

REFERENCES

Amacher A.L. (1996) :

Surgical management of meningocele and myelomeningoceles in operative neurosurgical technique; 165 : 2055-2068.

Amacher A.L., Drake C.G., McLaughlin A.D. (1968) :

Anterior sacral meningocele; surg Gynecol obstet; 126 : 986-994.

Barson A.J. (1970) :

Spina bifida : The significance of the level and the extent of the defect to the morphogenesis. Develop. Med. Child Neurol., 12 : 129-144.

Barkovich A.; Edwards M., and Cogen P. (1991) :

MRI evaluation of spinal dermal sinus tracts in children. Am. J. Neuroradiol 12 : 123.

Barnes P.D.; Reynolds A.F., and Gallaway D.C. (1984) :

Digital Myelography of Spinal dysraphism in infancy : Preliminary results. Am. J. Roentgenol., 142:1249.

Barry A.; Ratten B.M. and Stewart B.H. (1957) :

Possible factors in the development of the Arnold-chiori malformation. J. Neurosurg. 11 : 285 : 301.

Bell W.O.; Sumner T.E., and Volberg F.M. (1987) :

The significance of ventriculomegaly in the newborn with myelodysplasia. Childs Neyr. Syst.; 3:239-241.

Bellinger M.K. (1991) :

Neuropathic Bladder in adult and pediatric urology; 55 : 2055-2082.

Bentley JFE., and Smith JR. (1960) :

Developmental posterior enteric remnants and spinal malformations : the split notochord syndrome. Arch Dis. Child; 35 : 76-86.

REFERENCES

Amacher A.L. (1996) :

Surgical management of meningocele and myelomeningoceles in operative neurosurgical technique; 165 : 2055-2068.

Amacher A.L., Drake C.G., McLaughlin A.D. (1968) :

Anterior sacral meningocele; surg Gynecol obstet; 126 : 986-994.

Barson A.J. (1970) :

Spina bifida : The significance of the level and the extent of the defect to the morphogenesis. Develop. Med. Child Neurol., 12 : 129-144.

Barkovich A.; Edwards M., and Cogen P. (1991) :

MR evaluation of spinal dermal sinus tracts in children. Am. J. Neuroradiol 12 : 123.

Barnes P.D.; Reynolds A.F., and Gallaway D.C. (1984) :

Digital Myelography of Spinal dysraphism in infancy : Preliminary results. Am. J. Roentgenol., 142:1249.

Barry A.; Ratten B.M. and Stewart B.H. (1957) :

Possible factors in the development of the Arnold-chiori malformation. J. Neurosurg. 11 : 285 : 301.

Bell W.O.; Sumner T.E., and Volberg F.M. (1987) :

The significance of ventriculomegaly in the newborn with myelodysplasia. Childs Neyr. Syst.; 3:239-241.

Bellinger M.K. (1991) :

Neuropathic Bladder in adult and pediatric urology; 55 : 2055-2082.

Bentley JFE., and Smith JR. (1960) :

Developmental posterior enteric remnants and spinal malformations : the split notochord syndrome. Arch Dis. Child; 35 : 76-86.

Boone D.; Pansons, D., and Lochman S.M. et al. (1985) :

Spina bifida occulta; Lesion or anomaly? Clin Radiol 36 : 159.

Braun I.; Raghavendra B., and Kricheff (1983) :

Spinal cord imaging using real-time high-resolution ultrasound. Radiology, 147:459.

Breig A. (1970) :

Overstretching and pathological tension in the spinal cord-a basic cause of symptoms in cord disorders. J. biomech., 387-9.

Brunberg J.; Latchaw R., and Kanal E. (1988) :

Magnetic resonance imaging of spinal dysraphism. Radiol Clin. North. Am. 2 : 181.

Bruce D.A. and Schut L. (1979) :

Spinal lipoma in infancy and childhood. Childs Brain. 5-192.

Brun R.B., and Garson J.A. (1983) :

Neurulation in the Mexican salamander (Ambystoma mexicanum) : a drug study and cell shape analysis of the epidermis and the neural plate. J. Embryol Exp. Morphol. 74 : 275-295.

Carolli N. (1983) :

The orthopedic and orthotic management of the spina bifida child. Clin. Neurosurg., 30 : 413-435.

Carter C.O.; Roberts J.F.; Evans K., and Bude A.R. (1971) :

Genetic Clinic; A follow up. Lancet 1 : 281-285.

Cadduck, W.M. and Uthman, E.I. (1984) :

Squamous cell carcinoma and meningomyelocelw. Neurosurgery, 14 : 601-603.

Centers for Disease Control (1992) :

Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. MMWR; 41 : 1-7.

Champan P.H. (1996) :

Surgical management of occult spinal dysraphism in operative neurosurgical technique. 166 : 2069-2078.

Chapman P.H. (1982) :

Congenital intraspinal lipomas : anatomic considerations and surgical treatment. Childs Brain; 9:37-47.

Chan A.; Robertson E.F., and Ranieri E. (1995) :

The sensitivity of ultrasound and serum alpha-fetoprotein in population-based antenatal screening for neural tube defects. British Journal of Obstetrics and Gynaecology, Vol. 102, PP. 370-376.

Charney E.B.; Miller S.C., and Sutton L.N. (1985) :

Management of the newborn with myelomeningocele : Time for a decision-making process. Pediatrics 75 : 58.

Cheek W.R.; Laurent J.P. (1985) :

Dermal sinus tracts. In : Chapman PH, ed. Concepts in pediatric neurosurgery. Vol. 6. Basel, Switzerland : Karger, 63 -75.

Cocoyorin and Kieffer E.D., and Kieffer S.A. (1982) :

Application and limitations of computed tomography of the spine. Radiol. Clin. North. Am., 20:185.

Comarr A.E. (1977) :

Neurological disturbances of sexual function among patients with myelodysplasia. In McLaurin, R.L.; ed. : Myelomeningocele. New York. Grune & Stratton, PP. 797 : 808.

Dias M. and Pang D. (1995) :

Human neural embryogenesis : a description of neural morphogenesis and a review of embryonic mechanisms. In Pang D (ed) : Disorders of the pediatric Spine. New York : Raven Press. PP. 1-26.

Dickson R.H., and Leatherman K.D. (1977) :

The kyphotic spine in myelomeningocele. In McLaurin, R.L., ed. : Myelomeningocele. New York. Grune & Stratton, PP. 609-620.

Dremon J.B. (1970) :

The role of muscles in the development of human lumbar kyphosis. Develop. Med. Child. Neural., 121 Suppl. 22:33-38.

Duckworth T.; Sharrard W.J.; Lister J. and Seymour N. (1968) :

Hemimyelocoele. Develop. Med. Child Neurol. suppl., 16: 69-75.

Eckstein H.B. (1972) :

Myelomeningocele. Postgrad. Med. J., 481: 469-500.

Eckstein H.B. (1949) :

Neuropathic bladder. In Williams DI (ed) : Urology in Childhood. Berlin, Springer Verlag, P. 250.

Ehlers K.; Elmazar, M.M., and Nau, H. (1995) :

Biochemical and Molecular roles of nutrients, American Institute of nutrition. 6 : 67-75.

Emery J.L., and Lendon R.G. (1973) :

The local cord lesion in neurospinal dysraphism (meningomyelocoele). J. Pathol. 110: 80-96.

Elwood JH. (1972) :

Major central nervous system malformations notified in Northern Ireland, 1964-1968. Dev Med. Child. Neurol 14 : 731.

Fitz C.R., and Harwood-Nash D.C. (1975) :

The tethered conus. Am. J. Roentgenol., 125 : 515-523.

French B.N. (1982) :

Midline fusion defects and defects of formation in yomans, I.P. ed, : Text book of Neurological Surgery. Philadelphia, W.B. Saunders Co. Pub., Vol. 3, PP. 1236-1380.

French B.N. (1983) :

The embryology of spinal dysraphism clin. Neurosurg.
30 : 359-340.

Frerebeau Ph.; Dimegho A.; Grass M. and Harbi H. (1983) :

Diastematomyelia : Report of 2; cases surgically treated
by a neurosurgical and orthopedic team. child's Brain,
10 :328.

Furness M.E.; Barbary J.E. and Verco P.W. (1985):

A pointer to spina bifida : fetal head shape in the second
trimester. In World Federation of Ultrasound in
Medicine and Biology 1985 Proceedings (R.W. Gill and
M.J. Dadd, eds), Pergamon Press, Syndey, P. 296.

Gardner W.J. (1973) :

The dysraphic states from syringomyelia to anencephaly.
Amsterdam, Excerpta Medica. PP. 1-201.

Gardner W.J. (1977) :

Etiology and pathogenesis of the development of
myelomeningocele. In McLawrin, R.K.; ed. :
Myelomeningocele. New York, Grune & Stratton, PP. 3-
30.

Globus M.S.; Loughman W.D., and Epstein C.J. (1979) :

Prenatal genetic diagnosis in 3000 amniocenteses. N.
Engl J. Med. 300 : 157.

Greene W.B. Terry R.C., and DeMasi R.A. (1991) :

Effect of race and gender on neurological level in
myelomeningocele. Dev. Med. Child Neurol 33 : 110.

Guthkelch A.N. (1996) :

Diastematomyelia in wilkins R.H. and Regchary, S.S
eds. Neurosurgery; 353 : 3505-3519.

Guthkelch A.N. (1974) :

Diastematomyelia with median syptommm. Brain, 97 :
729-742.

Hall P.V.; Compbell R.L.S., and Kolarbeck J.E. (1975) :

Meningomyelocele and progressive hydromyelia.
Progressive paresis in myelodysraphism. J. Neurosurg.,
43 : 457-463.

Harood-Nash D.C. (1980):

Computed tomography of pediatric spine. A protocol for
the Radiol. clin. North. Am., 19 : 479.

Harris L.W.; Oakes W.J. (1996) :

Open Neural tube defects in practice of neurosurgeon;
184: 2779-2789.

**Hawass N.D.; El-Badawi M.G.; Fantani J.A.; Meshari A.A.; Abbas
F.S.; Edress Y.B., and Jabbar F.A. (1987) :**

Myelographic study of the spinal cord ascent during fetal
development. Am. J. Neurorad., 8 : 691-695.

Hendrick E.B.; Hoffman H.J., and Humphreys R.P. (1983) :

The tethered spinal cord clinic. Neurosurg., 30 : 457.

Hoffman HJ.; Hendrick E.B., and Humphreys R.P. (1976) :

The tethered spinal cord; its protean manifestations,
diagnosis and surgical correction. Childs Brain; 2:145-
155.

Hoffman HJ.; Hendrick E.B., and Humphreys R.P. (1982) :

Management of hydrocephalus. Monogr Neural Sci. 8 :
21-25.

**Hogge W.A.; Thiagarajah S.; Ferguson J.E.; Schnatterly P.T., and
Herbert G.M. (1989) :**

The role of ultrasonography and amniocentesis in the
evaluation of pregnancies at risk for neural tube defects.
Am J. Obstet. Gynecol 161, 520-524.

Huballah M.Y., and Hoffman H.J. (1987) :

Early repair of myelomeningocele and simultaneous
insertion of ventriculoperitoneal shunt : technique and
results. Neurosurgery; 20:21-23.

Humphreys R.P. (1996) :

Spinal dysraphism in wilkins, R.H. and Regachary, S. S.,
eds. : Neurosurgery. 359: 3453-3463.

Humphreys R.P.; Hendrick E.B., and Hoffman H.J. (1983) :

Diastematomyelia. Clin. Neurosurg, 30 :436.

Inoue H.K.; Kobayashi S., and Nakamura M. (1994) :

Treatment and prevention of tethered and retethered
spinal cord using a Gore-Tex surgical membrane. J.
Neurosurg, 80; 689-693.

James C.C.M., and Lassman L.P. (1972) :

Spinal Dysraphism. Spina Bifida occult. London, Butter-
worth and Co.

James H.E., and Walsh J.W. (1981) :

Current Problems in Pediatrics. spinal Dysraphism.
chicago, III : Yearbook Medical Publisher; XI : 8.

Jennings M.T.; Clarren S.K.; Kokich V.C., and Alvord E.C.Jr. (1982):

Neuroanatomical examination of spina bifida arperta and
Arnold chiari malformation in a 180 dog human fetus. J.
Neurol. sci., 54 : 325-338.

Jones P.H., and Love J.G. (1956) :

Tight filum terminale. Arch. Surg. (Chicago), 37:556-
566.

Kaplan J.O., and Quencer R.M. (1980) :

High resolution spinal sonography in infants. Am. J.
Roentgenol., 142 : 1243.

Kanev P.M.; Lemire R.J.; Loeser J.D., and Berger M.S. (1990) :

Management and long-term follow-up review of children
with lipomyelomeningocele, 1952-1987. J. Neurosurg.
73 : 48.

Katona F.; Berenyi M. (1975) :

Intravesical transurethral electrotherapy acta paediatr
Hung. 16 : 363.

Khoury A.; Hendrick E., and McLorie G. et al. (1990) :

Occult spinal dysraphism : Clinical and urodynamic outcome after division of the filum terminale. J. Urol; 144 : 426.

Kurzrock E.A.; Lowe, P., and Hardy, B.E. (1995) :

Bladder wall pedicle wraparound sling for neurogenic urinary incontinence in children. Journal of Urology, Vol. 155, 305-308.

Lassman L.P.; James CCM. (1977) :

Meningocele manque, Child's Brain 3 : 1.

Latchaw R.E. (1985) :

Congenital anomalies. In Latchaw, R.E. (ed.) : computerized tomography of the head, neck, and spine, Year-Book Medical Publishers, PP. 439-456.

Laurence K.M. (1977) :

The impact of antenatal diagnosis for central nervous system malformations in high risk pregnancies and pregnancy screening in a high risk community. Zeitschrift für Kinderchirurgie und Grenzgebiete 22 : 383-396.

Laurence K.M. (1989) :

A declining incidence of neural tube defects in the U.K. Z Kinderchir 44 : 51.

Laurence K.M. (1990) :

The genetics and prevention of neural tube defects and uncomplicated hydrocephalus in principles and practice of medical genetics. Vol. 1 : 22: 323-346.

Lemire R.L.; Loesser J.D.; Leech R.W., and Alvord Jr. E.C. (1975) :

Normal and Abnormal development of the human nervous system. New York Happel, Row. 11- 421.

Lemire R.J.; (1974) :

Embryology of the central nervous system. In Davis J.A. and Dobbing, J. Scientific foundation of pediatrics. London, William Medical books. 32 : 450-462.

Lindhout D., and Meinardi H. (1984) :

Spina bifida and in-utero exposure to valproate. Lancet II : 396.

Lindhout D., and Omtzigt J.G.C. (1994) :

Teratogenic effects of antiepileptic drugs : implications for the management of epilepsy in women of childbearing age. Epilepsia 35 (Suppl. 4) : 19-28.

Lindseth R.E. (1977) :

Valgus and rotational deformities of the knee, ankle and foot in myelomeningocele, incidence and development of hydrocephalus. Arch. Dis. Child., 36 : 381-389.

Lorber J. (1974) :

Selective treatment of myelomeningocele : To treat or not to treat ? Pediatrics, 53 : 307-309.

Luthy D.A.; Wardinsky T., and Shurtleff D.B. (1991) :

Cesarean section before the onset of labor and subsequent motor function in infants with myelomeningocele diagnosed antenatally. N Engl. J. Med. 10 : 662.

McAdams A.J. (1977) :

The Arnold-Chiari malformation : pathology and developmental considerations in McLone, P.K. ed. : Myelomeningocele. New York, Orune & Stratton, Inc., PP. 197-207.

McComb J.G., and Chen T.C. (1996) :

Closed spinal neural tube defects in practice of neurosurgery; 183 : 2753-2777.

McLone D. (1983) :

Results of treatment of children born with a myelomeningocele. Clin Neurosurg; 30: 407-112.

McLone D.G. (1992) :

Continuing concepts in the management of spina bifida. Pediatr Neurosurg, 18 : 254.

McLone D.G. (1995) :

Spina bifida aperta in disorders of paediatric spine;
9:137-157.

McLone DG., and Dias MS. (1994) :

Normal and abnormal early development of the nervous system. In: Cheek WR, ed. Pediatric neurosurgery, 3rd ed. Philadelphia : WB Saunders, 3-39.

McLone D.G., and Knepper P.A. (1986) :

Role of complex carbohydrates and nuerulation./ Pediatr Neurosci 1 : 2-9.

McLone D.G., and Naidich T.P. (1985) :

Terminal myelocystocele. Neurosurgery, 16:36.

McLone D.G., and Naidich T.P. (1985) :

Spinal dysraphism : experimental and clinica. In: Holtzman RNN, Stein BM (eds.), the Tehtered Spinal Cord. New York : Thieme-Stratton, 14-28.

Medical Research Council Vitamin Study Research Group (1991) :

Prevention of neural tube defects; results of the Medical Research Council Vitamin Study. Lancet; 338 : 131-137.

Meyer S., and Landau H. (1984) :

Precocious puberty in myelomeningocele patients. J. Pediat. Orthop., 4 : 28-31.

Merx J.; Niezen-Bakker S.; Thijssen H., and Walder H. (1989) :

The tethered spinal cord syndrome : A correlation of radiological features and preoperative findings in 30 patients. Neuroradiology, 31 : 63.

Miller P.; Smith D.W., and Shepard T.H. (1978) :

Possible teratogens of nural tube defect lancet II; 519.

Miller J.H.; Reid R.S., and Kemberling P.P. (1982) :

Utilization of ultrasound in the evaluation of spinal dysraphism in children. Radiology, 143 : 737-740.

Morris J., and Laurence KM. (1976) :

The effectiveness of genetic counselling for neural tube malformations. *Developmental Medicine and Child Neurology* 28 : Suppl. 37, 157-163.

Morriss-Kay G.M., and Crutch B. (1982) :

Culture of rat embryos with beta-Dxyloside : evidence of a role for proteoglycans in neurulation. *J. Anat.* 134 : 491-506.

Morrow R.J.; McNay M.B., and Whittle M.J. (1991) :

Ultrasound detection of neural tube defects in patients with elevated maternal serum alpha-fetoprotein. *Obstet Gynecol* 78, 1055-1057.

Naidich T.; Fernbach S.; McLone, D., and Shkolnik A. (1984) :

Sonography of the caudal spine and back : Congenital anomalies in children. *Am. J. Radiol.* 142 : 1229.

Naidich T.P.; Publowski R.M., and Naidich J.B. (1980) :

Computed tomographic signs of the Chiari II malformation. Part I: Skull and dural partitions. *Radiology*, 134 : 65.

Nathalie MJ Van der put.; Martin den Heyer, Rima Rozen, Henk, J. Blom, (1995):

Mutated methylenetetrahydrofolate reductase as a risk factor for spina bifida. *Lancet* Vol. 346 : 1070-1071.

Nicolaides K.H.; Campbell S.; Gabbe S.G., and Guidetti R. (1986):

Ultrasound screening for spina bifida : cranial and cerebellar signs. *Lancet* ii, 72-74.

Oakes W.J. (1985) :

Chiari malformations. Hydromyelia. Syringoscoliosis in myelomeningocele. In McLone.P.U. : *Myelomeningocele*. New York. Brunner Stratton. PP. 581-590.

Oakes W.J. (1996) :

Management of spinal cord lipoma and lipomyelomeningoceles in Wilkins R.H and Rengachary, S.S. eds. *Neurosurgery*; 353 : 3497-3504.

- Osaka K.; Tanimura T.; Hiravama A., and Matsumoto S. (1978) :**
Myelomeningocele before birth. *Neurosurg.* 49:711-724.
- Padget D.H. (1972) :**
Development of so called dysraptism, with embryonic evidence of clinical Arnold-Chiari and Dandgwalver malformation. *Johns Hofkines Med. J.* 180 : 127-165.
- Pang D. (1996) :**
Tethred cord syndrome in wilkins, R.H. and Regachary, S.S, eds. : *Neurosurgery*, 351 : 3465-3496.
- Pang D. (1995) :**
Spinal cord lipomas. In Pang D(ed) : *Disorders of the Pediatric Spine*, New York : Raven Press, PP. 203-251.
- Pang D.; Dias M. (1993) :**
Cervical myelomeningoceles. *Neurosurgery*, 33:363-373.
- Parke W.W. (1982) :**
Development of the spine in fothmam. E.H. : Simone., S.A. Eds. : *The spine philadelphia*, W.B. Saunders Co. Pub. Vol. 1 PP. 1-17.
- Petersen M.C.; Sherbondy A., and Wagener, J. (1995) :**
Abnormalities in control of ventilation in newborn infants with myelomeningocele. *Journal of Pediatrics* Vol. 126 : 1011-1015.
- Pttersson H.; Fitz C.P., and Harwood-Nash D.C. (1982) :**
Adverse reactions to Myelography with metrizamide in infants, children, and adolescents : General and CNs effects. *Acta. Radiol. Diag.*, 23 : 223.
- Ralis J., and Ralis H.M. (1972) :**
Morphology of peripheral nerves in children with spina bifida. *Dev. Med. Child Neurol.* 109 : 101.
- Reigel D.H. (1989) :**
Spina bifida. In *Pediatric Neurosurgery : Surgery of the Developing Nervous System*. Philadelphia : Saunders, PP. 35-52.

Reigel D.H. (1994) :

Spina bifida, pediat. neurosurgery, 3 : 51-75.

Reigel D.H., and McLone D.G. (1994) :

Tethred spinal cord, pediat. Neurosurgery; 9 : 77-95.

Reigel D.H.; Scarff T.B., and Woodford J. (1976) :

Surgery for tethered spinal cord in myelomeningocele patients (abstract). American Association of neurological surgeons. 26 : 1012-1019.

Rekate H.L. (1988) :

Management of hydrocephalus and the erroneous concept of shunt independence in spina bifida patients. BNIQ; 4 : 17-20.

Roberts C.J. Evans K.T.; Hibbard B.M.; Laurence K.M.; Roberts E.E., and Robertson I.B. (1983) :

Diagnostic effectiveness of ultrasound in detection of neural tube defect. The South Wales experience of 2509 scans (1977-1982) in high-risk mothers. Lancet ii, 1068-10069.

Robertson P. (1991) :

Operative and assisted delivery. Curr Opinion Obstet. Gynacol. 3 : 769.

Rozen R.L. (1977) :

Pathophysiology and spinal deformity in myelomeningocele Myelomeningocele. New York, Grune & Stration Inc., PP. 565-574.

Sandford M.K.; Kissling G.E., and Joubert P.E. (1992) :

Neural tube defect etiology : new evidence concerning maternal hyperthermia, health, and diet. Dev. Mved. Child. Neurol; 34 : 661-675.

Satoshi Shokti Yamada and Roberts (1987) :

Extensibility of the lumbar and sacral cord, pathophysiology of tethered spinal cord in cast. J.neurosurgery. 66 : 116-123.

Sayers M.D. (1977) :

The significance of the Arnold-Chiari malformation. In McLaurin. P.L. ed : Myelomeningocele. New York, 4 : 485-250.

Scarff T.B.; Toleikis J.R., and Bunck W.H. (1979) :

Dermatome somatosensory evoked potentials in children with myelomeningocele. Z. Kinderchir, 28 : 384.

Schneidau T.; Zebold K., and Kaplan W. (1995) :

Selective sacral rhizotomy for the management of neurogenic bladders in spina bifida patients. Journal of Urology, Vol. 154 : 766-768.

Schoenwolf G.C. (1988) :

Microsurgical analyses of avian neurulation : separation of medial and lateral tissues. J. Comp Neurol; 276 : 498-507.

Schoenwolf G.C., and Franks M.V. (1984) :

Quantitative analyses of changes in cell shapes during bending of the avian neural plate. Dev. Biol. 105 : 257-272.

Scheible W.; James H.E.; Lepold G.R., and Hilton S.V.W. (1983) :

Occult spinal dysraphism in infants : Screening with high-resolution real time ultrasound. Radiology, 146 : 743.

Sharred W.J.W. (1977) :

Assessment of the myelomeningocele child. In McLaurin RL (ed) : Myelomeningocele. New York, Grune & Staratton, PP. 389-410.

Smithells R.W.; Sheppard S., and Schorah C.J. (1980) :

Possible prevention of neural-tube defects by periconceptional vitamin supplementation. Lancet 1:647.

Stark G. (1971) :

Prediction of urinary incontinence in myelomeningocele. Devleop. Med. Child. Neurol., 13 : 388-389.

Stark G.D. (1971) :

Neonatal assessment of the child with a myelomeningocele Arch Dis. Child; 46:539-548.

Stark G.D. (1977) :

Spina bifida, problems and management. Oxford Blackwell Scientific publications. 15 : 435-439.

Steinbok P., and Cochrane, DD. (1991) :

The nature of congenital posterior cervical or cervicothoracic midline cutaneous mass lesions. Report of eight cases. J. Neurosurg; 75:206-212.

Storrs B.B. (1987) :

Selective posterior rhizotomy for treatment of progressive spasticity in patients with myelomeningocele. Preliminary report, ped Nuro. Sc. 13 : 135.

Sutton L.N. (1994) :

Spinal dysraphism, in wilkins, R.H. and Rengchary S.S eds : principles of neurosurgery; 5: 65-80.

Thompson M.W.; Willard H.F., and Mcinnes, R.R. (1991) :

Prenatal diagnosis Genetics in Medicine; 19 : 411-425.

Thompson M.W., and Rudd N.L. (1976) :

The genetics of spinal dysraphism. In Morley TP (ed) : current Controversies in Neurosurgery. Philadelphia, W.B. Saunders, PP. 126-146.

Toinson P., and Sugarman I.D. (1995) :

Complications with shunts in adults with spina bifida. British medical journal, Vol. 311 : 286-287.

Tskayannis D.E., and Shamperger R.C. (1995) :

Association of imperforate anus with occult spinal dysraphism. Journal of Pediatric surgery, Vol. 30 : 1010-1012.

Van Den Hof M.C.; Nicolaidis K.H.; Campbell J., and Campbell S. (1990) :

Evaluation of the lemon and banana signs in one hundred thirty fetuses with open spina bifida. Am. J. Obstet Gynecol. 161; 322-327.

Vandertop W.P.; Asai A., and Hoffman H.J. et al. (1992) :

Surgical decompression for symptomatic Chiaria II malformation in neonates with myelomeningocele. J. Neurosurg; 77: 541-544.

Vandresse J.H., and Cornelis, G. (1975) :

Diastematomyelia : report of eight observations. Neuroradiology, 10:87-93.

Venes J., and Stennings E.N. (1983) :

Surgical pathology in tethered cord secondary to myelomeningocele repair. Concepts pediatric neurosurgery. 4 : 165-185.

Venes J.L., and Stevens E.A. (1983) :

Surgical pathology in tethered cord secondary to myelomeningocele repair : Implications for initial closure technique. Concepts Ped Neurosurg; 4 : 165-185.

Venes J.L. (1984) :

Letter to the Editor. Pediatrics 74 : 948.

Venger B.; Laurent JP., and Cheek W.R. (1971) :

Congenital thoracic dermal sinus tract. In Marlin AE, ed. Concepts in pediatric neurosurgery, Vol. 9. basel, Switzerland; Karger, 161-172.

Wald N.J.; Cuckle H., and Brock JH. et al. (1977) :

Maternal serum alphafetoprotein measurement in antenatal screening for anencephaly and spina bifida in early pregnancy. (Report of United Kingdom collaborative study on alpha-fetoprotein in relation to neural tube defects.) Lancet 1:1323.

Wald N.J.; Cuckle H., and Nanchahal L. (1989) :

Amniotic fluid acetylcholinesterase measurement in the prenatal diagnosis of open neural tube defects. Second report of the Collaborative Acetylcholinesterase study. *Orenat Diagn.* 9 : 813.

Walker H.S.; Dietrich R.B.; Flannigan B.D.; Lufkin R.B.; Peaceck W.J., and Kangarlea H. (1987) :

Magnetic resonance imaging of the pediatric spine. *Radiograph.* 7 : 1129-1152.

Warder E.D., and Oakes W.J. (1993) :

Tethered cord syndrome and the conus in a normal position : *Neurosurgery* 33 :374-378.

Warder D.E., and Oakes W.J. (1994) :

Tethered cord syndrome : the low-lying and normally positioned conus. *Neurosurgery*, 34 : 597-600.

Warkany J. (1977) :

Morphogenesis of spine bifidia. In McLaurin, R.L., eds.: *Myelomeningocele*. New York, Grane and stratton, PP. 31-339.

Weaver D.D. (1996) :

Genetics of developmental defects in wilkins. R.H. Rengchary SS. eds.; 347 : 3419-3456.

Whittle L.R.; Johnston, I.H., and Besserm, M. (1985) :

Intraoperative monitoring of spinal somatosensory evoked potential during surgery for the tethered cord syndrome. *J. Pediatric neurosci.*, 1:178-186.

Wilkins R.H. (1996) :

Lateral and ant spinal meningocele in wilkins RH and Rengchary, S.S eds. *Neurosurgery*; 355 : 3521-3525.

Wilkins R.H., and Odom G.L. (1976) :

Spinal intradural cysts. In vinken PJ. Bruyn GW (eds.) : *Handbook of Clinical Neurology*, Vol 20. Tumors of the spine and spinal Cord, Part II. Amsterdam : North-Holland, PP. 55-102.

Wilkins R.H., and Rossitch E.Jr. (1995) :

Intraspinal cysts. In Pang D (ed) : Disorders of the pediatric Spine. New York : Raven Press, PP. 445-466.

Wright RL. (1971) :

Congenital dermal sinuses. In: Krayenbuhl H, Maspes PE, Sweet WH, eds. Progress in neurological surgery, Vol. 4. Basel, Switzerland : Karger, 175-191.

Yamada S., and Lacono R.P. (1995) :

Tethered cord syndrome, Disorders of pediatric spine; 10:159-173.

Yamada S.; Zinke D.E., and Sanders D. (1981) :

Patholphysiology of tethered cord syndrome. Y. Neurosurgery., 54 : 494-503.

Zimmerman K.; Weingorton C.E.; Johnson L.A.; Heier, and M.D.F. Deck (1990) :

Neuroradiology of the spine. In yomans neurological surgery 3rd edition, W.B. Sanders company. Vol. 1. 364-404.