

Reference

- A.A.P. (1992):* Emergency cardiac care committee and sub committee : Neonatal resuscitation program. J.A.M.A. 268: 2276.
- A.A.P. CO.N. (1993):* American Academy of pediatric committee on nutrition. Nutrition. Hound book Evanshion III.
- A.A.P. (1997):* American Academy of pediatric and American college of obstetric and gynecology (1997): Guidelines for perinatal care 4th Ed. Elk grove, Illinois.
- A.A.P. (1997):* Report of committee on heart dis. Elk.Grove 4th Ed Illinois.
- Aisenberg R.B., (1982):* Correlation between hypoxaemia and congestive heart failure. Ped. cardio l. 3:133.
- Allan, L.D., (1994):* Prospective diagnosis of 1000 cases of congenital heart dis. in the fetus J. Am. Coll of cardiol. 23:1452.
- Anand, KJS, Phil, D., Hickey, PR., (1992):* Halothane - morphine compared with high dose sufentanil for anaesthesia and post operative analgesia in neonatal cardiac surgery. N. Eng. J. Med., 326:1-9.
- Antol J.M., Davis, L.T., Erzurum S.C., et al., (1996):* Surfactant suppresses N. F_k B. activation in human monocytic cells. Am.J.Respir. Cell. Mol. Biol. 14:374.
- Avery, M.E., Gate wood O.B., Brumley G., (1996):* Transient tachypnea of newborn: Possible delayed reabsorption of fluid at birth. Am.J. dis. child. 111:380.

Ayers J., Ress J., Lee T., and cochrane G.M., (1982): British Med. Journal
vol. 284: 927:928.

Baden M, Bauer CR, calle E, Klein G, papageorgiou A, and stern(1973):
l., plasma corticosteroids in infants with respiratory distress
syndrome. Pediatrics; 52:782-7.

Bauchner H., May, A., and Coules E., (1992): The use of analgesic agents
for invasive medical procedures in pediatric and neonatal
I.C.U. J.Ped.121:647-9.

Bifano. EM., and Ehrenkranz, Z. Ed. (1995): Perinatal haematology,
clinical perinatology 23:3.

Boeck, C.De, Reempts P. Van, Rigattoh, chernick, (1984): Naloxone reduces
the decrease in ventilation induced by hypoxia in neonatal
infants. J. Appl. Physiol. 56:157-11.

Bohin. S., and Field, DJ., 1994): The epidemiology of neonatal respiratory
disease. Early Hum. Dev. 37:73.

Burke R., (1995): Video Assisted thoracoscopic surgery for C.H.D. J.
Thoracs. Cardiovasc. Sug. 109-3: 499:50.

Burnard. E.D., Todd, D.A., J. hone E., and Hindmarch, K.W., (1982): B.E.
levels in newborn cerebro spinal fluid. Aust. Pediatr, 18:258-
63.

Chernick. V., (1981): N. Eng. J. Med. Vol. 304:1227-28.

Clement J. (1996): Hot topics in neonatology, washington, Dc. Dec. 1-3.

Corcoran J.D., patterson C.C., Thomas P.A., and Halliday, H.L., (1993):
Reduction in the risk of bronchopulmonary dysplasia from

1980:1990. Results of a multi variate and logistic regression analysis. Eur. J. pediatr. 152:677.

Dnaford DA., Miskes, Headley Nelson KM, (1983): Effects of routine care procedures on transcutaneous oxygen in neonates
a quantitative approach arch. Dis. child. 58:20-3.

Davis, (1993): Davis DJ. (1993): How aggressive should the delivery room be cardiopulmonary resuscitation for ELBW pediatric,
92:447.

Dawes, (1968): Fetal and neonatal physiology; Chicago: Year Book Medical Publishers.

Donn, S.M., and Nicks, J.J., (1996): Assisted ventilation in the neonates
Philadelphia, W.B. Saunders P 215.

Edwin. J.B., Eric. M., Smith Debora Harbour and Mac Menamin, (1985):
Journal of immunology, 135:7795-7825.

Edwin J. Blalock, (1989): Physiological reviews. Vol, 69:No.1.

Emond, D., Lachance C., Gagnon J., Bard LD, (1993): Arterial partial pressure of oxygen required to achieve 90% saturation of
Haemoglobin in very low birth weight newborn. Pediatrics.
91:602.

Facchinetti, F., Bagnoli, Petraglia, et al., 1982: Feto-maternal opioid level
and parturition obs. J. Neuroendocrinol. 62:764-768.

Fauza D., Hines M., Fackler J., et al., : Continuous positive airway pressure
with perfluorocarbon accelerates postnatal lung growth.
Surg. from 46:666.