

References

- Akesson, B. (1975):** Occurrence of phospholipase A₁ and A₂ in human decidua. Prostaglandins. 9: 667-673. Quoted by Kerise (1978).
- Allot, H.A. and Palmer, C.R. (1993) :** Sweebling the membranes . A valid procedure in stimulating the onset of labor. Br. J. Obstet . Gynaecol 100:898.
- Anderson, A.B.M.and Turnbull, A .C. (1969) :** Relationship between length of gestation and cervical dilatation, uterine contractility and other factors during pregnancy. Am. J. Obstet .Gynecol .105: 1207.
- Anderson, G.G.; Hobbins, J.C. and Speroff, L. (1972) :** Intravenous prostaglandins E₂ and F₂ alpha for the induction of term labor. Am. J. Obstet. Gynecol. 112: 382-386.
- Anderson, G.G. ; Hobbins , J. C. ; Speroff, L. and Caldwell, B.V. (1972) :** Intravenous prostaglandins E₂ , F₂ and syntocinon for the induction of term labor. J. Repord. Med. 9: 287. Quoted by Wingerup and Ulmsten, 1981.
- Andersson , K. E. ; Forman , A. and Ulmsten , U. (1983) :** Pharmacology of labor. Clin. Obstet. Gynecol. 26:56-77.
- Andreasson ,B.; Bock, J.E.and Larsen ,J. (1983):** Induction of labor.

- suppositories compared with intranasal oxytocin and with sequential treatment. Acta. Obstet, Gynecol .Scand. 64:157-161.
- Baily , D. H.; Newman , C. and Ellinas , S. P. (1975) :** Use of prostaglandin E₂ vaginal suppositories in intrauterine fetal death and missed abortion. Obstet. Gynecol. 45: 110-113.
- Barr, W. and Graham, R. (1967):** Postgrad. med. J. Suppl. 433, 101.
- Barta , S. and Bengtsson, L.P. (1976b) :** Mechanism for increase production of prostaglandins in labor. Lancet. 1:1164-1165.
- Basu, S.B. and Feeney, J.G. (1980) :** The influence of pre-induction vaginal prostaglandin E₂ gel on subsequent labor. J. Ir. Med. Assoc. 73: 194-196.
- Bennett , A. ; Charlier , E.M. ; McDonald , A.M. ; Simpson, J.S.; Stamford , E. F. and Zebro , T. (1977) :** Prostaglandins and breast cancer. Lancet. (ii). 624 – 626.
- Bennett , A. ; McDonald , A.M.; Simpson , J.S. and Stamford , I.F.(1975):**Breast cancer, prostaglandins and metastases . Lancet. (I). 1218- 1220.
- Bergstrom, S. and Sjovalld (1960):** The isolation of prostaglandin from sheep prostate glands. Acta. Chemica .Scandinavia. 14: 1701-1705. Quoted by Population Reports Series G. 1980.
- Bernner , W. E. (1983) :** Reid's Controversy in Obstetrics and

- Gynecology-III. W.B. Sunnders Company. Philadelphia, London, Toronto, Mexico City, Rio de Janeiro, Sydney, Tokyo. pp. 371-385.
- Bernstein ,P.; Leyland, N. ;Gurland ,P.and Gare ,D. (1987):** Cervical ripening and labor induction with prostaglandin E₂ gel: A Placebo-controlled study. Am. J. Obstet .Gynecol .156: 336-340.
- Bibby, J.G.; Rnint , J.; Mitchell, M.D. ; Anderson A .B .M. and Turnbu, A . C. (1979) :** The effect of cervical encerclage on plasma prostagland concentration during early human pregnancy. Br. J. Obstet.Gynaecol. 86: 19.
- Bishop , E. H. (1964) :** Pelvic scoring for elective induction. Obstet. Gynecol 24: 266.
- Bitensky, L. and Cohen, S. (1965):** J. Obst. Gynaecol. Brit. Cwlth. 72: 769 – 774.
- Blumenthal, P.D.and Ramamauskar, R. (1990):** Randomized trial of dilapan and laminaria as cervical ripening agents before induction of labor. Obstet .Gynecol 5: 365.
- Brandel , R.; Bygdemen , M. and Weqvist, N.A. (1970):** Comparable study of the influence of prostaglandin E ,oxytocin and ergometrine on the pregnant human uterus. Obstet. Gynecol. Scand .49 supp .5.

- Browne, F. J. (1939) :** Antenatal and postnatal care , 3rd ed. London: Churchill.
- Bryant, W.M.; Greenwill ,J.E.and Weeks, P.M. (1968):** Alterations in collagen organization during dilatation of the cervix uteri. Surg .Obstet. Gynecol .126: 27.
- Buchan, P.C. (1979):** Pathogenesis of neonatal hyperbilirubinaemia after induction of labor with oxytocin. British Medical Journal. 2: 1255-1257.
- Buckingham, J.C.; Selden, R.andDanforth, D.N.; (1962):** Connective tissue changes in the cervix during pregnancy and labor. Ann .NY. Acad. Sci .97: 733.
- Bugalho , A. ; Bique , C. ; Almeida , L. and Fandes , A.; (1993):** The effectiveness of intravaginal Misoprostol (Cytotec) in inducing abortion after eleven weeks of pregnancy. Stud. Fam. Plann. Vol. 24 no. 5 pp. 319-23.
- Bugalho, A.; Bique, C.; Machungo, F. and Fandes, A. (1995):** Low-dose vaginal Misoprostol for induction of labor with a live fetus. Int. J. Gynecol. Obstet. vol 49, no. 2 pp. 149-55.
- Burnett, J.E. (1966):** Pre-induction scoring. An objective approach to induction of labor. Obstet .Gynecol. 28: 479.
- Bygdeman, M. (1964):** Acta. physiol. Scand. 63, Suppl. 242:1, Quoted

from Karim and Hillier., 1973.

Bygdeman , M . ; Bergstron , S. and Samuelsson , B. (1967) : (eds.)

Prostaglandins (Proceedings of the 2nd. Nobel Symposium,
Stockholm. June, 1966. Wiley-Interscience, New York, p. 71.

Quoted by Karim and Hillier., 1973.

Cabrol , D. ; Breton , M. ; Berrou , E. ; Visser , A. ; Sureau , C. and

Picard, J. (1980): Variations in the distribution of glycosaminoglycans

in the uterine cervix of the pregnant woman. Europ. J. Obstet.

Gynecol Reprod. Biol. 10: 281.

Calder , A . A. ;Embrey, M.P. and Hillier, K.(1974) : Extra-amniotic

prostaglandins E₂ for the induction of labor at term. J. Obstet.

Gynecol commonw. 81: 39.

Calder , A. A. (1990) : Control of parturition. Scientific and clinical

aspects in uterine contractility. In: Grafield ,R.E. Ed. Serono

Symposia. Norwalk. Massachusetts. p. 283-293.

Calder, A.A. (1980): Pharmacological management of the unripe cervix

in the human. In Naftolin ,F. and Stubblefield, P.G. (eds):

Dilatation of the uterine cervix. New York, Raven Press. pp.317

Calder , A.A. ; Embrey , M.P. and Hillier , K. (1974): Extra-amniotic

prostaglandin E₂ for the induction of labor at term.Br J. Obstet.

Gynecol. Br. Commonw. 81:39-46.

- Calder, A.A.; Moar, V.A.; Ounsted, M.K. and Turnbull, A.C. (1974b):** Increased bilirubin levels in neonates after induction of labor by intravenous prostaglandin E₂ or oxytocin. *Lancet*. 2: 1339-1342.
- Calder, A.A. ; Embrey, M.P. and Tait, T. (1977) :** Ripening of the cervix with extra-amniotic prostaglandin E₂ in viscous gel before induction of labor. *Br. J. Obstet. Gynaecol* .84: 264.
- Calderyo-Barcia, R. and Poserio, J. J. (1960) :** Physiology of the uterine contraction. *Clin. Obstet. Gynecol*. 386-408.
- Caldeyro-Barcia, R. ; Schwarcz, R. ; Belizan, J.M. ; Martell, M. ; Nieto, F.; Sabatino, H. and Tenzer, S.M. (1974) :** Adverse perinatal effects of early amniotomy during labor. In L. Gluck: *Modern Perinatal Medicine*. Chicago: Year Book Medical Publishers, Inc.p. 431- 449.
- Campell, J. M. (1984) :** Induction of labor using prostaglandin E₂ pessaries. *Clin. Exp. Obstet. Gynecol*. XI: 1-5.
- Carsten, M. E. (1974) :** Prostaglandins and oxytocin : Their effects on uterine smooth muscle. *Prostaglandins*. 5: 33-40.
- Carsten, M. E. (1979) :** Calcium accumulation by human uterine microsomal preparations: Effects of progesterone and oxytocin. *Am. J. Obstet. Gynecol*. 133: 598 – 601.

- Carsten, M.E. and Miller, J.D. (1979):** P.G. Specific binding to several microsomal fractions from bovine myometrium. 4th International Prostaglandin Conference. Prostaglandins.
- Chalmers, I.; Campbell, H. and Turnbull, A.C. (1975):** Brit. med. J. 2: 116.
- Chard, T. and Gibbons, G.L. D. (1983) :** Spurt release of oxytocin during surgical induction of labor in women. Am. J. obstet. Gynecol.:147: 678-680.
- Chayen, B.; Tejani, N. and Verma, U. (1986):** Induction of labor with an electric breast pump. J. Reprod. Med.31:116-118.
- Chuck, F. J. and Huffaker, B. J. (1995) :** Labor induction with intravaginal misoprostol versus intracervical prostaglandin E₂ gel (Prepidil gel):A randomized comparison. Am. J. Obstet. Gynecol. Vol. 173. no. 4. pp. 1137-42.
- Chvapil, M.; Droegemuller, W. and Meyer, T. et al. (1982):** New synthetic laminaria. Obstet. Gynecol. 60:729.
- Cisternino, A.; Panella, M.; Ettora, G.; Scollo, P. and Nocero, F. (1985):** Therapy to ripen the uterine cervix with intravenous infusion of prostaglandin E₂ before induction of labor in primigravidae. Archives in Gynecology. Abstracts XI. World Congress of Gynecology and Obstetrics. Berlin West Abstract N. 04. 42. 05.

- Coceani, F. ; Olley, P.M. and Bodach, E. (1976)** : Prostaglandins: A possible regulator of muscle tone in the ductus arteriosus. In Advances in Prostaglandin and Thromboxane Research. Vol.1. pp.417-420. Samuelsson, B. and Paoletti, R. (eds) New York: Raven Press
- Collins, G. ; Outerbridge, E. ; Manku, M.S. and Horrobin, D.F. (1976)** : Chloroquine as prostaglandin antagonist in treatment of patent ductus arteriosus. Lancet. (ii). 810.
- Conard, J.T. and Ueland, K. (1976)** : Reduction of the stretch modulus of human cervical tissue by prostaglandin E₂. Am. J. Obstet. Gynecol. 126: 218.
- Corsten, M. E. and Miller, J.D. (1977)** : Effects of PG and oxytocin on Calcium release from uterine microsomal fraction. Journal of Biological Chemistry. 252. (5). 1576 – 1581.
- Coudert, S. P. ; Winter, J. S. D. and Faiman, C. (1974)** : American J. of Obstet. & Gynecol. 119. (6). 755, 761.
- Coutinho, E.M. and Maia, H. S. (1971)** : The contractile response of the human uterus, fallopian tubes and ovary to prostaglandins in vivo. Fertility and Sterility. 22: 539 – 543.
- Craft, J.L.; Cullum, A.R.; May, D.T.L.; Noble, A.D. and Thomas, D.J. (1971)** : Prostaglandin E₂ compared with oxytocin for the induction

- of labor. Br. Med. J. 3: 276.
- Craft, I. (1972):** Amniotomy and oral prostaglandin E₂ for induction of labor. Br. Med. J. 11: 191. Quoted by Gordon Wright et al (1979a).
- Creinin, M. D. and Vittinghoff, E. (1994) :** Methotrexate versus Misoprostol alone for early abortion. A randomized control trial. JAMA. Vol. 272. no. 15 pp 1190-5.
- Creinin, M.D. ; Moyer, R. and Guido, R. (1997) :** Misoprostol for medical evacuation of early pregnancy failure. Obstet. Gynecol. 89.(5pt 1): 768-772.
- Cross, W.G. and Pitkin, R. M. (1978) :** Laminaria as an adjunct in induction of labor. Obstet. Gynecol. 51:606-608.
- Csapo, A. I. ; Jaffin, H. and Kerenyi, T. et al (1963) :** Volume and activity of the pregnant human uterus. Am .J. obstet. Gynecol.
- Cunningham, F.G.; Cox, K.; Hauth, J.C.; Strong, J.D. and Whalley, P.J. (1976):** Oral prostaglandin E₂ for labor induction in high risk pregnancy. Am.J. Obstet. Gynecol. 125: 881- 888.
- Cuthbert, M.F. (1969):** Effect on airway resistance of prostaglandin E₁ given by aerosol to healthy and asthmatic volunteers. Br. Med. J. (iv). 723 – 726.

- Danforth, D.N. (1980):** Early studies of the anatomy and physiology of the human cervix and implications of the future. In Naftolin, F. and Stubblefield, P.G. (eds): Dilatation of the uterine cervix. New York, Raven Press. p 3.
- Danforth, D.N. and Buckingham, J.C. (1973):** The effects of pregnancy and labor on the amino acid composition of the human cervix. In: Blandau, R.J. and Maghissi, K. (eds): The Biology of the cervix. Chicago, University of Chicago Press. p. 351-354
- Danforth, D.N.; Buckingham, J.C. and Roddick, J.W. (1960):** Connective tissue changes incident to cervical effacement. Am.J. Obstet. Gynecol. 80: 939.
- Danforth, D.N. (1954):** The distribution and functional activity of the cervical musculature. AM. J. Obstet. Gynecol. 68: 1261 – 1271.
- Danforth, D.N. (1983):** The morphology of the human cervix. Clin. Obstet. Gynecol. 26: 7-13.
- Danforth, D.N. and Buckingham, J.C. (1973):** The effects of pregnancy and labor on the amino acid composition of the human cervix. In R.J. Blandau and K. Moghissi: The Biology of the cervix. Chicago. University of Chicago Press. p.351 – 354.
- Danforth, D.N.; Veis, A.; Breen, M.; Weinstein, H.G.; Buckingham,**

- J.C. and Manalo, P. (1974):** The effect of pregnancy and labor on the human cervix: Changes in collagen, glycoproteins and glycosaminoglycans. *Am. J. Obstet. Gynecol.* 120: 641 - 651.
- Danforth, D.N. (1947):** The fibrous nature of the human cervix and its relation to the isthmic segment in gravid and non gravid uteri. *Am. J. Obstet. Gynecol.* 53: 541.
- Davey, D.A. (1980):** Induction of labor. *Clinics. Obstet. Gynecol.* 7:481-509.
- Davies, B.N.; Horton, E.W. and Withrington, P.G. (1968):** The occurrence of prostaglandin E₂ in splenic venous blood of the dog following nerve stimulation. *Br. J. Pharmac.* 32: 127-135.
- Dawood ,M.Y. ;Lauersen, N.H.; Trivedi ,D. andFuchs F (1979):** Studies on oxytocin in the baboon during pregnancy and delivery. *Acta. Endocrinol.* 91:704-18.
- Dawood, M.Y.; Raghavan ,K.S.; Pociask, C. andFuchs ,F. (1978):** Oxytocin in human pregnancy and parturition. *Obstet. Gynecol.* 51:138-43.
- Dawood, M.Y.; Wang ,C.F.; Gupta ,R.and Fuchs F (1978):** Fetal contribution to oxytocin in human labor. *Obstet. Gynecol.* 52:205-9.

- Deborah, A.; Wing; Ann Rahall; Margaret M. Jones; Murphy Goodwin, T. and Richard, H. Paul. (1995):** Misoprostol: An effective agent for cervical ripening and labor induction. Am. J. Obstet. Gynecol. pp. 1811-6.
- Dhont, M. ;Vanderheyden, K.; Vandekerckhove, D.; Vermeulen, A. and Thiery, M. (1974) :** PGE₂ : Effect on cortisol , growth hormone , prolactin, L.H, FSH & TSH levels before and after suppression with dexamethasone. I.R.C.S. 2: 1301. PGF_{2a}: I.R.C.S. 2: 1164.
- Dillon ,T.F. ;Douglas, R.G. ;Du Vigneaud ,V.and Barber, M. (1960):** Transbuccal administration of oxytocin for induction and stimulation of labor. Obstet .Gynecol. 15 : 587 – 92.
- Dimitrios, S.; Mastrogiannis; Robert, A.and Knuppel. (1995):** Labor induced using methods that do not Involve oxytocin.Clinic. Obstet. and Gynecol. Vol. 38. No. 2. p. 259-266.
- Domisse, J.; Davey, D.; Martin, B. and Cohen, M. (1981):** An Evaluation of prostaglandin E₂ administered intrarectally to induce labor.S.Afr. Med. J. 59: 817 – 818.
- Downie, J.; Poyser, N. L. and Wundererlich, M. (1974):** Level of PGs in human endometrium during the normal menstrual cycle. J. Physiol. 236, 465, 473.

Dunn, P.M. (1992): Dr. Thomas Denam of London (1733-1815):

Rupture of the membranes and management of the cord. Arch.
Dis. Child. 67:882-884.

Durlot, F.; Buboiss, C.; Brunerie, J. and Frydman, R. (1988): Efficacy

of progesterone preoperative cervical dilatation during first
trimester abortion. Human Reprod. 3:583-584.

Eglinton, G.; Raphael, R.A.; Smith, G.N.; Hall, W.J. and Pickles,

V.R. (1963): Nature (London). 200: 960 – 993. Quoted by Karim and
Hiller, 1973.

Ekman, G.; Forman, A.; Marshal, K. and Ulmsten, U. (1983):

Intravaginal versus intracervical application of prostaglandin E₂ in
viscous gel for cervical priming and induction of labor at term in
patients with an unfavorable cervical state. Am. J. Obstet.
Gynecol. 147: 657-61.

Ekman, G. Granstorm, L. and Ulmstem, U. (1986): Induction of labor

with intravenous oxytocin or vaginal PGE₂ suppositories. A
random ized study. Acta. Obstet. Gynecol. Scand. 65:857-859.

Ellwood, D.A.; Mitchell, M.D.; Andresson, A.B.M. and Turnbull,

A.C. (1979): Oestrogens, prostaglandins, and cervical ripening. Lancet.
1: 376 – 377.

- El-Refaey, H.; O'Brien, P.; Morafa, W.; Walder, J. and Rodeck, C. (1997):** Use of oral Misoprostol in the prevention of post-partum hemorrhage. Br. J. Obstet. Gynecol. 104. (3) : 336-339.
- El-Torkey, M. and Grant, J.M. (1992):** Sweeving of the membranes is an effective method of induction of labor in prolonged pregnancy: A report of randomized trial. Br. J. Obstet Gynaecol 99: 465.
- Embrey, M.P. (1966):** Mem.Soc. Endocr. 14: 89. Quoted by Embrey , 1977.
- Embrey, M.P. (1969):** The effect of prostaglandins on the human pregnant uterus. J. Obstet. Gynecol. Br. Common W. 76: 783 – 789.
- Embrey, M.P. (1977):** Oxytocin and the uterus: Scientific foundation of obstetrics and gynecology. London. William Heine Mann. Medical Books.LTD. pp. 798 – 810.
- Embrey, M.P. and Calder, A.A. (1975):** Induction of labor. In R.W. Beard; M. Brudenell; P. Dunn and D. Fairweather: The management of labor: College of Obstetricians and Gynecologists. London.p. 62. Quoted by Ulmsten et al., 1983.
- Embrey, M.P. and Mollison, B.G. (1967):** The unfavorabe cervix and induction of labor using a cervical balloon. J. Obstet. Gynaecol. Br. Common w. 74:44-48.

- Embrey, M.P. and Morrison, D.L. (1968):** J. Obstet.Gynecol. Br. Common W. 75: 829. Quoted by Karim and Hiller., 1973.
- Embrey, M.P.; Graham, N.B. and McNeill, M.E. (1980):** Induction of labor with a sustained –release prostaglandin E₂ vaginal pessary. Br. Med. J. 281: 901 – 902.
- Embry, M.P. and Anselmo, J.F. (1962):** The effects of intravenous oxytocin on uterine contractility. J. Obstet. Gynaecol .Br. Commonw 69: 910.
- Eskin; B.A. and Azarbal,S. (1973):** Advance. Bio -sciences. 9: 931 – 935.
- Ferguson, J.K.W.(1941):** A study of the motility of the intact uterus at term. Surg. Gynecol .Obstet.73: 359 – 366.
- Fitz Gibbon , G . (1923) :** Practical Midwifery . London : Churchill .
- Fitzpatrick, R.J. and Liggins, G.C. (1980):** Effects of prostaglandins on the cervix of pregnant women and sheep. In F. Naftolin and P.G. Stubblefield: Dilatation of the uterine cervix. New York: Raven Press.p. 287 – 300.
- Fletcher, H.M.; Mitchell, S.; Simeon, D.; Friderick, J. and Brown, D. (1993):** Intravaginal Misoprostol as a cervical ripeneing agent. Br. J. Obstet. Gynaecol. Vol. 100:. 641-4.

- Fletcher, H.M.; Mitchell ,S.; Simeon ,D.; Fredrick, J.and Brown, D. (1994) :** Intravaginal Misoprostol versus dinoprostone as cervical ripening and labor-inducing agents. *Obstet. Gynecol.* 83:244-7.
- Flynn, C.A. (1997):** Intravaginal Misoprostol :A new option for labor induction?. *J. Fam. Pract.* Vol. 44 no. 1. pp. 31.
- Forman, A.; Gandrup, P.; Andersson, K.E. and Ulmsten, U. (1983):** Effects of nifedipine on oxytocin and prostaglandin F₂ – alpha-induced activity in the post partum uterus. *Am. J. Gynecol.* 144: 665-670.
- Forman, A.; Ulmsten, U.; Banyai, J.; Wingerup, L. and Uldbjerg, N. (1982):** Evidence for a local effect of intracervical prostaglandin E₂-gel. *Am.J. Obstet. Gynecol.* 143:756-760.
- Francis, J.S.; Turnbull, A.C. and Thomas, F.F. (1970):** Automatic oxytocin infusion equipment for Induction of Labor. *J. Obstet. Gynaecol. Br. Commonw.* 77: 594-602.
- Friedman, E.A.; Niswander, K.R.; Bayonet-Rivera, N.P. and Sachtelben , M. R. (1966) :** Relation of pre-labor evaluation to inducibility and the course of labor. *Obstet .Gynecol* .28: 495.
- Frydman ,R.; Fernandez ,H.; Pons ,J.C.and Ulman A (1988):** Mifepristone (RU 486) and therapeutic late pregnancy

termination. A double blind study of two different doses. Human
.Reprod.3:803-806.

Frydman, R. ;Lelaidier ,C.and Baton-Saint-Mleux, C. et al (1992):

Women at term with mifepristone (RU 486) A double blind
randomized placebo controlled study. Obstet. Gynecol. 80:972-
975.

Fuchs , A.R. ;Fuch ,F. ; Husslein,P.and Soloff,M.S.(1984): Oxytocin

receptors in the human uterus during pregnancy and
parturition.Am.J.Obstet. Gynecol:150:734-41.

Graber, D.J. and Meier, K.H. (1991): Acute Misoprostol toxicity.

Ann. Emerg. Med. Vol.20 no. 5 :549-51.

Garette, W.J. (1960): Prognostic sign in surgical induction of labor.

Med .J .Aust. 49: 929.

Garfield, R.E.; Kannas, M.S. and Daniel, E.E. (1980): Gap. junction

formation in myometrium: Control by estrogen, progesterone and
prostaglandins. Am. J. Physiol. 238: c81 – c89. Quoted by

Husslein, P. and Sinzinger, H. (1984): British. J. of Obstet. and
Gynaecol. 91: 228 – 231.

Gerson Weiss (1995):Relaxin used to produce the cervical ripening of

labor.Clinic.Obstet.Gynecol.38:293-300.

- Ghosh, A. and Hudson, F.P. (1972):** Oxytocic agents and neonatal hyperbilirubinaemia. *Lancet*. 2: 823 – 825.
- Goldblatt, M.W. (1933):** *Chemistry and Industry*. 1052-56. Quoted by Karlson.: The influence of seminal fluid on the motility of the non-pregnant uterus. *Acta .Obstet. Gynecol. Scand*. 38: 503-521 (1959).
- Goldblatt, M.W. (1935):** *J. of Physiology*. 84: 208. Quoted by Karlson, S: The influence of seminal fluid on the motility of the non – pregnant human uterus. *Acta .Obstet. Gynecol. Scand*. 38: 503 – 521.
- Gordon –Wright ,A.P.and Elder, M.G. (1979):** Prostaglandin E₂ tablets used intravaginally for the induction of labor. *Br. J. Obstet. Gynaecol*. 86:32-36.
- Gordon, A.J. and Calder, A.A. (1977):** Estradiol applied locally to ripen the unfavorable cervix. *Lancet*. 2: 1319.
- Gordon-Wright, A.P.; Dutt, T.P. and Elder, M.G. (1979a):** The routine use of oral prostaglandin E₂ tablets for induction or augmentation of labor. *Acta. Obstet. Gynecol.Scand*. 58: 23-26.
- Graber, D.J. and Meier, K.H. (1991):** Acute Misoprostol toxicity. *Ann. Emerg. Med*. Vol.20 no. 5 :549-51.

- Grunberger, W.; Huber, I. And Husslein, P. (1984):** Local application of PGE₂ by means of a portio-adaptor. A new method for induction of labor. Acta Obstet. Gynecol. Scan. 63: 293-297.
- Gustavii, B. (1973):** Studies on the mode of action of intra-amniotically and extramniotically injected hypertonic saline in therapeutic abortion. Acta. Obstet. Gynecol. Scand. (Suppl.) 25: 5.
- Haberal, A ; Celikkanat, H. and Batioglu, S.(1996):** Oral Misoprostol use in early complicated pregnancy. Adv. Contracept. 12(2):139-143.
- Hedqvist, P. (1973):** Prostaglandin mediated control of sympathetic neuroeffector transmission. Adv. Bio-sci. 9: 461-473.
- Hedqvist, P. (1976):** Prostaglandin action on transmitter release at adrenergic neuroeffector junctions. In Advances in prostaglandin and thromboxane research. Vol.1: 357-363. Samuelsson, B. and Paoletti, R. (eds). New York: Raven Press.
- Henzel, M.R.; Smith, R.E. ;Boost, G. and Tyler, E.T. (1972):** Lysosomal concept of menstrual bleeding in humans. J. Clin. Endocrinol. 34: 860, 875.
- Herabutya, Y.and O-Prasertsawat, P.(1997):** Misoprostol in the management of missed abortion. Int. J. Gynecol. Obstet. 56(3): 263-6.

- Herxheimer, H. and Roetscher, I. (1971):** Effects of prostaglandin E₁, on lung function in bronchial asthma. Eur. J. Clin. Pharmac. 3 : 123-125.
- Heymann, M.A.; Rudolph, A.M. and Silverman, N.H. (1976):** Closure of the ductus arteriosus in premature infants by inhibition of prostaglandin synthesis. New Eng. J. Med. 295: 530-533.
- Hicks, J. B. (1969) :** On seatangle Tent . Practioner 3:83-90.
- Hillier, K. and Embrey, M.P. (1972):** High-dose i.v. administration of PGE₂ and F_{2a} for termination of mid-trimester pregnancies. J. Obster. Gynaecol. Brit. Cwlth. 79: 14-22.
- Hillier, K.; Button, A.; Corker, C.S.; Singer, A. and Eimbrey, M.P. (1972):** Brit. Med. J. 4: 333-336.
- Holmes, S.W.; Horton, E.W. and Main, I.M. H. (1963):** The effect of prostaglandin E₁ on response of smooth muscle to catecholamines, angiotensin and vasopressin. Br. J. Pharmac. 21: 538-543.
- Hughesdon, P.E. (1952):** The fibromuscular structure of the cervix and its changes during pregnancy and labor. J. Obstet. Gynaecol. Br. Commonw. 59: 763-776.

- Husslein ,P. ;Fuchs ,A.R.and Fuchs ,F .(1981):** Oxytocin and the initiation of human parturition: Prostglandin release during induction of labor by oxytocin. Am. J. Obstet. Gynecol. 141:688 – 93.
- Ian Donald. (1974):** Induced labor. Practical Obstetric Problems 480-512.
- Ito, A.; Kitamura, K.; Mori, Y. and Hirakawa, S. (1979):** The change in solubility of type I collagen in human uterine cervix in pregnancy at term. Biochem. Med. 21: 262-270.
- James Crossland; (1980):** Male prostaglandin and the female reproductive system. Lewis' Pharmacology. p.374.
- James Crossland; (1980):** Prostaglandin and fever. Lewis' Pharmacology.p.384.
- James Crossland; (1980):** Prostaglandin, hormones, and nucleotides, Lewis' Pharmacology. p.387.
- Jaye, M.; Shyken and Roy H. Petrie. (1995):** Oxytocin to induce labor. Clin. Obstet and Gynecol. Vol. 38. No. 2: 232-245.
- Jeeva, M.A. and Dommissse, J. (1982):** Laminaria tents or vaginal prostaglandins for cervical ripening. A comparative trial. S. Afr. Med. J. 61:402-403.

- John, D. ;Busowski and Michael, T. Parsons. (1995):** Amniotomy to induce labor. Clin. Obstet . Gynecol. Vol. 38. No. 2: 246-258.
- Johnson, N. (1989):** Intracervical tents usage and mode of action. Obstet. Gynecol. Survey.44:6.
- Judith Krammer and William, F. O'Brien. (1995):** Mechanical methods of cervical ripening. Clin. Obstet and Gynecol. Vol. 38, No. 2: 280-286.
- Kajanoja, P.; Ranta, T. and Seppala, M. (1978):** Prostaglandins. 16(2): 327-332
- Karim, S.M.M. (1970):** Once-month vaginal administration of prostaglandin E₂ and F₂ for fertility control. In Press. Quoted by Karim and Sharma, 1971b.
- Karim, S.M.M. (1970):** Use of prostaglandin E₂ in the management of missed abortion, missed labor, and hydatiform mole. Br. Med. J. 3: 196-197.
- Karim, S.M.M. (1971):** Action of prostaglandin in the pregnant woman. Ann. NY. Acad. Sci. 180: 483-498.
- Karim, S.M.M. (1972b):** Prostaglandin and human reproduction. Physiological role and clinical uses of prostaglandins in relation to human reproduction. In PGs: Progress in Research. 71-164.

- Karim, S.M.M. (1975a):** Advance in prostaglandin research. Prostaglandin and Reproduction ed. Karim S.M.M. (1975) p. 150.
- Karim, S.M.M. and Devlin, J. (1967):** Prostaglandin content of amniotic fluid during pregnancy and labor. J. Obstet. Gynaecol. Br. Commonw. 74: 230 – 234.
- Karim, S.M.M. and Hillier, K. (1973):** The role of prostaglandins in myometrial contraction, in J.B. Josimovich: Uterine contractions –side effects of steroidal contraceptives. New York: John Wiley and Sons, pp. 141-169.
- Karim, S.M.M. and Sharma, S.D. (1971):** Oral administration of prostaglandins for the induction of labor. Br. Med. J. 1: 260-262.
- Karim, S.M.M. and Sharma, S.D. (1971b):** Therapeutic abortion and induction of labor by the intravaginal administration of prostaglandins E₂ and F₂. J. Obstet. Gynaecol. Br. Commonw. 78: 294-300.
- Karim, S.M.M. and Sharma, S.D. (1972):** Termination of second trimester pregnancy with 15-methyl analogues of prostaglandins E₂ and F₂ α. J. Obstet. Gynaecol. Br. Commonw. 78: 737-743.
- Karim, S.M.M.; (1975):** Fetal –neonatal effect of PGF₂α used for induction of labor by i.v. route. In advances in prostaglandins

- research: Prostaglandins and reproduction. ed. Karim S.M.M. pp 124 – 167 Lancaster. Medical and Technical Publication.
- Karim, S.M.M.; Hillier, K.; Somers, K. and Trussell. (1971):** J. Obstet. Gynaecol. Br. Commonw. 78: 172. Quoted by Embrey, 1977.
- Karim, S.M.M.; Trussell, R.R.; Patel, R.C. and Hillier, K. (1968):** Response of pregnant human uterus to prostaglandin $F_{2\alpha}$. Induction of Labor. Br. Med. J. 4: 621-623.
- Karlson, S. (1959):** The influence of seminal fluid on the motility of the non-pregnant human uterus. Acta .Obstet. Gynecol. Scand. 38: 503.
- Karube, H.; Kanke, Y.; Mori, Y.; Hirakawa, S. and Hayashi, M. (1975):** Increase of structural glycoprotein during dilatation of human cervix in pregnancy at term. Endocrinol .Jpn .22: 445.
- Kase, N.G. (1983):** Principles and Practice of Clinical Gynecology. A Wiley Medical Publication. John Wiley and Sons. New York. Chichester. Brisbane. Toronto. Singapore. pp. 91-110.
- Kazzi, G.M.; Bottoms, S.T. and Rosen, M.G. (1982):** A closer look at the use of laminaria digitata for pre-induction ripening of the cervix: Its efficacy and safety. Society of Perinatal Obstetrics. San Antonio. Texas . p.23. Quoted by Steiner and Creasy .(1983).

- Keirse, M.J.N.C.(1993a):** Intravenous PGE₂ versus oxytocin for induction of labor. In Enkin, M.W. Keirse ,M.J.N.C.; Renfrew, MJ, Neilson ,I.P. (eds) Pregnancy and Childbirth Module. Conchrane Database of Systematic Reviews. Review no. 04529. 3 April, 1992. Disc Issue 2 Oxford: Update Software.
- Keirse, M.J.N.C (1993c):** Oral PGE₂ versus intravenous oxytocin for induction. In Eukin, M.W.; Keirse ,M.J.N.C.; Renfrew, M.J.; Neil son ,J.P. (eds). Pregnancy and childbirth module. Conchrane Database of Systematic Reviews. Review no. 03327, 3 April 1992. Disc Issue 2. Oxford Update Software.
- Keirse, M.J.N.C and Van Oppen, A.C.C. (1989):** Preparing the cervix for induction of labor. In Chalmers, I.; Enkin Mand Keirse, M.J.N.C (eds): Effective care in pregnancy and childbirth. Oxford University Press. p. 988.
- Keirse, M.J.N.C. (1975):** Studies on prostaglandin in relation to human parturition. D. Phil. Thesis University Oxford. Quoted by Keirse. 1978.
- Keirse, M.J.N.C. (1978):** Biosynthesis and metabolism of prostaglandins in the pregnant human uterus. Advances in Prostaglandin and Thromboxane Resarch. 4: 87-102.

Keirse, M.J.N.C.; Thiery, M.; Parewijck, W. and Mitchell. (1983):

Chronic stimulation of uterine prostaglandin synthesis during cervical ripening before onset of labor. Prostaglandin. 25: 671.

Kennedy, J.H.; Gordon-Wright, A.P.; Stewart, P.; Calder, A.A. and

Elder, M. G. (1982b) : Induction of Labor with a Stable-based Prostaglandin E₂ Vaginal Tablet. European Journal of Obstet. Gynecol. And Reproductive Biology. 13: 203-208.

Kennedy, J.H.; Stewrt, P.; Barlow, D.H.; Hillan E. and Cailder, A.A.

(1982d) : Induction of labor : Acomparison of a single prostaglandin E₂ vaginal tablet with amniotomy and intravenous oxytocin. Br.J. Obstet. Gynecol. 89: 704-707.

Kitamura, K.; Ito, A.; Mori, Y. and Hirakawa, S. (1980):

Glycosaminoglycans of human uterine cervix: Heparan sulfate increase with reference to cervical ripening. Biochem. Med. 23: 159-166.

Klein, L. and Yen, S.C.C. (1970): Urinary peptide hydroxyproline

before and during postpartum involution of human uterus. Metabolism 19:19.

Kleissl, H.P.; Van Der Rest, M.; Naftelin, F.; Glorieox, F.N. and De

Leon, A. (1978) : Collagen changes in the human uterine cervix at parturition. Am. J. obstet. Gynecol. 130: 748-753.

- Kurzrok, R. and Lieb, C.C. (1930):** Biochemical studies of human semen.II. The action of semen on the human uterus. Proc. Soc. Exp. Biol. Med. 28: 268-272.
- Lange, A.P.; Secher, N.J.; Westergaard, J.G. and Skovgard, L. (1982):** Pre- labor evolution of inducibility. Obstet. Gynecol. 20: 137.
- Legarth, J.; Guldbaek, E. and Secher, N.J. (1985):** The efficiency of prostaglandin E₂ vaginal suppository versus intracervical prostaglandin gel for induction of labor in patients with unfavorable Bishop score. Archieves in Gynecology Abstracts. XI World Congress of Gynecology and Obstetrics. Berlin West. Abstract. No. 04. 42. 11.
- Lennon , G. G. (1957) :** Proc . Roy . So c. Med. 50, 793.
- Leppert ,P.C.; Keller, S.; Cerreta, J.and Mandl,I. (1982):** Conclusive evidence of the presence of elastin in human and monkey cervix. Am. J. Obstet. Gynecol. 142: 179.
- Leppert, P.C. and YU SY (1991):** Three dimensional structures of uterine elastic fibers, Scanning electron microscopic studies. Connective Tissue Res 26: 15.
- Liggins, G.C. (1962):** The treatment of missed abortion by high dose of syntocinon intravenous infusion. J. Obstet. Gynaecol .Br. Commonw. 69 : 277-282.

- Liggins, G.C. (1978):** Ripening of the cervix. In Oliver ,T.K. and Kirschbaum, T.K. (eds). New York. Sem. Perinatol. 2: 261.
- Liggins, G.C. (1979):** Controlled trial of induction of labor by vaginal suppositories containing prostaglandin E₂. Prostaglandins. 18: 167-172.
- Liggins, G.C.; Fairclough, R.J.; Griever, S.A.; Kendall, J.Z. and Knox, B. S. (1973):**The mechanism of initiation of parturition in the ewe. Recent .Prog .Horm. Res. 29: 111.
- Lindmark, G.; Zador,G. and Nilsson, B.A. (1975):** The induction of labor with prostaglandin F₂ by intravenous nfusion. Acta .Obstet. Gynecol. Scand. (suppl.). 37: 17-26.
- MacDonald, D. (1970):** Surgical induction of labor. Am. J. Obstet. Gynecol .107: 908-911
- Macer, J.; Buthanon, D. and Yonekura ,M.L. (1984):** Induction of labor with prostaglandin E₂ vaginal suppositories. Obstet .Gynecol.63:664-668.
- Mackenzie, I.Z. ; Bradley, S. and Embrey, M. P. (1981) :** A simpler approach to labor induction using lipid-based prostaglandin E₂ vaginal suppository. Am .J. Obstet. Gynecol. 141:158-162.
- Mackenzie, I.Z. and Embrey, M.P. (1977):** Cervical ripening with intravaginal prostaglandin E₂ gel. Br. Med. J. 2: 1381-1384.

- Mackenzie, I.Z. (1988):*** The unripe cervix and its management for labor induction. *Ann Acad. Med. Singapore.* 151-7.
- MacVicar, J. (1971):*** *J. Obstet. Gynaec. Brit. Cwlth*, 78: 1007.
- Main, I.H.M. (1964):*** The inhibitory actions of prostaglandins on respiratory smooth muscle. *Br. J. Pharmac.* 22: 511-519.
- Malik, K.U. and McGiff, J.C. (1976):*** Cardiovascular actions of prostaglandins. In *Prostaglandins: physiological, pharmacological and pathological aspects.* pp. 103 – 200. Karim, S.M.M. (ed) Lancaster: M.T.P.
- Malmsten, C.; Hamberg, M.; Svensson, J. and Samuelsson, B. (1975) :*** Physiological role of an endoperoxide in human platelets . Haemostatic defect due to platelet cyclo-oxygenase deficiency. *Proc. Natn. Acad. Sci. Sa. USA.* 72: 1446-1450.
- Manabe, Y. (1971):*** Laminaria tent for gradual and safe cervical dilatation. *Am. J. Obstet. Gynecol.* 110:743-745.
- Morgulies, M.; Perez ,G.C. and Voto ,L.S.(1992):*** Misoprostol to induce labor. *Lancet.* 339, 64.
- Mathie, J.K.G. and Dawson, B.H. (1959):*** Effect of castor oil, soap enema and hot bath on the pregnant human uterus near term. *Br. Med. J.* i.1162.

- Mati ,J.K.G.; Horrobin, D.F.and Bramley, P.S. (1973):** Induction of labor in sheep and in humans by single doses of corticosteroids. B.M.J. 2:149-151.
- Myer, S. Bornstein and Don Shuwarger. (1996):** Misoprostol (Cytotec) for cervical ripening and induction of labor. www.obgyn.net a service of Elecomm Corp.
- McCallum, M. F. and Govan, A.D.T. (1963):** J. Obstet. Gynaec, Brit, Cwlth.70: 244.
- McColgin, S.W.; Hampson, H.L. and McCaul, J.F. et al. (1990):** Stripping membranes at term. Can it safely reduce the incidence of post-term pregnancy. Obstet. Gynecol. 76: 678.
- McInnes, D.R.; Naftolin, F. and Van Der Rest, M. (1980):** Cervical changes in pregnant women. In Naftolin ,F. and Stubblefield, P.G. (eds).Dilatation of the uterine cervix. New York. Raven Press. p.181.
- MacLennan, A.H. and Green, R.C. (1980):** A double blind dose trial of intravaginal prostaglandin F₂ α for cervical ripening and the induction of labor. Aust. N. Z.J. Obstet. Gynecol. 20: 80-83.
- MacLennan, A.H.; Green, R.C.and Grant ,B. (1985):** Ripening of the human cervix and induction of labor with intracervical purified porcine relaxin. Obstet. Gynecol. 92: 693.

- Mellows, H. J.; Sims, C.D. and Craft, I. (1977):** Prostaglandin E₂ in tylose for induction of labor in patients with a favorable cervix. Proc. Roy. Soc. Med. 70: 537-539.
- Mitchell; Flint, A.P.F. and Bibby, J. et al (1977):** Rapid increase in plasma prostaglandin concentrations after vaginal examination and amniotomy. B.M.J. 2:1183-1185.
- Mochizuki, M. and Tojo, S. (1980):** Effect of dehydroepiandrosterone sulfate on softening and dilatation of the uterine cervix in pregnant women. In Naftolin, F. and Stubblefield, P.G. (eds): Dilatation of the uterine cervix. New York. Raven Press. p. 267.
- Morrisson, A.R.; Nishikawa, K. and Needleman, P. (1977):** Unmasking of thromboxane A₂ synthesis by ureteral obstruction in the rabbit kidney. Nature.Lond. 267: 259-260.
- Munro, C.D.; Calder, A.A.; Fleming, R.; McMannus, T.J.; Anghey, B.; McPhee, G.; Hillier, K. and Coutts, J.R.T. (1978):** Paper presented to the society for the study of fertility. December 13. London, 1978. Quoted by Stewart et al (1981).
- Murota, S.; Abe, M. and Otsuka, K. (1977):** Stimulatory effect of prostaglandins on the production of hexosamine-containing substances by cultured fibroblasts. Induction of hyaluronic acid synthetase by prostaglandin F₂. Prostaglandins. 14: 983-991.

- Murphy, B.E.P. (1973):** Amer. J. Obstet. Gynec. 115: 521.
- Naftolin, F. and Stubblefield, P.G. (1981):** Dilatation of the uterine cervix. Connective tissue biology and clinical management. New York: Raven Press. Quoted by Uldbjerg et al.(1983).
- Naismith, W.C.M.K.; Barr, W. and MacVicar, J. (1972):** Induction of labor by simultaneous intravenous administration of prostaglandin E₂ and oxytocin. Br. Med. J. 4: 461-462.
- Najak, Z.; Hillier, K.; Karim, S.M.M. (1970):** The action of prostaglandins on the human isolated non pregnant cervix. J. Obstet. Gynaecol. Br. Commonw.77: 701.
- Newton, B.W. (1972):** Laminaria tent: Relic of the past of modern medical device? Am. J. Obstet. Gynecol. 113:442-448.
- Nicolaides, K.H.; Welsh, C.C.; Koullapis, E.N.and Filshiew, G.W. (1983) :** Cervical dilatation by lamicel-studies on mechanism of action. Br. J .Obstet. Gynecol.90:1060.
- Norchi, S.; Zanini, A.; Ragussa, A.; Maccario, L.and Valle, A. (1993):** Induction of labor with intravaginal prostaglandin E₂ gel. Int .J. Gynecol. Obstet. 42:103 – 7.
- Nugteren, D.H. and Hazelhof, E. (1973):** Isolation properties of intermediate in prostaglandin bio-synthesis. Biochim. Biophys. Acta. 326: 438. Quoted by Kase.(1983).

- O'Herlihy, C. and MacDonald, H.N. (1979):** Influence of pre-induction prostaglandin E₂ vaginal gel on cervical ripening and labor. *Obstet. Gynecol.* 54: 708-710.
- Orga, S.S.; Kirton, K. T.; Tomasi, T.B. and Lippes, J. (1974):** Prostaglandins in human fallopian tube. *Fertil. And steril.* 25: 250-255.
- Page, E.W. (1943):** Response of human pregnant uterus to pitocin tannate in oil . *Proceeding of the society for experimental biology and medicine* .52:195-197. Quoted from Davey.1980.
- Parikh, S.; Garud, M.; Parikh, V. and Tejani, T. (1985):** Use of a new oral oxytocin derivative in labor. *Archives in Gynecology Abstracts.* X1. World Congress of Gynecology and Obstetrics. Berlin West. Abstract No.13. 42. 06.
- Peel, J.H. (1936):** Induction of labor for disproportion. *Lancet.* I. 935.
- Pentecost, A.F. (1973):** The effect of buccal "pitocin" on the unripe cervix. *Curr. Med. Res. Opin.* 1: 482-488
- Physicians Desk Reference. 44 edition (1990).** Publisher: EDWARD. BARNHART.
- Pickles, V.R. and Ward, P.V.F. (1965):** *J. Physiology (London)* 178: 38, 99.

- Pickles, V.R.; Hall, W.H.; Best, F.A. and Smith, G.N. (1965):** J. Obstet. Gynecol. British. Commonwealth, 72: 185. Quoted from Karim and Hiller. 1973.
- Ponnath, H. and Weitzel, H. (1985):** Induction of labor with prostaglandin E₂. Archives in Gynecology. Abstracts XI World Congress of Gynecology and Obstetrics. Berlin West. Abstract No. 01. 14. 28.
- Praagh, I.V. and Hendricks, C.H. (1964):** The effect of amniotomy during labor in multiparas. Obstet. Gynecol. 24:258-265.
- Pritchard, J.A.; MacDonald, P.C. and Gant, N.F. (1985):** Williams Obstetrics. Seventeenth Edition. Appleton-Century-Crofts/Norwalk, Connection. pp. 295-321.
- Puig-Parellada, P. and Planas, J.M. (1977):** Action of selective inhibitor of thromboxane synthetase on experimental thrombosis induced by arachidonic acid in rabbits. Lancet. (ii). 40.
- Raisz, L.G.; Dietrich, J.W. and Simmons, H.A (1977):** Effect of prostaglandin endoperoxides and metabolites on bone resorption in vitro. Nature. Lond. 267: 532-534.
- Rall, T.W. and Schleifer, L.S. (1985):** Oxytocin, prostaglandins, ergot alkaloids and other drugs: tocolytic agents. In Gilman, A.G.; Goodman, L.S.; Rall, T.W.; Murad, F.; Goodman and Gilman's.

The Pharmacological Basis of Therapeutics. 7th edition. New York: Macmillan Publishing.

Ramwell, P.W.; Foegh, M.; Loebe, R. and Leovey, E.M.K. (1980):

Synthesis and metabolism of prostaglandins, prostacyclin and thromboxane. The arachidonic Cascade. Semin. Perinatol. 4: 3.

Quoted from Sperroff, et al., 1983.

Robert, A.; Nezamis, J.E. and Phillips, J.P. (1968): Effect of

prostaglandin E₁ on gastric secretion and ulcer formation in the rat. Gastroenterology. 55: 481-487.

Roberts, G. and Turnbull, A.C. (1971): Uterine hypertonus during

labor induced by prostaglandins. Br. Med. J. 1: 702-705.

Roden, L. (1980): Structure and metabolism of connective tissue

proteoglycans. In Lennarz, W.J. (ed): The biochemistry of glycoproteins and proteoglycans. New York and London. Plenum Press. p 267.

Rouben, D. and Arias, F. (1993): A randomized trial of extra-amniotic

saline infusion with intracervical Foley's catheter balloon

versus prostaglandin E₂ vaginal gel for ripening the cervix and

inducing labor in patients with unfavorable cervix. Obstet .

Gynecol .82: 290.

Samuelsson, B. (1977): The role of prostaglandin endoperoxides and

- thromboxanes in human platelets. In prostaglandins ,in hematology.pp. 1-10. Silver, M.J.; Smith, J.B. and Kocsis, J.J. (eds.) New York: Spectrum.
- Samuelsson, B.; Goldyne, M.; Granstrom, E.; Hamberg, M.; Hammarstrom, S. and Malmsten, C. (1978) :** Prostaglandins and thromboxanes. Ann. Rev.Biochem. 47: 997-1029.
- Sanborn, B.M. ; Held, B. and Kuo, H.S. (1976) :** Hormonal action in human cervix 11. Specific progestogen binding proteins in human cervix. J. Steroid. Biochem. 7: 665.
- Sanborn, B.M. ; Held, B. and Kuo, H. S. (1975) :** Specific estrogen binding proteins in human cervix. J. Steroid. Biochem. 6:1107.
- Sanchez-Ramos, L.; Kaunitz, A.M.; Del Valle Go; Delke, I.; Schroeder, P.A. and Briones, D.K. (1993) :** Labor induction with the prostaglandin E₁ methyl analogue Misoprostol: A randomized trial. Obstet. Gynecol. 81:332-6.
- Sanchez-Ramos, L.; Kaunitz, A. M. and Connor P.M. (1992):** Hygroscopic cervical dilators and prostaglandin E₂ gel for pre-induction cervical ripening: A randomized prospective comparison. J. Reprod. Med. 1: 335.
- Sandberg, L.B. (1976) :** Elastic structure in health and disease. Int. Rev. Connect. Tissue. Res. 7: 665.

- Saunders ,W.G. and Munsick , R.A . (1966):** Antidiuretic potency of oxytocin in women post-partum. Am. J. Obstet .Gynecol. 95:5-11.
- Shaala ,S.; Darwish, E.; Anwar, M.; Rocca, M.and Ismail, A.A. (1989):** Cervical prostaglandin injection : A novel method of administration for ripening the cervix and induction of labor. Int .J. Gynecol. Obstet. 30:221-3.
- Shepherd,J. H.; Bennett, M.J.; Laurence, D.; Moore, F.and Sims, D. (1981):** Prostaglandin vaginal suppositories: A single and safe approach to the induction of labor. Obstet. Gynecol. 58: 596.
- Shepherd, J.H.; Pearce, J.M.F. and Sims, C.D. (1979):** Induction of labor using prostaglandin E pessaries. Br. Med. J. 108-110.
- Shimizu, T.; Endo, M. and Yosizawa, Z. (1980):** Glyco-conjugates (glycosaminoglycans and glycoproteins) and glycogen in the human cervix uteri . Tohoku .J. Exp. Med. 131: 289.
- Sims, C.D. (1985):** The management of labor. Blackwell Scientific Publications Oxford. London Edinburgh. Boston .Palo. Aito. Melbourne. pp. 133-143.
- Sims, C.D.; Mellows, H.J.; Spencer ., P.J. and Craft, I.L. (1979):** Routine induction of labor with extra-amniotic prastaglandin E₂ in a viscous gel. Br. J. Obstet. Gynaecol. 86: 529-532.

- Spence, D.N. and Chalmers, J.A. (1964):** Buccal oxytocin in the induction of labor. *Lancet*. I. 633. Quoted by Davey. 1980.
- Speroff, L.; Glass, R.H. and Kase, N.C. (1983):** Clinical Gynecologic Endocrinology and Infertility. Third Ed. Williams and Wilkins. Baltimore/London : p. 307 – 333.
- Srisomboon, J.; Tongsong, T. and Tosiri, V. (1996):** Pre-induction cervical ripening with intravaginal prostaglandin E₁ methyl analogue Misoprostol: a randomized controlled trial. *J. Obstet. Gynecol. Res.* 22(2): 119-124.
- Steiner, A.L. and Creasy, R.K. (1983):** Methods of cervical priming. *Clin. Obstet. Gynecol.* 26:37-46.
- Steinetz, B.G.; O'Bryne, E.M. and Kroe, R.L. (1980):** The role of relaxin in cervical softening during pregnancy in mammals. In Naftolin, F. and Stubblefield, P.G. (eds): *Dilatation of the uterine cervix*. New York. Raven Press. p. 155.
- Stewart, P.; Kennedy, J.H.; Barlow, D.H. and Calder, A.A. (1981):** A comparison of estradiol and prostaglandin E₂ for ripening the cervix. *Br. J. Obstet. Gynaecol.* 88: 236.
- Stys, S.J.; Dresser, B.L.; Otte, T.E. and Clark, K.E. (1981):** Effect of prostaglandin E₂ on cervical compliance in pregnant ewes. *Am. J. Obsstet Gynecol.* 140: 415-419.

- Sweatman, W.J.F. and Collier, H.O.J. (1968):** Effect of prostaglandins on human bronchial muscle. *Nature.Lond.* 217: 69.
- Smythe, H. J. D. (1931):** Indications for induction of premature labor. *Br. Med. J. i.* 1018.
- Smythe, H. J. D. and Thompson, D.J. (1937):** Induction of labor by rupture or high puncture of the membranes. *J. Obstet. Gynaec, Br. Emp.* 44: 480.
- Szalay, S.; Husslein, P. and Grunberger W. (1981):** Local application of prostaglandin E₂ (PGE₂) and its influence on collagenolytic activity of: cervical tissue. *Singapore .J. Obstet. Gynecol.* 12: 15.
- Thiery, M. (1979):** Ripening of the cervix by prostaglandins at or near term. *International Congress Series No. 512 Gynecology and Obstetrics.* In Shoichi Sakamoto; Shimpei Tojo, and Tetsuya Nakayama: *Proccedings of The IX World Congress of Gynecology and Obstetrics.* Tokyo, October 25-31 (1979). Amsterdam-Oxford-Princeton: Excerpta Medica.
- Theobald, G. W. (1930):** Malaria, women and quinine. *Proc. R. Soc. Med.* 23, 31.
- Theobald, G. W. (1963):** Induction of labor and of premature labor. In *British Obstetric and Gynecological Practice.* ED. Claye, A. and Bourne, A. 3rd ed. p. 1055. London: Heinemann.

Theobald, G.W. (1973): The induction of labor. *Obstetric Therapeutic.*

Clinical Pharmacology and Therapeutics in Obstetric Practice
.341-379.

Theobald, G.W.; Graham, A.; Campell, J.; Gavage, P. and Driscoll,

W.J. (1948) : The use of posterior pituitary extract in physiological
amounts in obstetrics. *Br. Med. J.* ii. 123. Quoted by Clarke,
1975.

Thiery, M.; De Gezelle, H.; Van Kets, H.; Voorhoof, L. Verheugen,

C. ; Sims , R. Gerris , J. and Martin , G. (1978) : Extra-amniotic
oestrogen for the unfavorable cervix (Letter). *Lancet* 2: 835.

Thomas, F.V. and Dorflinger, C.T. (1978): Mechanism of rapid

antigonadotrophin action of PG on cultural luteal cell –
Proceeding of the National Academy of Sciences .75. (3). 1344 –
1348 March 1978.

Tiry Jones , P . and Malvern , J.(1973):Personal communication,

Queen Charlotte's Year Book.Quoted by Shepherd et al .(1979).

Toaff,M.F.; Hezroni,J. and Toaff,R.(1987):Induction of labor by

pharmacological and physiological dose of intra-venous oxytocin.

*Br.J.Obstet.Gynecol.*85:101-8.Quoted by Sims,1985.

- Tohan, N.; Tejani, N.A.; Varanasi, M. and Robins, J. (1979):**
Ripening of the term cervix with laminaria. *Obstet. Gynecol.*
54:588-590.
- Toplis, P.J. and Sims, C.D. (1979):** Prospective study of different
methods and routes of administration of prostaglandin E₂ to
improve the unripe cervix. *Prostaglandins*. 18: 127-136.
- Trofatter, K.F. (1993):** Endocervical prostaglandin E₂ for pre-induction
cervical ripening: Clinical trial results. *J. Reported. Med* 38:78-
82.
- Tromans, P.M.; Beazley, J.M. and Shenouda, P.I. (1981):**
Comparative study of estradiol and prostaglandin E₂ vaginal gel
for ripening the unfavorable cervix before induction of labor. *Br.*
Med. J. 282: 679.
- Turnbull, A.C. and Anderson, A.B.M. (1968a) :** Induction of labor .
Part II: Intravenous oxytocin infusion. *J. Obstet. Gynecol. Br.*
Commonw. 75: 24-31.
- Turnbull, A.C. and Anderson, A.B.M. (1968b):** Induction of labor.
Part III. Results with amniotomy and oxytocin "titration" .*J.*
Obstet. Gynecol. Br. Commonw 75: 24-31.
- Tweedy, E. H. and Wrench, G.T. (1919) :** *Practical Obstetrics*, 4th ed.
London: Hodder and Stoughton.

- 1- **Uldbjerg, N.; Ekman, G.; Malmstrom, A.; Olsson, K. and Ulmsten, U. (1983a)** : Ripening of the human uterine cervix related to changes in collagen, glycosaminoglycans, and collagenolytic activity. *Am. J. Obstet. Gynecol.* 147: 662-667.
- Uldbjerg, N.; Ekman, G.; Malmstrom, A.; Sporrang, B.; Ulmsten, U. and Wingerup, L. (1981)**: Biochemical and Morphological Changes of Human Cervix after Local Application of Prostaglandin E₂ in Pregnancy. *Lancet.*1: 267-268.
- Uldbjerg, N.; Malmstrom, A.; Ekman, G.; Sheehan, J.; Ulmsten, U. and Wingerup, I. (1983c)** : Isolation and characterization of dermatan sulphate proteoglycan from human uterine cervix. *Bio. Chem. J.* 209: 497. Quoted by Uldbjerg et al., 1983c.
- Uldbjerg, N.; Ulmsten, U. and Ekman, G. (1983b)**: The ripening of the human uterine cervix in terms of connective tissue biochemistry. *Clin. Obstet. Gynecol.* 26:14-26.
- Ulmsten, U.; Wingerup, L. and Andersson, K.E. (1979b)**: Comparison of prostaglandin E₂ and intravenous oxytocin for induction of labor. *Obstet. Gynecol.* 54:581:584.
- Ulmsten, U.; Wingerup, L.; Belfrage, P.; Ekman, G. and Wijkvist, N. (1982)** : Intra-cervical application of prostaglandin gel for induction of term labor. *Obstet. Gynecol.* 59: 336-339,

- Urquhart ,R. and Templeton, A. (1987):** Mifeprystone and second trimester termination. Lancet.ii:1405.
- Visscher, R.D.; Struyk, C.D. and Visscher, H.G. (1977):** Guidelines for the elective induction of labor with oral prostaglandin E₂. Obstet. Gynecol. 49: 15.
- Von, Euler, U.S. (1936):** J. of Physiology. 88: 213. Quoted by Karlson, S. The influence of seminal fluid on the motility of the non-pregnant human uterus. Acta. Obstet.Gynecol. Scand. 38: 503-521.
- Von Euler, U.S. (1939):** Scandinav. Arch. F.Physiol. 81: 61. Quoted by Karlson, S.: The influence of seminal fluid on the motility of the non-pregnant human uterus. Acts. Obstet. Gynecol. Scand. 38: 503-521.
- Von Malliot, K.V. and Zimmermann, B.K. (1976):** The solubility of collagen of the uterine cervix during pregnancy and labor. Arch .Gynakol. 220: 275.
- Von Malliot, K.V.; Stuhlsatzh, W.; Mohanarad hakrishnan, V. and Greiling , H. (1979):** Changes in glycosaminoglycans distribution pattern in the human uterine cervix during pregnancy and labor. Am.J.Obstet. Gynecol. 135: 503.

- Voutsas, N.; Beveridge, G.C. and Foraker, A.C. (1963):** The uterine cervix at term : Glycogen, mast cells and connective tissue elements. *Obstet. Gynecol.* 22: 108.
- Wallenberg, H.C.S.; Keirse, M.J.N.C.; Freie, H.M.P. and Blacquiere, J.F. (1980) :** Intramuscular administration of 15 (S) -15-methyl prostaglandin F₂ alpha for induction of labor in patients with fetal death. *Br.J. Obstet. Gynaecol.* 87: 203-209.
- Watson, B.P. (1922):** Further experience with pituitary extract in the induction of labor. *Am. J. Obstet. Gynecol.* 4: 603.
- Watson, B.P. (1920):** Induction of labor: indications and methods with special reference to the use of pituitary extract. *Am. J. Obstet. Gynecol.* 1: 70.
- Weiss, G.; Goldsmith, L.T.; Sachdev, R.; Von Hagen, S. and Lederer, K. (1993) :** Elevated first-trimester serum relaxin concentrations in pregnant women following ovarian stimulation, predict prematurity risk and preterm labor. *Obst. Gynecol.* 82:821-828.
- Wentz, A.C. and Jones, G.S. (1973):** Transient luteolytic effect of PGF₂a in the human . *Obstet. Gynecol (N.Y.)* 42: 172-181.
- Wheeler, R.J. and Schneider, K. (1983):** Properties and safety of cervical dilators. *Intelligence Rep. Obstet. Gynecol.* 2:24.

Wing, D.A.; Rahall, A.; Jones, M.M.; Goodwin, T.M. and Paul, R.H.

(1995): Misoprostol : an effective agent for cervical ripening and labor

induction. Am. J. Obstet. Gynecol. Vol.172. no.6 pp. 1811-6.

Wingerup, L.; Ulmsten, U. and Andersson, K.E. (1979): Ripening of

the cervix by intra-cervical application of PGE₂ gel before

termination of pregnancy with dilatation and evacuation. Acta.

Obstet. Gynecol. scand (suppl). 84: 15.

Wrigley, A. J. (1958): Lancet. 1. 1167.

Young, J.H. and James Hamilton (1963): Obstetrician and

controversialist. Med.Hist.:7:62-73.