



References



References

- Acton C, Davitt J and Ryan E (1988) :** Hystero-salpingo-graphy in infertility-an experience of 3631 examinations. Aust NZ J Obstet Gynaecol, 28:127.
- Adelusi B, Khashoggi T, Chowdhury N, Makanjuola D, Al-Nuaim L and Kangave D (1995) :** Accuracy of hysterosalpingography and laparoscopic hydrotubation in diagnosis of tubal patency. Fertil steril, 63:5.
- Ansari A.H (1979) :** Diagnostic procedures for assessment of tubal patency, Fertil Steril, 31:469.
- Ayida G, Kennedy S, Barlow D and Chamberlain P (1996):** A comparison of patient tolerance of hystero-salpingo-contrast-sonography (HY-CO-SY) with Echovist-200 and X-ray hysterosalpingo-graphy for outpatient investigation of infertile women. Ultras Obstet Gynecol, 7:201.

- Balen F.G, Allen C.M, Siddle N.C and Lees W.R**
(1993): Ultrasound contrast hysterosalpingography-
evaluation as an out-patient procedure. Br J Radiol,
66:592.
- Barbaric Z.L (1994) :** Radiographic contrast material In :
Principles of Genitourinary Radiology.
Edit Barbaric Z.L, 2nd ed. New York, 58:58.
- Barbot J (1995):** Hysteroscopy and hysteroigraphy.
Obstet Gynecol Clin North Am, 22:591.
- Beyth Y, Nowat D and Lax E (1985) :** A simple
improvement in the technique of hysterosalpin-
gography achieving optimal imaging and avoiding
possible complications. Fertil Steril, 44:543.
- Bonilla-Musoles F, Simon C and Serra V. (1992):**
An assessment of hysterosalpingo-
sonography (HSSG) as a diagnostic tool for uterine
cavity defects and tubal patency.J Clin Ultras,20:175.

Charles J. F (1988): Hysterosalpingography : Textbook of Diagnostic Imaging, Edit Putman C.E and Ravin C.E. Volume three, W.B Saunders company, Philadelphia, 145:2059.

Crequat J, Pennehouat G and Cornier E (1993): Intrauterine pathology and tubal patency evaluation by contrast ultrasonography. *Contracep Fertil Sex*, 861.

Daniel W.W (1991): Biostatistics : A foundation for analysis in the health science. 5th ed., Edit John Wiley and Sons. London, 3:70.

Davison G.B and Leeton J (1988): A case of female infertility investigated by contrast-enhanced echogynecography. *J Clin. Ultras.* 16:44.

De boer A.D, Vemer H.M, Willemsen W.N.P and Sanders F.M.B (1988): Oil or aqueous contrast media for hysterosalpingography : a prospective, randomized, clinical study. *Eur J Obstet Gynaecol Reprod Biol*, 28:65.

- Decherney A. H, Kort H, Barney J.B and Devore G.R (1980) :** Increased pregnancy rate with oil-soluble hysterosalpingography dye. *Fertil Steril* 33:407.
- Deichert U (1993):** Contrast hysterosalpingography (Hy-Co-Sy). *Contracep Fertil Sex*, 21(3):213.
- Deichert U, Schlieff R, Vandesandt M and Juhnke I (1989):** Transvaginal hysterosalpingo-contrast-sonography (Hy-Co-Sy) compared with conventional tubal diagnostics. *Hum Reprod*, 4:418.
- Deichert U, Schlieff R, Vandesandt M and Daume E (1992):** Transvaginal hysterosalpingo-contrast sonography for the assessment of tubal patency with gray scale imaging and additional use of pulsed wave doppler. *Fertil Steril*, 57:62.
- Feinstein S.R, Cheirif J, Ten cate E.J, Silverman P.R, Heidenreich P.A, Dick C, Desir R.M, Armstrong W. F, Quinones M.A and Shoihi P.M (1990):** Safety and efficacy of a new trans-pulmonary ultrasound agent : Initial multicenter clinical results. *J Am Coll Cardiol*, 16:316.

Friberg B and Joergensen C (1994): Tubal patency studied by ultrasonography. *Acta Obstet Gynecol Scand* 73:53.

Fritsch T.h, Mutzel W and Scharl M (1986): First experiences with a standardized contrast medium for sonography In: Otto Rch, Higgins ch.B (eds) *new developments in imaging*. Thieme-Verlag Stuttgart, New York.

Fritsch T.h, Scharl M and Siegert J (1988): Pre-clinical and clinical results with an ultrasonic contrast agent. *Invest Radiol*, 23(suppl I):302.

Fritsch T, Hilmann J, Kampfe M, Muller N, Schoobel C and Siegert J (1990): SHU 508, a transpulmonary echocontrast agent : Initial experience. *Invest Radiol*, 25:160.

Gavant M.L, Ellis J.V and Klesges L.M (1992) : Diagnostic efficacy of excretory urography with low-dose, non ionic contrast media. *Radiology*, 182:657.

Gleicher N, Thurmond A, Burry K and Coulam C

(1992): Gynecoradiology : a new approach to diagnosis and treatment of tubal disease, *Fertil Steril*, 58:885.

Heikkinen H, Martikainen H, Aydin T, Pentti J

and Volpi E (1995): Transvaginal salpingography for the assessment of tubal patency in infertile women : methodological and clinical experiences. *Fertil Steril*, 64:2.

Henkel B and Schlieff R (1987): Transvaginal contrast-

sonographic assessment of fallopian tube patency in : Euro Son 87 proceedings of the sixth congress of the European Federation of Societies for Ultrasound in Medicine and Biology, Helsinki, Finland, June 14-18. Helsinki: Finnish Society for Ultrasound in Medicine and Biology, 223.

Highman J.H and William R.L (1987): Gynaecological

Imaging : A textbook of Radiology and Imaging.

Edit Sutton D. 4th ed, volume 2, Churchill

Livingstone, London, New York. Part 5, 47:1246.

- Hoffman J (1992):** Anatomy and Physiology of the Fallopian Tube : Atlas of female infertility surgery. Edit Hunt B.R, 2nd ed, Mosby Year Book, 1: 3.
- Horwitz R.C, Morton P.C, Shaff M.J and Hugo P (1979) :** A radiological approach to infertility-hysterosalpingography. Br J Radiol 52: 255.
- Innocenti P, Pandimiglio A.M, Nicolucci A and Nori Bufalinin G (1989):** Use of non-ionic contrast medium (omnipaque 350) in hysterosalpingography. A comparison with ionic contrast media-Radiol Med, 77:4.
- Karasick S (1991):** Hysterosalpingography. Urologic Radiology, 13 : 67.
- Kasby C.B (1980):** Hysterosalpingography : An appraisal of current indications. Br J Radiol, 53 : 279.
- Kaufman R.H, Adam E, Noller K (1986) :**
Upper genital tract changes and infertility in diethylstilbestrol exposed women. Am J Obstet Gynecol 154:1312.

Kaufman F.R, Xu Y.K and Ng W.G (1989):

Gonadal function and ovarian galactose metabolism in classic galactosemia.

Acta Endocrinol. (Copenh), 120.

Kessler R.M and Altman D.H (1982): Real time

sonographic detection of vesicourethral reflux in children. A J R, 138:1053.

Komrower G.M (1982): Galactosaemia thirty years on :

The experience of a generation. J Inherited Metab. Dis., 5 (suppl) : 96.

Lange L, Fritsch T.H, Hilmann J, Kubowicz G

and Mutzel W (1986): Right heart echo-contrast in the anaesthetized dog after i.v administration of a new standardized sonographic contrast agent (II).

Arzneimittel for Sebung, 36 : 1034.

La Sala G.B, Sacchetti M.D, Deglincertitocci F,

Dessanti L and Torrelli M.G (1987): Complementary use of hysterosalpinography, hysteroscopy and laparoscopy in 100 infertile patients : Results and comparison of their diagnostic accuracy. Acta Eur Fertil, 18:369.

Lindequist S, Justesen P, Larsen C and Rasmussen F
(1991): Diagnostic quality and complications of
hystero-salpingography oil-versus water soluble
contrast media-a randomized prospective study.
Radiology, 179 : 69.

Loy R.A, Weinstein F.G and Seibel M.M (1989) :
Hysterosalpingography in perspective : the predictive
value of oil soluble versus water soluble contrast
media. Fertil Steril, 51 : 170.

Madanes A, Homer M, Turksoy R.N and Faber M
(1982): A single diverticulum of the fallopian tube.
Fertil Steril, 37:121.

Martin M.C (1991): Infertility : current obstetric &
gynecologic diagnosis & treatment. 7th ed,
Edit Pernoll M.L. Appleton and Lange, section
VI, 51:1031.

McComb P, Gomel V and Rowe T (1986):
Investigation of tuboperitoneal causes of female
infertility : Infertility male and female. Edit
Insler V and Lunenfeld B. Churchill Livingstone,
London and New York, 9: 215.

- Meyerovitz M.F (1991) :** Hysterosalpingography and fallopian tube cannulation : use of a double balloon introducing catheter. Radiology, 181: 901.
- Mitri F.F, Andronikou A.D, Perpinyal S, Hofmeyr G.J and Sonnendecker E.W.W (1991):** A clinical comparison of sonographic hydrotubation and hystero-salpingography, Br J Obstet Gynaecol, 98:1031.
- Ophir J and Paker K.J (1989):** Contrast agents in diagnostic ultrasound-Ultras Med Biol, 15:319.
- Parek M and Arronet G (1972):** Diagnostic procedures and methods in the assessment of pelvic organs with specific reference to infertility. Clinic Obstet Gynecol, 15 : 140.
- Pontifex G, Trichopoulos D and Karpathios S (1972) :** Hysterosalpingography in the diagnosis of infertility (statistical analysis of 3437 cases). Fertil Steril 23: 829.

Randolph J.R, Ying Y.K and Marier D.B (1986):

Comparison of real-time ultrasonography,
hystero-salpingography and laparoscopy/ hystero-
scopy in the evaluation of uterine abnormalities and
tubal patency, Fertil Steril, 46: 828.

Rasmussen F, Lindequist S, Larsen C and Justesen P

(1991): Therapeutic effect of hysterosalpingography :
Oil-versus water-soluble contrast media-a randomized
prospective study. Radiology, 179 : 75.

Rice J.P, London S.N and Olive D.L (1986):

Revaluation of hysterosalpingography in infertility
investigation. Obstet Gynaecol 67 : 718.

Richman T.S, Viscomi G.N, Decherney A, Polan M.L

and Olson Alcebo L (1984): Fallopian tubal
patency assessed by ultrasound following fluid
injection. Radiology, 152 : 507.

Schlieff R (1987): First steps in ultrasound contrast media

in : Felix R, ed. Contrast media from the past to the
future, Stuttgart : Thieme, 179:187.

Schlief R (1988): Echovist : physico-pharmacological properties, results of clinical studies and possible fields of use of a novel ultrasound contrast medium (in German). In Cockel HP, ed. Jahrbuch Der Radiologie. Regensburg & Biermann, 163:170.

Schlief R (1991): Ultrasound contrast agents. Current opinion in radiology, 3: 198.

Schlief R, Staks T, Mahler M, Rufer M, Fritzsche T and Seifert W (1990): Successful opacification of the left heart chambers on echocardiographic examination after intravenous injection of a new saccharide based contrast agent. Echocardiography, 7 : 61.

Schlief R and Deichert U (1991): Hysterosalpingo-contrast sonography of the uterus and fallopian tubes : Results of a clinical trial of a new contrast medium in 120 patients. Radiology, 178 :213.

Schlief R, Schürman R and Niendorf H.P (1993): Basic properties and results of clinical trials of ultrasound contrast agents based on galactose. Ann Acad Med Singapore, 22 : 762.

Schürmann R.J, Niendorf H.P and Schließ R (1991):

Efficacy and tolerance of hysterosalpingo-contrast sonography with a saccharide-based contrast agent in 425 patients. Radiology, 181 (suppl I) : 89.

Seigler A.M (1983): Hysterosalpingography, Fertil Steril, 40:2.

Siebert J, Fobbe F and Fritzsche T.H (1989):

Darstellung der arteriellen Durchblutung nach i.v. Gabe eines lungenkapillargängigen Ultraschallkontrastmittels in der farbkodierten Duplexsonographie. Ultraschall Klin. und Praxis suppl. 1:133.

Slezak P, and Tillinger K.G (1968): The occurrence and significance of a double outlined uterine cavity in the hystero-graphic picture. Radiology, 90: 756.

Slezak P and Tillinger K.G (1973): Broad longitudinal folds in the uterine cavity in hystero-graphy. Radiology, 106: 87.

Slezak P and Tillinger K.G (1973): The significance of the speculated outline of the uterine cavity in hystero-graphy. Radiology, 107 : 527.

Slezak P and Tillinger K.G (1976): The incidence and clinical importance of hystero-graphic evidence of cavities in the uterine wall. *Radiology*, 118:581.

Slezak P and Tillinger K.G (1978): Hystero-graphic evidence of polypoid filling defects in the uterine cavity. *Radiology*, 115 : 79.

Smith M.D, Kwan O.I, Reiser H.J and Demaria A.N (1984): Superior intensity and reproducibility of SHU 454, a new right heart contrast agent. *J Am Coll Cardiol*, 3 : 992.

Smith M.D, Elion J.L and McLure R.R (1989): Left heart opacification with peripheral venous injection of a new saccharide echo contrast agent in dogs. *J Am Coll Cardiol*, 13 : 1622.

Soules M.R and Spandoni L.R (1982): Oil versus aqueous media for hysterosalpingography : a continuing debate based on many opinions and few facts. *Fertil Steril*, 38:1.

Stewart A, Web J and Hewitt D (1968) : A survey of childhood malignancy. Br. Med J, 1:1495.

Tyrrell P.N.M, Mchugo J.M, and Hale M (1993):
Patients perception of the hysterosalpingogram : the initial stages of the audit cycle. Br J Radiol, 66:103.

Varma T.R (1991): Female infertility in clinical gynaecology : Edit Thankam R, Varma. Edward Arnold, 645.

Venezia R and Zangara C (1991): Echohysterosalpingography : new diagnostic possibilities with SHU 454. Acta Eur Fertil, 22 (5) : 279.

Vorwerk D, Gehl H.B, Schlieff R, Nelles A and Gunther R.W (1990): Dynamische kontrastmittelgestützte ultraschall cavo-graphie bei kavafilterpatienten. Ultraschall in Med 11 : 146.

Watson A, Vail A, Vandekerckhove P, Brosens I, Lilford R and Hughes E (1994): A meta-analysis of the therapeutic role of oil soluble contrast media a hysterosalpingography : a surprising result ? Fertil Steril, 61:3.

Wentz A.C (1988): Infertility : Novak's Text book of Gynaecology. Edit Jones H.W, Wentz A.C and Burnette, L.S : 11th ed, Williams and Wilkins, 10: 291.

Whitehouse G.H and Wright C.H (1992): The female reproductive system : Diagnostic Radiology, an Anglo-American text book of imaging. Edit Ronald G, David J, Alexander R and Robert E : 2nd ed, Volume three, Churchill Livingstone, London, New York. 87 :1826.

Yoder I.C (1991): Hysterosalpingography in the 1990s. Am J Roentgenol 157 (4) : 675.