

REFERENCES

REFERENCES

1. **Allen, D.W., Schroeder, W.A. and Balong, J. (1958):**
Observations on the chromatographic heterogeneity of normal adult and fetal human hemoglobin.
J. Am. Chem. Soc, 80:1628.
2. **Allen, I. Arief (1982):**
Diabetes Mellitus.
Kidney Int.; 7:342.
3. **Allgrove, J and Cockrill, B.L. (1988):**
Fructosamine or glycated hemoglobin as a measure of diabetic control?
Arch. Dis. Child., 63(4): 418-22.
4. **Andersen, D.W., Nicholas, A.V., Pan, S.S. and Lindgren, F.T. (1978):**
High density lipoprotein distribution. Resolution and determination of three major components in a normal population sample.
Atherosclerosis, 29 (2): 161-79.
5. **Averna, M.R, Carroccia, A., Barbagallo, C.M., Montalto, G., Sores, M. and Notarbartolo, A. (1988):**
Diagnostic use of fructosamine assay in the control of type II diabetes mellitus.
Acta Diabetol lat, 25 (1); 63-8.
6. **Baker, J.R., Oconnor, J.P., Metcalf, P.A., Lowson, M.R., and Johnson, R.N. (1983):**
Clinical usefulness of estimation of serum fructosamine concentration as a screening test for diabetes mellitus.
Br. Med. J., 287:863-867.

7. **Baker, J.R., Johnson, R.N. and Scott, D.J. (1984):**
Serum fructosamine concentrations in type 11 non insulin dependent diabetes mellitus.
Br. Med. J., 288 : 1484-1486.
8. **Baker, J.R., Metcalf, P.A., Holdaway, I.M. and Johnson, R.N. (1985):**
Serum fructosamine concentration as a measure of blood glucose control in type I (insulin dependent) diabetes mellitus.
Br. Med. J., 290 : 352-355.
9. **Berglund, J., Lins, L.E. and Lins, P.E. (1985):**
Predictability of diabetic nephropathy.
Act. Med. Scand., 215:263-70.
10. **Beyer-Mears, A. (1986):**
The polyol pathway, sorbinil and renal dysfunction.
Metabolism 35 (suppl. 1): 46-54.
11. **Bilbrey, G.L., Faloona, G.R., White, M.G. and Knochel, J.P. (1974):**
Hyperglucagonemia of renal failure.
J. Clin. Invest., 53:841-847.
12. **Bilbrey, G.L.; Faloona, G.R., White, M.C., Atkins, C.; Hull, A.R. and Knochel J.P. (1975):**
Hyperglucagonemia in uremia. Reversal by renal transplantation.
Ann. Int. Med, 82:825.
13. **Bisse, E., Berger, W. and Fluckiger, R. (1982):**
Elimination of labile glycohemoglobin during sample hemolysis at pH 5.
Diabetes 31:630-3.

14. Bloodworth, J.M.B., Jr. (1963):

Diabetic microangiopathy.

Diabetes 12:99.

15. Bloom, A. and Ireland, J. (1980):

The kidney and lower urinary tract.

In a colour Atlas of Diabetes. ed. by Carruthers, G.B. Wolfe Medical Atlases, London; 10:100-103.

16. Boden, G., Master, R.W., Gordon, S.A., Shuman, C.R. and Owen, O.E. (1980):

Monitoring metabolic control in diabetic outpatients with glycosylated hemoglobin.

Ann. Intern. Med., 92:357-360.

17. Bookchin, R.M. and Gallop, B.M. (1968):

Structure of HbA₁: Nature of the N-terminal B-chain blocking group.

Biochem. Biophys. Res. Commun., 32:86-93.

18. Boucher, B.J., Burrin, J.M. and Gould B.J. (1983):

A collaborative study of the measurement of glycosylated hemoglobin by several methods in seven laboratories in the United Kingdom.

Diabetologia 24:265-71.

19. Briggs, J.D., Buchanan, K.D., Luke, R.G. and Mchiddie, M.T. (1967):

Role of insulin in glucose intolerance in uremia.

Lancet 1:462

20. Bruns, D.E., Lobo, P.L., Savory, J. and Wills, M.R. (1984):

Specific affinity chromatographic measurement of glycated hemoglobins in uremic patients.

Clin. Chem. 30(4): 569-571.

21. Bryan, D. and Myers (1985):

Diabetes and the kidney.

Renal Diseases 616-618.

22. **Bunn, H.F., Briehl, R.W. and Kamin, S. (1970):**
The interaction of 2,3 diphosphoglycerate with various human hemoglobins.
J. Clin. Invest., 49:1088-1095.
23. **Bunn, H.F., Haney, D.N., Gabbay, K.H. and Gallop, P.M. (1975):**
Further identification of the nature and linkage of the carbohydrate in hemoglobin A_{1c}
Biochem. Biophys. Res. Commun., 67:103-109.
24. **Bunn, H.F., Haney, D.N. and Kamin, S. (1976):**
The biosynthesis of human hemoglobin A_{1c}. Slow glycosylation of hemoglobin in vivo.
J. Clin. Invest., 57:1652-59.
25. **Bunn, H.F., Gabbay, K.H. and Gallop, P.M. (1978):**
The glycosylation of hemoglobin relevance to diabetes mellitus.
Science 200:21.
26. **Bunn, H.F., Shapiro, R. and Mc Manus, M. (1979):**
Structural heterogeneity of human hemoglobin A due to non enzymatic glycosylation.
Biol. Chem., 254:3892-3898.
27. **Cartwright, G.E., Huguley, C.M., Jr. Ashenbruken, H., Fay, J., Wintrobe, M.M. (1948):**
"Studies on free erythrocyte protoporphyrin, plasma iron and plasma copper in normal and anaemic subjects" Quoted from wintrobe, M.M., Lee, G.R., Boggs, D.R., Bithell, T.C., Hthens, J.W. and Foerester, J. (1974). Clinical Hematology, 7th. ed. pp. 186.
28. **Casparie, A.J. and Meidema, K. (1977):**
Glycosylated hemoglobin in diabetes and renal failure.
Lancet 2:8041-8058.

29. **Cefalu, W.T., Prather, K.L., Murphy, W.A. and Parker, T.B. (1989):**
Clinical evaluation of serum fructosamine in monitoring elderly outpatient diabetics.
J. Am. Geriatr Soc., 37(9): 833-7.
30. **Cerletty, J.M. and Engbring, N.H. (1967):**
Azotemia and glucose intolerance in uremia.
Ann. Intern. Med., 66:1097.
31. **Chanutin, A. and Curnish, R.R. (1957):**
Effect of organic acid, inorganic phosphates on the oxygen equilibrium of human erythrocyte.
Arch. Biochem. Biophys.
32. **Churg, J. and Dogler, H. (1979):**
Diabetic renal disease. In: Earley, L.E, Gottscnolk, C.W., eds. Strouss and Welt's. Disease of the kidney, 3rd ed. Boston: Little, Brown and company, 2:1199-1215.
33. **Clegg, M.D. and Schroeder, W.A. (1959):**
A chromatographic study of the minor components of normal adult human hemoglobin including a comparison of hemoglobin from normal and phenylketonuric individuals.
J. Amer. Chem. Soc., 81: 6065- 6069.
34. **Cohen, I.M. and Lee, S. (1981):**
Diabetes, intraperitoneal insulin, and (APD).
Clin. Exp. Dialysis Apheresis 5, 269-275.
35. **Cohen, M.P. (1986):**
Aldose reductase, glomerular metabolism and diabetic nephropathy.
Metabolism 35 (Suppl., 1): 55-59.

36. **Cole, R.A., Soeldner, J.S., Dunn, P.H. and Bunn, H.F. (1978):**
A rapid method for the determination of glycosylated hemoglobins using high pressure liquid chromatography.
Metabolism 27:289-301.
37. **Compagnucci, P., Cartechini, M.G., Bolli, G., De Feo, P., Santeusano, F. and Brunetti, P (1981):**
The importance of determining irreversibly glycosylated hemoglobin in diabetes.
Diabetes 30:607-17.
38. **Coronel, A., Cidoncha, M., Macia, L and Hidalgo. (1990):**
Role of fructosamine in the glycemic status of diabetic patients on dialysis.
EDTA procedure, p. 77.
39. **Dachs, S., Churg, J., Mounier, W. and Grishman, E. (1964):**
Diabetic nephropathy.
Am. J. Patholol., 44:155.
40. **Dandona, P., Freedman, D. and Moorhead, J.F. (1979):**
Glycosylated hemoglobins in chronic renal failure.
Brit. Med. J., 1:1183-1184.
41. **David - Bar or Stern, Z. and Naparstek, Y. (1981):**
Glycosylated hemoglobin and late complications of diabetes mellitus.
N. Eng. J. Med., 304(26): 1607-1608.
42. **Davidson, M.B., Lowrie, E.G. and Hampers, G.L. (1969):**
Lack of dialysable insulin antagonist in uremia.
Metabolism 18:387.
43. **Davis, J.E., Mc Donald, J.M. and Jarrett, L. (1978):**
A high performance liquid chromatography method for hemoglobin A_{1c}.
Diabetes 27:102-7.

44. **Daubresse, J.C., Laurent, E., Ligny, C., Bailly, A., Lemy, C., Duchateau, A., and Meunier J.C. (1987):**
The usefulness of fructosamine determination in diabetic patients and its relation to metabolic control.
Diabet. Metab., 13(3): 217-21.
45. **Day, J.F., Thornburn, R.W., Thorpe, S.R. and Baynes, J.W. (1979):**
Non enzymatic glycosylation of rat albumin.
J. Biol. Chem. 254:934-940.
46. **de Boer, M.J., Meidema, K. and Casparie, A.F. (1980):**
Glycosylated hemoglobin in renal failure.
Diabetologia 18:437-440.
47. **Deckert, I. and Poulsen, J.E. (1968):**
Prognosis for juvenile diabetics with late diabetic manifestations.
Acta Med. Scand., 183:351-356.
48. **D'Elia, J.A., Kaldany, A., Miller, D.G., Abourizk, N.N. and Weinrauch, L.A. (1985):**
Diabetic nephropathy: In "Joslin's Diabetes Mellitus", 12th edition. A. Marble, L.P. Krall, R.F., Bradley, A.R., Christlieb, and J.S. Sorum (eds). Lea & Febiger Philadelphia, chapter 30, p. 635-644.
49. **De Gruchy, G.C. (1977):**
In clinical hematology in medical practices. Blackwell Scientific Publications. Oxford London. Normal Hemoglobin, 279-281.
50. **De Marchi, S., Cecchin E., Basile, A., et al. (1983):**
More on the increase of HbA₁ in C.R.F. The role of acidosis.
Nephron 35; 49-53.

51. **Ditzel, J., Anderson, H. and Peters, N.D. (1973):**
Increased Hb A_{1c} and 2,3 DPG in diabetes and their effects on red cell oxygen releasing capacity.
Lancet 11:1034.
52. **Ditzel, J. (1975):**
Eye and systemic disease, ed. F.A. Mausolf, ST. Lovis, Mosby, p. 179.
53. **Ditzel, J. and Jens-Jorgen kjaegaard (1978):**
Hemoglobin A_{1c} concentration after initial insulin treatment for newly discovered diabetes.
Br. Med. J., p. 741.
54. **Dix, D., Cohen, P., Kingsley, S., Senkbeil, J. and Sexton, K. (1979):**
Glycohemoglobin and glucose tolerance tests compared as indicators of borderline diabetes.
Clin. Chem. 25:877-79.
55. **Dods, R.F. and Bolmey, C. (1979):**
Glycosylated hemoglobin assay and oral glucose tolerance test compared for detection of diabetes mellitus.
Clin. Chem. 25:764-68.
56. **Dolhofer, R. and Wieland, O.H. (1979):**
Glycosylation of serum albumin: elevated glycosyl - albumin in diabetic patients.
FEBS (Lett) 103:282-6.
57. **Dolhofer, R. and Wieland, O.H.(1980):**
Increased glycosylation of serum albumin in diabetes mellitus.
Diabetes 29:417-422.

58. **Dominiczak, M.H., Orrell, J.M. and Finlay, W.E. (1989):**
The effect of hypoalbuminaemia, hyperbilirubinaemia and renal failure on serum fructosamine concentration in non diabetic individuals.
Clin. Chem Acta 182 (2):123-9.
59. **Drabkin, D.L. and Austin, J.H. (1932):**
Spectrophotometric studies: spectrometric constants for common hemoglobin derivatives in human, dog and rabbit blood.
J. Biol. Chem., 98:719.
60. **Drury, M.I. (1986):**
Diabetes Mellitus. Osney Mead (ed.)
Blackweel Scientific pub., London.
61. **El-Badry, A. and Soliman, M. (1985):**
Diabetic nephropathy, clinicopathology study of
Med. J. Cairo University 53(1):23034.
62. **Eschabach, J.W., Korn, D. and Finch, C.A. (1977):**
¹⁴C cyanate as a tag for red cell survival in normal and uremic man.
J. Lab. Clin. Med. 89:823.
63. **Fayed, I., Zaki, A.M., Hafiez, A.A., Tawadrous, G.A., Abou Zekry, M., Karas, M.S., Abdel Monem, A.M. and El-Miligy, D. (1986):**
Glycosylated hemoglobin in chronic liver disease and acute hemolytic anaemia.
The Gaz. of Egypt. Soc., Paed. Assoc. In press.
64. **Flock, E.V., Bennett, P.H., Savage, P.J., Webner, C.J., Howard, B.V., Ruch forth, N.B. and Miller, M. (1979):**
Bimodality of glycosylated hemoglobin distribution in pima indians: relationship to fasting hyperglycemia.
Diabetes care 7:147-50.

65. **Fluckiger, R., Harmon, W., Meiler, W. and Loo, S. (1981):**
Hemoglobin carbamylation in uremia.
New. Engl. J. Med., 304(14): 823.
66. **Fluckiger, R., Woodtli, T. and Berger, W. (1987):**
Evaluation of the fructosamine test for the measurement of plasma protein glycation.
Diabetologia 30:648-652.
67. **Freedman, E.A. (1982):**
Renal consequences of diabetes.
In: Diabetes Mellitus and obesity. B.N. Brodoff and S.J. Bleicher eds.
Williams and Wilkins, Baltimore, London, 72:667-69.
68. **Freedman, D., Dandona, P., Fernando, O. and Moorhead, J.F. (1982):**
Glycosylated hemoglobin in chronic renal failure and after renal transplantation.
J. Clin. Pathol., 35:737-739.
69. **Gabbay, K.H., Hasty, K., Berslow, J.L., Ellison, R.C., Bunn, H.F. and Gallop, P.M. (1977):**
Glycosylated hemoglobins and long term blood glucose control in diabetes mellitus.
J. Clin. Endocrinol Metabol., 44: 859.
70. **Gabbay, K.H., Sosenko, J.M., Banuchi, G.A. Mininsohu, M.J., and Fluckiger, R. (1979):**
Glycosylated hemoglobins. Increased glycosylation of hemoglobin A in diabetic patients.
Diabetes 28: 337-340.

71. **Gellman, D.D., Pirani, C.L., Soothill, J.P., Mu Hrcche, E.C. and Hark, R.M. (1959):**
Diabetic Nephropathy. A clinical and pathological study based on renal biopsies.
Medicine, 38: 321-367.
72. **Giovannetti, S., Cionil and Balestril (1974):**
Evidence that guanidines and some related compounds cause hemolysis in chronic renal failure.
Clin. Sci., 34:141.
73. **Glasaock, R.J. and Brenner, B.M. (1984):**
Glomerulopathies associated with multisystem diseases. In "Harrison's principles of internal medicine. Petersdort, R.G., Adams, R.D., Braumwald, E., Wilson, J.D., and Fauci, A.S. (eds). McGraw-Hill Book Company.
74. **Goldstein, D.E., Peth, S.B., England, J.D., Hess, R.L. and Da costa, J. (1980):**
Effect of acute changes in blood glucose on HbA_{1c}.
Diabetes 29:623-628.
75. **Goldstein, D.E., Parker, K.M., England, J.D., Wiedmeyer, J.R., Rowlings, S.S., Randle, R.M., Smonds, J.F. and Breyfogle, R.P. (1982):**
Clinical application of glycosylated hemoglobin measurements.
Diabetes 31 (Suppl. 3):70-8.
76. **Gonen, B., Rubenstein, A.H., Rochman, H., Tanega, S.P. and Horwitz, D.L. (1977):**
Hemoglobin A₁: An indicator of metabolic control of diabetic patients.
Lancet 11:734-736.

77. **Gorden, P., Sherman, B.M. and Simopoulos, A.P. (1972):**
 Glucose intolerance with hypokalemia and increased proportion of circulating proinsulin like component.
 J. Clin. Endocrinol. Metab., 34:235.

78. **Gottschalk, A. (1972):**
 Glycoprotein, El sevier, New York,
 part I. 141-157.

79. **Graf, R.J., Halter, J.B. and Porte, D., Jr. (1978):**
 Glycosylated hemoglobin in normal subjects and subjects with maturity onest diabetes: evidence for a saturable system in man
 Diabetes 27:834-839.

80. **Graf, H., Stummvol, H.K., Schernthaner, G. and Muller, M.M. (1980):**
 Glycosylated hemoglobin in renal failure.
 Diabetologia 19:555-6.

81. **Guthrow, C.E.; Morris, M.A.; Day, J.F.; Thorpe, S.R. and Baynes, J.W. (1979):**
 Enhanced non-enzymatic glycosylation of human serum albumin in diabetes mellitus.
 Proc. Natl. Acad. Sci (U.S.A.) 76:4258-61.

82. **Hall, G.F.M. (1952):**
 The significance of atheroma of the renal arteries in Kimmelstiel- Wilson syndrome.
 J. Path. Bact., 64:103-20.

83. **Hall, P.M., Cook, J.G.H., Sheldon, J., Rutherford, S.M. and Gould, B.J. (1984):**
 Glycosylated hemoglobins and glycosylated plasma proteins in the diagnosis of diabetes mellitus.
 Diabetes care 7:147-50.

- 84. Hampers, G.L., Soeldner, J.S., Doak, P.B. and Merrill, J.P. (1966):**
Effect of chronic renal failure and hemodialysis on carbohydrate metabolism.
J. Clin. Invest., 45:1719-1731.
- 85. Haney, D.N. and Bunn, H.F. (1976):**
Glycosylation of hemoglobin in vitro, affinity labelling of hemoglobin by glucose -6-phosphate.
Proc. Natl. Acad. Sci. (U.S.A.) 73:3534.
- 86. Hankin, W.D. and Hollady, L.A. (1980):**
A temperature conversion monogram for glycosylated hemoglobin analysis.
Clin. Chem. Acta 104:251-7.
- 87. Henry, J. (1984):**
Clinical diagnosis and management by laboratory methods. Todd, Sanford. Davidson WB, Saunders Company Philadelphia, London, Toronto, Vol I:175.
- 88. Higgins, P.J., Garlich, R.L. and Bunn, H.F. (1982):**
Glycosylated hemoglobin in human and animal red cells. Role of glucose permeability.
Diabetes 31:743.
- 89. Hill, A.B. (1977):**
In "A short Textbook of Medical statistics" 10th edition. The English language Book Society and Hodder and Stoughton London.
- 90. Hindle, E.J., Rostron, G.M. and Gatt, J.A. (1985):**
The estimation of serum fructosamine: an alternative measurement to glycated hemoglobin.
Ann. Clin. Biochem; 22:84-89.

91. **Hirszel, P. Galen, M.A., Happe, T., et al (1981):**
Glycosylated hemoglobin in patients treated by chronic dialysis.
Int. Urol Nephrol. 13: 185-191.
92. **Horton, E.S., Johnson, C. and Lebevitz, H.E. (1968):**
Carbohydrate metabolism in uremia.
Ann. Int. Med. 68:63-74.
93. **Huisman, T.H.J. and Dozy A.M. (1962):**
Studies on the heterogeneity of hemoglobin. Binding of hemoglobin with oxidized glutathione.
J. Lab. Clin. Med., 60:302-319.
94. **Ingram, V.M. and Stretton, A.O.W. (1962):**
Human hemoglobin A₂, Comparison of hemoglobin A₂ and A.
Biochem. Biophys. Acta; 62:456.
95. **Ireland, J.T., Viberti, G.C. and Watkins, P.J. (1982):**
The kidney and renal tract. In complications of diabetes 2nd ed., H. Keen and J. Jarrett. London, Edward Arnold.
96. **Jarrett, J. (1985):**
The natural history and prognosis of diabetes.
Medical Interna., 13:530-532.
97. **Javid, J., Pettis , P.K., Koenig, R.J., et al. (1978):**
Immunologic characterization and quantification of hemoglobin A_{1c}.
Br. J. Hematol., 38:329-337.
98. **Jermendy, G., Szelenyi, J., Bodonyi, A. and Foldi, J. (1988):**
Serum fructosamine (alkaline - reducing activity) for evaluation of glycemic control in diabetic patients.
Exp. Clin. Endocrinol, 92(2): 171-6.

99. John, W. Hare (1986):

Signs and symptoms in endocrine and metabolic disorders.
Medicine (Baltimore), 78:527.

100. John, W.G. and Richadson, R.W. (1986):

Glycosylated hemoglobin levels in patients referred for oral glucose tolerance tests.
Diabetic Med., 3(1):46-8.

101. Johnson, M.W., Dobrea, G.M. and Bendezu, R. (1980):

Temperature dependence of the chromatographic assay of hemoglobin A1 and application of a temperature controlled assay to clinical evaluation of diabetic control.
Clin. Chem. Acta., 104:319-28.

102. Johnson, R.N., Metcalf, P.A. and Baker, J.R. (1982):

Fructosamine a new approach to the estimation of serum glycosyl protein-
An index of diabetic control Clin.
Clin. Chem. Acta, 127:87-95

103. Johnson, R.N., Metcalf, P.A. and Baker, J.R. (1987):

Relationship between albumin and fructosamine concentration in diabetic and non diabetic sera.
Clin. Chem. Acta, 30, 164(2):151-62.

104. Jovanovic, L. and Peterson, C.M. (1981):

The clinical utility of glycosylated hemoglobin.
Am. J. Med., 70(12):331.

105. Kador, P.F. and Kinoshita, J.H. (1983):

Role of aldose reductase in the development of diabetes associated complications.
Am. J. Med., 79(Suppl. 5A): 8-12.

- 106. Kasai, K., Nakamura, T., Kase, N., Hirooka, T., Suzuki, R., Kogure, F., and Shimoda, I. (1983):**
Increased glycosylation of proteins from cataractous lenses in diabetes.
Diabetologia, 25:36-38.
- 107. Kennedy, A.L. and Merimee, T.J. (1981):**
Glycosylated serum protein and hemoglobin A levels to measure control of glycemia.
Ann. Intern. med., 95:56-58.
- 108. Kennedy, A.L., Mehl, T.D., Rieley, W.J. and Merimee, T.J. (1981):**
Non enzymatically glycosylated serum protein in diabetes mellitus an index of short term glycemia.
Diabetologia 21:94-98.
- 109. Kennedy, A.L. and Lyons, T.J. (1989):**
Non enzymatic glycosylation.
British Medical Bulletin 45(1): 174-190.
- 110. kern, T.S. and Engerman, R.L. (1987):**
Kidney morphology in-experimental hyperglycemia.
Diabetes 36:244-249.
- 111. Kerr, D.N.S. (1979):**
In "Cecil Textbook of Medicine" editors: Beeson, P.B., McDermott, W., and Wyngaarden, J.B. Saunder company 15th edition, 1351-1367.
- 112. Kim, H.S., Kalkoff, P.K., Costrini, N.V., Cerletty, J.M. and Jacobson, M. (1969):**
Effect of parathromone on plasma insulin response and pancreatic islets secretion of insulin.
J. Lab. Clin Med., 74:891.

- 113. Kimmelstiel, P. and Wilson, C. (1936):**
Complication in relation to diabetic control.
Am. J. Path., 12:83-97.
- 114. Kimmelstiel, P., Osawa, G. and Beres, J. (1966):**
Glomerular basement membrane in diabetics.
Am. J. Clin. Path, 45:21.
- 115. Klenk, D.C., Hermanson, G.T. and Krohn, R.I. (1982):**
Determination of glycosylated hemoglobin by affinity chromatography:
comparison with colorimetric and ion exchange methods, and effects of
common interferences.
Clin. Chem; 28:2088-94.
- 116. Koenig, R.J., Araujo, D.C. and Cerami, A. (1976):**
hemoglobin A_{1c} as an indicator of the degree of glucose intolerance in
diabetes.
Diabetes 25:230.
- 117. Koskinen, P., Irjala, K., Viikari, J., Panula-Ontto, R. and
Matikainen, M.T. (1987):**
Serum fructosamine in the assessment of glycemic control in diabetes
mellitus.
Scand. J. Clin. Lab. Invest., 47(3):285-92.
- 118. Kovarik, J., Stummvoll, H.K. and Graff, H. et al., (1981):**
Glucose intolerance and hemoglobin A₁ in chronic renal insufficiency.
Nephron, 28:209-212.
- 119. Lantz, B., Ochoa, C., Wajeman, H., et al. (1981):**
Minor hemoglobin components in diabetic and uremic patients.
Horm. Metab. Res., 13:660-663.

120. *Lapolla, A., Poli, T., Barison, A. and Fedele, D. (1988):*
Fructosamine assay: an index of medium term metabolic control parameters in diabetic disease.
Diabetes Res. Clin. Pract., 4(3):231-5.
121. *Lim, Y.S. and Staley, M.J. (1985):*
Measurement of plasma fructosamine evaluated for monitoring diabetes.
Clin. Chem., 31(5):731-33.
122. *Lindal, A., Carmena, R., Cohen, S. and Comty, C. (1971):*
Insulin hypersecretion in patients with chronic hemodialysis, role of parathyroid.
J. Clin. Endocrinol Metab., 32:653.
123. *Little, R.R., England, J.D., Wiedmeyer, H.M. et al. (1983):*
Glycosylated hemoglobin measured by affinity chromatography: microsample collection and room temperature storage.
Clin. Chem., 29:1080-2.
124. *Little, R.R., England, J.D., Wiedmeyer, H.M. et al. (1988):*
Relationship of glycosylated hemoglobin to oral glucose tolerance.
Diabetes, 37:60-64.
125. *Lowrie, E.G., Soeldner, J.S., Hampers, G.L. and Merrill, J.P. (1970):*
Glucose metabolism and insulin secretion in uremics, prediabetics and normal subjects.
J. Lab. Clin. Med., 76:603-515.
126. *Luke, R.G., Dinwoodie, A.J., Linton, A.L. and Kennedy, A.C. (1964):*
Fructose and glucose tolerance in uremia.
J. Lab. Clin. Med., 64:731.

127. **Mako, M., Block, M., Starr, J., Nielsen, K., Freedman, E. and Rubenstein, A. (1973):**
Proinsulin in chronic renal failure and hepatic failure, a reflection of the relative contribution of the liver and kidney to its metabolism.
Clin. Res., 21:631.
128. **Malaisse, W.J. and Methias, P.C.F. (1985):**
Stimulation of insulin release by an organic calcium agonist.
Diabetologia 28:153-156.
129. **Malins, J. (1968):**
Renal disease in diabetes.
Clinical Diabetes Mellitus. Eyre and Spottiswoode, 7:152-173.
130. **Mallia, A.K., Hermanson, G.T., Krohn, R.I., et al (1981):**
Preparation and use of a boronic acid affinity support for separation and quantitation of glycosylated hemoglobins.
Anal lett 14(B8), 649-661.
131. **Manicardi, V., Bonora, E. and portoli, I. (1981):**
Comparison of glycosylated hemoglobin and OGTT in the screening of diabetes (Abstract).
Diabetologia 21:301.
132. **McCance, D.R., Coulter, D., Smye, M., and Kennedy, L. (1987):**
Effect of fluctuations in albumin on serum fructosamine.
Diabetic Med., 4:434-36.
133. **McCance, D.R., Ritche, C.M. and Kennedy, L. (1988):**
Is hemoglobin A₁ superfluous in non insulin dependent diabetes mellitus?
Diabetes care, 11:512-514.
134. **McDonald., M.K. and Ireland J.T. (1964):**
The glomerular lesion in idiopathic and secondary diabetes.
Ciba Foundation colloquia, 15:301-14.

135. *McFarland, K.F., Catalona, E.W., Day, J.F., Thorpe, S.F. and Baynes, J.W. (1979):*
Non enzymatic glycosylation of serum proteins in diabetes mellitus.
Diabetes 28:1011-1014.
136. *Modan, M., Halkin, H., Karasik, A. and Lusky, A. (1984):*
Effectiveness of glycosylated hemoglobin, fasting plasma glucose, and a single post load plasma glucose level in population screening for glucose intolerance.
Am. J. Epidemiol 119:431-44.
137. *Mortensen, H.B. (1985):*
Glycated hemoglobin. Reaction and Biokinetic studies. Clinical application of hemoglobin A_{1c} in the assessment of metabolic control in children with diabetes mellitus.
Dan. Med. Bull., 32(6): 309-28.
138. *Nathan, D.M. (1981):*
Labile glycosylated hemoglobin contributes to hemoglobin A₁ as measured by liquid chromatography or electrophoresis.
Clin. Chem., 27:1261-63.
139. *Nathan, D.M., Singer, D.E., Hurxthal, K. and Goodson, J.D. (1984):*
The clinical information value of the glycosylated hemoglobin assay.
N. Engl. J. Med., 310:341-346.
140. *Navalesi, R., Pilo, A., Lenzis, S. and Donato, L. (1975):*
Insulin metabolism in chronic uremia and in the anephric state, effect of dialytic treatment.
J. Clin. Endocrin. Metab., 40:70-85.
141. *Negoro, H., Morley, J.E. and Rosenthal, M.J. (1988):*
Utility of serum fructosamine as a measure of glycemia in young and old diabetic and non diabetic subjects.
Am. J. Med., 85(3):360-4.

142. *Neuiman, D.R., Pierre, R.V. and Limman, J.W. (1973):*
Studies on the diagnostic significance of hemoglobin F levels.
Mayo Clin. Proc., 48:199.
143. *Oimomi, M., Ishikama, K., Kawasaki, T., Kubota, S.,
Yoshimwa, Y. and Baba, S. (1981):*
Glycosylated hemoglobin in renal failure.
Diabetologia 21:163.
144. *Okada, Y., Arima, T., Okazaki, S. and Nakated, R. (1982):*
Increased I^{125} labelled concentration binding to erythrocytes in diabetes
mellitus.
Diabetologia 22:212.
145. *Ottoman, R.E., Woodruff, J.H., Jr., Wilk, S. and Isooc, F.
(1956):*
The roentgen aspects of necrotizing renal papillitis.
Radiology 67:157.
146. *Paisy, R.B., McFarland, D.G., Sherrif, R.J., Hartog, M., Stade,
R.R. and White D.A.J. (1980):*
The relationship between glycosylated hemoglobin and home capillary
blood glucose levels in diabetics.
Diabetologia 19:31
147. *Parker, K.M., England, J.D., Dacosta, J., Hess, R.L. and
Goldstein, D.E. (1981):*
Improved colorimetric assay for glycosylated hemoglobin.
Clin. Chem., 27:669-72.
148. *Paulsen, E.P. and M. Koury (1976):*
Hemoglobin A_{1c} levels in insulin dependent and independent diabetes
mellitus.
Diabetes 25; 890-96.

149. *Pecoraro, R.E., Graf, R.J., Halter, J.B. and Porto, D. (1979):*
Comparison of a colorimetric assay for glycosylated hemoglobin with ion exchange chromatography.
Diabetes 28:1120-1125.
150. *Perez, G.T., Rabinovitch, A., Rietberg B., Owen, B. and Schiff, E.R. (1975):*
Impaired glucagon stimulated glucose output in livers of acutely uremic rats.
J. Lab. Clin. Med., 99:669.
151. *Perutz, M.F., Rassman, M.G., Cullis, A.F., Murihead, H., Will, G. and North, A.C.T. (1963):*
Structure of hemoglobin.
Nature, 199:633.
152. *Perutz, M.F. (1964):*
"The hemoglobin molecule"
Sci. Am., 211.
153. *Peters, T. (1968):*
Proposal for standardization of total protein assays.
Clin. Chem., 14:1147.
154. *Peters, J., and Watkins (1982):*
Nephropathy, ABC of Diabetes.
B.M.J., 285:627.
155. *Peterson, C.M., Jovanovic, L., Raskin, P. and Goldstein, D.E. (1984):*
A comparative evaluation of glycosylated hemoglobin assays: Feasibility of references and standards.
Diabetologia 26:214-7.

- 156. Pirat,, J. (1965):**
Diabetic Nephropathy. A metabolic or vascular disease
Diabetes 14:1-9.
- 157. Putten, J.P.M., Weringe, T.I. and Krans, H.M.J. (1985):**
Low pH and Ketoacids induce insulin receptor binding and postbinding
alternation in cultured 3T3 adipocytes.
Diabetologia 34:744.
- 158. Rahbar, S.S. (1968):**
An abnormal hemoglobin in red cell of diabetics.
Clin. Chem Acta, 22:296-298.
- 159. Rahbar, S.S., Blumenfeld, O. and Ranney, H.M. (1969):**
Studies of unusual hemoglobin in patients with diabetes mellitus.
Biochem. Biophys. Res Commun., 36:838.
- 160. Ramzy, M.F., Habib, M.R., Tawadrous, G.A., Hafiez, A.A,
Samney, S.G., Sadek, H. and El-Sheemy, N. (1984):**
Glycosylated hemoglobin in chronic renal failure.
Bulletin of Egypt. Soc. Physiol. Sci. 4(2):90-103.
- 161. Relman, A.S. (1972):**
Metabolic consequence of acid base-disorders.
Kidney Int., 1:347.
- 162. Rezk, H.S. (1981):**
Glycosylated hemoglobin in chronic renal failure.
M. Sc. Thesis, Faculty of Medicine, Cairo University.
- 163. Riggs, A. (1965):**
Functional properties of hemoglobins.
Physiol. Rev. 45:619.

- 164. Roberts, A.B., Court, D.J., Henley, P., Baker, J.R., James, A.G. and Ronayne, I.D. (1983):**
Fructosamine in diabetic pregnancy.
Lancet, 22 ii:998-999.
- 165. Roger, G. (1979):**
Post graduate nephrology.
Second edition, Butterworth London, Pages 27, 35, 68,119.
- 166. Rosenthal, M.A. (1979):**
The effect of temperature on the fast hemoglobin test system.
Hemoglobin 3:215-7.
- 167. Rosenthal, P.K., Vazquer, D.A. and Seckinger, D.L. (1981):**
Determination of glycohemoglobins by a rapid batch separation
Am. J. Clin. Pathol., 75:45-49.
- 168. Sabour, M.S., Mc Donald, M.K., and Robson, J.S. (1962):**
An electron microscopic study of the human kidney in young diabetic patients with normal renal function.
Diabetes, 11:291.
- 169. Saddi, R., Wajcman, H., Eschwege, E., levy, R. and Duchuteau, A. (1980):**
Glycosylated hemoglobin in patients on venesection therapy for hemochromatosis.
Lancet 11:141.
- 170. Saglid, U., Andersen, V. and Anderson, B.P. (1961):**
Glucose tolerance and insulin responsiveness in experimental potassium depletion.
Acta. Med. Scand., 169:243.

- 171. Salemans, T.H., Van Dieijen - visser, M.P. and Brombacher, P.J. (1987):**

The value of HbA₁ and fructosamine in predicting impaired glucose tolerance-an alternative to OGTT to detect diabetes mellitus or gestational diabetes.

Ann. Clin. Biochem., 24(Pt 5) (7):447-52.

- 172. Samaan, N.A. and Freedman, R.M. (1970):**

Growth hormone level in severe renal failure.

Metabolism 19:102.

- 173. Sandor, G. (1966):**

Serum proteins in health and disease.

Chapman and Hall, London.

- 174. Santiago, J.V., Davis, J.E. and Fisher, F. (1978):**

Hemoglobin A₁ levels in a diabetes detection program.

J. Clin. Endocrinol Metab., 47:578-80.

- 175. Schellekens, A.P.M., Sanders, G.T.B. and Thornton, W. (1981):**

Sources of variation in the column-chromatographic determination of glycohemoglobin.

Clin. Chem., 27:94-9.

- 176. Schernthaner, G., Stummvoll,, K.H. and Muller, M.M. (1979):**

Glycosylated hemoglobin in chronic renal failure.

Lancet 1:774.

- 177. Schleicher, E. and Wieland, O.H. (1981):**

Specific quantitation by HPLC of protein (Lysine) bound glucose human serum albumin and other glycosylated proteins.

J. Clin. Chem. Clin. Biochem., 19:81-7.

178. **Schwartz, H.C., King, K.C. and Schwartz, R. (1976):**
Effect of pregnancy on HbA_{1c} in normal, gestational diabetic and diabetic women.
Diabetes 25:1118.
179. **Shapiro, F.L., Leonard, A. and Comty, C.M. (1974):**
Mortality, morbidity and rehabilitation results in regularly dialysed patients with diabetes mellitus.
Kidney Int., 6 (Suppl 1): 5-8.
180. **Sharma, S.S., Solanki, R.L., Arora, H.L., Kumar, K. and Anand, V.K. (1989):**
Glycosylated hemoglobin (HbA_{1c}) and serum protein bound hexose in diabetes mellitus.
J. Assoc. Physicians India, 37(10): 639-41.
181. **Shaw, A.B. (1967):**
Hemolysis in chronic renal failure.
Br. Med. J., 2:213.
182. **Shimojo, N., Hirai, M., Naka, K. et al., (1988):**
Plasma fructosamine assay in children with insulin dependent diabetes mellitus.
Clin. Chem. Acta 176(1):101-6.
183. **Simon, D., Coignet, M.C., Thibault, N., Senan, G. and Eschwege, E. (1985):**
Comparison of glycosylated hemoglobin and fasting plasma glucose with two-hour post-load plasma glucose in the detection of diabetes mellitus.
Am. J. Epidemiol 122(4): 589-93.
184. **Spergel, G., Bleicher, J.J., Goldberg, M., Adesman, J. and Goldner, M.G. (1967):**
Effect of potassium on the impaired glucose tolerance in chronic uremia.
Metabolism 16:581.

- 185. Sridama, Thia, A.C. and Lui, K.F. (1990):**
Serum fructosamine determination as an index of glycemic control.
Ann. Acad. Med. Singapore, 18(4): 352-6.
- 186. Stanton, K.G., Davis, R. and Richmond, J. (1978):**
Glycosylated hemoglobin in renal failure.
Lancet 1:100.
- 187. Stark, A., Keck, E, Berger, M. and Zimmermann, H. (1981):**
Effect of calcium and calcitonin on circulating levels of glucagon and glucose in diabetes mellitus.
Diabetologia 20:547-552.
- 188. Svendsen, P.A., Christiansen, J.S., Welinder, B. and Nerup, J. (1979):**
Fast glycosylation of hemoglobin.
Lancet 1:603.
- 189. Svendsen, P.A., Christiansen, J.S., Soegard, U., Welinder, B.S. and Nerup, J. (1980):**
Rapid changes in chromatographically determined hemoglobin A₁ induced by short term changes in glucose concentration.
Diabetologia 19:130-136.
- 190. Tegos, R. and Beutler, E. (1980):**
Glycosylated hemoglobin A₂ components.
Blood 56:571.
- 191. Tesio, F., Cecchin, E., De Marchi, S. et al. (1982):**
Glycosylated or carbamylated hemoglobin in uremia.
N. Eng. J. Med., 306:1053.
- 192. Thomsen, A.C. (1965):**
The kidney in diabetes mellitus.
Copenhagen, E. Munksgaard, P. 124.

193. Tietz (1987):

Fundamentals of clinical chemistry. 3rd Edition

194. Trinder, P. (1969):

Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor.

Ann. Clin. Biochem., 6:24-27.

195. Trivelli, L.A., Ranney, H.M. and Lai, H. (1971):

Hemoglobin components in patients with diabetes mellitus.

N. Eng. J. Med., 284:355-357.

196. Varley (1988):

Practical clinical biochemistry. Sixth Edition.

197. Verillo, A., de Teresa, A., Golia, R. and Nunziata, V. (1983):

The relationship between glycosylated hemoglobin levels and various degrees of glucose intolerance.

Diabetologia 24:391-93.

198. Vermes, I., Zeyen, L.J., Van Roon, E. and Brandts, H. (1989):

The role of serum fructosamine as a screening test for gestational diabetes mellitus.

Horm. Metab. Res., 21(2): 73-6.

199. Viberti, G.C., Hill, R.D., Janett, R.J., et al. (1982):

Microalbuminuria as a predictor of clinical nephropathy in insulin dependent diabetes mellitus.

Lancet 1: 1430-1432.

200. Vincent, T. and Andriole (1985):

Urinary tract infections and pyelonephritis.

Renal Diseases, 619-630.

201. **Warren, S., Le Comble, P.M., Legg, M.A., eds. (1966):**
The pathology of diabetes mellitus 4th ed.
Philadelphia: Lea and Febiger, 227.
202. **Watkins, P.J. (1985):**
Diagnosis and treatment of diabetic nephropathy.
Med. Intern. (1"):554-55.
203. **Westervelt, F.B., Jr. and Schreiner, G. (1962):**
The carbohydrate intolerance of uremic patients.
Ann. Intern. Med., 57:266.
204. **White, J.M. (1972):**
Hematology, William, Heineman Medical Book "The structure of normal hemoglobin" LTD, London.
205. **Widness, J.A., Rogler, Brown, T.L. and McCormik, K.L. (1980):**
Rapid fluctuations in glycohemoglobin (A_{1c}) related to acute changes in glucose.
J. Lab. Clin. Med., 95:386-94.
206. **Wintrobe, M.M., Lee, C.R., Roggs, D.R., Bithell, T.C., Athens, J.W. and Lukens, J.N. (1981):**
Clinical hematology, eighth edition p. 90-96.
207. **Zager, R.A. and Johannes, G. (1982):**
Hyperphosphatemia, a factor that provokes severe experimental acute renal failure.
J. Lab. Clin. Med., 100:23.
208. **Zawada, E.T. and Virgi, A.Jr. (1980):**
Glycosylated hemoglobin test in diabetic and non diabetic hemodialysis patients.
Dialysis Transplant, 9:360-362.