

# REFERENCES

## REFERENCES

- Abbott, W.G.H. ; Lillioja, S. and Young, A.A. (1987) :* Relationships between plasma lipoprotein concentrations and insulin action in an obese hyper insulinemic population. *Diabetes*. 36: 897 - 904.
- Abdul-Karim, R. and Assal, N.S. (1961) :* Pressor response to angiotensin in pregnant and non pregnant women. *Am. J. Obstet. Gynecol.* 82 : 246.
- Anderson, E.A. and Mark, A.L. (1993) :* The vasodilator action of insulin: implications for the insulin hypothesis of hypertension. *Hypertension* 21: 136.
- Arduini, D. ; Rizzo G. ; Romanni C. ; Mancuso S. (1987) :* Uteroplacental blood flow velocity wave forms as predictors of pregnancy-induced hypertension. *Eur.J. Obstet. Gynecol. Reprod Biol.* 26: 335 - 41.
- Austin, M.A. ; Breslow, J.L. and Hennekens, C.H. (1988):* Low-density lipoprotein subclass patterns and risk myocardial infraction. *JAMA* 260: 1917.
- Austin, M.A. ; King, M.C. ; Vranizan, K.M. and Krauss R.M. ; (1990) :* Atherogenic lipoprotein phenotype : A proposed genetic marker for coronary heart disease risk circulation. 82: 495.
- Alexrod, M.D. (1991):* perspectives in diabetes insulin, prostaglandins and the pathogenesis of hypertension. *Diabetes*, vol. 40: 1223-1227.
- Axelrod, L. and Levine, L. (1981):* Prostacyclin production by isolated adipocytes. *Diabetes*. 30: 163-67.

- Ballegree, V. ; Spitz, B. ; Kieckens, L. ; Moreou, H. ; Assche A.van and Collen D. (1989) :** Predictive value of increased plasma levels of fibronectin in gestational hypertension. *Am. J. Obstet. Gynecol.* 161: 432 - 6.
- Barter, P.J. ; Hopkins, G.J. and Gorjatschkol, A. (1985) :** Lipoprotein substrates for plasma cholesterol esterification. Influence of particle size and composition of high density lipoprotein subfraction 3. *Atherosclerosis*. 58: 97.
- Bauman, M.R. ; Maiman, M. and Langer, O. (1982) :** An association between hyperinsulinemia and hypertension during the third trimester of pregnancy. *Am. J. Obstet. Gynecol.*; 159: 446 - 50.
- Beer, A.E. (1978) :** Possible immunologic bases of preeclampsia / eclampsia. *Semin perinatal*. 2: 39.
- Belizan, J.M. ; Villar, J. ; Gonzalez, I. ; Campodonico, I. and Bergel, E. (1991) :** Calcium supplementation to prevent hypertensive disorders of pregnancy. *N. Engl. J. Med.* 325: 1399.
- Belizan, J.M. ; Villar, J. and Repke, J. (1988) :** The relationship between calcium intake and pregnancy-induced hypertension : up-to-date evidence. *Am. J. Obstet. Gynecol.* 158: 898.
- Belizan, J.M. and Villar, J. (1980) :** The relationship between calcium intake and edema, proteinuria and hypertension gestosis : an hypothesis. *Am. J. of clinical Nutrition* 33: 2202 - 10.
- Bergstroms ; Carison, L.A. and Weeks, J.R. (1968) :** The prostaglandins : family of biologically active lipids. *Pharmacol. Rev.* 20: 1 - 48.

- Berkounitz, G.S. ; Shovron, M.L. ; Lapinski, P.H. and Berkowitz, R.L. (1990) :** Delayed child bearing and the outcome of pregnancy. New. Eng. J. Med. 322: 659 - 664.
- Bernstein, L. ; Pike, M.C. ; Lobo, R.A. ; Dupue, R.H. ; Ross, R.K. and Henderson, B.E. (1989) :** Cigarette smoking in pregnancy results in marked decrease in maternal HCG and oestradiol levels. Br. J. Obstet. Gynecol. 96: 92-6.
- Biezanski, J.J. (1974) :** Maternal lipid metabolism. Obstet Gynecol Ann. 33: 203 - 33.
- Bilheimer, D.W. ; Goldstein, J.L. ; Grandy, S.M. ; Starzl, T.E. and Brown, M.S. (1984) :** Liver transplantation to provide low density lipoprotein receptors and lower plasma cholesterol in a child with a homozygous familial hypercholesterolemia. Familial hypercholesterolemia. New. England J. Med. 311: 1658.
- Borok, Z.; Weiz, J. ; Owen, J. ; Auerbach, M. and Nossel, H.L. (1984) :** Fibrinogen proteolysis and platelet-granule release in preeclampsia / eclampsia. Blood. 63: 525 - 31.
- Brain, M.C. ; Dacie, J.V. and Hourihane, D.O. (1962) :** Microangiopathic haemolytic anaemia : the possible role of vascular lesions in pathogenesis. Br.J. Haematol. 8: 358 - 74.
- Branch, D.W.; Dudley, D.J. and Mitchell, M.D. (1991):** Preliminary evidence for homeostatic mechanism regulating endothelin production in preeclampsia. Lancet 337: 943-5.

- Branch, D.W.; Mitchell, M.D., Milker, E.; Paliniski, W. and Witzum, J.L. (1994):** Preeclampsia and serum antibodies to oxidised LDL. Lancet. 343: 645-6.
- Brands, M.S. ; Hildebrandt, D.A. ; Mizelle, H.L. and Hall, J.E. (1991) :** Sustained hyperinsulinemia increases arterial pressure in conscious rats. Am. J. Physiol. 260: R764.
- Brewer, H.B. ; Gregg, R.E. ; J.M. and Fojo, S.S. (1988) :** Apolipoprotein and lipoproteins in human plasma An overview. Clin. Chem. 34: 84.
- Brinton, E.A. ; Eisenberg, S. and Breslow, J.L. (1994) :** Human HDL cholesterol levels are determined by Apo A-I fractional catabolic rate which correlates inversely with estimates of HDL particle size. Arterio. Thromb. 14: 707.
- Brown, C.E.L. ; Cunningham, F.G. and Pritchard, J.A. (1987) :** Convulsions in hypertensive proteinuric primiparas more than 24 hours after delivery : Eclampsia or some other cause ?? J Reproduction Med. 32: 494.
- Brown, M.A. (1989) :** Pregnancy-induced hypertension. Curr. Concepts. Anaesth. Intens. Care. 17: 183 - 97.
- Brunner, H.R. and Gavras, H. (1975) :** Vascular damage in hypertension Hosp. Pract. 10: 97.
- Brustein, M. (1970) :** HDL cholesterol determination. Lipid Rest 11:583.
- Buchahan, T.A. ; Metzger, B.E. ; Freinkel, N. and Bergman, R.N. (1990):** Insulin sensitivity and B-cell responsiveness to glucose during late

- pregnancy in lean and moderately obese diabetes. Am. J. Obstet. Gynecol. 162: 1008 - 14.
- Byaruhanga, R.N. ; Chipato, T. and Rusakaniko, S. (1988) :** A randomized controlled trial of low-dose aspirin in women at risk from pre-eclampsia. International J. of Gynecology and Obstetrics. 60: 129-135.
- Campbell, D.M. and MacGillivray, I. (1985) :** Preeclampsia in second pregnancy. Br. J. Obstet. Gynecol. 92: 131 - 40.
- Campbell, D.M. and MacGillivray, I. (1980) :** The relevance of hypertension and oedema in pregnancy. Clin. Exp. Hypertens. 2 894 - 914.
- Campbell, M.J. and Machin, D. (1993) :** Medical statistics. A common sense approach. 2nd Ed. John Wiley & Sons. New York; Chichester; Brisbane; Toronto & Singapore.
- Campbell, S. ; Griffin, D.R. ; Pearce, J.M. ; Diaz-Pecasens, J. ; Cohen-over beek, T.E. ; Willson, K. and Teague, M.J. (1983) :** New Doppler technique for assessing uteroplacental blood flow. Lancet. 1.675-7.
- Campbell, S. ; Pearce, J.M.F. ; Hackett, G. ; Cohen-over beek, T. and Hernandez, G. (1986) :** Qualitative assessment of uteroplacental blood flow: early screening test for high - risk pregnancies. Obstet. Gynecol. 68 - 649 - 53.
- Carol, G. ; Duley, I. ; Belizan, J.M. and Villar, J. (1994) :** Calcium supplementation during pregnancy : A systemic review of randomized controlled trials. Br. J. Obstet. Gynecol. 101: 753 - 8.

- Catalano, P.M. ; Tyzbir, E.D. ; Roman, N.M. ; Amini, S.B. and Sims, E.A. H. (1991) :** longitudinal changes in insulin release and insulin resistance in nonobese pregnant women. *Am. J. Obstet. Gynecol.* 165: 1667 - 72.
- Chen, Y-D.I. ; Shew, W.H.-H. ; Swislock, A.L.M. and Reven, G.H. (1991):** High density lipoprotein turnover in patients with hypertension. *Hypertension.* 17 - 386.
- Chen, Y.D.L. ; Swami, S. and Skowronski, R. (1993) :** Differences in post prndial lipemia between patients with normal glucose tolerance and non-insulin-dependent diabetes mellitus. *J. Clin. Endocrinol. Metab.* 72: 172.
- Chesley, I.C. (1971) :** Hypertensive disorders in pregnancy. Williams obstetrics, 14<sup>th</sup> edition. Edited by: I.M. Hellman and J.A. Pritchard, New York, Appleton-Century-Crofts. pp 120-125.
- Chesley, I.C. (1974) :** A short history of eclampsia. *Obstet. Gynecol.* 43: 599.
- Chesley, I.C. (1985) :** Diagnosis of pre-eclampsia *Obstet. gynecol.* 65: 423.
- Chesley, L.C. (1972) :** Plasma and red cell volumes during pregnancy. *Am. J. obst. Gynecol.* 112 : 440 - 9.
- Chesley, L.C. (1984) :** History and Epidemiology of preeclampsia - eclampsia. *Clin. Obstet. Gynecol.* 27: 801 - 20.
- Chesley, L.C. and Sibai, B.M. (1988) :** Clinical significance and elevated mean arterial pressure in the second trimester. *Am. J. Obstet Gynecol.* 93: 176 - 81.

- Chesley, L.C. ; Talledo, O.E. ; Bohler, C.S. and Zuspan, F.P. (1975) :**  
Vascular reactivity to angiotensin II and norepinephrin in pregnant and non pregnant women. Am. J. Obstet. Gynecol. 91: 837 - 72.
- Chesley, L.C. and Cooper, D.W. (1986) :** Genetic of hypertension in pregnancy : Possible single gene control of preeclampsia and eclampsia in decendants of eclamptic women. Br. J. Obstet. Gynecol. 93 142.
- Cincotta, R.B. and Brennecke, S.P. (1988):** Family history of pre-eclampsia as a predictor for pre-eclampsia in primigravides. International Journal of Gynecology and Obstetrics 60: 23-27.
- Clark, B.A. ; Halvorson, I. ; Sachs, B. and Epstein, F.H. (1992) :** Plasma endothelin levels in preeclampsia : Elevation and correlation with uric acid levels and renal impairment. Am. J. Obstet. Gynecol. 166 : 962.
- Cooper, D.W. ; Brennecke, B.P. and Wilton, A.N. (1993):** Genetics of pre-eclampsia. Hypertens pregnancy 12: 1-23.
- Crouse, J.R. ; Parks, J.S. and Schey, H.M. (1985) :** Studies of low density lipoprotein molecular weight in human beings with coronary artery disease. J. Lipid. Res. 26: 566.
- Cunningham, F.G. ; MacDonald, P.C. and Gant, N.F. (1994) :** hypertensive disorders of pregnancy in willam obst. 19th. ed. Aplleton and lan p. 763 - 817.
- Daniel W.W. (1987) :** Biostatistics, A foundation for analysis in the health sciences. 4th Ed. John Wiley & Sons. New York; Chichester; Brisbane; Toronto & Singapore.

- Dati, F. ; Lammers, M. ; Metzmann, E. ; Marcovina, S. and Redaelli, G. (1989) :** Evaluation and assays for apolipoprotein A-I and B on turbidimeter. Clin. Chem. 35: 1072.
- Davey, D.A. (1995) :** Hypertensive disorders of pregnancy in Dewhurst's Textbook of Obstet. and Gynecol. for postgraduates 5th. (ed) London Backwell Scientific Publications.
- Davey, D.A. ; Dommissie, J. and Opie, L.H. (1984) :** A comparison of serum cystine aminopeptidase and urinary oestrogen secretion as placental function tests. In : Baldwin R. W. M. and Handon G.C. (eds) the critically III Obstetric patient. London : Farrand press p. 246.
- Davey, D.A. and MacGillivray, I. (1988) :** Sever Hypertension Preeclampsia and eclampsia. Am.J. Obstet. Gynecol. 158 982 - 8.
- Defronzo, R.A. and Ferrannini, E. (1991) :** Insulin resistance a multifaceted syndrome responsible for NIDDM obaty hypertension dyslipidemia and atherosclerotic cardiovascular disease. Diabetes. Care. 14: 173-194.
- Degani, S. ; Abinader, E. ; Eibschitz, I. ; Oettinger, M. ; Shapiro, I. and Sharf, M. (1985) :** Isometric exercise test for predicting gestational hypertension. Obstet. Gynecol. 65: 657 - 4.
- Dejager, S. and Turpin, G. (1996):** Hyperlipidemia in pregnancy Press-Med. 25 (37): 1839-45.
- Dekker, G.A. (1989) :** Prediction and prevention of pregnancy-induced hypertensive disorders : a clinical and pathophysiologic study [Thesis]. Rotterdam : Erasmus university.

- Dekker, G.A. and Sibai, M.D. (1991)** : Early detection of preeclampsia. Am. J. Obstet. Gynecol. 165: 160 - 70.
- Dieckmann, W.J. and Michel, H.L. (1973)** : Vascular - renal effects of posterior pituitary extracts in pregnant women. Am. J. Obstet. Gynecol. 33: 131.
- Differs, G.M. and MacGillivray, I. (1968)** : The incidence of pre-eclamptic toxemia in smokers and non-smokers. Lancet, 994-5.
- Donatella Santoro, M.D.; Alfredo Quinones Galvan, M.D.; Andrea Natali, M.D.; Eleuterio Ferrannini, M.D.; Pisa; Italy (1993)** : Some metabolic aspects of essential hypertension and its treatment. Am. J. 94 (Supp 4A) 32S - 39 S.
- Douglas, J.T. ; Shah, M. ; Lowe, G.D.O. ; Belch, J.J.F.I. ; Forbes, C.D. and Prentice, C.R.M. (1982)** : Plasma fibrinopeptide A and betathromboglobulin in preeclampsia and pregnancy hypertension. Thromb Haemost. 47: 54 - 5.
- Downing, I. ; Sheppard, G.L. and Lewis, P.J. (1982)** : Kinetic of prostaglandin synthetase in umbilical microsomes from normal and preeclamptic pregnancies. Br. J. Pharmacol. 13: 195 - 198.
- Eisenberg, S. (1984)** : High density lipoprotein metabolism. J. Lipid Res. 25: 1017.
- Endresen, H.J. ; Lorentzen, B. and Henriksen, T. (1992)** : Increased lipolytic activity and high ratio of free fatty acids to albumin in sera from women with preeclampsia leads to triglyceride accumulation in cultured endothelial cells. Am. J. Obstet. Gynecol. 167: 440 - 447.

- Eskenazi, B. ; Fenster, L. and Sidney, S. (1991) :** A multivariate analysis of risk factors for preeclampsia. *J.A.M.A.* 266: 237 - 41.
- Easterling, T.R. ; Benedetti, T.J. ; Schmucker, B.C. and Millard, S.P. (1990) :** Maternal hemodynamics in normal and preeclamptic pregnancies. a longitudinal study. *Obstet. Gynecol.* 76: 1061 - 9.
- Evans, D.H. ; Hoffman, R.G. and Kalkhoff, R.K. (1984):** Relationship of body fat to pography to insulin sensitivity and metabolic profiles in premenopausal women. *Metabolism.* 33: 68-72.
- Facchini, F. ; Chen, Y-D.I. and Clinkingbeard, C. (1992):** Insulin resistance hyperinsulinaemia and dislipidemia in individuals with a family history of hypertension.
- Facchini, F. ; Chen, Y-D.I. ; Hollenbeck, C.B. and Reaven, G.M. (1991):** Relationship between resistance to insulin-mediated glucose uptake, urinary uric acid clearance and plasma uric acid concentration. *J.A.M.A.* 266: 3008.
- Falkner, B. ; Hulman, S. and Kushner, H. (1993) :** Insulin-stimulated glucose utilization and borderline hypertension in young adult blacks. *Hypertension.* 22: 18.
- Falkner, B. ; Hulman, S. ; Tennenbaum, J. and Kushner, H. (1990) :** Insulin resistance and blood pressure in young black men. *Hypertension.* 16: 706.
- Ferrannini, E.; Buzzigoli, G. and Bonadonna, R. (1987) :** Insulin Resistance in essential hypertension. *N. Engl. J. Med.* 317: 350 - 7.

- Ferrannini, E. ; Haffner, S.M. ; Mitchell, B.D. and Stern, M.P. (1991) :**  
Hyperinsulinemia : The key feature of a cardiovascular and  
metabolic syndrome. *Diabetologia*. 34: 416 - 22.
- Ferrazzani, S. ; Caruso, A. ; Decarolis, S. ; Maltino, I.V. and Mancuso, S.  
(1990) :** Proteinuria and outcome of 444 pregnancies complicated  
by hypertension. *Am. J. Obstet. gynecol.* 162: 366.
- Fitzgerald, D.J. ; Entman, S.S. ; Mulloy, K. and Fitzgerald, G.A. (1987):**  
Decreased prostacyclin biosynthesis preceding the clinical  
manifestation of pregnancy-induced hypertension. *Circulation*. 75:  
956 - 63.
- Fonteyn, V.Z. and Isada, N.B. (1988) :** Non--genetic implication of child  
bearing after the age of thirty-five. *Obstet. Gynecol. Surv.* 4: 709 -  
720.
- Fossati, P. and Prencipe, L. (1982) :** Triglycerides Enzymatic  
determination. *Clin. Chem.* 28 2077.
- Fox, H. (1990) :** Placental structure of health and disease; In: *Modern  
Antenatal Care of the Fetus*; ed., Geoffrey Chamberlain, Blackwell  
Scientific Publications, Oxford, London, p. 52.
- Franz, H. and Wendler, D. (1992) :** A controlled study of maternal serum  
concentrations of lipoprotein in pregnancy-induced hypertension.  
*Arch. Gynecol. Obstet.* 252: 81 - 86.
- Friedwald, W.T.; Levy, R.I. and Fredrickson, D.S. (1972) :** Estimation of  
the concentration of low-density lipoprotein cholesterol without the  
use of the preoperative ultracentrifuge *Clin. Chem.*, 18: 499

- Friedman, E.A. and Neff, R.K. (1976)** : Pregnancy outcome as related to hypertension edema and proteinuria. In Lindheimer M.D. ; Katz A.I. ; Zuspan F.P. (eds) : Hypertension in pregnancy. New York wiley p. 13.
- Friedman , E.A. and Neff, R.K. (1977a)**: Pregnancy hypertension : A systolic evaluation of clinical diagnostic criteria. Littleton, M.A. PSG Publishing. p. 238.
- Friedman, S.A. (1988)** : Preeclampsia : A review of the role of prostaglandins. *Obstet Gynecol.* 71: 122 - 37.
- Friedman, S.A. ; Taylor, R.N. and Roberts, J.M. (1991)** : Pathophysiology of preeclampsia. *Clin. Perinatol.* 18 (4): 661 - 82.
- Gallery, E.D.M. ; Hunyor, S.N. and Györy, A.Z. (1979)** : Plasma volume concentration : a significant factor in both pregnancy associated hypertension (pre eclampsia) and chronic hypertension in pregnancy 48: 593 - 602.
- Gant, N.F. ; Daley, G.L. ; Chands ; Whalley, P.J. and MacDonald, P.G. (1973)** : A study of angiotensin II pressor response throughout primigravid pregnancy. *J. Clin. invest.* 52: 2682 - 9.
- Gant, N.F. ; Vhand, S. ; Worley, R.J. ; Whalley, P.J. ; Crosby, U.D. and MacDonald, P.G. (1974b)** : A clinical test useful for predicting the development of acute hypertension in pregnancy. *Am J Obstet. Gynecol.* 120: 1.
- Gibson, B. ; Hunter, D. ; Ncane, P.B. and Kelton, J.G. (1982)** : Thrombocytopenia in preeclampsia and eclampsia. *Semin thromb. Haemost.* 8: 234 - 47

- Giles, C. and Inglis, T.C.M. (1981) : Thrombocytopenia and macrothrombo-cytosis in gestational hypertension. Br. J. Obstet. Gynecol. 88: 1115 - 9.*
- Goode, G.K.; Miller, J.P. and Heagerty, A.M. (1995): Hyperlipidaemia, hypertension and coronary heart disease. Lancet. 345: 362-4.*
- Golay, A. ; Zech, L. and Shi, M.Z. (1987): High density lipoprotein (HDL) metabolism in non-insulin-dependent diabetes mellitus: Measurement of HDL turnover using tritiated HDL. J. Clin. Endocrinol. Metab. 65: 512.*
- Gimbrone, M.A. and Alexander, R. (1975) : Angiotensin II stimulation of prostaglandin production in cultured human vascular endothelium. G. A. ; Brown M. ; Lewis P.J. (1979) : Prostacyclin stimulates renin A II aldosterone in man. J. Clin. Endocrin. Meta. 49: 943 - 949.*
- Graninger, W. ; Tatra, G. ; Pirich, K. and Nas, Z.F. (1985) : Low antithrombin III and high fibronectin in preeclampsia. Eur. J. Obstet. Gynecol. Reprod. Biol. 19 : 223 - 9.*
- Gray, C.H. ; Howorth, P.J.N. and Rinsler, M.G. (1985) : Clinical chemical pathology 10th ed. Edward Arnold Ltd.ch. 9 pp. 152.*
- Greer, I.A. ; Walker, J.J. and McLaren, M. (1985): Immunoreactive prostacyclin and thromboxane metabolites in normal pregnancy and puerperium. Br. J. Obstet. Gynecol. 92: 381-385.*
- Haffner, S.M. ; Fong, D. ; Hazuda, H.P. ; Pugh, P.A. and Potterson, J. K. (1988) : Hyperinsulinemia upper body adiposity and cardiovascular risk factors in non-diabetics. Metabolism. 37: 338 - 45.*

- Haffner, S.M. ; Stern, M.P. and Hazuda, H.P. (1987):** Do upper body and centralization adiposity measure different aspect of regional body fat distribution.
- Hall, J.E. ; Brands, M.W. and Kivlighn, S.D. (1990):** Chronic hyperinsulinemia and blood pressure. *Hypertension*. 15: 519.
- Hamberg, M. ; Svensson, J. and Samuelson, B. (1975) :** Thromboxanes : new group of biologically active compounds derived from prostaglandin endoperoxides. *Proc. Natol. Acad. Sci. : USA* 71: 345 - 349.
- Hamsten, A. ; Wiman, B. ; Defaire, V. and Blombäck, M. (1985) :** Increased plasma level of a rapid inhibitor of tissue plasminogen activator in young survivors of myocardial infarction. *N. Engl J. Med.* 313: 1557.
- Hannele Laivuori Matti, J. ; Tikka, N.E.N. and Olavi Ylikork al, A. (1996) :** Hyperinsulinemia 17 years after preeclamptic first prgnancy. *Journal of Clinical Endocrinology and Metabolism* 81(8): 2908 - 2911.
- Hanretty, K.P. and Whittle, M.J. (1988) :** Doppler uteroplacental wave forms in pregnancy-induced hypertension : a reappraisal. *Lancet* 1. 850 - 5.
- Havel, R.J. ; Goldstein, J.L. and Brown, M.S. (1980) :** Lipoprotein and lipid transport. In : *Metabolic control and disease* 3rd edition Bondy P.K. and Rosenberg L.K. (eds) W.B. Saunder Co. Philadelphia p 393.

- Hays, P.M. and Gruikshank Dunn, L.J. (1985) :** Plasma volume determination in normal and preeclamptic pregnancies. *Am. J. Obstet. Gynecol.* 151: 958 - 66.
- Herbert, P.N. ; Asmann, G. ; Gotto, A.M. and Fredrickson, D.S. (1982) :** Familial lipoprotein deficiency in : The metabolic basis of inherited disease 2nd edition. Goldstein J.L. and Brown M.S. (eds). MacGraw Hill Inc. New York p. 589.
- Herrera, E. ; Lasuncion, M. A. ; Gomezcoronado, D.; Aranda, P. ; Lopen-Luna, P. and Maier, I. (1988) :** Role of lipoprotein lipase activity on lipoprotein metabolism and the fate of circulating triglycerides in pregnancy. *Am. J. Obstet. Gynecol.* 158: 1575 - 83.
- Hollenbeck, C.B. and Reaven, G.M. (1987) :** Variations in insulin-stimulated glucose uptake in healthy individuals with normal glucose tolerance. *J. Clin. Endocrinol. Metab.* 64: 1169.
- Hubel, C. A., Ph.D. ; McLaughlin, M. K. Ph.D. ; Evans, R.W. Ph.D. ; Beth, A. ; Hauth, B.S. ; Cynthia, J. ; Sims, M.D. ; James, M. and Roberts, M.D. (1996) :** Fasting serum triglycerides free fatty acids and malondialdehyde are increased in preeclampsia are positively correlated and decrease within 48 hours post partum 174: 975 - 982.
- Husby, H.; Roaled, B.; Schjetlein, R.; Nesheim, B.I. and Berg, K. (1996):** High levels of LP (a) lipoprotein in a family with cases of severe preeclampsia. *Clin. Genet.* 50 (1): 47-9.
- Jimenez, S.M. ; Pocovi, M. ; Ramon Cajal, J. ; Romero, M.A. ; Jones, D. D. ; Bahijri, S. ; Roberts, E.L. and Williams, G.F. (1988):**

- Activity of serum cytidine deaminase during pregnancy. Br.J. Obstet. Gynecol. 89: 314 - 7.
- Jones, D.D ; Bahijri, S. ; Roberts, E.L. and Williams, G.F (1982):** Activity of serum cytidine deaminase during pregnancy. Br. J. Obstet. Gynecol. 89: 314-7.
- Juhan-Vague, I. ; Alessi, M.C. and Vague, P. (1991) :** Increased plasma Plasminogen activator inhibitor 1 levels. A possible link between insulin resistance and atherothrombosis. Diabetologia 34: 457.
- Juhan-Vague, I. ; Thompson, S.G. ; Jespersen, J. ; on Behalf of the ECAT Angina Pectoris Study Group. (1993) :** Involvement of the hemostatic system in insulin resistance syndrome. Arterioscler. Thromb. 13: 1865.
- Juliasteinberger, M.D; Catherine Moorehead, M.S.; Victor Katch, Ed.D. and Albert, P. Rocchini, M.D. (1995) :** Relationship between insulin resistance and abnormal lipid profile in obese adolescents. The Journal of Pediatrics. Volume 126, Number 5, Part 1 page 690 – 695.
- Kaaja, R. ; Tikkanen, M.J. ; Viinikka, L. and Ylikorkala, O. (1995) :** Serum lipoproteins insulin and urinary prostanoid metabolites in normal and hypertensive women. Obstet. Gynecol. 85: 353 - 356
- Kalpan, L.A. and Pesce, A.J. (1989) :** Disorders of lipid metabolism. In : Clinical chemistry theory analysis and correlation; 2nd edition, CV Mosby Company st. louis Missouri p. 468.

- Kalpan, N.M. (1989) :** The deadly quartet : upper-body obesity glucose intolerance hypertriglyceridemia and hypertension. Arch Intern Med. 149: 1514 - 1520.
- Kaunitz, A.H. ; Hughes, J.H. ; Grimes, D.A. ; Smith, J.C. ; Rochat, R.W. and Kafrissen, M.E. (1985) :** Causes of maternal mortality in United States. Obstet. Gynecol. 65: 605 - 12.
- Kawasak, N. ; Matsui, K. ; Ito, M. ; Mushijima, H. ; Yoshimura, T. and Okamura, H. (1990) :** Effect of increased calcium intake during the third trimester on the vascular sensitivity of angiotensin II. Clin. and Exp. Hypertension in Pregnancy B9: 19 - 26.
- Khong, T.Y. ; Dewoff, F. ; Robertson, W.B. and Bronsens, I. (1986) :** Inadequate maternal vascular response to placentation in pregnancies complicated by preeclampsia and by small - for gestational - age infants. Br. J. Obstet. Gynecol. 93: 1049 - 1059.
- Kilby, M.D. ; Broughton, P. ; Pkin, F. and Symond, E. M. (1992) :** Calcium platelets and hypertension. J. hypertension. 10: 997 - 1003.
- Kissebah, A.H. ; Adams, P.W. and Wynn, V. (1982) :** Inter-relationship between insulin secretion and plasma free fatty acid and triglyceride transport kinetics in maturity onset diabetes and the effect of phenethylbiguanide (Phenformin). Diabetologia 10: 119 - 30.
- Kissebah, A.H. ; Vydellingum, N. and Murray R. (1982):** Relationship of body fat distribution to metabolic complications of obesity. J. Clin. Endocrinol. Metab. 54: 254-60.
- Knopp, R.H. ; Bergelin, R.O. ; Whal, P.W. ; Walden, C.E. ; Cahpman, M. and Irvin, S. (1982) :** Population-based lipoprotein lipid reference

values for pregnant women compared to non pregnant women classified by sex hormone usage. *Am. J. Obstet. Gynecol.* 143: 626 - 637.

**Knopp, R.H. ; Bonet, B. ; Lasuncion, M.A.; Montelongo, A. and Herrera, E. (1992) :** Lipoprotein metabolism in pregnancy. In : Herrera E. ; Knopp R.H. ; eds. *Perinatal biochemistry*. Boca Raton : CRC press 19 - 51.

**Knudsen, P. ; Eriksson, J. ; Lahdenpera, S. ; Kahri, J. ; Groop, L. and Taskinen, M.R. (1995) :** Changes of lipolytic enzymes cluster with insulin resistance syndrome. *Diabetologia*. 38: 344 - 50

**Kobayashi, S. ; Tanaka, M. ; Masaki, K. ; Hirakawa, S. and Momose, K. (1992) :** Apolipoprotein levels in preeclamptic pregnancies. *Nippon Sanka. Fujinka. Gakkai. Zasshi*. 44 (2): 223 - 8.

**Kokia, E. ; Barkai, G.; Reichman, B.; Segal, P.; Goldman, B. and Mashiach, S. (1990) :** Maternal serum lipid profile in pregnancies complicated by hypertensive disorders. *J. Perinat. Med* 18: 473-8.

**Kottke, B.M. (1986) :** Lipid markers for athero sclerosis. *Am J Cardiol* ; 57; 57: 11C.

**Krauss, R.M. (1982) :** Regulation of high density lipoprotein levels. *Med Clin. North Am.* 66: 403.

**Kreisberg, R.A. and Kasim, S. (1987):** Cholesterol metabolism and aging. *Am. J. of medicin*. 82: (Suppl 1B) 54-59.

**Krotkiewski, M. ; Bjorntop, P. and Sjostrom, C. (1983):** Impact of obesity on metabolism in men and women: importance of regional adipose distribution. *J. Clin. Invest.* 72: 1150-62

- Kuhai, C. (1975) :** Glucose metabolism during and after pregnancy in normal and gestational diabetic women. I. Influence of normal pregnancy on serum glucose and insulin concentration during basal fasting conditions and after a challenge with glucose. *Acta. Endocrinol* 79: 709 - 19.
- Laaksor, M. and Barrett-Connor, E. (1989):** A symptomatic hyperglycemia is associated with lipid and lipoprotein changes favoring atherosclerosis. *Arteriosclerosis*. 9: 665 - 72.
- Laivuor, H. ; Tikkanen, M.Z. and Ylikorkala, O. (1996) :** Hyperinsulinemia 17 years after preeclamptic first pregnancy. *J. Clin. End. And Metabol.* 81 (8) 29: 8-11.
- Landin, K. ; Tengrory, L. and Smith, U. (1990) :** Elevated fibrinogen and plasminogen activator inhibitor I (PAI-I) in hypertension are related to metabolic risk factors for cardiovascular disease. *J. Intern. Med.* 227: 273.
- Laws, A. ; King, A.C. ; Haskell, W.L. and Reaven, G.M. (1991) :** Relation of fasting plasma insulin concentration to high density lipoprotein cholesterol and triglyceride concentrations in men. *Arterio. Thromb.* 11: 1636.
- Lazarchick, J. ; Stubbas, T.M. ; Romein, L. ; Dorsteh, J.P-van and Loadholt, C.B. (1986) :** Predictive value of fibronectine levels in normotensive gravid women destined to become preeclamptic. *Am. J. Obstet. Gynecol.* 154: 1050 - 2.
- Leerink, C.B.; de-Varies, C.V. and Van-der-Killis, F.R. (1997):** Elevated levels of serum lipoprotein (a) and apolipoprotein (a) phenotype are

not related to preeclampsia. Acta. Obstet. Gynecol. Scand. 76 (7): 625-8.

**Lefevre, M. and Roheim, P.S. (1984) :** Metabolism and apo A-II. J. lipid Research 25: 1603.

**Leturque, A. ; Ferre, P. ; Burnal, A.F. ; Kande, J. ; Maulard, P. and Girard, J. (1986) :** Glucose utilization rates and insulin sensitivity in vivo in tissues of virgin and pregnant rates. Diabetes. 35 : 172 - 7.

**Levine, R.J. ; Raymond, E. ; Dersimonian, R. and Clemens, J.D. (1992) :** Preeclampsia prevention with calcium supplenation. Clin. Appl. nutr. 2: 30 - 8.

**Lewis, G.F. ; D'Meara, N.M. and Softys, P.A. (1990):** Postprandial lipoprotein metabolism in normal and obese subjects. comparison after vitamin A fat loading test. J. Clin. Endocrinol. Metab. 71: 1041.

**Lewis, P.J. ; Boyland, P. ; Friedman, L.A.; Hensby, C.N. and Downing, I. (1980) :** Prostacyclin in pregnancy. Br. Med. J. 282: 1581 - 1580.

**Lind, A.R. and McNicol, G.W. (1976) :** Circulatory responses to sustained hand grip contractions performed during other exercise both rhythmic and static. J. Physiol. 192: 595 - 607.

**Long, P.A. ; Abell, D.M. and Beischler, N.A. (1980) :** Fetal growth retardation and preeclampsia. Br. J. Obstet. Gynecol. 87: 13 - 18.

**Long, P.A. and Oats, J. (1987) :** Preeclampsia in twin pregnancy-severity and pathogenesis. Aust NZJ Obstet. Gynecol. 27: 1 - 5.

**Lopez-Epinoza, I. ; Dhar, H. ; Humphreys, S. and Redman, C.W.G.**

**(1986) :** Urinary albumin excretion in pregnancy. Br. J. Obstet. Gynecol. 176 - 81.

**López-Jaramillo ; Narvaez, M. ; Weigel, R.M. and Yepez, R. (1989) :**

Calcium supplementation reduces the risk of pregnancy-induced hypertension in an Andes population. Br. J. Obstet. Gynecol. 96: 648.

**Lorentzen, B. ; Endersen, M.J. ; Hovig, T. ; Haug, E. and Henriksen, T.**

**(1991) :** Sera from preeclamptic women increase the content of endothelial cells. Thromb. Res. 63: 363 - 372.

**Lorentzen, B. ; Endersen, M.J. ; Clausen, T. and Henriksen T. (1994) :**

Fasting serum free fatty acids and triglycerides are increased before 20 weeks of gestation in women who later develop preeclampsia. Hypertens. Preg. 13: 103 - 9.

**Louden, L.A. and Kilby, M.D. (1996) :** Low dose aspirin : The rationale for

preventing pre-eclampsia and intrauterine growth retardation : A role after CLASP? In recent advances. in Obstet. and Gynecol. pp. 15 - 33.

**Lorenzo, B.; Christian, A. ; Drevon ; Marit, J. ; Endresen and Tore**

**Henriksen (1995):** Fatty acid pattern of esterified and free fatty acids in sera of women with normal and preeclamptic pregnancy. Br. J. Obstetrics and Gynaecology 102: 530-537.

**MacGillivray, I. (1961) :** Hypertension in pregnancy and its consequences. J.

Obstet. Gynecol. Br. Connwlth 68 557 - 69.

- MacGillivray, I. (1977) :** Is mild pre-eclampsia harmful to the baby. *Ist J Med. Sci.* 12: 500 - 3.
- MacGillivray, I. (1983) :** Pre-eclampsia. The hypertensive disease of pregnancy. London : W B Saunders.
- MacGillivray, I. (1988) :** Some observation on the incidence of preeclampsia. *J. Obstet. Gynecol. Br. Emp.* 65: 536 - 9.
- Mahley, R.W. (1982) :** Atherogenic hyperlipoproteinemia : the cellular and molecular biology of plasma lipoproteins altered by dietary fat and cholesterol. *Med. Clin. North Am.* 66: 375.
- Maki, M. (1983) :** Coagulation fibrinolysis platelet and kinin forming systems during toxemia of pregnancy. *Biol. Res. Pregn. Perinatol.* 4: 152 - 4.
- Manas, K.J. ; Welsh, J.D. ; Rankin, R.A. and Miller, D.D. (1985) :** Hepatic haemorrhage without rupture in preeclampsia. *N. Engl. J. Med.* 312: 426.
- Marcoux, S. ; Brisson, J. and Fabia, J. (1991) :** Calcium intake from diary products and supplements and the risks of preeclampsia and gestational hypertension. *Am. j. Epidemial.* 133: 1266.
- Marshall, W.J. (1990) :** Lipid and lipoproteins. In : illustrated text book of clinical chemistery. 1st. edition. Goodfellow and Moehrole (eds) J.B. lippin cott company Philadelphia p 210.
- Martinez, H. and Grande, F. (1988) :** Longitudinal study of plasma lipids and lipoprotein cholesterol in normal pregnancy and puerperium. *Gynecol. Obstet. Invest.* 25: 158 - 164.

- Marya, R.K. ; Rathee, S. and Mittel R. (1988) :** Evaluation of three clinical tests for predicting pregnancy-induced hypertension. *Am. J. Obstet. Gynecol.* 158: 683 - 4.
- Maseki, M. ; Nishigaki, I. ; Hagihara, M. and Tomodak Yagi, K. (1981) :** Lipid peroxide levels and lipid content of serum lipoprotein fractions of pregnant subjects with or without preeclampsia. *Clin. Chim. Acta.* 115: 155 - 161.
- Mastrogiannis, D.S. ; O'Briem, W.F. ; Krammer, J. ; Benoit, R. (1991):** Potential role of endothelin I in normal and hypertensive pregnancies. *Am. J. Obstet. Gynecol.* 165: 1711.
- Mccarron, D.A. and Morris, C.D. (1985) :** Blood pressure response to oral calcium in persons with mild to moderate hypertension. A randomized double-blind placebo controlled crossover trial.
- Mckay, D.G. (1965) :** Disseminated intravascular coagulation. New York Harper and Row.
- Mehata, J. ; Mehta, P. ; Lawson, D. and Saldeen, T. (1987) ;** Plasma tissue plasminogen activator inhibitor levels in coronary artery disease : Correlation with age and serum triglyceride concentration. *J. Am. Coll. Cardiol.* 9: 263.
- Michael, A.; Weber, M.D.; Irvine, California (1993) :** Hypertension as a risk factor syndrome. Therapeutic Implications. *Am. J.* 94 (Suppl 4A) 24S - 31S.
- Minkes, M. ; Stanford, N. and Chim, M.Y. (1979) :** Cyclic 3' 5' AMP inhibits the availability of arachidonate to prostaglandin synthetase in human platelet suspensions. *J. Clin. Invest.* 59: 449.

- Miyamori, I. ; Fitzgerald, G.A. ; Brown, M. and Lewis, P.J. (1979) :**  
Prostacyclin stimulates renin A II aldosterone in man. J. Clin. Endocrine Metab. 49: 943 - 949.
- Modan, M.; Halkin, H.; Almog, S.; Lusky, A.; Eshkol, A.; Shefi, M.; Shitrit, A. and Fuschs, Z. (1988):** Hyperinsulinemia a link between hypertension. Obesity and Glucose Intolerance J. Clin. Invest. 75: 809-17.
- Molnár, M. and Hertelendy, F. (1992) :** NW - Nitro - L - arginine an inhibitor of nitric oxidase synthesis increases blood pressure in rats and reverses the pregnancy-induced refractoriness to vasopressor agents. Am.J. Obstet. Gynecol. 166: 1560.
- Moncada, S. ; Gryglewski, R. ; Bunting, S. and Van, J.R. (1976) :** An enzyme isolated from arteries transforms -prostaglandin endoperoxides to an unstable substance that inhibits platelet activation. Nature. 263: 663 - 665.
- Moncada, S. and Vane, J.R. (1979) :** Arachidonic acid metabolism and the interaction between platelets and the endothelium. N. Engl. J. Med 300: 1142 - 1147.
- Montaquin, J.M. ; Rainville, G. ; Giroux, L. ; Raynauld, P. ; Amyot, G. ; Bilodeau, R. and Pelland, N. (1985) :** A prospective study of blood pressure in pregnancy : prediction of preeclampsia. Am. J. Obstet. Gynecol. 151: 191 - 6.
- Monelongo, A. ; Lasuncoín, M.A. ; Pallarado, L.F. and Herrera, E. (1992):** Longitudinal study of plasma lipoproteins and hormones

- during pregnancy in normal and diabetic women. *Diabetes*. 41: 1651-1659.
- Muller, D. C. ; Dariush E. ; Richard, E. P.; Jordan, D. T. and Reubin, A. (1993):** An epidemiological test of the hyperinsulinemia-hypertension hypothesis. *Journal of Clinical Endocrinology and Metabolism* 76 (3): 544 – 48.
- Muller, F. ; Savey, L. ; Lifiblec, B. ; Bussieres, L. ; Nolayizamba, G. ; Claude, J. and Giraudet, P. (1996):** Maternal serum human chorionic gonadotropin level at fifteen weeks is a predictor for preeclampsia. *Am. J. Obstet. Gynecol.* 175: 37-40.
- Munoz, A.; Uberos, J. Molina, A.; Valenzuela, A.; Cano, D.; Ruiz, C. and Molina Font, J.A. (1995):** Relationship of blood rheology to lipoprotein profile during normal pregnancies and those with intrauterine growth retardation. *3-Clin. Pathol.* 48 (6): 571-4.
- Myant, N.B. (1983) :** The metabolic basis of familial hypercholesterolaemia. *Klinische Wochenschrift*. 61: 383.
- Myatt, L. (1986) :** Eicosanoids and blood pressure regulation. In : *Hypertension in pregnancy. Proceedings of the 16th study group for the RCOG Vol. (3) USA: Perinatology Press* pp 167 - 187.
- Myatt, L. and Elder, M.G. (1977) :** Inhibition of platelet aggregation by a placental substance with prostacyclin-like activity. *Nature*. 268 - 159 - 160.
- Nakamura, T. ; Ito, M. ; Mastui, K. ; Yushimura, I. ; Kawasaki, N. and Maeyana (1968):** Significance of angiotensin sensitivity test for

- prediction of pregnancy-induced hypertension. *Obstet. Gynecol.* 67: 388-94.
- Nikkilä, E.A. (1984)** : Plasma lipid and lipoprotein abnormalities in diabetes. In : Miller NE, Miller GJ eds. *Clinical and metabolic aspects of high-density lipoproteins*. Amsterdam: Elsevier. Science Publishers BV, 217 - 45.
- Nilson, E. ; Carfinkel, A.S. and Schotz, M. (1980)** : Lipolytic enzymes and plasma lipoproteins metabolism. *Ann. Rev. Biochem.* 49: 668.
- Nisell, H. ; Hjendahl, P. and Linde, B. (1965)** : Cardiovascular response to circulating catecholamines in normal pregnancy and in pregnancy-induced hypertension. *Clin. Physiol.* 5: 479 - 93.
- Nova, A. ; Sibai, B.M. ; Barton, J.R. ; Mercer, B.M. and Mitchell, M.D. (1991)** : Maternal plasma level of endothelin is increased in preeclampsia. *Am. J. Obstet. Gynecol.* 165: 724.
- O'Brein, W.F. (1990)** : Prediction of preeclampsia. *Obstet. Gynecol.* 75: 445.
- Odendaal, H.J. ; Pattinson, R.C. and Bam, R. (1990)**: Aggressive or expectant management for patients with severe preeclampsia between 28-34 weeks' gestation: a randomized controlled trial. *Obstet. Gynecol.* 76: 1070-5.
- Olefsky, J.M. ; Farquhar, J.W. and Reaven, G.M. (1974)** : Role of insulin in hypertriglyceridemia. *Am. J. Med.* 46: 1756.
- Olefsky, J.M. ; Reaven, G.M. and Farquhar, J.W. (1974)** : Effect of weight reduction on obesity : studies of carbohydrate and lipid metabolism. *J. Clin. Invest.* 53: 64.

- Öney, T. and Kaulhausen, H. (1982)** : The value of the angiotensin sensitivity test in the early diagnosis of hypertensive disorders of pregnancy. *Am. J. Obstet. Gynecol.* 142: 17.
- Orchad, T.J.; Recker, D.J.; Bates, M.; Kuller, L.H. and Drach, A.L. (1983)** : Plasma insulin and lipoprotein concentration: an atherogenic association? *Am. J. Epidemiol.* 118: 326 – 37.
- Otani Usuki, S. ; Saitoh, T. ; Yauagisawa, M. ; Iwasaki, H. ; Tanaka, J.; Suzuki, N. ; Fujino, M. ; Goto, K. and Masaki, T. (1991):** Comparison of endothelin-1 concentrations in normal and complicated pregnancies. *J. Cardiovascular Pharmacol.* 17: 5308.
- Ounsted, M. ; Cockburn, J. ; Moarv, A. and Redman, C.W. (1983) :** Maternal hypertension with superimposed preeclampsia : Effect on child development at 7 years. *Br. J. Obstet. Gynecol.* 90: 644.
- Page, E.W. and Christianson, R. (1976)** : Influence of blood pressure with and without proteinuria upon outcome of pregnancy. *Am. J. Obstet. Gynecol.* 126: 821- 33.
- Page, E.W. and Christianson, R. (1976)** : The impact of mean arterial blood pressure in the middle trimester upon the outcome of pregnancy. *Am. J. Obstet. Gynecol.* 125: 740 - 6.
- Pulmer, R.M.J. ; Ferrige, A.G. and Moncada, S. (1987)** : Nitric oxide release account for the biological activity of endothelium-derived relaxing factor. *Nature.* 327 - 524.
- Patterson, J.K. (1988)** : Hyperinsulinemia upper body adiposity and cardiovascular risk factors in non diabetics. *Metabolism.* 37: 338 - 45.

- Piechota, W. and Staszewski, A. (1992)** : Reference range of lipid and apolipoproteins in pregnancy. Eur. J. Obstet. Gynecol. Reprod. Med. 45: 27 - 35.
- Piper, P.J. and Van, J.R. (1969)** : The release of additional factors in anaphylaxis and their antagonism by anti-inflammatory drugs. Nature: 223: 39 - 35.
- Potter, J.M. and Nestel, P.J. (1979)** : The hyperlipidemia of pregnancy in normal and complicated pregnancies. Am. J. Obstet. Gynecol. 133: 165 - 70.
- Pritchard, J.A. ; Cunningham, F.G. and Mason, R.A. (1976)** : Coagulation changes in eclampsia : their frequency and pathogenesis. Am. J. Obstet. Gynecol. 124 855.
- Raah, W. ; Schroeder, G. ; Wagner, R. and Gige, W. (1956)** : Vascular reactivity and electrolytes in normal and toxemic pregnancy. J. Clin. Endocrinol. 16: 1196.
- Randeree, H.A. ; Omar, M.A.K. ; Motala, A.A. and Seedat, M.A. (1992)** : Effect of insulin therapy on blood pressure in NIDDM patients with secondary failure. Diabetes care. 15: 1258.
- Reaven, G.M. ; Lerner, R.L. ; Stern, M.P. and Farquhar, J. W. (1967)** : Role of insulin in endogenous hypertriglyceridemia. J. Clin. Invest. 46: 1756.
- Reaven, G.M. and Greenfield, M.S. (1981)** : Diabetic hypertriglyceridemia. evidence of three clinical syndromes. Diabetes. 30: 66 - 75.
- Reaven, G.M. (1988)** : Role of insulin resistance in human disease. Banting lecture (1988). Diabetes. 37: 1595 - 1607.

- Reaven, G.M. and Chen, Y-D.I. (1988)** : Role of insulin in regulation of lipoprotein metabolism in diabetes. *Diabetes/Metab. Rev.* 4: 639.
- Reaven, G.M. (1991)** : Insulin resistance hyperinsulinaemia hypertriglyceridemia and hypertension : parallels between human disease and rodent. *Diabetes. Care*; 14: 195.
- Reaven, G.M. ; Brand, R.J. and Chen, Y.D. (1993)**: Insulin resistance and insulin secretion are determinants of oral glucose tolerance in normal individuals. *Diabetes* 24:1324.
- Reaven, G.M. ; Chen, Y.D. and Jeppesen, J. (1993)** : Insulin resistance and hyperinsulinemia in individuals with small dense low density lipoprotein particles. *J. Clin. Invest.* 92: 141.
- Redman, C.W.G. ; Beilin, L.J. ; Bonnar, J. and Wilkinson, R.H. (1976)**: Fetal outcome in trial of ant: hypertensive treatment in pregnancy. *Lancet*, 1: 1370 -3.
- Redman, C.W.G. ; Beilin, L.J. ; Bonnar, J. and Wilkinson, R.H. (1976)**: Plasma-urate measurement in predicting fetal death in hypertensive pregnancy. *Lancet*. 1: 1370-3.
- Redman, C.W.G. ; Bonnar, J. and Beilin, L.J. (1978)** : Early platelet consumption in preeclampsia. *B. M. J.* 1: 467 - 9.
- Redman, C.W.G. and Jefferies (1988)** : Revised definition of preeclampsia. *Lancet*. 1: 809.
- Redman, C.W.G. (1991)** : Current topic: preeclampsia and the placenta. *Placenta* 12: 301-308.
- Redman, C.W.G. and Robert, J.M. (1993)** : Management of pre-eclampsia. *Lancet* 97: 762-769.

- Remuzzi, G.; Marchesi, D. and Zoja, C. (1980):** Reduced umbelical and placental vascular prostacyclin in sever preeclampsia . Prostaglandin 20: 105-110.
- Repke, J.T. and Villar, J. (1991) :** Pregnancy-induced hypertension and low birth weight : The role of calcium. Am. J. Clin. Nutr. 54: 2375.
- Richmond, W. (1993) :** Cholesterol determination. Clin. Chem. 19: 1350 - 1356.
- Rifai, N. (1986) :** Lipoproteins and apolipoproteins. Arch. Pathol. Lab. Med. 110: 694.
- Risto Kaa, J.A. ; M.D. Ph. D. ; Matti, J. Tikkanen M.D. Ph.D. ; LasseViinikka, M.D. Ph.D. and Olaviylikorkala, M.D. Ph. D. (1995) :** Serum lipoproteins insulin, and urinary prostanoid metabolites in normal and hypertensive pregnant women. Obstet. Gynecol. 185: 353 - 56.
- Rittenhouse-Simmonds, S. and Deykin, D. (1978) :** The activation of platelet phopholipase A<sub>2</sub>. The effect of CAMP. Biochem. Biophys. Acta : 543: 409 - 413.
- Robert, A. and Kreisberg, M.D.(1987) :** Cholesterol metabolism and Aging Am. J. 82 (Supp 1B) : 54 – 59.
- Robertson, E.G. (1971) :** The natural history of oedema during pregnancy. J. Obstet. Gynecol. Br. Common. 78: 520.
- Robertson, W.B. and Khong, T.V. (1987) :** Pathology of uteroplacental bed in : Sharp F.; Symonds E.M. ; (eds) : Hypertension in pregnancy. Ithaca New York : Perinatology Press. 101 - 18.

- Roberts, J.M.; Taylor, R.N.; Musci, T.J.; Rodgers, G.M.; Hubel, G.A. and Mclaughlin, M.K. (1989) :** Preeclampsia : an endothelial cell disorder. *Am. J. Obstet. Gynaecol.* 163: 1365 – 6.
- Romero, R. ; Mazor, M. ; Lockwood, C.J. ; Emamian, M.; Belanger, K.P.; Hobbins, J.C. and Duffy, T. (1989) :** Clinical significance ,prevalence and natural history of thrombocytopenia in pregnancy-induced hypertension. *Am. J. Perinatol.* 6: 32.
- Röschlau, P. ; Bernt, E. and Gruber, W. (1974) :** Enzymatische bestimmung des gesamt cholesterines im serum. *J. Clinchem. Clin. Biochem.* 12: 405 - 8.
- Rosing, U. ; Samsiol, G. ; Olund, A. ; Johansson, B. and Kallner, A. (1989) :** Serum levels of apolipoprotein A-I, AII and HDL-cholesterol in second half of normal pregnancy and in pregnancy complicated by preeclampsia. *Horm. Metab. Res.* 21 (7): 376 - 82 (Abstract).
- Ryan, E.A. ; O'Sullivan, M.Z. and Skyler, J.S. (1985) :** Insulin action during pregnancy : Studies with euglycemic clamp technique. *Diabetes.* 34: 380 - 9.
- Saad, M. ; Lillioza, S. and Myomba, B. L. (1991):** Racial differences in the relation between blood pressure and insulin resistance. *N. Engl. Med.* 324: 733.
- Sagen, N. ; Koller, O. and Haram, K. (1982) :** Haemoconcentration in sever preeclampsia. *Br. J. Obstet. Gynecol.* 89: 802 - 5.
- Sanchez-Remos, L. ; Sandroni, S. ; Andres, F.J. and Kaunitz A.M. (1991):** Calcium excretion in preecalmpsia. *Obstet. Gynecol.* 77: 510 - 3.

- Sanchez-Torres, F. and Santamaria, A. (1965) :** Review of Obstet and Gynecologie Venezuela 25 657 - 69. Quonted by Chesley L.C. : (1978) in : Hypertensive disorder of pregnancy. New York : Appleton - Century Crofts.
- Sattar, N.; Greer, I.A.; Loudon, J.; Lindsay, G.; McConnel, M.; Shepherd, J. and Packard (1977):** Lipoprotein subfraction changes in normal pregnancy: threshold effect of plasma triglyceride on appearance of small dense low density lipoprotein J Clin. Endocrinal. Metab. 82(8): 24: 83-91.
- Satter, N.; Mbchb; Astrid, B.; Mrco, G.; Colin, B.; James, S.; Greer, I. MD and Chris, J.P. DSC (1997):** Lipoprotein subfraction concentrations in pre-eclampsia : pathogenic parallels to Atherosclerosis. Obstetrics and Gynecology 89: 403-8.
- Savitz, D.A. and Zhang, J. (1992) :** Pregnancy- induced hypertension. Am. J. Public health - 82; 765 - 9.
- Schiff Ben-Baruch, G. ; Peleg, E. ; Rosenthol, T. ; Alcolay, M. ; Devir, M. and Mashiach, S.C. (1992) :** Immunoreactive circulation endothelin-I. in normal and hypertensive pregnancies. Am. J. obstet Gynecol. 166: 624.
- Schulman, H. (1987) :** The clinical implications of doppler ultrasound analysis of the uterine and umbilical arteries. Am. J. Obstet Gynecol. 156: 889 - 93.
- Schuster, E. and Wepelmann, B. (1981) :** Palsma urate measurements and fetal outcome in preecalampsia. Gynecol. Obstet. Invest 21: 162 - 7.

- Schwarz, Z.R. and Retzke, U. (1971) :** Cardiovascular response to infusion of angiotensin II in pregnant women. *Obstet. Gynecol.* 38: 714 - 8.
- Scott, J.S. (1968) :** Pregnancy toxemia associated with hydrops fetalis, Hydatidiform mole and hydramnios. *J. Obst. Gynecol. Br. Emp.* 65: 689-701.
- Seidmen, D.S. ; Ever-Hadani, P. ; Stevenson, D.K. and Gale, R. (1989) :** The effect of abortion on the incidence of preeclampsia. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 33: 109 - 14.
- Shamiss, A. ; Carroll, J. and Rosenthal, T. (1992) :** Insulin resistance in secondary hypertension. *Am. J. Hypertens.* 5: 26.
- Sheppard, B.L. and Bonnar, J. (1981) :** An ultrastructured study of the uteroplacental spiral arteries in hypertensive and normotensive pregnancy and fetal growth retardation. *Br. J. Obstet. Gynecol.* 88 695 - 705.
- Sibai, B.M. ; Anderson, G.D. and McCubbin, J.H. (1982) :** Eclampsia II. Clinical significance of laboratory findings. *Obstet. Gynecol.* 59: 153 - 7.
- Sibai, B.M. ; Anderson, G.D. ; Spinnata, J.A. and Shaver, D.C. (1983) :** Plasma volume findings in patients with mild pregnancy-induced hypertension. *Am. J. Obstet. Gynecol.* 147: 16-19.
- Sibai, B.M. (1990) :** Preeclampsia - eclampsia. *Currprob. Obstet. Gynecol. Fertil.* 13: 1 - 45.
- Sibai, B.M. ; Mercer, B.M. ; Schiff, E. and Friedman, S.A. (1994) :** Aggressive versus expectant management of severe preeclampsia at

- 28 to 32 weeks gestation: A randomized controlled trial. *Am. J. Obstet. Gynecol.* 171: 818-22.
- Smith, J.B. ; Ingerman, C. ; Kociss, J.J. and Silver, M.J. (1974) :** Formation of an intermediate in prostaglandin biosynthesis and its association with platelet release reaction. *J. Clin. Invest.* 53: 1468 - 1572.
- Smolarczyk, R.; Romejoko, E.; Wojcickajagodzinśka, J.; Czajkowski, K.; Teligacza; Jkowska, J. and Piekaraski, P. (1996):** Effect of total estrogens and luteal hormone on lipid metabolism in women during normal pregnancy. 67: (9): 438-42.
- Socal, M.L.; Weiner, C.P.; Louis, G.; Rehnberg, K. and Ross, E.G. (1985):** Platelet activation in preeclampsia. *Am. J. Obstet. Gynecol.* 151: 494 - 7.
- Stalder, M.; Pometta, D. and Suenram, A. (1981) :** Relationship between plasma insulin levels and high density lipoprotein cholesterol levels in healthy men. *Diabetologia.* 21: 279 - 87.
- Steel, S.A. ; Pearce, J.M. and Chamberlain, G.V. (1988) :** Doppler ultrasound of the uteroplacental circulation as a screening test for severe preeclampsia with intrauterine growth retardation. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 28: 279 - 87.
- Steel, S.A.; Pearce, J.M.; McParland, P. and Chamberlain, G.V.P. (1990):** Early doppler ultrasound screening in prediction of hypertensive disorders of pregnancy. *Lancet.* I: 1548 - 51.
- Stein, Z.A. (1985) :** A women's age : Child bearing and child rearing. *Am. J. Epidemiol.* 121: 327 - 342.

- Steinberger, J. ; Moorehead, C. ; Katch, V. and Rocchini, A.P. (1995) :**  
Relationship between insulin resistance and abnormal lipid profile in obese adolescents.
- Steven, M. Haffner; Leena Mykkanen; Rodolfo, A. Valdez, and Michals Katz (1993) :** Relationship of sex hormones to lipids and lipoproteins in nondiabetic Men. J. Clin. End. And Metabol. 77 (6) : 1610 – 15.
- Stevena-Friedman. (1988) :** Preeclampsia; A review of the role of prostaglandins. Obstet. Gynecol. 71: 22.
- Stuart, M.J.; Clark, D.A. and Sunderji, S.G. (1981):** Decreased prostacyclin production : a characteristic of chronic placental insufficiency . Lancet (I): 1126 : 1128.
- Suhonen, L. and Teramo, K. (1993) :** Hypertension and preeclampsia in women with gestational glucose intolerance. Acta. J. Obstet. Gynecol. Scand. 72: 269 - 72.
- Swenson, T.L. (1991) :** The role of the cholesterylester transfer protein in lipoprotein metabolism. Diab. Metab. Rev. 7: 139.
- Tall, A.R. and Small, D.M. (1978) :** Plasma high density lipoproteins. N. Engl. J. Med. 299: 1232.
- Talledo, O.E. (1966) :** Renin- angiotensin system in normal and toxemic pregnancies. I . Angiotensin infusion test. Am. J. Obstet. Gynecol. 96: 141 - 3.
- Talledo, O.E. ; Chesley, I.C. and Zuspan, F.P. (1968) :** Renin-angiotensin system in normal and toxemic pregnancies; III. Differential

- sensitivity to angiotensin II and norepinephrine in toxemia of pregnancy. *Am. J. Obstet. Gynecol.* 100: 218.
- Tan, C.E.; Squire, L.; Caslake, M.J.; Bedford, D.; Watson, T.D.G. and McCamnell, M. (1995):** Relationship between VLDL and LDL subfraction in normotensive men and women. *Arterioscler. Metab. Vasc. Biol.* 13, 839-48.
- Taskinen, M.R. (1990) :** Hyperlipidemia in diabetes. In : Betteridge DJ, ed. lipid and lipoprotein disorders. Beilieres Clin. Endocrinol. Metab ; 4: 743 - 75.
- Tatami, R. ; Mabuchi, H. and Ueda, K. (1981) :** IDL and cholesterol rich VLDL in angiographically determined CAD. *Circulation*; 64: 1174.
- Tateson, J.G. ; Moncada, S. and Vane, J.R. (1977) :** Effect of prostacyclin. *J. Obstet. Gynecol.* 10: 467-482.
- Tedde, R. ; Sechi, L.A. and Marigliano, A. (1989):** Antihypertensive effect of insulin reduction in diabetic hypertensive patients. *Am. J. Hypertens.* 2: 163.
- Teitz, N.W. (1987) :** Lipids; Lipoproteins; In : fundamentals of clinical chemistry. 3rd edition. W.B. Saunders; Philadelphia; p 448
- Teitz, N.W. ; Aldrich, J. ; Bhagavan, N.V. ; Coiner, D. ; Conn, R.B. ; Kowalczyk, R.S. ; Lechmann, H.P. ; Pruden, E.L. and Whitley, R.J. (1987) :** "Fundamentals of clinical chemistry". 3rd ed W.B. Saunders Company; Philadelphia; London; Toronto. Ch 14; pp. 448.
- Thaler, I. ; Monor, D. ; Itskovitz, J. ; Rotten, S. ; Levit, N. ; Timor-Tritsch, I. and Brandes, J.M. (1990) :** Changes in uterine blood flow during human pregnancy. *Am. J. Obstet. Gynecol.* 162, 121-5.

- Thomson, A.M. ; Hytten, F.E. and Billunicz, M.Z. (1967):** The epidemiology of oedema during pregnancy. J. Obstet. Gynecol. Commonwlth 74: 1-10.
- Tikkanen, M.J. (1989) :** intravascular lipoprotein remodeling : lipolytic enzymes. In : Fruchart JC, Shepherd J, eds. Human plasma lipoproteins: Chemistry, Physiology and Pathology. Berlin, New York : De Gryuter. 143 - 51.
- Tobey, T.A. ; Greenfield, m. ; Kraemer, F. and Reaven, G.M. (1981) :** Relationship between insulin resistance; insulin secretion; very low density lipoprotein kinetics and plasma triglyceride levels in normotriglyceridemic man. Metabolism. 30 - 165 - 71.
- Toop, K. and Klopper, A. (1981) :** Concentrations of pregnancy associated plasma protein A (PAPP-A) in aptients with preeclamptic toxemia. Placenta. 3 (Suppl) 167 - 73.
- Trudinger, B.J. ; Giles, W.B. and Cook, C.M. (1985) :** Uteroplacental blood flow velocity-time waveforms in normal and complicated preganacy. Br. J. Obstet. Gynecol. 92: 39 - 45.
- Van den Elzen H.J. ; Juriy, W. ; Wladimiroff ; Titia, E. ; Cohen-over beek ; Anton, J. ; de Bruijn and Diederick, E Grobbee. (1996) :** Serum lipids in early pregnancy and risk of preeclampsia. Briti.J. Obstet.Gynaecol. (103). pp 117 - 122.
- Velentages, P. ; Benga-De, E. and Williams, M.A. (1994) :** Chronic hypertension; pregancy-induced hypertension; and low birth weight. Epideminology. 5: 345 - 8.

- Victor, R. ; Suarez, M.D. ; Juan, G. ; Trelles, M.D. ; Juan M. and Miyahira, M.D. (1996) :** Urinary calcium in asymptomatic primigravides who later developed preeclampsia. *Obstet. Gynecol.* 87: 79 - 82.
- Villar, M.A. ; Sibai, B.M. ; Moretti, M.L.; Mundy, D.C.; Tabb TaN. and Anderson, G.D. (1989b) :** The clinical significance of elevated mean arterial blood pressure in the second trimester and a threshold increase in systolic and diastolic blood pressure during the third trimester *Am. J. Obstet. Gynecol.* 160: 419.
- Walker, J.J. (1987) :** The case for early recognition and intervention in pregnancy induced hypertension. In: Sharp F.; Symonds E.M. ; eds. *Hypertension in pregnancy.* Ithaca; New York: Perinatology. Press 289 - 300.
- Wallenburg, H.C.S. and Rotmans, N. (1983) :** Circulation large platelets and platelet turnover in normotensive and hypertensive pregnancies. *Ain Shams University Press; Cairo;* pp 283 - 288.
- Walsh, S.W. ; Behr, M.J. and Allen, N.H. (1985) :** Placental prostaglandin production in normal and toxemic pateints. *AM. J. Obstet. Gynecol.* 151: 110 - 115.
- Wang, Y. ; Walsh, S.W. ; Guo, J. and Zhang, J. (1991b) :** Maternal levels of prostacyclin; thromboxane; vitamin E. and lipid peroxides- throughout normal pregnancy. *Am. J. Obstet. gynecol.* 165: 1690
- Weenink, G.H. ; Treffers, P.E. ; Vijn, P. ; Smorenberg-Schoorl, M.E. and Tencate, J.W. (1984):** Antithrombin III levels in preeclampsia

- correlate with maternal and fetal morbidity. *Am. J. Obstet. Gynecol.* 148: 1092 - 7.
- Weiner, C.P. and Brandt, J. (1982)** : Plasma antithrombin III activity an aid in the diagnosis of preeclampsia - eclampsia. *Am. J. Obstet. Gynecol.* 142: 275 - 81.
- Weinstein, L. (1982)** : Syndrome of hemolysis; elevated liver enzymes; and low platelet count : a severe consequence of hypertension in pregnancy. *Am. J. Obstet. Gynecol.* 142: 159 - 67.
- Whitby, O.H. ; Deufel, T.; and Paetzke-Brunner, I. (1985)** : Methods of enzymatic analysis. Third edition. VCH verlagsgesellschaft mbH; D- 6940. Weinheim . Vol. VIII. pp 481.
- Willams, G.F. and Jones, D.D. (1982)** : Serum deoxycytidylate deaminase as an index of high-risk pregnancy. *Br. J. Obstet. Gynecol.* 89: 309 - 13.
- Wilson, D.E.; Chan, I. ; Buchik, N. and Horton, S.C. (1985)** : Postchallenge plasma lipoprotein remnants : chylomicron remnants in endogenous hypertriglyceridemia. *Metabolism.* 34: 551.
- Wood, P.L. and Durham, B.H. (1988)** : Change in plasma cystyl aminopeptidase (Oxytocinase) between 30 - 34 w weeks' gestation as a predictor of pregnancy-induced hypertension. *Obstet. Gynecol.* 72: 850 - 2.
- Yen, S.S.C. (1973)** : Endocrine regulation of metabolic homeostasis during pregnancy. *Clin. Obstet. Gynecol.* 16: 130 - 47.

- Ylikorkala, O. ; Perkonen, F. and Viinkka, L. (1986) :** Renal prostacyclin and thromboxane in normotensive and preeclamptic women and their infants. *J. Clin. Endocrinol. Metab.* 63: 1307 - 12.
- Zaltnik, F.J. and Burmeister, L.F. (1983) :** *Dietary protein and preeclampsia. Am. J. Obstet. Gynecol.* 147: 345.
- Zavaroni, I.; Dall'Aglio, E. and Alpio (1985):** Evidence for an independent relationship between plasma insulin and concentration of high density lipoprotein cholesterol and triglyceride. *Atherosclerosis* 55: 259.
- Zavaroni, I.; Mazza, S. ; Dall'Aglio, E. (1992):** Prevalence of hyperinsulinaemia in patients with high blood pressure. *J. Intern. Med.* 231-235.
- Zavaroni, I.; Mazza, S. and Fantuzzi, M. (1993):** Changes in insulin and lipid metabolism in males with symptomatic hyperuricaemia. *J. Intern. Med.* 234: 24.
- Zemel, M. B. ; Zemel, P.C. ; Berrys ; Norman, G. (1990):** Altered platelet calcium metabolism as an early predictor of increased peripheral vascular resistance and preeclampsia in urban black women. *N. Engl. J. Med.* 323: 434-8.
- Zilva, J.F. and Pannall, P.R. (1984) :** "Clinical Chemistry in Diagnosis and treatment" . 4th ed. LLOYD-Luke (Medical Books) LTD, London. Ch. X. pp 232.