

REFERENCES

A

Abi-Samra, F. and Tarzi, R.C. (1983):

Determinations of left ventricular hypertrophy and function in hypertensive patients.

Am. J. Med., 75 (Suppl. 34): 26-33.

Adams, M.A.; Bobik, A. and Korner, P.I. (1989):

Differential development of vascular and cardiac hypertrophy in genetic hypertension.

Hypertension, 14: 191-202.

Adams, T.D.; Yanowitz, F.G. and Fisher, A.G. (1985):

Heritability of cardiac size: an echocardiographic and electrocardiographic study of monozygotic and dizygotic twins. Circulation, 71: 39-44.

Albert, B. and Davids, S. (1997):

Twenty-four hour ambulatory blood pressure monitoring: now that technology had come of age-we need to catch up. J. Pediatrics, 130: 167-9.

Albert, P. and Rocchini, M.D. (1993):

Adolescent obesity and hypertension. Pediatrics clinics of North America, V. 40, No. (1).

Albert, W. Pruitt (1996):

Nelson text book of pediatrics - 15th edition by Richard E. Behrman, Robert M. Kliegman, Ann. M. Arvin., Senior editor, Waldo E. Nelson. P. 404-415.

Albert, W.; Pruitt, Behrman, R.E. and Vaughan, V.C. (1992):

Nelson text book of pediatrics - 14th edition by WB saunders company, philadelphia. P. 345-360.

Alli, C.; Di-Twllio, M.; Mariotti, G.; Taioli, E. and Bellic and Radice, M. (1992):

Effect of long term treatment with prazosin on left ventricular diastolic function in mild to moderate hypertension.

Chest 101 (1): 181-186.

Anderson, N.B. (1989):

Racial differences in stress induced cardiovascular reactivity and hypertension: current status and substantive issues. Psychol. Bull., 105: 89-105.

Anversa, P.; Ricci, R. and Olivetti, G. (1986):

Quantitative structural analysis of myocardium during physiologic growth and induced cardiac hypertrophy. A review. J. Am. Coll. Cardiol., 7: 1140.

Ashizawa, N.; Seto, S.; Kitano, K.; Toyama, K. and Sasaki, H. (1989):

Effect of blood pressure changes on development and regression of electrocardiographic left ventricular hypertrophy. A 26 year longitudinal study. J. Am. Coll. Cardiol., 13: 165-72.

Astrand, I. and Rodahl, K. (1986):

Textbook of work physiology: physiological bases of exercise. McCraw-Hill, Inc: New York.

Astrand, P.O. and Saltin, B. (1969):

Plasma and red cell volume after prolonged exercise. J. Appl. Physiol., 19: 819-823.

Aviv, A. and Gardner, J. (1989):

Racial differences in ion regulation and their possible links to hypertension in blacks. Hypertension, 14: 584.

B

Balady, G.J.; Weiner, D.A.; Rose, L. and Ryan, T.J. (1990):

Physiological responses to arm ergometry exercise relative to age and gender. J. Am. Coll. Cardiol., 16: 130.

Barlow, R.J.; Connel, M.A. and Mil. F.J. (1986):

A study of 48-hour faecal and urinary electrolyte excretion in normotensive black and white south African males. J. Hypertens. 401: 197.

Benbassat, J. and Froom, P. (1986):

Blood pressure response to exercise predictor of hypertension. Arch. Intern. Med., 146: 2053-2055.

Blumenthal, S. (1987):

Report of the task force on blood pressure control in children.
Pediatric 59 (Suppl.): 797. Copy right American Academy of
Pediatric, 1987.

Bohm, M.; Mende, V.; Schmitz, W. and Scholz, (1986):

Cardiac alpha receptors and cardiac hypertrophy in genetic
predisposition to hypertension.
Am. Heart, J., 112: 1347-1349.

Braunwald, E. (1992):

Clinical aspects of heart failure Braunwald E.; A textbook of
cardiovascular medicine A. HBJ. I., 3rd Edition W.B.
Saunders, 446.

Breslin, D.J.; Cifford, R.W.Jr. and Bairbrawn, J.F. (1986):

Essential hypertension: a twenty year follow-up study.
Circulation, 33: 87-93.

Bruno, E.L. (1988):

Clinical aspects of endocrine hypertension.
Med. Clin. North, 71: 907.

***Brutsaert, D.L.; Rademakers, F.E.; Sys, S.V.; Gillbert, T.C. and
Housmans, P.R. (1985):***

Analysis of relaxation in the evaluation of ventricular function
of the heart.
Prog. Cardiovasc. Dis., 28: 142 (Abstract).

***Burke, G.L.; Arcilla, R.A.; Cullpepper, W.S.; Webber, L.S.; Chiangy,
Berenson, G.S. (1997):***

Blood pressure and echocardiographic measurements in
children: the Bogalusa heart study.
Circulation, 75: 106-114.

C

Cahill, N.; Sodt, P. and Lester, L. (1987):

Cardiac echo-angio. Cor-relation in children (abstr.).
Am. J. Cardiol., 37: 125.

- Camble, W.H.; Shaver, J.A.; Alvares, R.F.; Salerni, R. and Rddy, P.S. (1993):***
A critical appraisal of diastolic time intervals as a measure of relaxation in left ventricular hypertrophy.
Circulation, 68: 76-87.
- Casale, P.N.; Devereux, R.B. and Milner, M. (1990):***
Value of echocardiographic measurement of left ventricular mass in predicting cardiovascular morbidity events in hypertensive men.
Ann. Intern. Med., 105: 173-178.
- Chase, H.P.; Garg, S.K. and Harris, S. (1990):***
High normal blood pressure and early diabetic nephropathy.
Arch. Intern. Med., 150: 639.
- Choong, C.Y.; Abascal, V.M. and Thomas, J.D. (1988):***
Combined influence of ventricular loading and relaxation on transmitral flow velocity profile in dogs measured by doppler echocardiography.
Circulation, 78: 672-683.
- Clarke, W.R.; Schrott, H.G.; Laeverton, P.E.; Connor, W.E. and Laver, R.M. (1988):***
Tracking of blood lipids and blood pressure in school age children:
Muscantine study. Circulation, 58: 626-634.
- Craettinger, M.; Alli, C. and Avanzini, F. (1991):***
Left ventricular structure and function in normotensive adolescents with genetic predisposition to hypertension.
Am. Heart J., 111: 115-20.
- Criqui, M.H.; Haskell, W.L.; Heiss, G.; Tyroler, H.A.; Green, P. and Rubenstein, C.J. (1993):***
Predictors of systolic blood pressures responses to trail mill exercise. The lipid research clinics program prevalence study.
Circulation, 68: 225-233.
- Culpepper, W.S.; Sodi, P.C.; Messerli, F.H.; Ruschhaupt, D.G. and Arcilla, R.A. (1997):***
Cardiac status in juvenile borderline hypertension.
Ann. Intern. Med., 98: 1-7.

Cumming, D.J.; Irving, J.B.; Kerr, F. and Kirby, B.J. (1988):

Static exercise in untreated systemic hypertension.
Br. Med. J., 35: 413-421.

D

Danglond, D.A.; James, C.H. and Daniel, J.M. (1986):

Doppler echocardiographic approaches to ventricular diastolic function.
Echocardiography. A review of cardiovascular ultrasound 3:1,
P. 33-40.

Daniels, S.D.; Meyer, R.A. and Loggie, J.M.H. (1990):

Determinants of cardiac involvement in children and adolescents with essential hypertension. Circulation, 82: 1243-8.

De-Simone, R.; Irac, L.; Perna, B.; Caruso, C.; Guda, M. and Iacono, A. (1993):

Effect of cilazapril therapy on diastolic filling and left ventricular hypertrophy in patients with arterial hypertension. Minerva-Cardioangiol., 41(5): 205-209.

De-Swiet, R.; MacMahons, L. and Narcisse, G. (1989):

Blood pressure, stroke and coronary heart disease. Part 2, short-term reductions in blood pressure: Overview of randomized drug trials in their epidemiological context. Lancet, 335: 827.

Devereux, R.B.; Phillips, M.C.; Casale, P.N.; Eisenberg, R.K. and Kligfied, P. (1983):

Geometric determinants of electrocardiographic left ventricular hypertrophy.
Circulation, 67: 907-11.

Devereux, R.B. and Reichek, N. (1984):

Reploration abnormalities of LVH, clinical, echocardiographic and hemodynamic correlates.
Br. Heart J., 52: 516-23.

Flamm, S.D.; Taki, J. and Moore, R. (1990):

Redistribution of regional and organ blood volume and effect of cardiac function in relation to upright exercise intensity in healthy human subjects.
Circulation, 81: 1550.

Fouad, F.M. (1987):

Left ventricular diastolic finding in hypertensive patient
Circulation 75 (Suppl. I). 1-48.

Fouad, F.M.; Tarazi, R.C. and Galagher, J.H. (1980):

Abnormal left ventricular relaxation in hypertensive aptients.
Clin. Sci., 59: 4115.

Freedman, D.S.; Newman, W.P. and Tracy, R.E. (1988):

Black and white differences in aortic fatty streaks in adolescence and early adulthood: The Bogalusa heart study.
Circulation, 77: 856-864.

Frohlich, E.D.; Kozal, V.J. and Torazi, R.C. (1983):

Physiological comparison of labile and essintial hypertensive.
Circ. Res., 46: Suppl. 1: 55-69.

Froelicher, V.F. (1987):

Exercise and the heart clinical concepts. 2nd ed. Chicago, yearbook medical publishers, Inc., 1987.

Froelicher, V.F. and Dlarcordes, G.D. (1989):

Manual of exercise testing. Chicago, yearbook of medical publishus, Inc., 1989.

G

Gerard, P.; Aurgimma, M.D.; Pamelan, Douglas, M.D.; William, H. and Gaasch, M.D. (1997):

Quantitative evaluation of left ventricular structure, wall stress, and systolic function. The practical of clinical echocardiography (Otto) 2nd ed., P. 1-24.

Gibson, D.G. and Brown, D. (1973):

Measurement of instantaneous left ventricular dimension and filling rate in man using echocardiography.
Br. Heart J., 35: 114.

- Gibson, D.G. and Brown, D.J. (1975):**
Measurement of peak rates of left ventricular wall movement in man: Comparison of echocardiography with angiography. *Br. Heart J.*, 37: 677.
- Godfrey, S.; Davies, C.T.M.; Wozinak, E. and Bornes, C.A. (1991):**
Cardiorespiratory response to exercise in normal children. *Clin. Sci.*, 40: 419-431.
- Gottdiener, J.S.; Brown, J.; Zoltick, J. and Fletcher, R.D. (1990):**
Left ventricular hypertrophy in men with normal blood pressure: relation to exaggerated blood pressure response to exercise. *Ann. Intern. Med.*, 112: 161-6.
- Graettinger, W.F.; Neutel, J.M.; Smith, D.H.G. and Weber, M.A. (1991):**
Left ventricular diastolic filling alternations in normotensive young adults with a family history of systemic hypertension. *Am. j. Cardiol.*, 68: 51-6.
- Granger, C.B.; Karimeddini, H.K. and mith, Y.E. (1985):**
Rapid ventricular hilling in left ventricular hypertrophy: physiologic hypertrophy. *Am. J. Cardiol.*, 5: 862-868.
- Graudi, A.; Poletti, L.; Tettamanti, F. and Solda, P. (1989):**
The left ventricle in normotensive young adult with hypertensive parents. *Am. Heart J.*, 10 (Suppl) 189.
- Grossman, W. (1980):**
Cardiac hypertrophy: useful adaptation for pathological process. *Am. J. Med.*, 69: 576-584.
- Grossman, W. (1985):**
Wall stress and patterns of hypertrophy in the human left ventricle. *J. Clin. Invest.*, 56: 56.
- Grossman, W. and McLaurin, L.P. (1975):**
Diastolic properties of the left ventricle. *Ann. Intern. Med.*, 84: 316.

Gunther, S. and Grossman, W. (1979):

Determination of ventricular function in pressure overload hypertrophy in man. *Circulation*, 59: 679.

Guntheroth, W.G. (1965):

Pediatric electrocardiography, Philadelphia W.B.; Saunders Coated from Guntheroth W.G. and Park, M.K.; How to read pediatric ECG, year book, medical: publishers, Inc., Chicago, London, 2nd edition, 560.

Guyton, A.C. (1986):

Textbook of medical physiology.
7th ed. Philadelphia, W.B. Saunders Company, 376.

Guzzetta, P.C.; Davis, C.F. and Ruley, E.J. (1991):

Experience with bilateral renal artery stenosis as a cause of hypertension in childhood. *J. Pediatric. Surg.*, 26: 532.

H

Hanrath, P.; Mathy, D.C. and Kremer, P. (1980):

Effect of verapamil on left ventricular isovolumic relaxation time and a regional left ventricular filling in hypertrophic cardiomyopathy. *Am. J. Cardiol.*, 45: 1258.

Harizi, R.C.; Bianco, J.A. and Alpert, J.S. (1988):

Diastolic function of the heart in clinical cardiology. *Arch intern. Med.*, 148: 99-109.

Harshfield, G.A.; Pulliam, D.A.; Somes, G.W. and Alpert, B.S. (1993):

Ambulatory blood pressure pattern in Youth Am. J. Hypertens., 6: 968-73.

Hartford, M.; Wikstrand, J.; Wallentin, I.; Jungman, S.; Wilhensen, L. and Braglund, G. (1984):

Diastolic function of the heart in untreated primary hypertension. *Hypertension*, 6: 329-338.

Hayashida, W.; Kumada, T. and Nohara, R. (1990):

Left ventricular regional wall stress in dilated cardiomyopathy. *Circulation*, 82:2075.

- Henry, W.L.; Ware, J.; Gardin, J.M.; Hepner, S.I.; McKay, J. and Weiner, M. (1987):**
 Echocardiographic measurements in normal subjects: Growth related changes that occur between infancy and early adult hood.
 Circulation, 57: 278-285.
- Himmelmann, A. (1994):**
 Blood pressure and left ventricular mass in children and adolescents. Blood Press. Suppl. 1994, 3:1-46.
- Hoenecke, H.R.; Goldberg, S.J.; Shan, D.J.; Allen, H.D. and Valdes-Cruz, L.M. (1982):**
 Effect of beam directional alterations on left ventricular shortening fraction.
 Am. J. Cardiol., 50: 1120.
- Hohn, A.R.; Riopel, D.A.; Keil, J.E.; Loadholt, C.B.; Margolius, H.S.; Halushka, P.V.; Privitera, P.J.; Webb, J.G.; Medley, E.S.; Schuman, S.H.; Rubin, M.I.; Pantell, R.H. and Braunstein, M.L. (1994):**
 Childhood familial and racial differences in physiologic and biochemical factors related to hypertension. Hypertension, 5: 56-70.
- Hoof, I.M.S. van (1993):**
 Alteration in calcium metabolism in young people at risk of hypertension; the Dutch hypertension and offspring study. Hypertension, 21: 267-272.
- Hoof, I.M.S.; Grobbee, D.E.; van-Manning, H.J. and Hofman, A. (1997):**
 Hemodynamic characteristics of the early phase of primary hypertension. Circulation, 87: 1006-1012.
- Horan, M.J.; Falkner, B. and Kin, S.Y. (1987):**
 Report of the second task force on blood pressure control in children. Pediatrics, 79 (1): 1-25.
- Horshfield, G.A.; Pulliam, D.A. and Splert, B.S. (1991):**
 Patterns of sodium excretion during sympathetic nervous system arousal. Hypertension, 17: 1156-1160.

Houtman, P.N. (1994):

The physiology and treatment of hypertension. Current pediatric textbook, 4: 77-82.

I

Inouye, I.K.; Massie, B.H.; Loge, D. and Simpson, P. (1984):

Failure of antihypertensive therapy with diuretic, beta blocking, and calcium channel blocking drug to consistently reverse left ventricular filling abnormalities. Am. J. Cardiol., 53: 1583-1587.

Inouye, I.K.; Massie, B.H.; Loge, D. and Topic, N. (1994):

Abnormal left ventricular filling, an early finding in mild to moderate systemic hypertension. Am. J. Cardiol., 53: 120-126(a).

J

Jennings, G.L.; Deakin, G. and Dewar, E. (1989):

Exercise cardiovascular disease and blood pressure. Clin. Exp. Hypertens. A 11: 1035.

Jonas, D.; Del Rosario, M.D.; Frank, A.; Treiber, Ph.D.; Gregory, A.; Harashfield, Ph.D.; Harry, S.; Davis, M.S. and William, B.; Strng, M.D. (1998):

Predictors of future ambulatory blood pressure in youth. J. Pediatric, 132: 693-8.

Jones, M.L. (1988):

Hemodynamic studies in patients with borderline pressure elevation. Circulation, 38: 282-288.

Julien, I.F. Hoffman (1996):

Systemic hypertension.
Rudolph's pediatrics, 20th edition.
Appleton & lange Asimon of Sccluster company. P: 1543-1551.

Julins, S.H.; Krause, L. and Bnda, A.J. (1989):

Neurogenic pressor episods to cause hypertension but do not induce cardiac hypertrophy.
Hypertension, 13: 422-29.

Julius, S. (1991):

Autonomic nervous system dysregulation in human hypertension. *Am. J. Cardiol.*, 67: 3B.

Julius, S. and Schork, M.A. (1996):

Predictors of hypertension. In: Perry H.U.Jr., Smith W.N.; eds. *Mild hypertension: to treat or not to treat*. New York Academy of Sciences, 38: 601-9.

K

Kannel, W.B. (1984):

Role of blood pressure in cardiovascular morbidity and mortality. *Prog. Cardiovascular Dis.*, 17:5.

Kannel, W.B.; Castelli, W.P. and McNomara, P.M. (1989):

Role of blood pressure in the development of congestive heart failure. The franingham study *N. Engl. J. Med.*, 287: 781-787.

Kapukw, G.K.; Setos, Moli, H. and Mori, M. (1993):

Impaired left ventricular filling in borderline hypertensive patients without cardiac structural changes. *Am. Heart J.*, 125 (6): 1710-1716.

Keren, G.; Meisner, J.S. and Mori, M. (1986):

Interrelationship of mid diastolic mitral valve motion, pulmonary venous flow, and transmitral flow. *Circulation*, 74: 36-44.

Klein, J.M. (1987):

Simultaneous direct and indirect blood pressure measurements in man at rest and during work. *J. Appl. Physiol.*, 6: 506-507.

L

Lagadie-Grossman, D.; Chesnais, J.M. and Fewray, D. (1988):

Intracellular pH regulation in capillary muscle cells from streptozotocin diabetic rats: Anion-sensitive microelectrode study. *Pflugers Arch.*, 412: 613-617.

References

- Larry, T.; Mahoney, Richard, M.; Schieken, William, P.; Clarke, and Ronald, M. Lower (1998):*
Left ventricular mass and exercise responses predict future blood pressure. (The Muscative study). Hypertension, 12: 206-213.
- Lieberman, M.; Chopin, L.F. and Carter, E. (1990):*
Factors related to blood pressure in a biracial adolescent female population. Hypertension, 8: 843-850.
- Light, K.C.; Koepke, J.P.; Obrist, P.A. and Willis, P.W. (1983):*
Psychological stress induces sodium and fluid retention in men at high risk for hypertension. Science, 220: 429.
- Littenberg, B.; Garber, A.M. and Sox, H.C. Jr. (1990):*
Screening for hypertension. Ann. Intern. Med., 112:192.
- Little, R.C. (1979):*
The mechanism of closure of mitral valve: A continuing controversy. Circulation, 59: 615-618.
- Little and Downes, T.R. (1990):*
Clinical evaluation of LV diastolic performance. Prog. Cardiovasc. Dis., 32 (4): 273-290.
- Lund, Johanson, P. (1986):*
Hemodynamic patterns in natural history or borderline hypertension.
J. Cardiovascular Pharmacol., 8 (suppl. 5): S8-S 14.
- Lund-Johansen, P. (1995):*
Hemodynamics in essential hypertension. Clin. Sci., 59: 343s-354s.
- Lutas, E.M.; Devereux, R.B.; Reis, G. and Alderman (1985):*
Increased cardiac performance in mild essential hypertension. Left ventricular mechanisms. Hypertension, 7: 979-988.

Macor, F.; Bongiovi, S. and Martina, S. (1991):

Left ventricular filling studied by echo-Doppler in hypertensive and normotensive subjects. G. Ital Cardiol., 21 (2): 151-161.

Mahoney, L.T.; Lauer, R.M. and Lee, J. (1988):

Factors affecting tracking of coronary heart disease risk factors in children: the muscatine study. Ann. NY Acad. Sci., 623: 129.

Marban, E. and Kortesune, Y. (1990):

Cell calcium, oncogenes and hypertrophy. Hypertension, 15: 652-658.

Marchandise, B. Schroder, E. and Bosl, A. (1989):

Early detection of doxorubicin cardiotoxicity. Am. Heart J., 11: 892-898.

Marco, E.J.; Cianciulli, T. and Espinosa, J.H. (1993):

Ambulatory blood pressure monitoring and diastolic left ventricular abnormalities in established and borderline hypertension. Medicine, B-Aires, 53 (3), 202-206.

McCloskey, D.I. and Streatfield, K.A. (1975):

Muscular reflex stimuli to the cardiovascular system during isometric contraction of muscle groups of different mass. J. Physiol., 250: 431-441.

McCullagh, W.H.; Covell, J.W. and Ross, J.Jr. (1972):

Left ventricular dilatation and diastolic compliance changes during chronic volume overloading. Circulation, 45: 943.

McIlhany, M.L.; Schaffer, J.W. and Hines, E.A.Jr. (1985):

The heritability of blood pressure: an investigations of 200 pairs of twins using the cold pressor test. Johns Hopkins Medical Journal, 136: 57.

Messerli, F.H.; Sundergard, R.K.; Reisen, E.D. and Dreslinski, G.R. (1983):

Dimorphic cardiac adaptation to obesity and arterial hypertension. Ann. Intern. Med., 99: 955-761.

Messerli, F.H.; Aristizabal, D. and Soria, F. (1993):

Reduction of left ventricular hypertrophy: How beneficial?
Am. Heart J., 125: 1520-40.

Mikel, D. and Smith, M.D. (1997):

Left ventricular diastolic function-clinical utility of Doppler echocardiography. The Practical of clinical echocardiography (Otto) 2nd ed., P. 49-74.

Molineux, D. and Steptoe, A. (1997):

Exaggerated blood pressure responses to submaximal exercise in normotensive adolescents with a family history of hypertension. J. Hypertens., 6: 361-5.

Morgan, H.E. and Baker, K.M. (1991):

Cardiac hypertrophy. Mechanical, neural and endocrine dependence. Circulation, 83: 13.

Mulvagh, S.; Quinones, M.A. and Kleiman, N.S. (1992):

Estimation of left ventricular end-diastolic pressure from Doppler transmitral flow velocity in cardiac patients independent of systolic performance.
J. Am. Coll. Cardiol., 20: 112-119.

N

Nada, N.C. (1993):

Doppler echocardiography.
2nd edition, Lea and Febiger, Philadelphia, London P. 197-215.

Naggar, C.Z.; Stemley, K. and Downing, L.L. (1982):

Effects of amyle nitrate on left ventricular relaxation patients with borderline hypertensive. Am. J. Cardiol., 50: 979-984.

Neutel, J.M.; Smith, D.H.G.; Graettinger, W.F.; Winer, R.L. and Weber, M.A. (1993):

Metabolic characteristics of hypertension: importance of positive family history.
Am. Heart J. 126: 924-9.

Niederle, P.; Widimsky, J.; Jandova, R.; Ressler, J. and Grosopic, A. (1982):

Echocardiographic assessment of the left ventricle in juvenile hypertension. *Int. J. Cardiol.*, 2: 91-101.

Nikolic, S.; Yellin, E.L. and Tamura, K. (1988):

Passive properties of left ventricle: Diastolic stiffness and restoring forces: *Circ. Res.*, 62: 1210-1222.

O

Orourke, M. (1990):

Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension*, 15: 339.

P

Park, M.K. (1996):

Textbook of pediatric cardiology. 3rd edition St. Louis, Mosby-Year book. Hypertension.

Pearson, A.G.; Arthur, J. and Denisem Oseky, L. (1987):

Assessment of diastolic function in normal and hypertrophied hearts. Comparison of Doppler echocardiography and M-mode echocardiography. *Am. Heart J.* 11 (3): 1417.

Pearson, A.C.; Pasierski, T. and Labovitz, A.J. (1991):

Left ventricular hypertrophy: Diagnosis, prognosis and management. *Am. H. J.*, 121: 148.

Pela, I.; Modesti, P.A.; Cocchic, Cecioni, I.; Gensini, G.F. and Bartolozzi, G. (1990):

Changes in the ampulatory arterial pressure of normotensive obese children. *Pediatric Med. Chir.*, 12: 495-7.

Pickering, S.; Davies, C.T.M.; Wozniak, E. and Barnes, C.A. (1990):

Cardiorespiratory response to exercise in normal children. *Clin. Sci.*, 40: 419-431.

Port, S. and Cobb, F.R. (1980):

Effect of age on the response of the left ventricular ejection fraction to exercise. *N. Engl. J. Med.*, 303: 1133.

R

Rahimootalla, S.H.; Ehasni, A. and Sinno, M.Z. (1985):

Left atrial transport function in myocardial infarction importance of it's booster function. Am. J. Med., 59: 686-594.

Reaven, G.M. (1991):

Insulin resistance, hyperinsulinemia, and hypertriglyceridemia in the etiology and clinical course of hypertension. Am. J. Med., 90 (2A): 7S-12S.

Rebeck, T.R.; Turner, S.T. and Sing, C.E. (1996):

Probability of having hypertension: effects of sex, history of hypertension in parents, and other risk factors. J. Clin. Epidemiol., 1996 Jul., 49 (7): 727-34.

Reicheck, N. and Devereux, R.B. (1981):

Left ventricular hypertrophy: relationship of anatomic, echocardiographic and electrocardiographic findings. Circulation, 63: 1391-1404.

Ren, J.; Hakki, A.; Kotler, M.N. and Iskandrian, A.S. (1985):

Exercise systolic blood pressure a powerful determinant of increased left ventricular mass in patients with hypertension. J. Am. Coll. Cardiol., 5: 1224-1231.

Ren, J.; Oancholy, S.B.; Iskandrian, A.S.; Lightly, G.W.; Mallavarapu, C. and Segal, B.L. (1994):

Doppler echocardiographic evaluation of the spectrum of left ventricular diastolic dysfunction in essential hypertension. Am. Heart J., 127: 906-13.

Richard, M. and Schieken, M.D. (1993):

Genetic factors that predispose the child to develop hypertension. Pediatric clinic of North America; volume, 40. No. 1.

Rocchini, A.P. (1989):

Childhood hypertension: Etiology, diagnosis and treatment Pediatric clin. North Am. 31 (6): 1259-73.

Rokey, R.; Kno, L.C.; Zoghbi, W.A.; Limacher, M.C. and Quinones, M.A. (1985):
Determination of parameters of left ventricular filling with pulsed Doppler echocardiography. Comparison with cineangiography. Circulation, 71: 543-550.

S

Safar, A.E.; Lehner, J.P.; Wincent, M.F.; Palinfosse, M.T. and Simon, A.C.H. (1989):
Echocardiographic dimensions in borderline and sustained hypertension. Am. J. Cardiol., 59: 623-632.

Sahn, D.J.; De Maria, A.; Kisslo, J. and Heyman, A. (1988):
Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiography measurements. Circulation, 48: 1072-1079.

Sanad, O.A.; Ramzy, A.; Taher, M.A. and Abo-El-Enein, H.M. (1995):
Right ventricular diastolic performance in essential hypertension. Benha Med. J., Vol. 11, P. 73-81.

Santamore, W.P. and Little, W.C. (1986):
Left ventricular diastolic performance. Cardiology, 3: 43-46.

Sasayama, S.; Ross, J. and Franklin, D. (1976):
Adaptation of the left ventricle to chronic pressure over load. Circu. Res., 38: 172-180.

Savage, D.D.; Drayer, J.M. and Henry, W.L. (1989):
Echocardiographic assessment of cardiac anatomy and function in hypertensive subjects. Circulation, 59: 623-632.

Schieken, R.M.; Clarke, W.R. and Lauer, R.M. (1998):
Left ventricular hypertrophy in children with blood pressures in the upper quintile of the distribution of blood pressure. The Muscative study. Hypertension, 3: 699-675.

- Seguro, C.; Sau, F.; Zedda, N.; Mura, O.; Montaldo, C.; Scano, G.; Arru, A.; Leo, M.R.; Aru, A. and Cherchi, A. (1995):**
Hemodynamic assessment at rest and during dynamic physical exercise in young subjects with and without hypertensive parents. *Cardiologia*, 1995 Jun., 40 (6): 391-7.
- Sen, S.; Tarazi, R.C.; Khairallah, P.a. and Bumpus, F.M. (1994):**
Cardiac hypertrophy in spontaneously hypertensive rates. *Circu. Res.*, 35: 775-781.
- Shapiro, L.M.; Macklinnon, J. and Beavers, D.G. (1981):**
Echocardiographic features of malignant hypertension. *Br. Heart. J.*, 46: 374-379.
- Shar, C.L.; Barke, G.L. and Freedman, D.S. (1986):**
Value of childhood blood pressure measurements and family history in predicting future blood pressure status: Results from 8 years of follow-up in the Boglausa Heart study. *Pediatrics*, 77: 862.
- Silverman, N.H. (1993):**
Quantitative methods to enhance morphological information using M-mode and Doppler, and cross sectional ultrasound in Silverman N.H. (ed): *Pediatric Echocardiography*. Baltimore, Williams and Wilkeins, P. 35-108.
- Sinha, M.K.; Buchanan, C. and Leggett, N. (1989):**
Mechanism of IGF-I-stimulated glucose transport in human adipocytes: Demonstration of specific IGF-I receptors not involved in stimulation of glucose transport. *Diabetes*, 38: 1217-1225.
- Smith, V.E.; White, W.B.; Mecran, M.K. and Karimeddin, M.K. (1986):**
Improved left ventricular filling accompanies reduced left ventricular mass during therapy of essential hypertension. *J. Am. Cardiol.*, 81: 1449-1454.
- Snider, A.R.; Gidding, S.S. and Rocchini, A.P. (1996):**
Doppler evaluation of left ventricular diastolic filling in children with systemic hypertension. *Am. J. Cardiol.*, 56: 921-926.

Sodi-Pallares, D. (1958):

Electrocardiography in infancy and childhood. Ped. Clin. North Amer. (Quoted in: Essa et al., 1980) follow up study of electrocardiography patterns from birth till the age of four months. Gas Egypt Ped. Ass., 28: 100.

Soergel, M.; Kirchstein, M.; Busch, C.; Danne, T.; Gallerman, J. and Holl, R. (1997):

Oscillometric twenty-four hour ambulatory BP values in healthy children and adolescents: a multicenter trial including 1141 subjects. J. Pediatric., 130: 178-084.

Soto, L.F.; Kikuchi, D.A. and Arcilla, R.A. (1997):

Echocardiographic functions and blood pressure in children and young adults from a Biracial population: The Bagalusa heart study.

American Journal of Medical Science, 297 (5): 271-279.

Spann, J.F.; Bove, A.A.; Natarajan, G. and Kreulen, T. (1980):

Ventricular performance, pump function and compensatory mechanisms in patients with aortic stenosis. Circulation, 62:576.

Sparrow, H.R.; Burggrof, G.W.; Cuddy, E.; Milliken, J.A. and Warren, J. (1983):

Prediction of left ventricular mass from the electrocardiogram. Am. J. Med., 24: 121-127.

Sudhir, K.; Dennis, M.; Roberta, L. and Anderson, R.N. (1996):

Parental hypertension and cardiac alterations in normotensive children and adolescents.

Am. Heart J., January: 81-87.

Suga, M.; Alli, C. and Avanzini, F. (1988):

Echocardiographic variables in progeny of hypertensive and normotensive parents. Acta Med. Scand., 693: 61-64.

Swinscow, T.D.V. (ed) (1994):

Statistics of square one, 9th edition.

London, Latimer Trend & Company.

T

Takahashi, M.; Sasayama, S. and Kawai, C. (1980):

Contractile performance of the hypertrophoid ventricle in patients with systemic hypertension.
Circulation, 62: 116-126.

The Fourth National Committee (1993):

The fourth report of the Joint National Committee on Detection, Evaluation and Treatment of high blood pressure.
Arch Intern. Med., 153: 154-183.

Treiber, F.A.; Musante, L. and Riley, W. (1989):

The relationship between hostility and blood pressure in children. Behav. Med., 15: 173.

Trevi, G.P.; Gallina, S. and Di-Giovanni (1993):

The clinical and echocardiographic parameters at rest and during mental stress in preadolescent subjects with a hypertensive parents. Cardiologia (Abstract). Jul; 38 (7): 437-44.

U

Urschel, C.W.; Covell, J.W. and Sonnenblick, E.H. (1978):

Myocardial mechanics in aortic and mitral valvular regurgitation: The concept of instantaneous impedance as a determinant of the performance of the intact heart. J. Clin. Invest., 47: 867.

V

Vittolio, de Leonardis, Maurizid, De Scalzi, Alberto Falchetti, Piero Cinelli, Emanuele, Croppi, Ricardo, Livi, Luca Scorpilli and Poolo Tommaso Scarpelli (1992):

Echocardiographic evaluation of children with and without family history of essential hypertension: American J. of Hypert., 1: 305-308.

Wadsworth, R.M. (1990):

Calcium and vascular reactivity in ageing and hypertension. J. Hypertens., 8: 975-983.

Wahr, D.W. and Schiller, N.B. (1983):

Left ventricular volumes determined by two-dimensional echocardiography in normal adult population. J. Am. Coll. Cardiol., 1: 863-868.

Walter, H.J. (1989):

Primary prevention of chronic disease among children. The school-based "known your body" intervention trials. Health Educ. Q., 16:201.

Wasserman, K.; Hansen, J.E. and Whipp, B.J. (1987):

Principles of exercise testing and interpretation. Philadelphia, Lee and Fibiger.

Weber, M.A.; Smith, D.H.; Neutel, J.M. and Greatting, I.G. (1991):

Arterial properties of early hypertension. J. Hum. Hypertens., 5: 417-423.

Whelton, P.K. and Klag, M.J. (1989):

Hypertension as a risk factor for renal disease. Review of clinical and epidemiological evidence. Hypertension., 13 (Suppl. I), : 1.

Widgren, B.R.; Berglund, G.; Wikstrand, J. and Andersson, O.K. (1992):

Reduced venous compliance in normotensive men with positive family histories of hypertension. J. Hypertens., 10: 459-465.

Wikstrand, J.; Berglund, G. and Wilhelmsen, L. (1978):

Value of systolic and diastolic time intervals: studies in normotensive and hypertension 50-years old man and women. Br. Heart Jr., 40: 256-269.

Wilka, N.A.; Sheldahl, L.M. and Levandoski, S.G. (1989):

Weight carrying versus handgrip exercise testing in man with coronary artery disease. Am. J. Cardiol., 64: 736.

Williams, R.R. (1989):

Will gene markers predict hypertension?
Hypertension, 14: 610.

Williams, E.F.; Sonnenblick, E.H. and Bramwald, E. (1989):

Determination of atrial contractile force of intact heart. Am. J. Physiol., 209: 1061-1068.

Williams, R.R.; Hung, S.C. and Hasstedt, S.J. (1990):

Multigenic human hypertension: evidence for subtypes and hop for haplo-types. J. Hypertens., 8 (Suppl. 7): S39.

Clinical data of normotensive children of hypertensive parents.

Case No.	Age (years)	Sex	Family H. of Hypertension	Weight (kg)	Height (cm)	Pulse	S.B.P. (mmHg)	D.B.P. (mmHg)
1	8	F	M	21	135	90/M	120	80
2	10	M	M	24	135	95/M	110	80
3	11	M	M	23	134	85/M	110	80
4	15	M	(M.F)	45	155	85/M	110	80
5	16	M	F	70	165	100/M	120	85
6	14	F	M	35	160	80/M	110	80
7	13	M	M	35	165	87/M	110	80
8	16	M	F	80	160	98/M	120	85
9	13	M	M	30	160	92/M	110	80
10	15	M	(M.F)	42	163	98/M	120	90
11	15	F	M	50	160	101/M	110	80
12	14	F	M	35	135	87/M	120	80
13	10	F	F	23	140	82/M	110	80
14	13	F	M	30	148	93/M	110	80
15	9	F	M	37	140	98/M	110	80
16	8	M	F	25	130	96/M	110	80
17	13	F	M	34	162	90/M	110	80
18	11	M	M	25	137	98/M	120	80
19	11	M	M	30	150	94/M	110	80
20	14	F	M	35	164	85/M	100	80
21	14	M	M	44	153	95/M	120	70
22	13	M	F	37	145	95/M	110	80
23	15	F	M	47	165	98/M	120	80
24	12	M	(M.F)	35	166	96/M	110	80
25	16	F	M	60	168	100/M	130	85
26	14	M	M	40	156	94/M	110	80
27	13	M	F	40	160	100/M	110	80
28	11	F	F	29	140	95/M	100	70
29	15	F	F	48	166	100/M	110	70
30	12	M	M	37	162	95/M	110	80
31	16	M	F	72	173	90/M	120	90
32	14	M	M	45	159	90/M	110	80
33	13	M	M	42	159	96/M	100	80
34	14	F	M	41	155	96/M	100	75
35	12	M	F	31	150	96/M	110	80
36	11	F	M	23	135	90/M	105	80
37	13	F	(M.F.)	32	152	97/M	120	85
38	15	M	F	46	160	85/M	120	85
39	13	M	F	49	162	90/M	120	85
40	12	F	M	33	152	87/M	100	80
41	11	M	M	26	137	88/M	100	80
42	12	M	M	42	150	103/M	120	85
43	15	M	(M.F.)	50	166	100/M	120	85
44	12	M	F	40	151	96/M	120	80
45	10	F	M	26	141	90/M	120	80
46	8	F	(M.F.)	20	136	100/M	110	80
47	13	M	M	37	160	96/M	120	80
48	10	F	M	24	140	89/M	100	80
49	12	F	M	33	156	92/M	100	80
50	11	M	(M.F)	3	140	100/M	120	85
51	16	M	M	7	172	102/M	120	85
52	9	M	M	22	138	80/M	110	80
53	13	F	F	30	159	85/M	100	75
54	14	M	M	33	163	90/M	110	80

Investigations of normotensive children of hypertensive parents.

Case No.	ECHOCARDIOGRAPHIC MEASUREMENTS					EXERCISE TEST			
	Systolic function (E,F)	Diastolic function			LVM gm	Mid exercise		Peak exercise	
		E wave (cm/S ²)	A wave (cm/S ²)	E/A ratio		HR/min.	B.P mmHg	HR. /min.	B.P mmHg
1	65%	110	50.1	2.2	31.4	137	130/80	146	130/85
2	72%	95.9	56.2	1.7	41.9	130	120/90	150	130/90
3	71%	105	60	1.8	40.9	140	120/85	160	130/85
4	69%	104	104.3	0.1	137	120	120/90	140	130/95
5	62%	83.5	85.7	0.1	250	140	130/85	165	150/90
6	71%	114	45.1	3.2	53.8	100	120/80	130	120/85
7	62%	110.5	66.8	1.7	154.9	130	120/85	160	130/90
8	57%	73	73.7	0.1	230	140	150/90	Fatigue	Fatigue
9	67%	73.6	48.4		58.4	120	120/80	140	130/80
10	74%	74.3	84.2		168	120	130/90	160	150/95
11	70%	88.5	73.3	1.2	108	140	120/85	170	130/90
12	80%	98.7	43.8	2.3	64.3	130	120/85	165	130/85
13	72%	116	53.6	2.2	62.4	116	120/80	130	120/85
14	62%	73.6	61.5	1.2	71.2	107	120/80	130	120/85
15	53%	74	45.3	1.6	51.3	130	120/80	150	130/85
16	54%	90	52.3	1.7	64	130	120/80	150	120/85
17	37%	152	55.6	2.7	58.6	120	120/80	150	130/85
18	71%	120.3	60.7	2	113.4	130	120/85	160	130/90
19	63%	106	65.2	1.6	84.2	96	120/80	110	120/85
20	70%	106	60.3	1.8	50.8	110	120/80	140	130/80
21	60%	83.3	59.5	1.4	74	120	120/80	160	150/90
22	74%	100	60.7	1.6	85	130	120/80	162	140/90
23	60%	87.6	63.5	1.4	64.5	120	130/80	167	130/90
24	70%	87	41.7	2	90.6	120	120/80	160	150/90
25	80%	60.3	70.2	0.9	100	150	140/85	170	140/90
26	74%	108	74.2	1	114	130	120/80	160	130/85
27	67%	100.2	66.1	1.5	75	130	120/80	150	120/85
28	74%	108	74.2	1.5	114	110	120/80	150	130/80
29	75%	105.1	66.2	1.6	92.1	160	120/80	180	130/85
30	63%	108.2	73.1	1.5	149.2	120	120/80	160	130/85
31	75%	103	104.1	0.1	244.2	130	130/90	Fatigue	Fatigue
32	80%	112	80.6	1.4	92.3	120	120/85	150	120/85
33	66%	111.1	60.1	1.8	89.3	120	120/85	160	140/85
34	58%	98.7	60.8	1.6	178	130	110/80	160	130/85
35	54%	84.6	68.4	1.2	41.3	125	120/80	144	130/80
36	80%	113	99.8	1.1	105	130	120/85	160	130/90
37	75%	110.2	99.9	1.1	102.5	140	130/90	170	140/95
38	70%	87.2	90.5	0.1	97.5	130	120/90	170	130/90
39	81%	90.1	91.6	0.1	104	140	130/85	160	130/90
40	37%	150	50.1	3	59.2	103	120/80	120	120/85
41	30%	120.1	56.2	2.1	60.3	120	120/80	140	120/85
42	89%	86.2	95	0.9	108.2	150	120/90	170	130/90
43	77%	90.2	50.6	1.8	115.2	130	130/85	160	140/90
44	70%	110	80.2	1.4	120	150	130/85	160	140/90
45	72%	96.1	58.3	1.6	99.1	130	130/80	160	130/95
46	67%	105.2	56.4	1.9	80.9	130	120/85	160	130/90
47	76%	100.1	71.3	1.4	105.4	125	130/85	170	140/90
48	50%	105	43.3	2.4	67.1	120	120/80	135	120/85
49	74%	108	74.2	1.5	59.3	120	110/85	130	120/85
50	80%	100.1	102	0.1	120	140	130/85	180	145/95
51	80%	98.6	82.6	1.2	130	130	130/90	Fatigue	Fatigue
52	38%	140.2	53	2	60.1	100	120/80	140	120/85
53	32%	131.1	57.3	2.3	150.3	105	110/80	130	120/85
54	40%	102	57.3	1.8	140.2	120	120/80	140	120/85