




REFERENCES

REFERENCES

Aggarwal BB; (1985):

Human tumour necrosis factor.
J. Biol. Chem. 260: 2345-48.

Arjan A., Kraayenbrink., et al. (1993):

Endothelial vasoactive mediators in preeclampsia.
Am. J. Obstet Gynecol; 169: 160- 5 .

Atkinson MW., Maher JF., Owen J; (1994):

The predictive value of umbilical artery doppler studies for preeclampsia or fetal growth retardation in preeclampsia.
Obstet & Gynecol; 83: 609 -12.

Axelrod L., Levine L; (1983):

Inhibitory effect of insulin on prostacyclin production by isolated rat adipocytes.
Prostaglandins; 25: 571- 79 .

Axelrod L; (1986)

Prostacyclin production by isolated rat adipocytes: evidence for cyclic adenosine 3',5' monophosphate-dependent and independent mechanisms and for selective effect of insulin.
Endocrinology; 119: 2233- 39 .

Axelrod L; (1991):

Insulin, prostaglandins, and the pathogenesis of hypertension.

Diabetes; 40: 1223- 27.

Baker V.V., Berek J.S. and Maza O.M; (1996):

Cytokines, Lymphokines and immune mediators
In molecular Biology and Genetics.

Novak's Gynecology 12th ed; 6:138-142.

Bauman MR., Maiman M., and Langer O; (1982):

An association between hyperinsulinemia and hypertension during the third trimester of pregnancy.

Am. J. Obstet & Gynecol; 159: 446- 50 .

Belizan JM., Villar J., Gonzalez L., et al; (1991):

Calcium supplementation to prevent hypertensive disorders of pregnancy.

N. Engl. J. Med; 325: 1399-1404.

Bender HS., Chickering WR., et al; (1985):

Pregnancy and diabetes: the maternal response.

Life sci 37: 1-5.

Berntorp K., et al; (1983):

Relation between plasma insulin and blood glucose in a cross- sectional population study of the oral glucose tolerance test.

Acta Endocrinol; 102: 549-56.

Benson SA., and Yalow RS; (1970):

Insulin "antagonists" and insulin resistance in diabetes mellitus. Theory and practice. Ellenberg. M., and Rifkin, H., Eds.

New York. MC Graw- Hill; 388-423.

Beutler B., Melsark I., Wand Ceram A; (1985):

Cachectin (tumour necrosis factor) production, distribution and metabolic fate in vivo.

J. Immunol; 135: 3972- 77 .

Beutler B., Cerami A, et al: (1987):

Cachectin: more than a tumour necrosis factor.

N. Engl, J Med; 316: 379-385.

Bomri A; (1980)

Histological technique for placental examination
5th ed.

Oxford, New York: Oxford University Press, 195.

Bray MA; (1983):

The pharmacology and pathophysiology of leukotriene B₄.

Br. Med. Bull; 39: 249-254.

Brosens I., Robertson WB., Dixan HG; (1967):

The physiological response of the vessels of the placental bed to normal pregnancy.

J. Pathol. Bacteriol; 93: 569 - 73.

Brosens I., Robertson WB., Dixan HG; (1972):

The role of the spiral arteries in the pathogenesis of pre eclampsia.

Obstet & Gynecol. Annu; 1: 177 - 79.

Bry K; (1993):

Epidermal growth factor and transforming growth factor- alpha enhance the interleukin- 1 and tumour necrosis factor- stimulated prostaglandin E₂ production on amnion cells.

Prostaglandins Leukotrienes & Essential fatty acids; 49: 923-8.

Burt RL., and Davidson IWF; (1974):

Insulin half-life and utilization in normal pregnancy.

Obstet & Gynecol; 43: 161-70.

Campbell MJ and Mechin D; (1993):

Medical statistics- A common sense approach- 2nd
ed: John Wiley & Sons. New York; Chichester;
Brisbane;
Toronto & Singapore.

Catalano PM., Tyzbir E. D, Roman NM, et al; (1991):

Longitudinal changes in insulin release and insulin
resistance in non obese pregnant women.
Am J Obstet & Gynecol; 165: 1667-72.

Chajek- Schaul T., Friedman G., et al; (1989):

Mechanism of the hypertriglyceridaemia induced
by TNF administration to rats.
Biochem. Biophys. Acta; 1001: 316- 24 .

Chang J., Gilman SC., and Lewis AJ; (1986):

Interleukin- 1 activates phospholipase A₂ in rabbit
chondrocytes: a possible signal for IL-1 action.
J Immunol; 136: 1283-7.

Chen Y.D., Shew W.H., Swislock A.L.M; (1991):

High density lipoprotein Turnover in patients with
hypertension.
Hypertension; 17- 386 .

Chesley IC; (1971):

Hypertensive disorders in pregnancy.

Williams Obstetrics, 14th edition, New York.

P.P 120- 125 .

Chesley IC; (1974):

A short history of eclampsia.

Obstet & Gynecol; 43 : 599-603.

Chesley LC., and Cooper DW; (1986):

Genetics of hypertension in pregnancy. Possible single gene control of pre eclampsia and eclampsia in the descendants of eclamptic women

Br.J obstet & Gynecol; 93: 898-903.

Clauss M., Ryan J., Stern D; (1992):

Modulation of endothelial cell hemostatic properties by TNF: insights into the role of endothelium in the host response to inflammatory stimuli.

Raven press New York; 49-63.

Cohen S. and Yoshida D; (1979):

Regulation of cytokine function.

A.N.Y.: Acd. Sci; 332 : 356-362.

Comens P., Wolf BA., Lacy PE., et al; (1987):

Interleukin 1 is a potent modulator of insulin secretion from isolated rat islets of Langerhans.

Diabetes; 36: 963-970.

Cooper DW and Liston; (1979):

Genetic control of severe pre eclampsia.

J. Med. Genet. 16: 409 -13.

Cotran RS; (1987):

New roles for the endothelium in inflammation and immunity.

Am J. pathol; 129: 407 -11.

Cousins L; (1987):

Pregnancy complications among diabetic women: Review 1965-1985.

Obstet & Gynecol. Surv; 42: 140-45.

Cunningham FG., et al; (1993):

Hypertensive disorders in pregnancy.

Williams Obstetrics 19th ed; 2: 767 -70.

Davies JA and prentice CRM; (1992):

Coagulation changes in pregnancy induced hypertension and growth retardation. In: Greer IA, Turpie AGG, Forbes CD (eds). Haemostasis and thrombosis in obstetrics and Gynecology.

Chapman & Hall, London, PP 143.

De Fronzo RA., Ferrannini E; (1991):

Insulin resistance: a multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia and atherosclerotic cardiovascular disease.

Diabetes care; 14: 174-94.

De Graaf J.,Har.Lemmers HL., Hoctars MPC et al.,(1991):

Enhanced susceptibility to in- vitro oxidation of the dense low- density lipoprotein subfraction in healthy subjects.

Arterioscler. Thromb; 11: 298- 306 .

Dakker G., Morris N. and Eaton BM (1996):

Nitric oxide, the endothelium, pregnancy and pre eclampsia.

Br. J. Obstet & Gynecol; 103 : 4-15

De Wolf F., De wolf., pecters C., Brosen SI; (1980):

The human placental bed: electron microscopic study of trophoblastic invasion of spiral arteries.

Am J. Obstet Gynecol; 137: 58-70 .

DiGiovine FS., and Duff GW; (1990):

Interleukin 1: The first interleukin.

Immunol today; 11: 13 -15 .

Douglas R., Baha M., et al; (1989):

Ultrastructural aspects of pre eclampsia.

Am J Obstet Gynecol; 161: 735-41.

Dusting GJ., Moncada S., Vane JR; (1982):

Prostacyclin its biosynthesis, actions, and clinical

potential in prostaglandins and the cardiovascular system.

Oates JA, Ed. New York, Raven: 59 .

Faure A., Sutter- Dub MT., et al; (1983):

Ovarian- adrenal interactions in regulation of endocrine pancreatic function in the rat.

Diabetologia; 24: 122 -27.

Fay RA., Hughes AO., Farron; (1983):

Platelets in pregnancy: hyperdestruction in pregnancy.

Obstet & Gynecol; 61: 238-240.

Feingold K.R., Serio M.K., Adis, Moser AH., and Grunfeld C; (1989):

TNF stimulates hepatic lipid synthesis and Secretion.

Endocrinol; 124: 2336- 42 .

Ferrannini E., Buzzigol G., Bonadonna R; (1987):

Insulin resistance in essential hypertension.
N. Engl. J. Med.; 317: 350-7.

Ferrannini E., Haffner S.M., Mitchell B.D. et al; (1991):

Hyperinsulinemia: The key feature of a
cardiovascular and metabolic syndrome.
Diabetologia; 34: 416- 22 .

Ferris TF; (1991):

Pregnancy, pre eclampsia and the endothelial cell..
N. Engl. J. Med; 325: 1439 - 43.

Fiers W; (1991):

Tumour necrosis factor, characterization at the
molecular, cellular and in vivo level.
FEBS; 285: 199-212.

Fox H., (1990):

Placental structure of health and disease; In
modern Antenatal care of the fetus; Geoffrey
chamberlain, Blackwell.
Scientific publications, Oxford, London, P. 52 .

Freinkel N., Metzger BE; (1979)

The critical implication of maternal metabolism for
fetal development.
Excerpta Medica; 3: 28-33.

Freinkel N., Dooley SL., Metzger BE; (1985):

Care of the pregnant women with insulin dependent diabetes mellitus.

N. Engl. J. Med; 313: 96-99.

Friedman S.A; Taylor R.N. and Roberts JM; (1991):

Pathophysiology of pre eclampsia.

Clin perinatal; 18: 661- 82 .

Friedmen S.A., Lubarsky SL., Ahokas RA, et al; (1995):

Pre eclampsia and related disorders. Clinical aspects and relevance of endothelin and nitric oxide.

Clinics in Perinatology; 22: 343-55.

Frolich M.A., Datta S., and Corn S.B; (1998):

Thrombopoietin in normal pregnancy and pre eclampsia.

Am. J. Obstet & Gynecol; 179: 100- 4 .

Furchgott RF., Zawadski; (1980):

The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholin.

Nature; 288: 373- 376 .

Gerretsen G., and Elema J.D; (1981):

Morphological changes of spiral arteries in the placental bed in relation to pre eclampsia and fetal growth retardation.

Br.J. Obstet & Gynecol; 88: 876-881.

Giles WB., truding B.J., Baird PJ; (1985):

Fetal umbilical artery flow velocity waveforms and placental resistance: pathological correlation.

Br. J. Obstet & Gynecol; 92: 31-35.

Gill S., Vince., Phyllis M., et al; (1995):

Interleukin-6, tumour necrosis factor and soluble tumour necrosis factor receptors in women with pre eclampsia.

Br. J. Obstet & Gynecol; 102: 20-25.

Giroir BP., Peppel K., Silva M., et al; (1992):

The biosynthesis of tumour necrosis factor during pregnancy.

European Cytokine Network; 3: 533-8.

Gomori G; (1980):

Rapidone step trichrome stain. 5th ed. Oxford, New York.

Oxford, University press; 185: 189-92.

Good GK., Miller J.P., and Heugerty A.M; (1995):

Hyperlipidaemia, hypertension and coronary heart disease.

Lancet; 345: 362- 4.

Goodman RP., Killam AP., Brush AR., et al; (1982):

Prostacyclin production during pregnancy:
Comparison of production during normal
pregnancy and pregnancy complicated by
hypertension.

Am. J. Obstet & Gynecol; 142: 817 – 23.

Granner DK; (1996)

Hormones of the pancreas and gastrointestinal
tract.

Harper's Biochemistry, 24th ed; 51: 581-91.

Grau GE; (1987)

Tumour necrosis factor (cachectin) as an essential
mediator in murine cerebral Malaria.

Science, 237: 1210 –14.

Greer L.A., Haddad NG., et al; (1989)

Neutrophil activation in pregnancy induced
hypertension.

Br. J. Obstet & Gynecol; 96 : 978-82.

Greer MD., Flona Ph D., et al; (1994)

Increased concentration of cytokines interleukin-1 receptor antagonist in plasma of women with pre eclampsia.

Obstet & Gynecol; 84: 937-40.

Haeger M., Unander M., and Bengtsson A; (1991):

Complement activation in relation to development of pre eclampsia.

Obstet & Gynecol; 78: 46-9.

Haeger M., Unander M., and Norder- Hanson B; (1992):

Complement, neutrophil and macrophage activation in women with severe pre eclampsia, and HELLP syndrome.

Obstet & Gynecol; 79: 19-26.

Halligan A., Bonnar J., Sheppard B; (1994):

Haemostatic, fibrinolytic and endothelial variables in normal pregnancies and pre eclampsia.

Br. J. Obstet & Gynecol; 101: 488-492.

Harlan JD; (1987):

Neutrophil- mediated vascular injury.

Acta Med Scand Suppl.; 715: 123-129.

Houwert- de- Jong MH., claas FH., et al; (1985):

Humoral, immunity in normal and complicated pregnancy.

Eur J. Obstet & Gynecol Reprod Biol; 19:205-14.

Hunt JS; (1993):

Expression and regulation of the tumour necrosis factor- alpha gene in the female reproductive tract.

Reproduction, Fertility&Development; 5: 141-53.

Janoff A; (1985):

Elastase in tissue injury.

Ann Rev Med; 36: 207-216.

Jovanovic L., Druzin M., and Peterson GM; (1981):

Effect of euglycemia on the outcome of pregnancy in insulin- dependent diabetic women as compared with normal control subjects.

Am. J. Med; 71: 921 -25 .

Kaaja R., Tikkanen MJ., Viinikka L; (1995):

Serum lipoproteins. Insulin, and urinary prostanoid metabolites in normal and hypertensive pregnant women.

Obstet & Gynecol; 85: 353-6.

Kamoi K., sudo N., Ishibashi, et al; (1990):

Plasma endothelin-1 levels in patients with pregnancy induced hypertension.

N., Engl. J. Med; 323: 1486-1487.

Karaayenbrink AA., Dekker GA.,and Van Kamp G;(1993):

Endothelial vasoactive mediators in pre eclampsia.

Am J. Obstet & Gynecol; 169: 160-165

Khong TY., De Wolf F., et al; (1986):

Inadequate maternal vascular response to placentation in pregnancies complicated by pre eclampsia and by small for gestational age infants.

Br. J. obstet Gynecol; 93: 1049-59.

klpatrick DC., Gibson F., Liston WA., et al; (1989):

Association between susceptibility to pre eclmapsia within families and HLADR₄.

Lancet; 1: 1063 -65.

Kupfermine MJ., peaceman AM, Wigton TR., et al; (1994):

Tumour necrosis factor-alpha is elevated in plasma and amniotic fluid of patients with severe pre eclampsia.

Am. J. Obstet & Gynecol; 170: 1752-1757.

Marschner I., et al., (1974):

Group experiments on the radioimmunological insulin determination.

Horm Metabol Res; 6: 293-6.

Mc Manus IFA; (1980):

Carlton's histological techniques. 5th ed. Oxford.
New York. Oxford University press, 237.

MC Namara JR., Venner JL., Zengling L., et al; (1992):

Change in LDL particle size is associated with change in plasma triglyceride concentration.

Arterio Scler. Thromb; 12: 1284- 90 .

Meekins JW., Pijnenborg R., Hanssens M., et al; (1994):

A study of placenta bed spiral arteries and trophoblast invasion in normal and severe pre eclamptic pregnancies.

Br. J. Obstet & Gynecol; 101: 669-674.

Memon RA., Feingold KR., Moster AH. et al; (1992):

Differential effects of interleukin 1 and TNF on ketogenesis.

Am. J. Physiol; 263: E 301- E 309 .

Mestman JH; (1980):

Outcome of diabetes screening in pregnancy and perinatal morbidity in infants of mothers with mild impairment in glucose tolerance.

Diabetes care 3: 447-50.

Micheal A., and Weber M.D; (1993):

Hypertension as a risk factor syndrome. Therapeutic Implications.

Am. J. Obstet & Gynecol; 94: 24- 31.

Minuz P., et al; (1987):

Reduced excretion of vasodilator prostaglandin in pre eclampsia.

Basel PP; 175-181.

Modan M., Halkin H., Almog S., et al; (1985):

Hyperinsulinemia- a link between hypertension, obesity and glucose intolerance.

J clin invest; 75: 809-17.

Molnar M., Romero R., and Hertelendy F; (1993):

Interleukin-1 and tumour necrosis factor stimulate arachidonic acid release and phospholipid metabolism in human myometrial cells.

Am J of Obstet & Gynecol; 169: 825-9.

Muller D.C., Richard E., Jordan D., et al; (1993):

An epidemiological test of the hyperinsulinemia-hypertension hypothesis.

J. Clinical Endocrinol and metabolism; 76: 544-48

Naeye RL, (1989):

Pregnancy hypertension, placental evidence of low uteroplacental blood flow and spontaneous premature delivery.

Hum Pathol; 20: 441-45.

National Committee for clinical laboratory standards (1991):

Procedures for the collection of diagnostic blood specimens by venipuncture, 3rd ed., NCCLS .

Document H₃-A₃ .

Nicholas H., and Bryan M, (1996):

Nitric oxide, the endothelium, pregnancy and pre eclampsia.

Br J Obstet & Gynecol; 103: 4-15.

O'Brien R., and Granner DK; (1991):

Regulation of gene expression by insulin.

Biochem. J; 278: 609-13.

Odendaal H.J., Pattinson R.C., and Ram R.; (1990):

Aggressive or expectant management for patients with severe pre eclampsia between 28- 34 weeks' gestation.

Obstet & Gynecol; 76: 1070- 5.

Opsjon S.L., and Austgulen R.; (1995):

Interleukin-1, interleukin-6 and tumour necrosis factor at delivery in pre eclamptic disorders.

Acta Obstet & Gynecol scand; 74: 19-26.

Perez C; (1990):

A non secretable cell surface mutant of tumour necrosis factor (TNF) kills by cell- to- cell contact.

Cell 63: 251-55.

Peristeris P., Clark B., D., Gatti S., et al; (1992):

N., Acctylcystein and glutathion as inhibitors of tumour necrosis factor production.

Cell Immunol; 140: 390-399.

Pijnenborg R., Dixon G., et al; (1980):

Trophoblastic invasion of human decidua from 8 to 18 weeks of pregnancy.

Placenta 13-19.

Pober JS., Gimbrone MA., Lapierre LA., et al; (1986):

Overlapping patterns of activation of human endothelial cells by interleukin- 1, tumour necrosis factor and immune interferon.

J Immunol; 137: 1893-96.

Porteou F., and Nathan C; (1990):

Shedding of tumour necrosis factor receptors by activated human neutrophils.

J. Exp Med; 172: 599-607.

Potter MF., and Nestel PJ; (1979):

Hyperlipidemia of pregnancy in normal and complicated pregnancies.

Am. J. Obstet & Gynecol; 133: 165- 70 .

Reaven GM., and Banting Lecture (1988):

Role of insulin resistance in human disease.

Diabetes; 37: 1595- 607.

Redman C.W. G; (1991):

Current topic: pre eclampsia and the placenta.

Placenta, 12: 301-308.

Reeves WG; (1983):

Insulin antibody determination: theoretical and practical consideration.

Diabetologia, 24: 399-403.

Roberts JM, (1989):

Pregnancy related hypertension maternal-fetal medicine: principles and practices, 2nd ed
Philadelphia, saunders; 777.

Roberts JM., Taylor RN., Muscit., et al., (1989):

Pre eclampsia: An endothelial cell disorder.
Am J. Obstet & Gynecol; 161: 1200-1204.

Roberts JM., Taylor RN., Goldfien A; (1991):

Clinical and biochemical evidence of endothelial cell dysfunction in pregnancy syndrome pre eclampsia.

Am. J. of Hypertension; 4 (8): 700 -705.

Roberts JMRN., Taylor SA., Friedman; (1993):

New development in pre eclampsia. In fetal medical Reveiw.

Am J. Obstet & Gynecol; 169: 166-9.

Robertson WB., Khong TY., Brosens L., et al; (1986):

The placental bed biopsy: review from three European center.

Am J obstet & Gynecol; 155: 401-12.

Rodgers GM., Taylor RN., and Roberts JM., (1988):

Pre eclampsia is associated with serum factor
cytotoxic to human endothelial cells.

Am J Obstet & Gynecol; 159: 908-914.

Ryan EA., O'sullivan MJ., and Skyler JS; (1985):

Insulin action during pregnancy. Studies with the
euglycemic clamp technique.

Diabetes; 34: 380-389.

Sacks G.P., Studena K., Sargent IL., et al; (1998):

Normal pregnancy and pre eclampsia both produce
inflammatory changes in peripheral blood
leucocytes akin to those of sepsis.

Am. J. Obstet & Gynecol; 179: 80- 6 .

Saleh AA., Bottoms SF., Norman G., et al; (1988):

Hemostasis in hypertensive disorders of
pregnancy.

Obstet & Gynecol; 71: 719-22.

Schafer W., Tielsch., and Gasper FW; (1993):

Urinary excretion of 6-keto-PGF₁ alpha TXB₂ and
PGE₂ in a rat animal model for pre eclampsia like
syndrome.

J. Clin. Invest; 91: 2744-46.

Spellacy WN., Miller SJ., and Wineger A; (1986):

Pregnancy after 40 years of age.

Obstet & Gynecol; 68: 452 -55.

Spiedel MT., Booyse FM., Abrams A., et al; (1990):

Lipolysed hypertriglyceridaemic serum and triglyceride rich lipoprotein cause lipid accumulation in and are cytotoxic to cultured human endothelial cells. High- density lipoprotein in hibit this cytotoxicity.

Thromb. Res; 58: 251- 264 .

Stanley K., Fraser R., Bruce C; (1998):

Physiological changes in insulin resistance in human pregnancy: Longitudinal study with the hyperinsulinaemic euglycaemic clamp technique.

Br. J. Obstet & Gynecol; 105: 756- 59 .

Stark J., M; ((1993):

Pre eclampsia cytokine induced oxidative stress.

Br. J. Obstet & Gynecol; 100: 105- 109 .

stewart DJ., and Monge JC; (1993):

Hyperlipidaemia and endothelial dysfunction.

Curr. Opin. Lipidol; 4: 319- 324 .

Stirrat G.M.; (1986):

The immunology of hypertension in pregnancy
perinatology press, Ithaca, New York
PP; 249-259.

Suhonen L., and Teramo K; (1993):

Hypertension and pre eclampsia in women with
gestational glucose intolerance.
Acta. Obstet & Gynecol; 72: 269- 72 .

Tanfield PA., Ales KL., Resnick L.M., et al; (1987):

Hypocalcuria in pre eclampsia.
N. Engl. J Med; 316: 725-9.

Terranova PF., Hunter VJ., Roby KF., et al; (1995):

Tumour necrosis factor-alpha in the female
reproductive tract.
Proc. So. Exp. Biol Med; 209: 325-42.

Trinchieri G; (1992):

Tumour necrosis factors: the molecules and their
emerging role in medicine.
New York: Raven Press; 1992: 515-30.

Van JR., Anggard EE., Botting RM; (1990):

Regulatory functions of the vascular endothelium.
N. Engl J. Med; 323: 27-36.

Van der poll T., Sander J., and Van Deventer H; (1991):

Comparison of the early dynamics of systemic prostacyclin release after administration of tumour necrosis factor and endotoxin to healthy humans.
J Infect. Dis.; 164: 599-601.

Verhoeff FH; (1980):

Histological technique for placental examination
5th ed. Oxford, New York: Oxford University press; 195.

Vilcek J., Lee TH; (1991):

Tumour necrosis factor.
J. Biol. Chem. 266: 7313-15.

Vince G., Starkey P., Sargent I; (1992):

The expression of TNF- R at the maternofetal interface in normal human pregnancy.
Placenta, 12: 442- 443 .

Vince GS., Starkey PM, Austgulen R., et al; (1995):

Interleukin- 6, tumour necrosis factor and soluble tumour necrosis factor receptors in women with pre eclampsia.
Br. J. Obstet & Gynecol; 102: 20-5.

Visser W., Beckman L., Brenner HA; (1994):

Bioactive TNF- α in pre eclamptic women with and without the HELLP syndrome

Br. J. Obstet & Gynecol; 101: 1081- 82.

Wallach D., Peyser MR., Lessing JB., et al; (1995):

Soluble tumour necrosis factor receptors in maternal plasma and second trimester amniotic fluid.

Am J. Obstet & Gynecol; 173: 900-5.

Wang Y., Walsh SW., Guo J., et al; (1991):

Maternal levels of prostacyclin, thromboxane, vitamin E, and lipid peroxides throughout normal pregnancy.

Am J Obstet & Gynecol; 165: 1690-4.

Wang Y., Walsh SW., Kay HH; (1992):

Placental lipid peroxides and thromboxane are increased and prostacyclin decreased in women with pre eclampsia.

Am. J. Obstet & Gynecol; 167: 946- 49.

Warso MA, and Lands WEM; (1993):

Lipid peroxidation in relation to prostacyclin and thromboxane physiology and pathophysiology.

Br J Med Bull; 277-80.

White MF., Kahn CR; (1994):

The insulin signaling system.
J. Biol Chem; 269: 1-5.

Whorton AR., Montgomery MF., and Kent RS; (1985):

Effects of hydrogen peroxide on prostaglandin production and cellular integrity in cultured porcine aortic endothelial cells.
J Clin invest; 76: 295-302.

Wisdom SJ., et al; (1991):

Antioxidant system in normal pregnancy and in pregnancy induced hypertension.
Am. J. Obstet & Gynecol; 165: 1701 -1705.

Yanush Polsky EH., Ozyurk M., Polgar K., et al; (1993):

The effects of cytokines on human chorionic gonadotrophin (HCG) production by a trophoblast cell line.
J. of Reproductive. Immunology; 25: 235-47.

Yui J., Garcia LL., et al; (1994)

Cytotoxicity of tumour necrosis factor – alpha and gamma interferon against primary human placental trophoblasts.
Placenta, 15: 814-35.

Zlatnik E.J. and Burmester LF; (1983)

Dietary protein and pre. Eclampsia.
Am. J. Obstet & Gynecol; 147: 345-48.