

REFERENCES

- Abdel Sattar, N.A.; Sadek, S.H.; Sallam, M.S. and Abdel Dayem, M.K.:
Apolipoprotein B (Apo-B) and certain other lipid parameters in coronary atherosclerosis.
Thesis submitted for partial fulfilment of M.D. of Clinical Pathology, 1986.
- Alpert, J.S. and Braunwald, E.:
Acute myocardial infarction (MI): Pathological, pathophysiological and clinical manifestations; Heart Diseases. In: A Textbook of cardiovascular Medicine; E. Braunwald, (2nd)ed.; 1262-1300, 1984. W.B. Saunders Co. Philadelphia.
- Alpert, J.S. and Rippe, J.M.:
Ischaemic heart disease; In: Manual of Cardiovascular diagnosis and Therapy; Alpert, J.S., (2nd)ed.; 161-190, 1981. Little and Brown Co. Boston.
- American Heart Association:
Planning fat controlled meals for approximately 2000-3600 calories.
American Heart Association, New York, 1967.
- Avogaro, P.; Billalo Bon, G. and Cazzalato, G.:
Are apolipoproteins better discriminators than lipids for atherosclerosis?
Lancet, 1:901, 1979.
- Bailar, J.C.:
Cause and effect in epidemiology. What do we know about hypertriglyceridemia?
N. Engl. J. Med., 302: 1417, 1980.
- Baker, N.N.; Delahunty, T. and Jackson, R.L.:
The primary structure of human plasma HDL Glutamine I (Apo-A-I).
Proc. Natl. Acad. Sci. USA, 71: 3631, 1974.
- Balcon, R.:
Detection of ischaemic heart disease; In: Oxford textbook of Medicine; D.J., Weatherall; J.G.G., Ledingham and D.A., Warrel, (1st)ed.; 13-168, 1983.
Oxford University Press. Oxford, New York and Toronto.
- Barboriak, J.J.; Anderson, A.J. and Hoffman, R.G.:
Inter relationship between coronary artery occlusion, HDL-c and alcohol intake.
J. Lab. Clin. Med., 94: 348, 1979.

- Barham, D. and Tinder, P.:
An investigation of the determination of serum
uric acid by enzymatic method.
Analyst, 97: 142, 1972.
- Bersot, T.P.; Mahley, R.W.; Brown, M.S. and Goldstein,
J.L.:
Interaction of swine lipoproteins with the LDL
receptor in human fibroblast.
J. Biol. Chem., 251: 2395, 1976.
- Blum, C.B.; Levy, R.I.; Eisenberg, S.; Hall, H.;
Geobel, R.H. and Berman, M.:
High density lipoprotein (HDL) metabolism in man.
J. Clin. Invest., 60: 795, 1977.
- Boyles, J.K.; Pitas, R.E.; Wilson, E.; Mahley, R.W. and
Taylor, J.M.:
Apo-E associated with astrocytic glia of the
C.N.S. and with non-myelinating glia of the
peripheral nervous system.
J. Clin. Invest., 76: 1501, 1985.
- Breier, C.; Drexel, H. and Ranon, R.:
Effect of post-heparin lipoprotein lipase activity
and of plasma testosterone in CAD.
Lancet, 1:1242, 1985.
- Breslow, J.L.; Ross, D. and Mc. Pherson, J.:
Isolation and characterization of cDNA clones for
human Apo-A-I.
Proc. Natl. Acad. Sci. USA, 79: 6861, 1982.
- Brewer, H.B.; Fairwell, T.; Larue, A.; Ranon, R.;
Houser, A. and Bronzert, T.J.:
The amino acid sequence of human Apo-A, on
apolipoprotein isolated from HDL.
Biochem. Res. Commun., 80: 623, 1978.
- Brewer, H.B.; Lux, S.E.; Ranon, R. and John, K.M.:
Amino acid sequence of human Apo-LP-Gln-II (Apo-A-
II), apolipoprotein isolated from HDL complex.
Proc. Natl. Acad. Sci. USA, 69: 1304, 1972.
- Brook, J.G.; Aviran, M. and Viener, A.:
HDL subfractions in normolipidemic patients with
coronary atherosclerosis.
Circulation, 66: 923, 1982.
- Brown, M.S. and Goldstein, J.L.:
Lipoprotein metabolism in the macrophage:
Implications for cholesterol deposition in
atherosclerosis.
Ann. Rev. Biochem., 52:223, 1983.

- Brunnell, I.D.; Sniderman, A.D.; Albers, J.J. and Kwiterovich, P.O.Jr.:
Apo-B and Apo-A-I in CAD in humans.
Atherosclerosis, 4: 79, 1984.
- Campeau, L.:
The relation of risk factor to the development of atherosclerosis in saphenous vein bypass grafts.
N.Engl. J. Med., 311: 1329, 1984.
- Carlson, L.A.:
Ischaemic heart disease in relation to fasting values of plasma triglycerides and cholesterol.
Stockholm prospective study.
Lancet, 1: 865, 1972.
- Castelli, W.P.:
Epidemiology of CAD: The Framingham Study.
Am. J. Med., 76: 4, 1984.
- Castelli, W.P.; Abbott, R.D. and Mc Namara, P.M.:
Summary estimates of cholesterol used to predict CAD.
Circulation, 67: 730, 1983.
- Castelli, W.P.; Gordon, T. and Hjortland, M.C.:
Alcohol and blood lipids: The cooperative lipoprotein phenotyping study.
Lancet, 2: 153, 1977.
- Cooper, A.D.:
Role of the liver in the degradation of lipoproteins.
Gastroenterology, 88: 192, 1985.
- Dedonder-Decoopman, E.; Fievet-Desreumaux, C.; Campos, E.; Moulin, S.; Deailly, P.; Sezille, G. and Jaillaro, J.:
Plasma levels of VLDL, LDL-c, HDL-c, triglycerides and Apo-B and Apo-A-I in a healthy population.
Atherosclerosis, 37: 559, 1980.
- Durrington, P.N.; Hunt, I. and Ishola, M.:
Serum Apo-A-I and B and lipoproteins in middle-aged men with and without previous MI.
Br.H.J., 56: 206, 1986.
- Durrington, P.N.; Ishola, M. and Hunt, L.:
Apo-A, A-I and B and parental history in men with early onset ischaemic heart disease.
Lancet, 14: 1070, 1988.
- Dyer, A.; Stamler, J. and Shokelle, R.H.:
The relationship of education to blood pressure.
Circulation, 54: 987, 1976.

- Eisenberg, S.:
Lipoproteins and lipoprotein metabolism.
Klin. Wochenschr., 61: 119, 1983.
- Eisenberg, S.; Bilheimer, D.W. and Levy, R.I.:
The metabolism of VLDL. Studies on the transference of apolipoproteins between plasma lipoproteins.
Biophys. Acta, 280: 94, 1972.
- Eisenberg, S. and Levy, R.I.:
Lipoprotein metabolism.
Adv. Lipid Res., 13:1, 1975.
- Eisenberg, S.; Windmuller, H.G. and Levy, R.I.:
Metabolic fate of rat and human lipoproteins and apolipoproteins in the rat.
J. Lipid Res., 14: 446, 1973.
- Eliot, R.S. and Edwards, J.E.:
Pathology of coronary atherosclerosis and its complications; In: The Heart; J.W. Hurst.
(4th)ed.; 1121-1134, 1978.
Mc Grow-Hill Book Co. New York.
- El Rabat, K.A.; Tawadrous, G.A.; Hifny, A.; El-Sheikh, A.A. and Tabl, A.S.M.:
Lipoprotein and apolipoprotein pattern in patients with CAD.
Thesis submitted for partial fulfilment of M.Sc. degree in Cardiology, Benha Faculty of Medicine, 1987.
- Enger, S.C.; Herjornsen, K.; Erikssen, K. and Fretland, A.:
HDL and physical activity. The influence of physical exercise, age and smoking on HDL-c and the HDL-c/total cholesterol ratio.
Scand. J. Clin. Lab. Invest., 37: 251, 1977.
- Esders, W. and Michrina, C.A.:
Purification and properties of L- α -glycerophosphate oxidase from streptococcus faecium ATCC 12755.
J. Biol. Chem., 254: 2710, 1979.
- Evan, A.S.:
Lipids, lipoproteins and apolipoproteins;
In: Fundamentals of Clinical Chemistry; Norbert W. Tietz, (3rd) ed.; 448-460, 1987.
W.B. Saunders Co. Philadelphia, London, Toronto, Tokyo and Hong Kong.

- Felker, T.E.; Fainaru, M.; Hamilton, R.L. and Havel, R.J.:
Secretion of the arginine-rich and A-I apolipoproteins by the isolated perfused rat liver.
J. Lipid Res., 18: 465, 1977.
- Fessel, W.J.:
High uric acid as an indicator of cardiovascular disease independence from obesity.
Am. J. Med., 68: 401, 1980.
- Frantz, I.D.Jr. and Moore, R.B.:
The sterol hypothesis in atherogenesis.
Am. J. Med., 46: 484, 1969.
- Franzen, J. and Fex, G.:
Low serum Apo-A-I in acute MI survivors with normal HDL.
Atherosclerosis, 59: 37, 1986.
- Fredrickson, D.S.; Levy, R.I. and Lees, R.S.:
Fat transport in lipoproteins: An integrated approach.
N. Engl. J. Med., 276: 34, 1967.
- French, R.H.; King III, S.B. and Douglas, J.S.:
Techniques of cardiac catheterization including coronary arteriography; In: The Heart; J.W. Hurst, (5th)ed.; 1843-1850, 1982.
Mc Grow-Hill Book Co. New York.
- Friedberg, C.K.:
Pathogenesis of atherosclerosis; In: Diseases of the Heart; Friedberg, C.K., (3rd)ed.; 651-675, 1966.
W.B. Saunder Co. Philadelphia.
- Friedman, G.G.; Petitti, D.B. and Bawol, R.D.:
Mortality in cigarette smokers and quitters effect of base-line differences.
N. Engl. J. Med., 304: 1407, 1981.
- Friedman, H.H.:
Problem-Oriented Medical Diagnosis; In: Medical Diagnosis; Friedman, H.H., (3rd)ed.; 35-39, 1983.
Little and Brown Co. Boston and Toronto.
- Fruchart, J.C.:
Estimation of HDL-c and LDL-c by enzymatic method.
Rev. Fr. des Laboratoires, 103: 7, 1982.
- Fuller, J.M.; Shipley, M.J. and Rose, G.:
CAD risk and impaired glucose tolerance.
Lancet, 1: 1373, 1980.

- Geddes, J.S.; Adgey, A.A. and Pantridge, J.F.:
Prevention of cardiogenic shock.
Am. Heart J., 99(2): 243, 1980.
- Gibbons, L.W.; Blair, S.N.; Cooper, K.H. and Smith, M.:
Association between CAD risk factors and physical
fitness in healthy adult women.
Circulation, 67: 977, 1983.
- Gibson, D.G.; Precoitt, T.A. and Brown, D.J.:
Analysis of left ventricular wall movement during
isovolumic relaxation and its relation to CAD.
Br. Heart J., 38: 1010, 1976.
- Glickman, R.M. and Green, P.H.R.:
The intestine as a source of Apo-A-I.
Proc. Natl. Acad. Sci. USA, 74: 2569, 1977.
- Glueck, C.J.; Laskarzewski, P.M.; Suchindran, C.M.;
Chambless, L.E.; Barrett-Connor, E.; Stewart, P.;
Heiss, G. and Tyroler, H.A.:
Progeny's lipid and lipoprotein levels by parental
mortality.
The Lipid Research Clinics Program Prevalence
Study.
Circulation, 73: 151, 1986.
- Glueck, C.J. and Tsang, R.:
Longevity syndromes: familial hypobeta-and
familial hyperalpha-lipoproteinemia.
J. Lab. Clin. Med., 88: 941, 1976.
- Goldstein, J.L. and Brown, M.S.:
The LDL pathway and its relation to
atherosclerosis.
Ann. Rev. Biochem., 46: 897, 1977.
- Goldstein, J.L. and Brown, M.S.:
Lipoprotein receptors, genetic defect against
atherosclerosis.
Clin. Res., 30: 417, 1982.
- Gordon, T.; Castelli, W.P. and Hjortland, M.C.:
HDL as a protective factor against CAD.
Am. J. Med., 62: 707, 1977.
- Gotto, A.M.:
HDL: Biochemical and metabolic factors.
Am. J. of Card., 52: 4, 1983.
- Hamby, R.I.:
Hereditary aspects of CAD.
Am. Heart J., 101: 639, 1981.

- Havel, R.J.:
Familial dysbetalipoproteinemia.
The Medical Clinics of North America, 66(2): 319.
1982.
- Henry, J.B.:
Clinical diagnosis and management by laboratory
methods; In: Clinical Diagnosis; Todd, Sanford
and Davidson, (17th)ed.; 184-283, 1984.
W.B. Saunders Co. Philadelphia.
- Heyden, S.; Heiss, G.; Manegold, C.; Tyroler, H.A.;
Hames, C.G.; Bartel, A.G. and Copper, G.:
The combined effect of smoking and coffee drinking
on LDL-c and HDL-c.
Circulation, 66: 22, 1979.
- Hill, A.B.:
Statistical analysis; In: A Short Textbook of
Medical Statistics; Hill, A.B., (10th)ed.; 1-120,
1979.
The English Language Book Society, Hodder and
Stoughton, London.
- Hopf, U.; Assman, G.; Schaefer, H.E. and Capurso, A.:
The intestine is a source of Apo-A-I.
Gut, 20: 219, 1979.
- Huff, M.W.; Fidge, N.H.; Nestel, P.J.; Billington, T.
and Watson, B.:
Metabolism of Apo-C: Kinetics of Apo-C-II, C-III
and C-III2 and VLDL-Apo-B in normal and
hyperlipoproteinemic subjects.
J. Lipid Res., 22: 1235, 1981.
- Hui, D.Y.; Innerarity, T.L. and Mahley, R.W.:
Lipoprotein binding to canine hepatic membranes,
metabolically distinct Apo-E and Apo-B, E-
receptors.
J. Biol. Chem., 256: 5646, 1981.
- Hulley, S.B.; Rosenman, R.H.; Bawol, R.D. and Brand,
R.J.:
Epidemiology as a guide to clinical decision: The
association between triglycerides and CAD.
N. Engl. J. Med., 302: 1383, 1980.
- Innerarity, T.L.; Pitas, R.E. and Mahley, R.W.:
Receptor binding of cholesterol induced HDL
containing predominantly Apo-E to cultured
fibroblasts with mutation at the LDL receptor
locus.
Biochemistry, 19: 4359, 1980.

- Jackson, R.; Morris, M.A.; Haynes, B. and Eisenbarth, G.S.:
Increased circulating Ia-antigen bearing T-cells in Type I diabetes mellitus.
N. Engl. J. Med., 306: 785, 1982.
- Jackson, R.; Morrisett, J.D. and Gotto, J.:
Lipoprotein structure and metabolism.
Physiol. Rev., 56: 259, 1976.
- Jennings, R.B.; Sommers, H.M. and Smyth, G.A.:
Myocardial necrosis induced by temporary occlusion of a coronary artery in the dog.
Arch Pathol., 70: 68, 1960.
- Kane, J.P.; Hardman, D.A. and Paulus, H.E.:
Heterogeneity of Apo-B: Isolation of a new species from human chylomicrons.
Proc. Natl. Acad. Sci. USA, 77: 2465, 1980.
- Kane, J.P.; Sata, T.; Hamilton, R.L. and Havel, R.J.:
Apolipoprotein composition of VLDL of human serum.
J. Clin. Invest., 56: 1622, 1975.
- Kannel, W.B.; Castelli, W.P. and Gordon, T.:
Cholesterol in the prediction of atherosclerotic disease.
Ann. Intern. Med., 90: 85, 1979.
- Kannel, W.B.; Castelli, W.P.; Gordon, T. and Mc Namara, P.M.:
Serum cholesterol, lipoproteins and the risk of CAD. The Framingham Study.
Ann. Intern. Med., 74: 1, 1971.
- Kannel, W.B.; Hjortland, M.C.; Mc Namara, P.M. and Gordon, T.:
Menopause and risk of cardiovascular disease. The Framingham Study.
Ann. Intern. Med., 85: 447, 1976.
- Karathanasis, S.K.; Mc Pherson, J.; Zannis, V.I. and Breslow, J.L.:
Linkage of human Apo-A-I and Apo-C-III genes.
Nature, 304: 371, 1983.
- Keys, A.:
CAD in seven countries.
Circulation, 41: 211, 1970.
- Khan, M.I.G.:
Management of acute MI: In: Manual of Cardiac Drug Therapy; Khan, M.I.G., (2nd)ed.; 125-155, 1988.
Bailliere Tindall/W.B. Saunders. London, Philadelphia, Toronto, Sydney and Tokyo.

- King, E.J.; Pillai, S.S. and Beal, D.:
Modification of Folin and Wu for blood sugar
estimation.
Lancet, 1: 310, 1941.
Quoted from, Varely, H.; Cowenlack, A.H. and Bell,
M.:
Estimation of blood sugar. In: Practical Clinical
Biochemistry; Vol. I, 86-88, 1980.
Churchill Livingstone Medical Division of Longman
Group Limited. London and Philadelphia.
- Kostner, G. and Holasek, A.:
Characterization and quantification of the
apolipoproteins from human chyle chylomicrons.
Biochemistry, 11: 1217, 1972.
- Kozarevic, D.; Mc Gee, D.; Vojardic, N.; Race, Z.;
Dawler, T.R. and Gordon, T.:
Frequency of alcohol consumption and morbidity and
mortality. The Yugoslavia Cardiovascular Disease
Study.
Lancet, 1: 613, 1980.
- Kukita, H.; Hamada, M.; Hiwada, K. and Kodubu, T.:
Clinical significance of measurement of serum Apo-
A-I, A-II and B in hypertriglyceridemic male
patients with and without CAD.
Atherosclerosis, 55: 143, 1985.
- Kukita, H.; Hiwada, K. and Kokuba, T.:
Serum Apo-A-I, A-II and B levels and their
discriminative values in relatives of patients
with CAD.
Atherosclerosis, 51: 261, 1984.
- Kukita, H.; Imamura, Y.; Hamada, M.; John, T. and
Kokubu, T.C.:
Plasma lipids and lipoproteins in Japanese male
patients with CAD and their relatives.
Atherosclerosis, 42: 21, 1982.
- La Rosa, J.C.; Levy, R.I.; Herbert, R.; Lux, S.E. and
Fredrickson, D.S.:
A specific apolipoprotein activator for
lipoprotein lipase.
Biochem. Biophys. Res. Commun., 41: 57, 1970.
- Layde, P.M.; Beral, V. and Kay, C.R.:
Further analysis of mortality in oral
contraceptive users.
Royal College of General Practitioner Oral
Contraception.
Lancet, 1: 549, 1981.

- Lee, T.H. and Goldman, L.:
Serum enzyme assays in the diagnosis of acute MI.
Annals of Internal Medicine, 105: 221, 1986.
- Levy, R.I.:
Cholesterol, lipoproteins, apolipoproteins and heart disease:
Present status a future prospect.
Clin. Chem., 27(5): 623, 1981.
- Levy, R.I. and Feinleib, M.:
Risk factors for CAD and their management;
In: Coronary Heart Diseases. Prevention, Complications and Treatment; Conner, W.E. and Bristow, J.D., (1st)ed.: 1205-1234, 1985.
Lippincott, Philadelphia, London, Mexico City, New York, St. Louis, Sao Paulo and Sydney.
- Lew, E.A. and Garfinkel, L.:
Variation in mortality by weight among 750,000 men and women.
J. Chronic dis., 32: 563, 1979.
- Lipid Research Clinical Program:
The lipid research clinics coronary primary prevention trial results. The relationship of reduction in incidence of CAD to cholesterol lowering.
JAMA, 251: 365, 1984.
- Lowe, G.D.; Machado, S.G.; Krol, W.F; Barton, B.A. and Forbes, C.D.:
White blood cell count and haematocrite as predictors of coronary recurrence after MI.
Thromb. Haemost., 54(3): 700, 1985.
- Maciejko, J.J.; Halwes, D.R. and Kottke, M.D.:
Apo-A-I as a marker of angiographically assessed CAD.
N. Engl. J. Med., 309: 385, 1983.
- Mahley, R.W.; Hui, D.Y.; Innerarity, T.L. and Weisgraber, K.H.:
Two independent lipoprotein receptors on hepatic membranes of dog, swine and man. Apo-B, E and Apo-E receptors.
J. Clin. Invest., 68: 1197, 1981.
- Mahley, R.W.; Weisgraber, K.H. and Innerarity, T.L.:
Atherogenic hyperlipoproteinemia induced by cholesterol feeding in the patas monkey.
Biochemistry, 15: 2979, 1976.

Mao, S.J.T.; Jackson, R.L.; Gotto, A.M. and Sparrow, J.T.:

Mechanism of lipid-protein interaction in the plasma lipoproteins: Identification of a lipid-binding site in Apo-A-II.
Biochemistry, 20: 1676, 1981.

Marcus, A.K.; Milton, J.C. and Lawrence, M.T.J.R.:
Acute MI; In: Current Medical Diagnosis and Treatment; Marcus, A.K. Middle East edition; 225-234, 1985.
Lange Medical Publications. California.

Marriott, H.J.L.:
Myocardial Infarction; In: Practical Electrocardiography; Marriott, H.J.L., (6th)ed.; 233-259, 1980.
Williams and Wilkins Co. Baltimore and London.

Maseri, A.; Chierchias and L'Abbate, A.:
Pathogenetic mechanisms underlying the clinical events associated with atherosclerotic heart disease.
Circulation, 62: 5, 1980.

Mc Conathy, W.J. and Alaupovic, P.:
Studies on the isolation and partial characterization of Apo-D and lipoprotein D in human plasma.
Biochemistry, 15: 515, 1976.

Menzel, H.J.; Kladetzky, H.G. and Assman, G.:
Apo-A-IV isoform, isoelectric focusing identification.
J. Lipid Res., 23: 915, 1982.

Micheli, H.; Pometta, D.; Jornot, C. and Scherre, J.R.:
HDL-c in male relatives of patients with CAD.
Atherosclerosis, 32: 269, 1979.

Miller, G.J. and Miller, N.E.:
Plasma HDL concentration and development of CAD.
Lancet, 1: 16, 1975.

Miller, N.E.:
HDL: A major risk factor for coronary atherosclerosis.
Bailliere's Clinical Endocrinology and Metabolism, 1(3): 603, 1987.

Miller, N.E.; Hammett, F. and Saltissi, S.:
Relation of angiographically defined CAD to plasma lipoprotein subfractions and apolipoproteins.
Br. Med. J., 282: 1741, 1981.

- Morrisett, J.D.; Jackson, R.L. and Gotto, A.M.:
Lipoprotein structure and function.
Ann. Rev. Biochem.. 44: 183. 1975.
- Morrison, J.A.; Mellaes, M.; De Groot, J.; Khoury, D.;
Gortside, P.S. and Glueck, C.J.:
Cigarette smoking, alcohol intake and oral
contraceptive relationships to lipids and
lipoproteins in adolescent school-children.
Metabolism, 28: 1166. 1979.
- Morton, N.E.; Gulbrandsen, C.L.; Rao, D.C.; Rhoads,
G.G. and Kagan, A.:
Determinants of blood pressure in Japanese-
American families.
Hum. Genet.. 53: 261. 1980.
- Newman, W.P. III; Freedman, D.S. and Voors, A.W.:
Relation of serum lipoprotein levels and systolic
blood pressure to early atherosclerosis: Boglusa
Heart Study.
N. Engl. J. Med.. 314: 138. 1986.
- Nishimura, R.A.; Reeder, G.S. and Miller, F.A.:
Prognostic value of predischage 2-dimensional
echocardiogram after acute MI.
Am. J. Cardiol.. 53: 429. 1983.
- Noma, A.; Yokosuka, T. and Kitamura, K.:
Plasma lipids and apolipoproteins as
discriminators for the presence and severity of
angiographically CAD.
Atherosclerosis, 49(1): 1. 1983.
- Nupuf, M.S. and Sutherland, W.H.F.:
HDL levels in children of young men with ischaemic
heart disease.
Atherosclerosis, 33: 365. 1979.
- Okazaki, M.; Ohro, Y. and Hara, I.:
Rapid method for the quantitation of cholesterol
in human serum lipoproteins by high performance
liquid chromatography.
J. Biochem.. 89: 879. 1981.
- Oram, S.:
Ischaemic heart disease; In: Clinical Heart
Diseases; Oram, S., (2nd)ed.; 413-465. 1981.
William Heineman Medical Books LTD. London.
- Persky, V.W.; Dyer, A.R. and Shekelle, K.B.:
Uric acid: A risk factor for CAD.
Circulation, 59: 969. 1979.

- Phillips, R.L.; Lilienfeld, A.M. and Diamond, E.L.:
Frequency of CAD and cerebrovascular accidents in
parents and sons of CAD index cases and controls.
Am. J. Epidemiol., 100: 87, 1974.
- Pometta, D.; Micheli, H.; Raymond, L.; Oberhaensli, I.
and Suenram, A.:
Decreased HDL-c in prepubertal and pubertal
children of CAD patients.
Atherosclerosis, 36: 101, 1980.
- Pometta, D.; Micheli, H.; Suenram, A. and Jornot, C.:
HDL in close relatives of CAD patients.
Atherosclerosis, 34: 419, 1979.
- Pometta, D.; Suenram, A.; Sheybani, E.; Brab, B. and
James, R.:
HDL-c levels in patients with MI and their
families.
Atherosclerosis, 59: 21, 1986.
- Rees, A.; Shoulder, C.C.; Stocks, J.; Galton, D.J. and
Barallel, F.E.:
DNA polymorphism adjacent to human Apo-A-I gene.
Lancet, 1: 444, 1983.
- Reichlin, M.; Visco, J.P. and Kocke, F.J.:
Radioimmunoassay for human myoglobin. Initial
experience in patients with CAD.
Circulation, 57: 52, 1978.
- Rifkind, B.M. and Segal, P.:
Lipid Research Clinics Program Reference Values
for hyperlipidemia and hypolipidemia.
JAMA, 250(14): 1869, 1983.
- Rissanen, A.M.:
Familial aggregation of CAD in a high incidence
area (North Karelia, Finland).
Br. Heart J., 42: 294, 1979.
- Rissanen, A.M. and Nikkila, E.A.:
CAD and its risk factors in families of young men
with angina pectoris and controls.
Br. Heart J., 39: 875, 1977.
- Rissanen, A.M. and Nikkila, E.A.:
Aggregation of coronary risk factors in families
of men with fatal and non-fatal CAD.
Br. Heart J., 42: 373, 1979.
- Robbins, S.L.:
Pathologic basis of CAD; In: Pathologic Basis of
Diseases; Robbins, S.L., (1st)ed.: 21-30, 1974.
W.B. Saunders Co. Philadelphia.

- Robbins, S.L. and Cortan, R.S.:
CAD pathology; In: Pathologic Basis of Diseases;
Robbins, S.L., (2nd)ed.; 1226-1238, 1979.
W.B. Saunders Co. Philadelphia, London and
Toronto.
- Ross, G.:
Familial patterns in ischaemic heart disease.
Br. J. Prev. Soc. Med., 18: 75, 1964.
- Rubenstein, B.:
Protein content and composition of human HDL
subfractions.
Atherosclerosis, 33: 415, 1979.
- Schaefer, E.J.; Eisenberg, S. and Levy, R.I.:
Lipoprotein-apolipoprotein metabolism.
J. Lipid Res., 19: 667, 1978.
- Scheidt, S.; Ascheim, R. and Killip III, T.:
Shock after acute MI.
Am. J. Cardiol., 26: 556, 1973.
- Schmidt, S.B.; Wasserman, A.G.; Muesing, R.A.;
Schlesselman, S.E.; La Rosa, J.C. and Ross, A.M.:
Lipoprotein and apolipoprotein levels in
angiographically defined coronary atherosclerosis.
Am. J. Cardiol., 55: 1459, 1985.
- Schmitz, G. and Assman, G.:
Biosynthesis of HDL subfractions.
J. Lipid Res., 23: 865, 1982.
- Schmitz, G.; Assman, G. and Melnik, B.:
Role of LCAT in HDL3/HDL interconversion.
Clin. Chim. Acta, 119: 225, 1982.
- Shirai, K.; Barnhart, R.L. and Jackson, R.L.:
Hydrolysis of human plasma HDL phospholipids and
triglycerides by hepatic lipase.
Biochem. Biophys. Res. Commun., 100: 591, 1981.
- Simmons, A.V.; Sheppard, M.A. and Cox, A.F.:
Deep venous thrombosis after MI.
Br. Heart J., 35: 623, 1973.
- Slack, J.:
Risks of ischaemic heart disease in familial
hyperlipoproteinemic states.
Lancet, 2: 1380, 1969.

- Slack, J. and Nevin, N.C.:
Hyperlipidemic xanthomatoses. Increased risk of death from ischaemic heart disease in first degree relatives of 53 patients with essential hyperlipidemia and xanthomatosis.
J. Med. Genet., 5:4, 1968.
- Sniderman, A.; Shapiro, S. and Marpole, D.:
Association of coronary atherosclerosis with hyperapobetalipoproteinemia.
Proc. Natl. Acad. Sci. USA. 77: 604, 1980.
- Sobel, B.E. and Shell, W.E.:
Serum enzyme determination in the diagnosis and assessment of MI.
Circulation, 45: 472, 1972.
- Stamler, J. Stamler, R. and Liu, K.:
High blood pressure: Role in CAD and implications for prevention and control.
Geneva, Cardiovascular Unit, W.H.O./C.V.D., 8: 835, 1983.
- Steiner, R.E.:
Ischaemic heart disease a challenge to radiology.
Br. J. Radiology, 56: 626, 1983.
- Stephen, M.F.:
Pathophysiology of myocardial ischaemia; In: The Heart; J. Willis Hurst; Robert C. Schlant; Charles E. Rachley; Edmund H. Sonnenblick and Nanette Kass Wenger, (7th)ed.; 940-960, 1990.
Mc Graw-Hill Information Services Company Health Professions Division, New York.
- Stone, D.; Shapiro, S.; Kaufman, D.W.; Rosenberg, L.; Nietnen, O.S. and Stolley, P.W.:
Risk of MI in relation to current and discontinued use of oral contraceptives.
N.Engl. J. Med., 305: 420, 1981.
- Tanaka, A.:
Role and metabolism of HDL subfractions. Analysis of serum LDL2-c and HDL3-c in patients with various diseases by high performance liquid chromatography.
Bull. Tokyo Med. Den. University, 29: 131, 1982.
- Tatami, R.; Mabuchi, H.; Veda, K.; Sveda, R.; Hava, T.; Kametani, T.; Ito, S.; Koizumi, J.U.; Ohata, M.; Miyamoto, S.; Nakayama, A. and Kanaya, M.:
Intermediate density lipoprotein and cholesterol rich VLDL in angiographically determined CAD.
Circulation, 64: 1174, 1981.

Taylor, K.G.; Carter, T.J.; Valente, A.J.; Wright, A.D.; Smith, J.H. and Matthews, K.A.:
Sex difference in the relationships between obesity, alcohol consumption and cigarette smoking and serum lipid and apolipoprotein concentration in a normal population.
Atherosclerosis, 38: 11, 1981.

Tyroler, H.A.:
Epidemiology of plasma HDL-c levels.
Lipid Research Clinics Program Prevalance Study.
Circulation, 62: 4, 1980.

Ten Kate, L.P.:
Familial aggregation of CAD and its relation to known genetic risk factors.
Am. J. Cardiol., 50: 945, 1982.

Thompson, G. and Soutar, A.:
Genetic polymorphism and plasma lipoproteins.
Nature, 301: 658, 1983.

Tibbutt, D.A.:
True left ventricular aneurysm.
Br. Med. J., 289: 450, 1984.

Timmis, A.D.:
Post myocardial infarction syndrome.
Br. Med. J., 289: 636, 1984.

Trahen, C.A.; Gere, J.B. and Bigham, D.A.:
Clinical assessment of serum myosin light chain in the diagnosis of acute MI.
Am. J. Cardiol., 41: 641, 1978.

Uterman, G. and Beispiegel, U.:
Apo-A-IV, a protein occurring in human mesenteric lymph chylomicrons and free in plasma, isolation and quantification.
Eur. J. Biochem., 99: 333, 1979.

Uterman, G.; Feussner, G.; Franceschini, G.; Hass, J. and Steinmetz, A.:
Genetic variants of group A apolipoproteins.
J. Biol. Chem., 257(1): 501, 1982.

Warnick, G.R.:
Apolipoproteins.
Clinical Chemistry News, 9:8, 1985.

- Warnick, G.R.; Mayfield, C.; Albers, J.J. and Hazzard, W.R.:
Gel isoelectric focusing method for specific diagnosis of familial hyperlipoproteinemia type III.
Clin. Chem., 25: 279, 1979.
- Warren, J.V. and Lewis, R.P.:
Beneficial effects of atropine in the pre-hospital phase of coronary care.
Am. J. Cardiol., 37: 68, 1976.
- Watson, D.:
Direct estimation of total cholesterol.
Clin. Chem. Acta, 5: 637, 1960.
- Weisgraber, K.H.; Bersot, T.P.; Mahley, R.W.; Franceschini, G. and Sirotori, C.R.:
Apolipoproteins.
J. Clin. Invest., 66: 901, 1980.
- Weisgraber, K.H. and Mahley, R.W.:
Apo-E-A-II complex of plasma lipoproteins. Characterization of this mixed disulfide and its identification in a HDL subfraction.
J. Biol. Chem., 253: 6281, 1978.
- Weisgraber, K.H.; Roll, S.C. and Mahley, R.W.:
Human Apo-E heterogeneity.
J. Biol. Chem., 256: 9077, 1981.
- Whayne, T.F.; Alaupovic, P.; Curry, M.D.; Lee, E.T.; Anderson, P.S. and Schechter, E.:
Plasma Apo-B and VLDL, LDL and HDL-c as risk factors in the development of CAD in male patients examined by angiography.
Atherosclerosis, 29: 411, 1981.
- Willerson, J.T.:
Acute MI; In: Cecil Textbook of Medicine; Wyngaarder, J.B. and Smith, L.H., (16th)ed.; 247-256, 1982.
W.B. Saunders Co. Philadelphia, London and Toronto.
- Willetts, W.; Hennekens, C.H.; Siegel, A.J.; Adner, M.M. and Castelli, W.P.:
Alcohol consumption and HDL-c in marathon runners.
N. Engl. J. Med., 303: 1159, 1980.
- Williams, R.S.; Louge, E.E.; Lewis, J.L.; Parton, T.; Stread, N.W.; Wall Ace, A.G. and Pizzo, S.V.:
Physical conditioning augments the fibrinolytic response to venous occlusion in healthy adults.
N. Engl. J. Med., 302: 987, 1980.

Windmueller, H.G.; Herbert, P.N. and Levy, R.I.:
Biosynthesis of lymph and plasma lipoprotein
apolipoprotein by isolated perfused rat liver and
intestine.
J. Lipid Res., 14: 215, 1973.

World Health Organization (WHO):
World health statistics. Vital statistics and
causes of death.
Annual, 1:1, 1973.
World Health Organization. Geneva.

Zannis, V.I. and Breslow, J.L.:
Characterization of a unique human Apo-E variant
associated with type III hyperlipoproteinemia.
J. Biol. Chem., 255(5): 1759, 1980.

Zannis, V.I. and Berslow, J.L.:
Human VLDL-Apo-E isoprotein polymorphism is
explained by genetic variation and post-
translational modification.
Biochemistry, 20: 1033, 1981.