

REFERENCES

[A]

Abd El - Hady, S. (1996) :

Clinical application of urine antigen detection in group B streptococcal neonatal sepsis. Thesis, submitted for partial fulfillment of M.Sc. degree in pediatrics. Benha Faculty of Medicine.

Abd El-Maksoud, Y.H. (1997) :

Some metabolic changes in neonatal sepsis. Thesis, submitted for partial fulfillment of M.D. degree in pediatrics. Benha Faculty of Medicine.

Abu-Farrag, M.S.; Rowisha, M.A.; Mahfouz, S.A. and El-Sheikh, M.M. (1996) :

Interleukin-6 in neonatal sepsis. The Gaz. Egypt. Pediatr. Assoc., 53 : 63.

Adriaanse, A.H. (1996) :

Prevention of neonatal septicemia due to group B streptococci. Clin. Obstet. Gynaecol., 9 (3) : 545.

Alpert, G .; Campos, M.; Preblud, S.R. and Plotkin, S.A. (1984):

Vancomycin dosage in pediatrics.
Am. J. Pharmacol., 63 : 190 .

Alt, R.; Willars, D.; Messer, J. et al. (1982) :

Interet de la C-reactive protein dans Les infections bacteriens neonatales.
Arch. Fr. Pediatr.; 39 : 811-813.

- Amadori, A.; Zamarchi, R.; Del Mistro, A. et al. (1989) :*
HIV-specific B cell activation. A major constituent of spontaneous B cell activation during HIV- infection. J. Immunol. 143, 2146-2152.
- Amara, E. E. (1995) :*
A study of calcium homeostasis in septicemic neonates. Thesis, submitted for partial fulfillment of M. D. degree in pediatrics. Benha Faculty of Medicine.
- Amato, M.; Howald, H. and Von moralt, G. (1988) :*
Quantitative changes of white blood cells and perinatal diagnosis of infection in high risk preterm infants. Pediat., 23 (2) : 129- 34.
- Ambrosino, D.M.; Delaney, N.R. and Shamberger, R.C., (1990) :*
Human polysaccharide specific B cells are responsive to IL-6. J.Immunol. 144 : 1221-1226.
- Andreas, H.; Groll Anja Meiser, Martina Weise, et al. (1992):*
IL-6 as early mediator in the neonatal sepsis pediatric infect Dis. J. 11 (6) : 496-498.
- Andus, H.; Groll, Anja Meiser, Martina Weise, et al., (1987) :*
IL-6 as early mediator in neonatal sepsis. Pediatric. Infect. Dis. J. 11, (6), 496-498.
- Asano, S.; Okano, A. and Ozawa, K. et al. (1990) :*
In vivo effects of recombinant human IL-6 in primates. Stimulated production of platelets. Blood 75, 1602-1605.

Avery, G.B. (1994) :

Bacterial and viral infection of the newborn. Arch. Dis. Child., 71:112.

[B]

Baker, C. J. and Edward, J. (1990) :

Noscomial septicemia and meningitis in neonates. Am. J. Med., 70 : 698.

Bakr, M.B. (1994) :

Adrenal functions in neonatal septicemia, Thesis submitted for partial fulfillment of M.D. degree in pediatrics, Benha Faculty of Medicine, Zagazig University.

Baltimore, R.S.; Kasper, D.L. and Baker, C.J. (1977) :

Antigenic specificity of opsonophagocytic antibodies in rabbit antisera to GBS. J. Immunol., 118 : 673.

Baner, J. and Hermann, F. (1991) :

IL-6 in clinical medicine. Ann. Hematol. 62, 203 -210.

Battisti, O.; Mitchinson, R. and Davies, P.A. (1991) :

Changing blood culture isolates in a referral neonatal intensive care . Arch . Dis . Child., 86 : 775.

Beagley, K.W.; Kiyono, H. and Kishimoto, T. (1989) :

Interleukins and IgA synthesis.
J. Exp. Md. 169 : 2133-2148.

Bell, J.Y. and McGuiness (1982) :

Serum level of IL-6 as a prognosis factor in metastatic renal cell carcinoma. Cancer Res. 52, 3317-3322.

Bennet, R.; Bergdahl, S.; Erikson, M. and Zetterstorm, R. (1989):

The outcome of neonatal septicemia during fifteen years. Acta. Pediatr. Scand. , 78 : 40 .

Bennet, P.; Eriksson, M. and Zetterstrom, R. (1981) :

Increasing incidence of neonatal septicemia. Causative organisms and predisposing risk factors.
Acta. Pediatr. Scand, 70 : 207-10.

Bhardway, N.; Tatter, S.B. and Rivelis, M. et al., (1989) :

IL-6/IFN- γ synovial effusion of patients with rheumatoid arthritis. J. Immunol. 143, 2153-2159.

Bhutta, Z.A.; Naqui, S.H. and Muzaffar, T. (1991) :

Neonatal septicemia in Pakistan : Presentation and Pathogens. Acta. Pediatr. Scand; 80 : 596-601.

Bont, E.S.J.M.; Martens, A. and Van Raanij. (1994) :

Diagnostic value of plasma levels of tumour necrosis factor- α (TNF- α) and IL-6 in newborns with sepsis. Acta. Pediatr., 83 : 696-9.

Boyer, K.M.; Samuel, P. and Gotoff, S. (1986) :

Prevention of early onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis . N. Engl.J.Med. , 314 : 1665.

Brown, T.J.; Lie, J.; Shoyab, M. and Rowe, J.M. (1990) :

Regulation of IL-6 expression by oncostatin.
M. J. Immunol. 147 : 2175-21800.

Buck, C.; Bundschu, J.; Gallati, H.; Battmann, P. and pohlandt, F. (1994):

Interleukin-6 : a sensitive parameter for the early diagnosis of neonatal bacterial infection. Pediatrics, 93:54-8

Burstein, S.A.; Downs, T. and Friese, P., et al., (1992):

Thrombocytosis in normal and sublethally irradiated dogs. Responses to human IL-6. Blood 80, 420-428.

[C]

Cairo, M.S. (1989):

Neonatal neutrophil host defence . AJDC., 143 : 40.

Cairo, M.S.; Plunkett, J.M.; Nguyen, A. and Van de Ven, C. (1991) :

Sequential administration of IL-6 and G-CSF in newborn rats : modulation of newborn granulopoiesis and thrombopoiesis. Pediatr. Res; 30 : 554-559.

Cairo, M.S.; Rucker, R.; Bennetts, G.A.; Hicks, D.; Katz, J. and Johnson, S. (1984) :

Improved survival of newborns receiving leucocyte transfusions for sepsis. Pediatr., 74 (5) : 887.

Caldron, T.M.; Scherman, J. and Bermann, J.W. et al. (1992):

IL-6 modulates gene expression in cultured human endothelial cells. Cell immunol. 143, 118-126.

Castell GV.; Geiger, T.; Gross V. et al (1988) :

Plasma Clearance organ distribution and target cells of IL -6 / hepatocyte stimulating factor in the rat. Eur J, Biochem, 177 :375-361.

Catherine, M.H.; Costarino, A.T.; Sullivan, J.S. and Dulkerian, S. (1994) :

Cytokine elevations in critically ill infants with sepsis and necrotizing enterocolitis. J. Pediatr., 124: 105-111.

Chen, I.; Mory, Y. and Revel, M. (1988) :

Growth inhibition of human breast carcinoma and leukemia/lymphoma cell lines by recombinant interferon B2. Proc. Natt. Acad. Sci. U.S.A. 85, 8037-8041.

Christensen, R.D.; Bradly, R. and Rothstein, G. (1981) :

The leukocyte left shift in clinical and experimental neonatal sepsis. J.Pediatr, 98 :101.

Christensen, R.D. and Rethstein, G. (1979) :

Granulocyte transfusion in neonates with bacterial infection, neutropenia and depletion of mature marrow neutrophils. Pediatr., 70 (1) : 1.

Clapp, D.W. ; Kligman, R.M.; Baley, J.E.; Shenker, N.; Fanaroff, A.A. and Berger, M. (1989) :

Use of intravenously administered immunoglobulin to prevent nosocomial sepsis in low birth weight infants : Report of a pilot study . J . Pediatr. , 115 (6) : 973.

Cole, F.S. (1991) :

Bacterial infection of the newborn. In Schaffer and Avery's (ed). Diseases of the newborn. 6th Edition W.B. Saunders company Philadelphia., P. 350.

Cole, F.S. and Cloherty, J.P. (1985) :

Infection, prevention and treatment in manual of neoant care by little, Brown and company, Bosoton P. 193, 153 : 156.

Coovadia, Y.M. ; Mayosi, B. and Adhikari, M.Z. (1989):

Hospital acquired neonatal bacterial meningitis : the impact of cefotaxime usage on mortality and of Amikacin usage on incidence . Ann. Trop. Pediatr. , 9 (4) : 233.

Corrigan, J.J. (1984) :

Thrombocytopenia. A laboratory sign of septicemia in infants and children. J. Pediatr, 105 : 219.

[D]

Dalvi, R.; Rao, S. and Rangnoker, J. (1991) :

Exchange transfusion in neonatal sepsis.
Indian. J. Pediatr., 28 (1) : 39.

Damas, P.; Ledaux, D.; DeGroote, D. and Lamy, M. (1992) :

Cytokine serum level during sever sepsis in human IL-6 as amarker of severity. Ann. Surg.; 215 (4): 356-62.

Dawodu, A.H. and Alausa, O.K. (1980) :

Neonatal septicemia in the tropics.
Afr. J. Med. sci., 2 : 1-6.

Debets, R.; Rao, S. and Rangnoker, J. (1989) :

Exchange transfusion in neonatal sepsis .
Indian .J. Pediatr., 28 (1):39.

Dietzman, D.E.; Fischer, G.W. and Schoenkencht, D. (1994):

Neonatal Escherichia coli septicaemia.
J. Pediatr., 185 : 128.

Dinareello, C.A. (1991):

The role of cytokines in the pathogenesis of fever :
basic mechanisms and management. New York :
Raven Press; 23-47.

Dollagramatic, H.D. and Tapratizitim, S.(1993) :

Comparison of 4 tests used in diagnosis of neonatal
septicemia. Arch. Dis. Child., 69 :429.

Donato, H.; Gebara, E.; De Cosen, R.H. and Gioseffio, M. (1992) :

Leukocyte alkaline phosphatase activity in the
diagnosis of neonatal bacterial infection . J . Pediatr . ,
121 (2) : 242

Dowton, S.B. and Calton, H.R. (1988) :

Acute phase reactants in inflammation and infection.
Hematol., 25 : 84.

El-Dadah, M. ; Frenkel, L.D. ; Hiatt, I.M. and Hegy, T. [E] (1987) :

Evaluation of routine lumbar punctures in newborn
infants with respiratory distress syndrome . Pediatr .
Infect . Dis . J., 6 : 243 .

El-Diwani, K.M.; Tayeb, S.; Eissa, A.M.; El-Fell, S. and Eltohamy, F. (1987) :

Neonatal septicemia. The Gaz. Egypt. Pediatr. Assoc., 35 : 37.

El-Maraghi, S. ; El-Saifi, A. and Abou-Alam. M. (1984) :

Diagnosis of neonatal septicemia by gastric aspirate examination . J.Egypt Soc . Ob/Gyn . , 10 (1) : 43 .

El-Radi, A.S.; Jawad, M.; Mansour, N.; Jamil, I. and Ibrahim, M. (1983) :

Sepsis and hypothermia in the newborn infant . J. Pediatr., 103 : 300.

El-Sallab, S.M.; Abdel Aal, A.; El-Tonbary, Y. and El-Kappany, M.A. (1987) :

Some aspects of neonatal septicemia of Mansoura Neonatal care Unit. Thesis submitted in partial fulfillment for the master degree in pediatrics, P.74-94.

Entman, M.I.; Bossen, R.D. and Anderson, D.C. (1991) :

Inflammation in the course of early myocardial ischemia.

FASEB. J. 5, 2529 -2537.

Eshali, H. ; Ringertz, S. and Nystrom, S. (1989):

Septicaemia with coagulase negative staphylococci in a neonatal intensive care unit . Acta . Pediatr . Scand. , 360: 127.

[F]

Feigin, R.D. ; Abramson, S.L. and Edwards, M.S. (1987) :
Infection in the fetus and neonate. In Franaroff, A.A.
and Martin, R.J.(eds.): Neonatal and Perinatal
Medicine. ed.4. Mosby Co., Baltimore, Boston.,
P:763.

**Freedman, R.M.; Ingram, D.L.; Gross. L.; Warshav, J.B.
and Baltimore, M. (1981) :**
A half century of neonatal sepsis at Yale (1928-1978).
Am. J. Dis. Child., 135 : 140.

[G]

Gamberzade, S. ;Toppare, M.F. and Senses, D.A. ; (1996) :
Neonatal sepsis and meningits, Pak.
Med. Assoc., 46 (2): 43.

Gaur, S.; Kesarwala, H. and Gupta, M. (1994) :
Clinical immunology and infectious diseases.
Pediatr. Clin. N. Am.; 41 (4) : 745-759.

Gerdes, J.S. (1991) :
Clinicopathologic approach to diagnosis of neonatal
sepsis. Clin. Perinatology, 18 (2) : 361.

**Gilstrap, L.G.; Leveno, K.J.; Cox, S.M.; Burris, J.S. and
Rosenfeld, C.R. (1988) :**
Intrapartum treatment of acute chorio-amnionitis :
Impact on neonatal sepsis. Am . J. Obstet. Gynecol .,
159 : 579.

Givon, T.; Salvin, S.; Revel, M. et al. (1992) :

Anatitumor effects of human recombinant IL-6 on acute myeloid leukemia in mice and in cell cultures. Blood 79, 2392-2398.

Gomella, T.L.; Cunningham, M.D. and Eyal, F.G.(eds). (1994):

Neonatology: Management, Procedures, on - call problems. prentice- Hall International Inc.2nd. ed. pp339.

Gotoff, S.P. (1996) :

Sepsis of newborn. In Behrman, R.E.; Kliegman, R.M. and Arvin, A.M. (eds.) : Nelson Textbook of Pediatrics. ed. 15, W.B. Saunders, Philadelphia, London, Toronto Montreal, Sydney, Tokyo, p : 528 .

Grael, El .; Halle . E.; Bollmann , R. and Buch, H. (1989):

Neonatal septicaemia - incidence , etiology and outcome . Acta . Pediatr. Scand . , 360 : 113.

Groll, A.H.; Meiser, A.; Weise M. et. al. (1992) :

IL-6 as early mediator in neonatal sepsis. Pediar. Infect. Dis. J., 11:496-8.

Gronross, P.; Vesikari, T. and Janas, M. (1985) :

Neonatal septicemia. Arch. Dis. Child.; 60 : 542.

Grossman, R.M.; Krueger, J.; Yourish, D. et al. (1989) :

IL-6 is expressed in high levelin psoriasis skin and stimulates proliferation of cultured human kratinocytes. Proc. Natl. Acad. Sci. USA, 86 : 6367 - 6371.

Guerina, N.C. (1991) :

Bacterial and fungal infection. In Cloherty J. Pand Stork. AR (ed). Manual of neonatal care. 3rd edition. Little Brown and company London. 146.

Guillermín, R.; Vargo, T.M. and Rossier, J. (1977) :

Beta-endorphin and adrenal corticotropin and secreted concomitantly by the pituitary gland. Science; 197 : 1367-70.

[H]

Hall, R.T.; Shigheoka, A.O. and Hill, H.R. (1983) :

Serum opsonic activity and peripheral neutrophil count before and after exchange transfusion in infants with early onset group B streptococcal septicemia. Pediatr. Infect. Dis., 2 (5) : 356.

Halliday, H.L. (1989) :

When to do a lumbar puncture in a neonate. Arch. Dis. Child, 64 : 313 .

Hama, T.; Miyamoto, M.; Tsukui, H. et al. (1989) :

IL-6 as a neutropic factor for promoting the survival of cultured basal.

Haque, K.H. (1992) :

Infection and immunity in the newborn. Textbook of Pediatr. P. 302 -316.

Haque, K.N.; Ziadi, M.H. and Bahakin, H. (1988):

IgM enriched intravenous immunoglobulin therapy in neonatal sepsis. Am. J. Dis. Child., 142 : 1293.

Harris MC, Costarino AT, Sullivan JS, et al. (1994) :

Cytokine elevations in clinically ill infants with sepsis and necrotizing enterocolitis. J. Pediatr., 124(1): 105-111.

Helgott, D.C.; May, L.T. and Sthoeger, Z. (1987):

Bacterial lipopolysaccharide (endotoxin) enhances expression & secretion of B₂ interferon by human fibroblasts G Exp Med., 166 : 1300

Heinrich, P.C. (1990) :

Interleukin-6 (IL-6) and the acute phase response. Biochem. J. 265 : 621-36.

Heney, D.; Lewis, J.; Evans, S.W. et al. (1992) :

IL-6: Its relationship to CRP and fever in children with febrile neutropenia. J. Infec. Dis., 165 : 886-890.

Hermann, E.; Gruss, H.J. and Lubbert, M. (1991) :

IL-4 inhibits growth for multiple myelomas by suppressing IL-6 expression. Blood 78 : 2070 - 2074.

Hill, R.J.; Warren, M.K. and Levin, J. et al. (1992) :

Evidence that IL-6 does not play a role in the stimulation of platelet production after induction of acute thrombocytopenia. Blood 80 : 346 - 351.

Hirano, T. and Matsuda, T. (1986) :

Complementary DNA for a novel human interleukin that induces B-lymphocytes to produce immunoglobulin. Nature (London) 324, 73-76.

Hirohata, S. and Miyamoto, T. (1990) :

Elevated levels of IL-6 in cerebrospinal fluid from patients with systemic lupus erythematosus and central nervous system involvement. *Arthritis Rheum.* 33, 644 - 649.

Hocher, J.R. ; Simposn, P.M. ; Rabaliais, G. and Stewart , L. ; (1992) :

Extracorporeal membrane oxygenation and early onset group B streptococcal sepsis . *Pediatr.*, 89 (1) : 3 .

Honda, M.; Kitamura, K. and Okura, T. (1990) :

Quantitative analysis of serum IL-6 and its correlation with increased levels serum IL-2R in HIV-induced diseases. *J. Immunol.* 145, 4059 - 4064.

Horowitz, B. ; Chang , M.D. and Lippin , A. (1984) :

Preparation of antihemophilic factor and fibronectin from human plasma cryoprecipitate. *Clin. Bioch.*, 24: 357.

Hosford, D. and Braquet, P. (1990) :

The potential role of platelet-activating factor in shock and ischemia. *J. Crit Care*; 5 : 115.

Houston, J.B. (1983) :

Pus cells and bacterial counts in the diagnosis of urinary tract infections. *Arch. Dis. Child.*, 38 : 600.

Houssian, F.A.; Olive, D.; Van Snick, J. et al. (1988) :

Synergetic activation of human T-cell by IL-6 and IL-1
Eur. J. Immunol. 18 : 653 - 656.

[I]

Ikebuchi, K.; Wong, G.G.; Clark, S. et al. (1987) :

IL-6 enhancement of IL-3 dependent proliferation of
multipotential hemopoietic progenitors. Proc. Natl.
Acad. Sci. U.S.A. 84 : 9035-9039.

Ischimi, Y.; Akatsu, T. and Yamguchi, A. (1990) :

IL-6 is produced by osteoblasts and induces bone
resorption. J. Immunol. 145 : 3297 - 3303.

Ishibashi, T.; Kimura, H. and Shikawa, Y. (1989) :

IL-6 is a potent thrombopoietic factor in vivo in mice.
Blood 74 : 1241-1244.

Isomaki, H.A.; Hakulinen, T. and Joustsenlahti, U. (1978) :

Excess risk of lymphomas, leukemias and myelomas in
patients with rheumatoid arthritis. J. Chron. Dis. 31 :
591-696.

[J]

Jacobes, R.F.; Darville, T. and Parks, A.J. (1992) :

Safety profile and efficacy of cefotaxime for the
treatment of hospitalized children. Clin. Infect. Dis.,
56 : 65.

Jourdan, M.; Battaille, B.; Segnin, J. et al. (1990) :

Constitutive production of IL-6 and immunologic
features in cardiac myxomas. Arthritis rheum. 33 : 398
- 402.

[K]

Kafetzis, D.A.; Brater, A. and Kapiki, A.N. (1982) :

Treatment of severe neonatal infection with cefotaxime: Efficacy and pharmacokinetics. J. Pediatr., 100 : 483.

Kawano, M.; Hirano, T. and Taga, T. (1988) :

Autocrine generation and requirement B F2/IL-6 for human multiple myeloma. Blood 73 : 2145 - 2148.

Kern, P.; Van Damme, J.; Dietrich, M. et al. (1989) :

Elevated TNF and IL-6 serum levels as markers for complicated plasmodium falciform malaria. Am. J. Med. 87 : 139 -143.

Khalifa, A.M.; Allam, M. ; Nawar, F. and Sabry, S.I. (1989):

Quantitative and Qualitative changes in neonatal bacterial infection . The Gazet Egypt. Pediatr. Associa. , 37 (1) .

Kirnbaur, R.; Kock, A.; forster, E. et al. (1991) :

Regulation of epidermal cell IL-6 production by UV light and corticosteroid.
J. Invest. Dermatol. 96 : 484 - 489.

Kitabara, M.; Kishimoto, S. and Hirano, T. (1990) :

The in vivo antitumours effect of human recomb. Inant IL-6 Jpn. J. Cancer Res. 81 : 1032 - 1038.

Kite, P.; Millar, M.R.; Gohram, P. and Congdon, P. (1988) :

Comparison of five tests used in diagnosis of neoantal bacteremia. Arch. Dis. Child., 63 : 39.

- Kite, P.; Todd, N.; Mackay, P. and Miller, M.R. (1990) :**
 Plasma neutrophil elastase alpha-1-proteinase inhibitor level in premature neonates with coagulase-negative staphylococcal bacteremia. *J. Pediatr*; 117 (4) : 630-1.
- Klein, J.O.; Dashafesky, B. and Mayer, A. (1983) :**
 Selection of antimicrobial agents for treatment of neonatal sepsis. *Rev. Infect. Dis.*, 5 : 55.
- Kleiman, M.B.; Reynold, J.K. and Shreiner, R.L. (1994):**
 Rapid diagnosis of neonatal bacteremia with acridine orange-stained buffy coat smears . *J. Pediatr* ; 125 : 419.
- Koj, A. (1985) :**
 In : The acute-phase response to injury and infection (A.H. Gordon and A. Kop, eds 9 Vol. 10 : P. 139, Amstradam/ New York.
- Kurlat , L. ; Stoll , B.J. and Mc Gowan , J.E. (1989) :**
 Time to positivity for detection of bacteremia in neonates . *J. Clin . Microbiol .*, 27 (5) : 1068 .
- Kushner, I. (1982) :**
 The pheonmenon of acute phase response.
Ann. NY Acad. Sci.; 389 : 39 - 48.
- Kyllonen, K.S. ; Clapp, W. ; Robert, M. ; Nancy, S. and Fanaroff, A. (1989):**
 Dosage of intravenously administered immune globulin and dosing interval required to maintain target levels of immunoglobulin G in low birth weight infants. *J. Pediatr.*, 115 (6) : 1013.

[L]

Label, M.H. and McCracken, G.H. (1989) :

Delayed cerebrospinal fluid sterilization and adverse outcome of bacterial meningitis in infants. Pediatrics, 83 : 161.

Lagrutta, E.F.; Drusin, I.M. and Mackles, A.W. et. al. (1993):

Neonatal infection . An J. Dis. child, 137:838-41

Lancefield, R.C. ; McCarty, M. and Everly, W. (1975):

Multiple mouse protective antibodies directed against Group B Streptococci. J. Exp . Med ., 142 : 165 .

Leeming, J.P.; Sutton, T.M. and Fleming, P.Y. (1995) :

Neonatal skin as a reservoir of infection in neonatal sepsis. Pediatr. Inf. Dis. J., 14 (8) : 719.

Lehrnbecher, T.;Schord, T.; Krous, D. and Troos, Martius, J. (1995) :

Interleukin-6 and soluble interleukin -6 receptor in cord blood in the diagnosis of early onset sepsis in neonates. Acta. Pediatr., 84 : 806-8.

Lerman, S.J. (1986) :

Infections of the newborn infant.
Biol. Neonates, 49 : 58.

Littlewood, J.M. (1982) :

Sixty six infants with urinary tract infection in the first month of life . Arch. Dis . Child . , 47 : 218 .

Liu, C.H. ; Lehan. C. ; Speer, M.E. ; Smith , E.O. ; Fernbach , D.J. and Rudolph, A.J. (1985):

Early detection of bacteremia in an outpatient clinic .
Pediatr. , 75 (5) : 827 .

[M]

Maberry, M.C. and Gilstrap, L.C. (1991) :

Intrapartum antibiotic therapy for suspected intraamniotic infection : Impact on the fetus and neonate. Clin. Obstet-Gynaecol., 34 (2) : 345.

Macintoch, K. (1984) :

Bacterial infections of the newborn . Biol . Neonates, 38 : 114 .

Manroe, B.L.; Weinberg, A.G.; Rosenfeld, C.R. and Browre, R. (1979):

The neonatal blood count in health and disease.
J. Pediatr. , 95 : 89 .

Marodi, L.; Kalmar, A. and Szabo, I. (1989) :

Opsonic activity in serum from septic infants treated with intravenous immunoglobulin . Arch. Dis. Child ., 64 530.

Martens, J.; Samson, W.P.F. and Fetter, A. (1993) :

TNF- α , IL-6 and IL-1B plasma level in neonatal sepsis. Pediatr. Res. 33 (4) :380 - 383.

Maruo, N.; Morita, I. and Matota, S. (1992) :

IL-6 increases endothelial permeability in vitro. J. Indocrinology 134 : 710 - 714.

Mathers, N.J. (1987) :

Diagnostic audit of C-reactive protein in neonatal infection. Eur. J. Pediatr., 146 : 147-51.

McCabe, W.R.; Treadwell, T.L. and Maria, A.D. (1983) :

Pathophysiology of bacteremia.
Am. J. Med.; 28 : 7-16.

McCracken , G.H. and Freij , B.J. (1987) :

Perinatal bacterial diseases. In Freij, R.D. and Chervy, J.D. (eds): Textbook of Pediatric Infectious Disease. ed.2. W.B. Saunders Co .Philadelphia., p : 940.
Cephalohematoma infection in neonatal septicaemia .
Pediatr., 44 (1) : 27.

McIntosh, N. (1992) :

The Newborn. In : Chanphell AGM and McIntosh, N (eds.). Forfar and Arneil's Textbook of Pediatrics, 4th ed. WB Saunders. Company, Philadelphia, London,Toronto, P. 109.

Meser, J.; Eyer, D.; Donato, L. et al. (1996) :

Evaluation of interleukin-6 and soluble receptors of tumor necrosis factor for early diagnosis of neonatal infection. J. Pediatr. 129 : 574- 580.

Miki, S.; Yamamoto, M. Sonoda, T. et al. (1989) :

IL-6 functions as in vitro autocrine growth factor in renal cell carcinoma. FEBS Lett. 250 : 607 - 891.

Miles, S.A.; Stevens, R.H.; Longan, D.M. et al. (1990) :

Kaposi Sarcoma derived cells produce and respond to IL-6. Proc. Natt. Acad. Sci. U.S.A. 87 : 4068 - 4072.

- Miller, L.C.; Isa, S.; Loprestd G. et. al. (1990) :**
Neonatal interleukin- 1B, interlukin -6 and tumour recrosisfactor :Cord blood levels and Cellular production. J. Pediatr, 22:281-285
- Mock, B.A.; Kozak, C. and Clark, S.C. (1989) :**
The murine IL-6 gene maps to the proximal region of chromosomes. J. Immunol. 142 : 1372-1376.
- Moffet, L.H. (1989) :**
Pediatric infection disease neonatal sepsis and meningitis., P. 496-507.
- Molyneux, R.; Schleiffenbaum, B. and Fehr, J. (1991):**
IL-1 and TNF stimulate human vascular endothelial cells to promate transendotheliol neutrephil passage. J. Clin. Invest, 83 : 444.
- Monga, K.; FERMAMDEZ, A. and DEADHAR, L. (1986) :**
Changing bacterialological septicemia. Indian J Pediatr., 53 : 505-508.
- Morinage, Y.; Suzuki, H. and Takasuki, F. (1989) :**
Contribution of IL-6 to the anti-proliferative effect of IL-1 and TNF- α on the tumor cell lines. J. Immunol. 143 : 4538 - 4542.
- Mul's, J.J.; Jablons, D.M.; Robenberg, A. et al. (1990) :**
Antitumour activity of recombinant IL-6 in Mice. J. Immunol. 148 : 2622 - 2629.

Mustafa, M.M.; Lebel, M.H. and Ramilo, O. (1989) :

Correlation of IL-1B and TNF concentrations in CSF and outcome from bacterial meningitis. J. Pediatr.; 115 : 208 - 13.

[N]

Naitoh, Y.; Tominaga, T.; Imura, H. et al. (1988) :

IL-6 stimulates the secretion of adrenocorticotrophic hormone in conscious, freely moving rats. Bioch. Biophys. Res. Commun. 155 : 1459 - 1463.

Nakajima, K.; Martinez, O.; Hirano, T. et al. (1989) :

Induction of IL-6 production by HIV. J. Immunol. 142, 144 - 147.

Neumann, S.; Hennrich, N.; Gunzer, G. and Lang, H. (1983):

Enzyme-linked immunoassay for complexes of human granulocyte elastase with α_1 -proteinase inhibitor in plasma. In Goldberg, D.M. and Werner, M. (eds.) : Progress in Clinical Enzymology. ed. 2, Masson Co, New York, p : 293.

Neurath, A.R.; Strik, N. and Sprot, P. (1992) :

Search for hepatitis B virus cell receptors reveals binding sites for IL-6 on the virus envelope protein. J. Exp. Med. 175 : 164 - 169.

Nicholas, G. (1991) :

Bacterial and fungal infection. Ped. Infect. Dis. J., 16: 39.

Nijsten, N.W.N.; DeGroot, R.; Ten Duis, H.J. et al. (1987) :
Serum levels of IL-6 and acute phase responses.
Lancet 2 : 921.

Nishimoto, N.; Yoshizaki, K.; Tagoh, M. et al. (1989) :
Elevation of serum IL-6 prior to acute phase proteins
on the inflammatory by surgical conditions. Clin.
Immunopath. 50 : 399-401.

[O]

O'Donnell, J.P. and Merenstein, G.B. (1993) :
Infection in the neonate . In Merenstein and Gardner
(eds.) : Handbook of Neonatal Intensive Care. ed. 3.
Mosby Year Book Boston Chicago London
Philadelphia Sydney, p : 287.

Odio, C.M. (1995) :
Cefotaxime for treatment of neonatal sepsis and
meningitis. Diagn. Microbiol. Infect. Dis., 11 : 85.

Ohabo, K.; Sugamura, K. and Kogune, K. (1991) :
IL-6 in cerebrospinal fluid of HTLV-1 associated
myelopathy, Neurology 41 : 594 - 595.

Okusawa, S.; Gelfand, J.A. and Ikejima, T. (1988) :
IL-1 induces a shock -like state in rabbits synergism
with TNF and the effect of cyclooxygenase inhibitors.
J. Clin. Invest., 81 : 1162 - 72.

Oster, W.; Klein, H. and Hirano, T. (1989) :
Participation of the cytokines IL-6, TNF α and TNF- β
secreted by acute myelogenous leukemia blasts in
autocrine and paracrine leukemia growth control. J.
Clin. Invest. 84 : 451-457.

Overall, J.C. (1987):

Infection of the newborn . In Behrman , R.F. and
Vaugan, V.C. (eds.) : Nelson Textbook of Pediatrics.
ed. 13 W.B. Saunders, Philadelphia, London, Toronto,
Tokyo, p : 422.

[P]

Padmanbhan, J.; Risembery, H.M. and Rowe, D. (1993) :

Howell-Jolly bodies in the peripheral blood of full
term and premature neonates. John. Hopkins. Med. J.,
132 : 146 .

Patchen, M.L.; McVittie, T.J. and Schwartz, G.N. (1991) :

Administration of IL-6 stimulates multilineage
hematopoiesis and accelerates recovery from radiation
induced hematopoietic depression. Blood. 77 : 472 -
280.

Pelton, B.K.; Hylton, W. and Demman, A.M. (1992) :

Activation of IL-6 production by U.V. irradiation of
blood mononuclear cells from patients with SLE. Clin.
Exp. Immunol. 89 : 251-254.

Philip, A.G. (1981) :

Decreased use of antibiotics using neonatal sepsis
screening technique . J . Pediatr. , 98 : 795 .

Philip, A.G. and Hewitt, J.R. (1980) :

Early diagnosis of neonatal sepsis. Pediatr., 65 (5) :
1036.

Pizzo, P.A. (1992) :

Modulation of host defenses by cytokines evolving adjuncts in prevention and treatment of serious infections in immunocopromised hosts. Clin. Infect. Dis., 15 : 508 - 524.

Plow, E.F. (1982) :

Leukocyte elastase during blood coagulation. J. Clin. Invest., 69 : 564 .

Poli, G.; Bressler, P.; Kinter, A. et al. (1990) :

IL-6 induces human immunodeficiency virus expression in infected monocytic cells alone and in synergy with TNF- α by transcriptional and posttranscriptional mechanisms. J. Exp. 172 : 151 - 158.

Pommier, C.G. ; O'Shea , J. and Chused , T. (1984) :

Studies on fibronectin receptors of human peripheral blood leucocytes . J. Exp . Med ., 159 : 137 .

Pourcyrons, M. ; Korohees, S.B. ; Bada, H.S. and Patterson, T. (1988) :

Indwelling umbilical arterial catheter catheter : A preferred sampling site for blood culture . Pediatr ., 81 (6) : 821.

Pyati, S.P.; R.S.; Jacobes, M.M.; Raval, D.S.; Amma, P. and Lillen, L. (1983) :

Penicillin in infants weighing two kilograms or less with early oncet group B streptococcal disease . N . Engl . J . Med ., 308 : 1383 .

- [R]
Raff, H.V.; Siscoe, P.J.; Wolf, E.A.; Maloney, G. and Shuford, W. (1988) :
 Human monoclonal antibodies to GBS : Reactivity and in vivo protection against multiple serotypes . J . Exp. Med ., 168 : 905 .
- Rekha, H.U ; Vase S. and Reyes, M. (1980) :**
 Premature rupture of membranes and neonatal infections. Indian. J. Pediatr. 37 : 137.
- Renauld, I.G.; Vink, A. and Van Snick, J. (1989) :**
 IL-4 and IL-6 in CTL induction accessory signals in murine cytolytic T-cell responses : Dual requirment for IL-4 and IL-6. J. Immunol. 143 : 1894 -1898.
- Riedy, M. and Stewart, C.C. (1992) :**
 Inhibitory role of interleukin-6 in macrophage proliferation. J. Leucocyte Biol. 52 : 125-127.
- Roberton, N.R. (1992):**
 Neonatal infections. Ped. Infect. Dis. J., 13: 65.
- Robillard, P.Y.; Nabeth, P. and Hulsey, T.C. (1993) :**
 Neonatal bacterial septicemia in a tropical area : Four-year experience in Guadeloupe (French West Indies). Acta. Pediatr., 82 : 687 - 9.
- Rodwell, L.R. ; Anton, L.L. and David. T. (1988) :**
 Early diagnosis of neonatal sepsis using a haematological scoring system. J. Pediatr., 112 : 761.

- Rozycki, H.J. ; Stahl, G.E. and Baumgart, S. (1987) :***
 Impaired sensitivity of a single early leucocytic count
 in screening for neonatal sepsis . *Pediatr. Inf. Dis. J* , 6
 : 440.
- Rusconi, F.; Parizzi, F.; Garlaschi, et al. (1991) :***
 IL-6 activity in infants and children with bacterial
 meningitis. *Pediatr. Infect. Dis. J.*; 10 : 117-21.
- Russel, G.A.; Symth, A. and Cooko, R.W. (1992) :***
 Receiver operating characteristic curves for comarison
 of serial neutrophil band from and CRP in neonates at
 risk of infection. *Arch. Dis. Child.*, 67 : 808.
- /S/***
- Saez-Liorens,X. and McCracken, G.H. (1993):***
 Sepsis syndrome and septic shock in pediatrics:
 current concepts of terminology, Pathophysiology, and
 management. *J. Pediatr.* 123:497-508.
- Samuel, P.G. (1992) :***
 Sepsis in the newborn. In : Krugman S, Kat SL,
 Gershon SL and Wilfert CM. (eds.). *Texbook of*
Infectious Disease of children, 9th ed. The CV &
 Mosby Company, Saint Louis, P. 402.
- Sanchez, P. ; Siegel, J.D. ; Cushion, N.B. and Norma, T.***
(1990):
 Significance of a positive urine group B streptococcal
 latex agglutination test in neonates . *J. Pediatr.* , 116 :
 601.

- Sehgal, P.B.; Helfgott, D.C.; Tatter, S.B. et al. (1988) :***
 Regulation of acute phase and immune response in viral disease enhanced expression of IL-6 gene in virus infected fibroblasts. J. Exp. Med. 167 : 1951-1956.
- Sherman, M.P.; Chance, K.H. and Goetzman, B.W. (1989) :***
 Gram's stains of tracheal secretions predict neonatal bacteremia. Am. J. Dis. Child. 138 : 848.
- Sherman, M.P.; Goetzman, B.W.; Ahlfors, C.E. and Weinberg, R.P. (1980) :***
 Tracheal aspiration and its clinical correlates in the diagnosis of congenital pneumonia. Pediatr., 65 : 258.
- Shimizu, S.; Hirano, Y.; Yoshiok, B. et al. (1988) :***
 IL-6 dependent growth of alennert's lymphoma-derived T-cell line Blood 72 : 1826-1828.
- Siegel, J.D. (1985) :***
 Neonatal septicaemia. Perinatol., 9 : 20.
- Simon, C.; Schroder, H.; Beyer, C. and Zerbst, T. (1991):***
 Neonatal sepsis in an intensive care unit and results of treatment. Infection, 19 (3) : 146.
- Skavunu, T.; Gatzola, K. and Hatzioannidis, K. (1991) :***
 Efficacy of -aztreonam in the treatment of neonatal sepsis . Rev . Infect . Dis., 13 (7) : 591 .
- Spangelo, B.L.; Isakson, P. and MacLead, R.M. (1989) :***
 IL-6 stimulates anterior pituitary hormone release in vitro Indocrinology 125 : 575 - 577.

- Rozycki, H.J. ; Stahl, G.E. and Baumgart, S. (1987) :***
 Impaired sensitivity of a single early leucocytic count
 in screening for neonatal sepsis . *Pediatr. Inf. Dis. J* , 6
 : 440.
- Rusconi, F.; Parizzi, F.; Garlaschi, et al. (1991) :***
 IL-6 activity in infants and children with bacterial
 meningitis. *Pediatr. Infect. Dis. J.*; 10 : 117-21.
- Russel, G.A.; Symth, A. and Cooko, R.W. (1992) :***
 Receiver operating characteristic curves for comarison
 of serial neutrophil band from and CRP in neonates at
 risk of infection. *Arch. Dis. Child.*, 67 : 808.
- /S/***
- Saez-Liorens,X. and McCracken, G.H. (1993):***
 Sepsis syndrome and septic shock in pediatrics:
 current concepts of terminology, Pathophysiology, and
 management. *J. Pediatr.* 123:497-508.
- Samuel, P.G. (1992) :***
 Sepsis in the newborn. In : Krugman S, Kat SL,
 Gershon SL and Wilfert CM. (eds.). *Texbook of
 Infectious Disease of children*, 9th ed. The CV &
 Mosby Company, Saint Louis, P. 402.
- Sanchez, P. ; Siegel, J.D. ; Cushion, N.B. and Norma, T.
 (1990):***
 Significance of a positive urine group B streptococcal
 latex agglutination test in neonates . *J. Pediatr.* , 116 :
 601.

- Sehgal, P.B.; Helfgott, D.C.; Tatter, S.B. et al. (1988) :**
Regulation of acute phase and immune response in viral disease enhanced expression of IL-6 gene in virus infected fibroblasts. J. Exp. Med. 167 : 1951-1956.
- Sherman, M.P.; Chance, K.H. and Goetzman, B.W. (1989) :**
Gram's stains of tracheal secretions predict neonatal bacteremia. Am. J. Dis. Child. 138 : 848.
- Sherman, M.P.; Goetzman, B.W.; Ahlfors, C.E. and Weinberg, R.P. (1980) :**
Tracheal aspiration and its clinical correlates in the diagnosis of congenital pneumonia. Pediatr., 65 : 258.
- Shimizu, S.; Hirano, Y.; Yoshiok, B. et al. (1988) :**
IL-6 dependent growth of alennert's lymphoma-derived T-cell line Blood 72 : 1826-1828.
- Siegel, J.D. (1985) :**
Neonatal septicaemia. Perinatol., 9 : 20.
- Simon, C.; Schroder, H.; Beyer, C. and Zerbst, T. (1991):**
Neonatal sepsis in an intensive care unit and results of treatment. Infection, 19 (3) : 146.
- Skavunu, T.; Gatzola, K. and Hatzioannidis, K. (1991) :**
Efficacy of -aztreonam in the treatment of neonatal sepsis . Rev . Infect . Dis., 13 (7) : 591 .
- Spangelo, B.L.; Isakson, P. and MacLead, R.M. (1989) :**
IL-6 stimulates anterior pituitary hormone release in vitro Indocrinology 125 : 575 - 577.

Togari, H.; Iwanaga, T. and Kito, H. (1983):

Endotoxin clearance by exchange blood transfusion in septic shock neonates. *Acta. Pediatr. Scand.*, 72 : 87.

Tollner, U.; Pohlandt and Rike, S.R. (1982) :

Septicemia in the newborn due to gram-negative bacilli : Risk factors, clinical symptoms and hematological changes. *Europ J. Pediatr.*; 123 : 243-254.

Tracey, K.J.; Beutler, B. and Lowry, S.F. (1986):

Shock and tissue injury induced by recombinant human cachectin-*Science*, 234: 470-4.

Trugo, W.E.; Gibson, R.L. and Henderson, W.R. (1990):

Tumour necrosis factor - induced neonatal pulmonary hypertension. Effects of dazmegrel pretreatment. *Pediatr. Res.*, 27 : 466.

Tsang, K.Y.; Primusf, J. and Schlom, J.(1991):

Human recombinant IL-6 enhance antibody dependent cellular cytotoxicity. *Clin. Obstet. Gynaecol.*, 8 (3) :545.

Tullus, B.; Lundstrom, G. and Burman, G. (1991) :

Antibiotic susceptibility of 629 bacterial blood and CSF isolates from swedish infants and the therapeutic implication. *Acta. Pediatr. Scand.*, 80 : 205.

[V]

Vain, N.E.; Mazlumian, J.R. and Swarner, O.W. (1980):

Role of exchange transfusion in the treatment severe septicaemia. *Pediatr.*, 66 : 693.

- Van Damme, J.; Fibbe, W.F. and Flkenbury, L.H. (1989):***
 Simultaneous production of IL-6, interferon - B and colony colony stimulating activity by fibroblasts after viral and bacterial infection. *Dur. Immunol.* 19, 163 - 168.
- Van Deventer, S.J.H.; Buller, H.R. and Catem, J.W. (1990):***
 Experimental endotoxemia in human : analysis of cytokine release and coagulation, fibrinolytic and complement pathways. *Blood*; 76 : 2520.
- Vesikari, T.; Isolauri, E.; Tuppurainen, N.; Renlund, M. ; Koivisto, M. and Kunnas, M. (1985) :***
 Neonatal septicaemia in Finland 1981-1985. *Acta. Pediatr. Scand.*, 78 : 44.
- Vink, A.; Coulie, P.G. and Stevens, M. (1990):***
 Mouse plasmacytoma growth in vivo : Enhancement by IL-6 and inhibition by antibodies direct against IL-6 or its receptor. *J. Exp. Med.* 172, 997 - 1000.
- Visser , V.E. and Hall, R.T. (1980) :***
 Lumbar puncture in the evaluation of suspected neonatal Sepsis. *J. Pediatr.*, 96 : 1063.
- [W]***
- Walchiers, D.; Van-Loock, F.; Yeanfils, G. and Vandepitte, J. (1995) :***
 Epidemiological study on neonatal septicemia and meningitis in Belgium . *Acta - Clin - Belg .*, 50 (6) : 326 .

Weisbart, P.H.; Gasson, J.C. and Golde, D.W. (1989):

Colon - stimulating factors and host defence. Ann Inter. Med., 110 : 297 - 303.

Whitfield, J.M. (1993):

Neonatal sepsis. In : Holbrook PR (eds.). Textbook of pediatric critical care 3rd ed. WB Saunders Company, Philadelphia, London, Toronto, P. 98.

Whitelaw, A. (1990) :

Treatment of sepsis with IgG in very low birth weight infants . Arch . Dis Child .; 65 : 347.

Winston, D.J. ; McGrattan, M.A. and Busuttill, R.W. (1984) :

Imipenem therapy of pseudomonas aeruginosa and other serious bacterial infections. Antimicrob. Agents. Chemother ., 26 : 673 .

Wright SD and Jong M.T.C. (1986) :

Adhesion Promoting reception on human macraphages recagize E-coli by binding to polysaccharides.
J. Exp. Med.; 164:1876-88.

[X]

Xanthou, M.; Xypolyta, A. and Anagnostakis, D. (1975):

Exchange transfusion in severe neonatal infection with sclerema . Arch . Dis . Child , 50 : 901 .

[Y]

Yachie, A.; Takano, N.; Ohta K. et al. (1992) :

Defective Production of IL-6 in very small premature infants in respanse to bacterial Pathogens. infect immunol, 60:749-53.

Yoder, P.R. ; Gibbs, R.S. and Balanco, J.D. (1983) :

A prospective, controlled study of maternal and perinatal outcome after intra- amniotic infection at term . Am . J . Obstet . Gyn ., 145 : 695.

Yoder, P.C. and Polin, R.A. (1986) :

Immunotherapy of neonatal septicemia. Ped. Clin. Nor. Am., 33 (3) : 481.

[Z]

Zipursky, A.; and Jaber, H. (1978) :

The Haematology of bacterial infection in newborn infants. Clin. Haematol., 7 : 175.