

References

- Abd-Alla, M.D. and Ravdin, J.I. (2002):** Diagnosis of amoebic colitis by antigen capture ELISA in patients presenting with acute diarrhoea in Cairo, Egypt. *Trop Med Int Health.*, 7(4):365-70.
- Abd-Alla, M. D.; Jackson,T. F.; Reddy, S., and Ravdin, J. I. (2000):** Diagnosis of invasive amoebiasis by enzyme linked immunosorbent assay of saliva to detect amoebic lectin antigen and antilectin immunoglobulin G antibodies. *J. Clin. Microbiol.*, 38:2344-2347.
- Acuna-Soto, R.J.; Samuelson, P.D.; Girolami, L.; Zarate, F.; Schoolnick,G. and wirth, D. (1993):** Application of polymerase chain reaction to the epidemiology of pathogenic and non pathogenic *E. histolytica*. *Am. J. Trop. Med. Hyg.*, 48:58-70.
- Adams, A.R.D. and Maegraith, B.G. (1980):** Fulminant amebic colitis. *Am. J. Trop. Med. Hyg.*, 22:56-63.
- Adams, S.A; Robson, S.C; and Gathiram, V. (1993):** Immunological similarity between the 170 KD amoebic adherence glycoprotein and human beta 2 integrins *Lancet.*, 341:17-19.
- Ahmad, M.; El-Hady, M. and Morsy, T. (1990):** Parasitic infection and hemoglobin level among school children of different socio-economic classes in Abha, Saudi Arabia. *J. Egypt. Soc. Parasitol.*, 20: 61-67.
- Ali, I. K.; Hossain, M.B.; Roy, S; Ayeh-Kumi, P.F.; Petri, W.A.; Haque, R. and Clark, C.G. (2003):** *Entamoeba*

moshkovskii infections in children, Bangladesh. *Emerg. Infect. Dis.*, 9:580-584.

Arrowood, M.J. (1998): Diagnosis of *Cryptosporidium* and Cryptosporidiosis .Editor: R.Fayer, Publisher, CRC Press, Inc., Boca. Raton, Fla., 43-64.

Ashrat, B. K.; Hossin, H.; and Erag, M. (1997): Assessment of human intestinal parasites in Iran during the last 50 years. Tehran, Islamic Republic of Iran., 137:19–22

Atchley, V.O.; Auerenheimer, A.H. and Wasley, M.A. (1963) Precipitate pattern in agar gel with sera from human amoebiasis and *E. histolytica* antigen. *J. Parasitol.*,249:313.

Ayeh-Kumi, P. F.; Ali, I.M.; L. A. Lockhart, L.A.; C. A. Gilchrist, C.A.; Petri, W.A. and Haque, R. (2001): *Entamoeba histolytica*: genetic diversity of clinical isolates from Bangladesh as demonstrated by polymorphisms in the serine-rich gene. *Exp. Parasitol.*, 99:80-88.

Azam, A.; Paul, J.; and Sehgal, D. (1996): Identification of novel genes from *Entamoeba histolytica* by expressed sequence tag analysis. *Gene.*, 181:113-116.

Barnes, P. F.; DeCock, K.M.; Reynolds, T.N. and Tralla, P.W. (1987): A comparison of amebic and pyogenic abscess of the liver. *Medicine (Baltimore).*, 66:472-483.

Benetton, M.L.; Goncalves, A.V.; Meneghini, M.E.; Silva, E.F. and Carneiro, M. (2005): Risk factors for infection by *Entamoeba histolytica*/*E.dispar* complex: an epidemiological study conducted in outpatients clinics in the city of Manaus,

Amazon Region, Brazil. Trans R Soc. Trop Med Hyg., 99:532-540.

Bercu, T.E.; Petri, W.A.; and Behm, J.W. (2007): Amoebic colitis: new insights into pathogenesis and treatment. Curr Gastroenterol Rep., 9(5):429-433.

Bessa, C.; Braga, M.; Gomes, M.W.; Andrea, S. and Barbara, J. (2001): *Entamoeba histolytica* and *Entamoeba dispar* infections as detected by monoclonal antibody in an urban slum in Fortaleza, North eastern Brazil. Trop., 34: 5437-5443.

Bhattacharya, S.; Bhattacharya, A.; Diamond, L.S. and Soldo, A.T. (1989): Circular DNA of *Entamoeba histolytica* encodes ribosomal RNA. J. Protozool., 36:455-458.

Blessmann, J. and Tannich, E. (2002): Treatment of asymptomatic intestinal *Entamoeba histolytica* infection. N.Engl.J. Med., 347(17): 1384.

Blessmann, J.; Buss, H.; Dinh, B.T.; Ngo, Q.T.; Van, A.L.; Alla, M.D.; Jackson, T.F.; Ravdin, J.I. and Tannich, E. (2002): Real-time PCR for detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* in fecal samples. J. Clin. Microbiol., 40:4413-4417.

Blessmann, J.; Van, A. and Tannich, E. (2006): Epidemiology and treatment of amoebiasis in Hue, Vietnam. Arch. Med. Res., 37:270-2.

Blessmann, J.; Ali, P.A.; Nu, B. T.; Dinh, T. Q.; Viet, A. L.; Van, C.G.; and Tannich, E. (2003): Longitudinal study of intestinal *Entamoeba histolytica* infections in asymptomatic adult adult carriers. J. Clin. Microbiol., 41:4745-4750.

- Bobbi, S.; Pritt, M.D. and Clark, C. G. (2008):** Amoebiasis. Mayo. Clin. Proc., 83(10):1145-1160.
- Britten, D.; Wilson, S.M.; McNerney, R; Moody, A.H.; Chiodini, P.L. and Ackers, J.P. (1997):** An improved colorimetric PCR-based method for detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar*. J. Clin. Microbiol., 35:1108-1111.
- Calderaro, A.; Gorrini, C.; Bommezzadri, S.; Piccolo, G.; Dettori, G. and Chezzi, C. (2006):** *Entamoeba histolytica* and *Entamoeba dispar*: comparison of two PCR assays for diagnosis in a non-endemic setting. Trans. R. Soc. Trop. Med. Hyg., 100:450-457.
- Cantellano, M.E. and Palomo, P.A. (2000):** *Entamoeba histolytica*: Mechanism of surface receptor capping. Exp. Parasitol., 79: 424-35.
- Casemore, D.P; Sands, R.T. and Curry, A. (1985):** Cryptosporidium species a new human pathogen. J.Clin. Pathol., 38:1321-1326.
- Cavalier-Smith, T. (1998):** A revised six-Kingdom system of life. Biol. Rev. Camb. Philos. Soc., 73:203-266.
- Cheesbrough, M. (1987):** Medical laboratory manual for tropical countries. Tropical Health Technology, Butterworth-Heinemann., 88:365-389.
- Cheesbrough, M. (1991):** Techniques used to identify parasite In: Medical Laboratory Manual for Tropical Countries. Britain, Butterworth-Heinemann Ltd., 175-185.

- Chien, A. ; Edgar, D.B. and Trela, J.M. (1976):** "Deoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*". J. Bacteriol., 174: 1550–1557.
- Clark, C. G. and Diamond, L.S. (2002):** Methods for cultivation of luminal parasitic protists of clinical importance. Clin. Microbiol. Rev., 15:329-341.
- David, M.; Cook, R. C.; Dennis, S. L. and Gary, M. (2009):** A Retrospective Analysis of Prevalence of Gastrointestinal Parasites among School Children in the Palajunoy Valley of Guatemala. J. Health. Popul. Nutr., 27(1):31-40.
- De Leon, A. (1970):** Pronostico tardio en el absceso hepatico amibiano. Arch Invest Med., 1:205-206.
- De Long, E.R.; De Long, D.M. and Clark- Pearson, D.L. (1988):** Comparing the areas under two or more correlated Receiver Operating Characteristic Curves: A non Parametric Approach, Biometrics, 44, 837-845.
- Diamond, L. S. (1982):** A new liquid medium for xenic cultivation of *Entamoeba histolytica* and other lumen dwelling protozoa. J. Parasitol., 68:958-959.
- Diamond, L. S. and Clark, C.G. (1993):** A redescription of *Entamoeba histolytica* separating it from *Entamoeba dispar*. J. Eukaryot. Microbiol., 40:340-344.
- Edman, U.; Meraz, M.A.; S. Rausser, S.; Agabian, N. and Meza, I. (1990):** Characterization of an immuno-dominant variable surface antigen from pathogenic and nonpathogenic *Entamoeba histolytica*. J. Exp. Med., 172:879-888

- Erlich, H.A.; Gelfand, D. and Sninsky, J.J. (1991):** Recent advances in polymerase chain reaction. *Science*, 252(5013):1643-51.
- Evangelopoulos, A.; Spanakos, G.; Patsoula, E.; Vakalis, N. and Legakis, N.A. (2000):** A nested, multiplex, PCR assay for the simultaneous detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* in faeces. *Ann. Trop. Med. Parasitol.*, 94:233-240.
- Fernades, H.; Clement, R.S.; Dsouza, G.K.; Swethadri, C.N. and Naik, R. (2009):** Ameboma of the colon with amoebic liver abscess mimicking colon cancer. *Indian. Journal of Parasitology And Microbiology.*, 52(2):122-124.
- Fotedar, R.; Stark, D.; Beebe, N.; Marriott, D.; Ellis, J. and Harkness, J. (2007):** PCR detection of *Entamoeba histolytica*, *Entamoeba dispar*, and *Entamoeba moshkovskii* in stool samples from Sydney, Australia. *J. Clin. Microbiol.*, 45:1035-1037.
- Franklin, A. and Harolad, W. (1994):** *Basic Clin. Parasitol.*, 21, 22
- Freitas, M. A.; Vianna, E.N.; Martins, A.S.; Silva, E.F.; Pesquero, J.I. and Gomes, M.A. (2004):** A single step duplex PCR to distinguish *Entamoeba histolytica* from *Entamoeba dispar*. *Parasitology.*, 128:625-628.
- Furrows, S. J.; Moody, A.H. and Chiodini, P.L. (2004):** Comparison of PCR and antigen detection methods for diagnosis of *Entamoeba histolytica* infection. *J. Clin. Pathol.*, 57:264-266.

- Garcia, L.S. and Bruckner, D.A. (1997):** Intestinal Protozoa Flagellates and Ciliates. In: Diagnostic Medical Parasitology, 2nd ed., 13-39.
- Gatti, S.; Swierczynski, G.; Robinson, F.; Anselmi, M.; Corrales, J.; Moreira, J.; Montalvo, G.; Bruno, A.; Maserati, R.; Bisoffi, Z. and Scaglia, M. (2002):** Amoebic infections due to the *Entamoeba histolytica*-*Entamoeba dispar* complex: a study of the incidence in a remote rural area of Ecuador. Am. J. Trop. Med. Hyg., 67:123-127.
- Ghosh, S.; Field, J.; and Rogers, R. (2000):** The *Entamoeba histolytica* mitochondrion derived organelle (crypton) contains double-stranded DNA and appears to be bound by a double membrane. Infect. Immune., 68:4319-4322.
- Gill, G.V.; and Beeching, N.J. (2004):** Tropical Medicine. 5th ed. Oxford, UK: Blackwell Publishing., 153-159.
- Goncalves, M. L.; da Silva, V.L.; de Andrade, C.M.; Reinhard, K.; da Rocha, G.C.; Le Bailly, M.; Bouchet, F.; Ferreira, L.F. and Araujo, A. (2004):** Amoebiasis distribution in the past: first steps using an immunoassay technique. Trans. R. Soc. Trop. Med. Hyg., 98:88-91.
- Gonin, P. and Trudel, L. (2003):** Detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* isolates in clinical samples by PCR and enzyme-linked immunosorbent assay. J. Clin. Microbiol., 41:237-241.
- Gonzales, M.; Leonila, F. D. and Elizabeth, G.M. (2006):** Antiamoebic drugs for treating amoebic colitis. Arch. Med. Res., 27:250-3.

- Gonzalez, C. R. ; Isibasi, A. ; Ortiz-Navarrete, V. ; Paniagua, J. ; Garcia, J.A. ; Ramirez, A. ; Salvatierra, B. ; Tapia, R. ; Sepulveda, J. and Gutierrez, G. (1995):** Prevalence of antibodies against *Entamoeba histolytica* in Mexico measured by ELISA. *Epidemiol. Infect.*, 115:535-543.
- Hamzah, Z.; Petmitr, S.; Mungthin, M.; Leelayoova, S. and Chavalitsheewinkoon, P. (2006):** Different detection of *Entamoeba histolytica* and *Entamoeba dispar* by a single- round PCR assay *J. Clin. Microbiol.*, 44:3196-3200.
- Haque, R. and Petri. W.A. (2006):** Diagnosis of amebiasis in Bangladesh. *Arch. Med. Res.*, 37:273-276.
- Haque, R.; Neville, L.M.; Hahn, P. and Petri, J.r. (1995):** Rapid diagnosis of *Entamoeba* infection by using *Entamoeba histolytica* stool antigen detection kits *J.Clin. Microbiol.*, 33:2558-2561.
- Haque, R.; Faruque, A.S.; Hahn, P.; Lyerly, D. and Petri, W.A. (1997):** *Entamoeba histolytica* and *Entamoeba dispar* infection in children in Bangladesh. *J. Infect. Dis.*, 175:734-736.
- Haque, R.; Mollah, N.U.; Ali, I.K.; Alam, K.; Eubank, A.; Luerly, D. and Petri, W.A. (2000):** Diagnosis of amebic liver abscess and intestinal infection with the TechLab *Entamoeba histolytica* II antigen detection and antibody tests. *J. Clin. Microbiol.*, 38:3235-3239.
- Haque, R.; Huston, C.D. and Hughes, M. (2003):** Amebiasis. *N Engl. J. Med.*, 348:1565.
- Haque, R.; Mondal, D. and Duggal, P. (2006):** *Entamoeba histolytica* infection in children and protection from subsequent amoebiasis. *Infect. Immune.*, 74:904-909.

- Heckendorn, R.; N’Goran, E.K.; Felger, I.; Vounatsou, P.; Yapi, A.; Oettli, A.; Marti, H.P.; Dobler, M.; Traore, M.; Lohourignon, K.L. and Lengeler, C. (2002):** Species-specific field tests of *Entamoeba* spp. in an area of high endemicity. *Trans. R. Soc. Trop. Med. Hyg.*, 97:305–307.
- Helmy, M.M.; Rashed, L.A. and Abdel-Fattah, H.S. (2007):** Detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* isolates in clinical samples by PCR. *J. Egypt. Soc. Parasitol.*, 37(1):257-74.
- Higgins, J. A.; Fayer, R.; Trout, J.M.; Xiao, L.; Lal, A.A.; Kerby, S. and Jenkins, M.C. (2001):** Real-time PCR for the detection of *Cryptosporidium parvum*. *J. Microbiol.*, 47:323-337.
- Hira, P. R.; Iqbal, J.; Al-Ali, F.; Philip, R.; Grover, S.; D’Almeida, E. and Al-Eneizi, A.A. (2001):** Invasive amebiasis: challenges in diagnosis in a non-endemic country (Kuwait). *Am. J. Trop. Med. Hyg.*, 65:341-345.
- Holland, J. L.; Louie, L.; Simor, A.E. and Louie, M. (2000):** PCR detection of *Escherichia coli* directly from stools: evaluation of commercial extraction methods for purifying fecal DNA. *J. Clin. Microbiol.*, 38:4108-4113.
- Hooshyar, H.; Rezaian, M.; Mahmood, M.; Farnial, S. and Solaymani, M. (2004):** A field study of the distribution of *Entamoeba histolytica*/ *dispar* cyst passers in Northern, Central, and Southern Iran. *Iranian J. Publ. Health.*, 33(2):28-32.
- Hung, C. C.; Deng, H.Y.; Hsiao, W.H.; Hsieh, S.M.; Hsiao, C.F.; Chen, M.Y. and Chang, S.C. (2005):** Invasive amebiasis as an emerging parasitic disease in patients with human

immunodeficiency virus type 1 infection in Taiwan. Arch. Intern. Med., 165:409-415.

Hunter, G.; Swartzwelder, J.C. and Clyde, D.F. (1976): Trop. Med. Hyg., 34:261-268.

Huston, C. D. and Petri, J.r. (1999): Amebiasis: clinical implications of the recognition of *Entamoeba dispar*. Curr. Infect. Dis. Rep., 1:441-447.

Jackson, T. F.; Anderson, C.B. and Simjee, A.E. (1984): Serological differentiation between past and present infection in hepatic amoebiasis. Trans. R. Soc. Trop. Med. Hyg., 78:342-345.

Jaco, J.; Roy, A.; Eric. A.T.; Brienens, T.; Marianne, A. and Anton, M. (2004): Simultaneous detection of *Entamoeba histolytica*, *Giardia Lamblia* and *Cryptosporidium Parvum* in fecal samples by using multiplex real time PCR. Journal of Clinical Microbiology., 1220-1223.

Josef, S. and David, W. R. (2001): In vitro Amplification of DNA by the Polymerase Chain Reaction. Trends Mol. Med., 6:334-338.

Julie, L.; Kirstin, E. and Nick, S. (2009): Multiplex- PCR: Current Technology and Applications. Trend. Mol. Med., 3:221-229.

Katzenstein, D.; Richerson, V. and Braude, A. (1982): New concept of amoebic liver abscess derived from hepatic imaging serodiagnosis, and hepatic enzymes in 67 consecutive cases in San Diego. Medicine., 61: 237.

Kebede, A.; Verweij, J.J.; Endeshaw, T.; Messele, T.; Tasew, G.; Petros, B. and Polderman, A.M. (2004): The use of real-time PCR to identify *Entamoeba histolytica* and *E. dispar*

- infections in prisoners and primary-school children in Ethiopia. Ann. Trop. Med. Parasitol., 98:43-48.
- Kellogg, D.E.; Rybalkin, I.; Chen, S.; Mukhamedova, N.; Vlasik, T.; Siebert, P.D. and Chenchik, A. (2002):** PCR facilitated by a neutralizing monoclonal antibody directed against Taq DNA polymerase. Biotechniques., 16(6):1134-7.
- Kelsall, B.L. and Ravdin, J.I. (1993):** Degradation of human IgA by *Entamoeba histolytica* J. Infect. Dis., 168:1319-1322.
- King, M. (1975):** A medical laboratory for developing countries; ELBS, ELBS, Oxford University Press.vol;15: 45-55.
- Knapp, R.G. and Miller, M.C. (1992):** Clinical Epidemiology and biostatistics. National Medical Series from Williams and Wilkins, Baltimore, Hong Kong, London, Sydney, Egyptian Edition. P:209.
- Lawyer, F.C.; Stoffel, S.; Saiki, R.K.; Chang, S.Y.; Landre, P.A.; Abramson, R.D. and Gelfand, D.H. (1993):** "High-level expression, purification, and enzymatic characterization of full-length *Thermus aquaticus* DNA polymerase and a truncated form deficient in 5' to 3' exonuclease activity". PCR Methods Appl., 2: 275–287.
- Lebbad, M. and Svard S.G.. (2005):** PCR differentiation of *Entamoeba histolytica* and *Entamoeba dispar* from patients with amoeba infection initially diagnosed by microscopy. Scand. J. Infect. Dis., 37:680-685.
- Leippe, M.; Bruhn, H. and Hecht, O. (2005):** Ancient weapons: the three dimensional structure of amoebapore A.. Trends. Parasitol., 21:5-7.

- Leon, G.; Fiori, C. and Das, P. (1997):** Electron Probe analysis and biochemical characterization of electron dense granules secreted by *Entamoeba histolytica* Mol. Biochem. Parasitol., 85:233-242.
- Li, E. and Stanley, J.r. (1996):** Protozoa. Amoebiasis. Gastroenterol. Clin. N. Am., 25:471-492.
- Lichtenstein, A.; Kondo, A.T.; Visvesvara, G.S.; Fernandez, A.; Paiva, E.F.; Mauad, T.; Dolhnikoff, M. and Martins, M.A. (2005):** Pulmonary amoebiasis presenting as superior vena cava syndrome. Thorax., 60:530-2.
- Logan, J.; Edwards, K. and Saunders, N. (2009):** Real-Time PCR: Current Technology and Applications. Nat. Protoc., 2:655-672.
- Lohourignon, N. and Lengeler, C. (2004):** Species-specific field tests of *Entamoeba* spp. in an area of high endemicity. Trans. R. Soc. Trop. Med. Hyg., 97:305–307.
- Lottre, H.; Russmann, H.; Heesemann, J. and Tannich, E. (2008):** Attenuated recombinant yersinia as live oral vaccine carrier to protect against amoebiasis. Int. J. Med. Microbiol., 12:79-86.
- Lucas, R. and Upcroft, J.A. (2001):** Clinical significance of the redefinition of the agent of amoebiasis. Rev. Latinoam. Microbiol., 43(4):183-187.
- Magana, M.L.; Fernandez, D .J.; and Magana, M. (2008):** Cutaneous amoebiasis in pediatrics. Arch Dermatol., 144(10):1369-72.
- Mansour, N.S.; Youssef, F.G.; Mikhail, E.M. and Mohamed, E.W. (1997):** Amoebiasis in schistomiasis endemic and non endemic areas of Egypt. J.Egypt. Soc. Parasitol., 27:617-628.

- Martinez, P. A. (1982):** The biology of *Entamoeba histolytica*. J Parasitol., 54:1047-1056.
- Millan, A. and Neillage, G.J. (1994):** Comparison of the sensitivity of microscopy and culture in the laboratory diagnosis of intestinal protozoal infection. J Clin Pathol., 37:809-811.
- Mirelman, D.; Nuchamowitz, Y. and Stolarsky, T. (1997):** Comparison of use of enzyme-linked immunosorbent assay-based kits and PCR amplification of rRNA genes for simultaneous detection of *Entamoeba histolytica* and *E. dispar*. J. Clin. Microbiol., 35:2405-2407.
- Misra, S.P.; Misra, V. and Dwivedi, M. (2006):** Ileocaecal masses in patients with amoebic liver abscess: Etiology and management. World. J. Gastroenterol., 12:1933-1936.
- Monteiro, L.; Bonnemaïson, D.; Vekris, A.; Petry, K.G.; Bonnet, Vidal, J.R.; Cabrita, J. and Megraud, F. (1997):** Complex polysaccharides as PCR inhibitors in feces: *Helicobacter pylori* model. J. Clin. Microbiol., 35:995-998.
- Moran, P. ; Ramos, F. ; Ramiro, M. ; Curiel, O. ; Gonzalez, E. ; Valadez, A. ; Gomez, A. ; Garcia, G. ; Melendro, E.I. and Ximenez, C. (2005):** Infection by human immunodeficiency virus-1 is not a risk factor for amoebiasis. Am. J. Trop. Med. Hyg., 73:296-300.
- Morimoto, N.; Komatsu, C.; Nishida, M. and Sugiura, T. (2000):** Detection of *E. histolytica* cyst in non fixed long term stored human faeces by direct immunofluorescent stain. J. Infect. Disease., 54(2): 72-74.

- Nolan, T.; Hands, R.E. and Bustin, S.A. (2006):** "Quantification of mRNA using real-time RT-PCR". Nat. Protoc., 1: 1559–1582.
- Nunez, Y. O.; Fernandez, M.A.; Torres, N. D.; Silva, J. A.; Montano, I.; Maestre, J. L. and Fonte, L. (2001):** Multiplex polymerase chain reaction amplification and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* DNA from stool samples. Am. J. Trop. Med. Hyg., 64:293-297.
- Nyenke, C.; Chukwujekwu, D. C.; Stanley, h. and Awoibi, N. K. (2008):** Prevalence of Intestinal Amoebiasis in Infant and Junior School Children in Degema . J. Appl. Sci. Environ. Manage., 12(3): 83 – 87.
- Paglia, M. G. and Visca. P. (2004):** An improved PCR-based method for detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* in formalin-fixed stools. Acta .Trop. 92:273-277.
- Patterson, M. and Schoppe, L.E. (1982):** The presentation of amoebiasis. Med. Clin. N. Am., 66:689-705.
- Paul, J.; Srivastava, S. and Bhattacharya, S. (2007):** Molecular methods for diagnosis of *Entamoeba histolytica* in a clinical setting: an overview. Exp Parasitol., 116(1):35-43.
- Pavlov, A.R.; Pavlova, N.V.; Kozyavkin, S.A.; and Slesarev, A.I. (2004):** "Recent developments in the optimization of thermostable DNA polymerases for efficient applications". Trends Biotechnol., 22: 253–260.
- Petri, W.A. (1996):** Recent advances in amebiasis. Crit Rev Clin Lab Sci., 33(1):1-37.

- Petri, W.A. and Singh, U. (1999):** Diagnosis and management of amoebiasis. Clin Infect Dis., 29(5):1117-25.
- Petri, W.A. and Singh, U. (2006):** Enteric Amoebiasis: Principles, Pathogens, & Practice. 2nd ed. Philadelphia, PA: Elsevier Churchill Livingstone; P. 967-983.
- Pillai, D. R., Keystone, J.S.; Sheppard, D.C.; MacLean, J.D.; MacPherson, D.W. and Kain, K.C. (1999):** *Entamoeba histolytica* and *Entamoeba dispar*: epidemiology and comparison of diagnostic methods in a setting of non endemicity. Clin. Infect. Dis., 29:1315-1318.
- Prathap, K. and Gilman, R. (1970):** The histopathology of acute intestinal amebiasis. A rectal biopsy study. Am. J. Pathol., 60:229–245.
- Pritt, B. S. and Clark, C.G. (2002):** Amoebiasis. Mayo. Clin. Pro., 83(10): 1154-9.
- Proctor, E. M. (1991):** Laboratory diagnosis of amebiasis. Clin. Lab. Med., 11:829-859.
- Que, X. and Reed, S.L. (1991):** Nucleotide sequence of a small subunit ribosomal RNA (16S-like rRNA) gene from *Entamoeba histolytica*: differentiation of pathogenic from non-pathogenic isolates. Nucleic Acids Res., 11:5438.
- Que, X. and Reed, S.L. (2000):** Cysteine Proteinases and the pathogenesis of amebiasis. Clin. Microbiol. Rev., 13:196-206.
- Qvarnstrom, Y.; James, C.; Xayavong, M.; Holloway, B.P.; Visvesvara, G.S.; Sriram, R. and Da Silva, A.J. (2005):** Comparison of real-time PCR protocols for differential laboratory diagnosis of amebiasis. J. Clin. Microbiol., 43:5491-5497.

- Rachid, H.; Alaoui Yazidi, A.; Loudadssi, F.; El Biaze, M.; Bakhatar, A.; Yassine, N.; El Meziane, A. and Bahlaoui, A. (2005):** Amoebic infections of the lung and pleura. *Rev Mal Respir.*, 22:1035-7.
- Ramos, F.; Moran, P.; Gonzalez, E.; Garcia, G.; Ramiro, M.; Gomez, A.; Leon M.C.; Melendro, E.I.; Valadez, A. and Ximenez, C. (2005):** *Entamoeba histolytica* and *Entamoeba dispar*: prevalence infection in a rural Mexican community. *Exp. Parasitol.*, 110:327-330.
- Ramos, F.; Valdez, E. and Morán, P. (2000):** Prevalence of *Entamoeba histolytica* and *Entamoeba dispar* in a Highly Endemic Rural Population. *Arch. Med. Res.*, 31:534-5.
- Ramos, F.; Zurabian, R.; Moran, P.; Ramiro, M.; Gomez, A.; Clark, C.G.; Melendro, E.I.; Garcia, G. and Ximenez, C. (1999):** The effect of formalin fixation on the polymerase chain reaction characterization of *Entamoeba histolytica*. *Trans. R. Soc. Trop. Med. Hyg.*, 93:335-336.
- Ravdin, J.I. (1995):** amoebiasis. *Clin. Infect Dis.*, 20:1453-1466.
- Ravdin, J.I. (2000):** Amoebiasis. Imperial College Press. London., 3:47-66.
- Ravdin, J.I. and Stauffer, W.M. (2005):** *Entamoeba histolytica* (Amoebiasis): Principles and Practice of Infectious Diseases. 6th ed. Philadelphia, Pa: Churchill Livingstone; 3097-3111.
- Ravdin, J. I.; Jackson, T.F.; Petri, W.A.; Murphy, C.F.; Ungar, B.L.; Gathiram, V.; Skilogiannis, J. and Simjee, A.E. (1990):** Association of serum antibodies to adherence lectin with

invasive amebiasis and asymptomatic infection with pathogenic *Entamoeba histolytica*. J. Infect. Dis., 162:768-772

Read, S.J.; Mitchell, J.L., and Fink, C.G. (2001): LightCycler multiplex PCR for the laboratory diagnosis of common viral infections of the central nervous system. J. Clin. Microbiol., 39(9):3056-9.

Redondo, R.B.; Laura, G.; Martinez, M. and George, A. (2004): *Entamoeba histolytica* and *Entamoeba dispar*: differentiation by Enzyme-Linked Immunosorbent (ELISA) and its clinical correlation in pediatric patients. Rev. inf. dis., 238:239.

Reed, S.L.; Ember, J.A. and Herdman, D.S. (1995): The extracellular neutral cysteine proteinase of *Entamoeba histolytica* degrades anaphylatoxins C3a and C5a. J. Immunol., 155:266-274.

Rickert, A.M.; Lehrach, H. and Sperling, S. (2004): Multiplexed real-time PCR using universal reporters. Clin Chem., 50(9):1680-3.

Rivera, W. L.; Tachibana, H. and Kanbara, H. (1998): Field study on the distribution of *Entamoeba histolytica* and *Entamoeba dispar* in the northern Philippines as detected by the polymerase chain reaction. Am. J. Trop. Med. Hyg., 59:916-921.

Rivera, W. L.; Santos, S.R. and Kanbara, H. (2006): Prevalence and genetic diversity of *Entamoeba histolytica* in an institution for the mentally retarded in the Philippines. Parasitol. Res., 98:106-110.

- Rivera, W.L.; Yason, J.A. and Adao, D.E. (2010):** *Entamoeba histolytica* and *E. dispar* infections in captive macaques (*Macaca fascicularis*) in the Philippines. *J. Parasitol.*, 51(1):69-74
- Roberts, L.S. and Janovy, J.J.R. (2002):** Foundations of parasitology 5th ed. W.M.C.Brown publ., 82-86.
- Romero, J. L.; Descoteaux, S.; Reed, S.; Orozco, E.; J. Santos, J. and Samuelson, J. (1992):** Use of polymerase chain reaction and nonradioactive DNA probes to diagnose *Entamoeba histolytica* in clinical samples. *Arch. Med. Res.*, 23:277-279.
- Roy, S.; Kabir, M.; Mondal, D.; Ali, I.K.; Petri, W.A. and Haque, R. (2005):** Real-time-PCR assay for diagnosis of *Entamoeba histolytica* infection. *Clin. Microbiol.*, 43:2168-2172.
- Ryan, K.J. and Ray, C.G. (2004):** Epidemiology of amoebiasis. *Sherris Medical Microbiology* (4th ed. Ed.). Mc.Graw Hill., 733-8
- Rychlik, W.; Spencer, W.J. and Rhoads, R.E. (1990):** "Optimization of the annealing temperature for DNA amplification in vitro". *Nucl. Acids Res.*, 18: 6409–6412.
- Sails, A.D. (2009):** "Applications in Clinical Microbiology". *Real-Time PCR. Nucleic Acids Res.*, 13:4635.
- Saklatvala, T. (1993):** Milestones in parasitology. *Parasitol.*, 9:347-348.
- Salles, J.M.; Moraes, L.A. and Salles, M.C. (2003):** Hepatic amebiasis. *Braz J. infect Dis.*, 7 (2):96-110.
- Sanchez, Giillen, M. C.; Velazquez-Rojas, M.; Salgado, R. H.; Torres, R.E.; Perez, F.R.; Martinez, M.J. and Talamas, R.P. (2000):** Seroprevalence of anti-*Entamoeba histolytica* antibodies by IHA and ELISA assay in blood donors from Puebla, Mexico. *Arch. Med. Res.*, 31:553-554.

- Sayyari, A.A.; Imanzadeh, F.; Bagheri, S.A.; Karami¹, H. and Yaghoobi, M. (2005):** prevalence of intestinal parasitic infections in the Islamic Republic of Iran. *J. Parasitol.*, 54-61.
- Sebastiaan, J.; Damien, J.; Fotedar, R.; Marriott, D.; John, T. and Jock, L. (2007):** Amoebiasis: Current status in Australia. *MJA.*, 186(8): 412-416.
- Sepulveda, B.; Lee, E.; Torre, M. and Landa, L. (1971):** EL diagnostico serologico de La amoebiasis invasara con La Techica de immunoelectroforesis cruzada. *Arch. de Invest. Medica.*, 2:263.
- Seydel, K.B.; Li, E. and Swanson, P.E. (1997):** Human intestinal epithelial cells produce proinflammatory cytokines in response to infection in a SCID mouse-human intestinal xenograft model of amebiasis . *Infect Immune.*, 65:1631-1639.
- Shaikh, G. S.; Begum, R.; Hussain, A. and Shaikh, R. (2009):** prevalence of intestinal protozoan and helminth parasites in sukkur, sindh. *Sindh univ. Res. Jour.*, 41 (2) :53-58.
- Shamsuzzaman, S.M. and Hashiguchi, Y. (2002):** Thoracic amoebiasis. *Clin Chest Med.*, 23:479-92.
- Shamsuzzaman, S.M.; Haque, R.; Hasin, S.K. and Hashiguchi, Y. (2000):** Evaluation of indirect fluorescent antibody test and enzyme-linked immunosorbent assay for diagnosis of hepatic amoebiasis in Bangladesh. *J. Parasitol.*, 86:611-615.
- Shandera, W. X.; Bollam, P.; Hashmey, R.H.; Athey, P.A.; Greenberg, S.B. and White, A.C. (1998):** Hepatic amoebiasis among patients in a public teaching hospital. *South Med. J.*, 91:829-837.

- Sharkey, E.R.; Scalice, K.G.; Christy, J.r.; Atwood, S.M. and Daiss J.L. (1994):** "Antibodies as Thermolabile Switches: High Temperature Triggering for the Polymerase Chain Reaction". Bio/Technology., 12: 506–509.
- Sharp, S. E.; Suarez, C.A.; Duran, Y. and Poppiti, R.J. (2001):** Evaluation of the Triage Micro Parasite Panel for detection of *Giardia lamblia*, *Entamoeba histolytica*/*Entamoeba dispar*, and *Cryptosporidium parvum* in patient stool specimens. J. Clin. Microbiol., 39:332-334.
- Sheiban, F. and Rezaian, M. (1998):** Study on intestinal protozoa in seven villages of Bandar- Abbas. Iranian. J. Publ. Healt., 10:45-55.
- Spackman, E. and Suarez, D.L. (2008):** "Type A influenza virus detection and quantitation by real-time RT-PCR". Mol. Biol., 436: 19–26.
- Stanley, S. L.; Becker, A.; Jenkins, C.K.; Foster, L. and Li, E. (1990):** Cloning and expression of a membrane antigen of *Entamoeba histolytica* possessing multiple tandem repeats. Proc. Natl. Acad. Sci. USA. 87:4976-4980.
- Stanley, S.L. (2003):** Amoebiasis. The Lancet., 361: 1025-1034.
- Stark, S.; Fotedar, R.; Butcher, A.; Marriott, D.; Elis, L. and Harkness, J. (2008):** Comparison of stool antigen detection kits to PCR for diagnosis of amoebiasis . J. Clin Microbiol., 46:1678-1681.
- Stauffer, W. and Ravdin, J.I. (2003):** *Entamoeba histolytica*: an update. Curr Opin Infect Dis., 16(5):479-485.

- Stauffer, W.; Abd-Alla, M. and Ravdin, J.I. (2006):** Prevalence and incidence of *E. histolytica* infection in South Africa and Egypt. Arch Med Res., 37:266-9.
- Takahashi, T.; A. Gamboa, D.A.; Gomez, T.J.; Remes, J.M.; Gonzalez, D.M.; Morales, J.C.; Granados, J. and J. Sierra, M. (1997):** Fulminant amoebic colitis: analysis of 55 cases. Dis. Colon Rectum., 40:1362-1367.
- Tan, W.; Wang, K. and Drake, T.J. (2004):** Molecular beacons. Curr Opin Chem Biol., 8(5):547-53.
- Tannich, E. and Burchard, G.D. (1991):** Differentiation of pathogenic from nonpathogenic *Entamoeba histolytica* by restriction fragment analysis of a single gene amplified in vitro. J. Clin. Microbiol., 29:250-255.
- Tanyuksel, M. and Petri, W.A. (2003):** Laboratory diagnosis of amoebiasis. Clin. Microbiol. Rev., 16:713-729.
- Taysser, A.; Iman, Y.M. And Youssef, F.G. (1997):** Epidemiology of amoebiasis. J. Egypt. Soc. Parasitol., 27 (1):291-297.
- Thornton, S.A.; West, A.H. and Dupont, H.L. (1992):** Comparison of methods for identification of *E.histolytica* Am. J. Clin. Pathol., 80:755-757.
- Tran, V.Q.; Herdman, D.S. and Torian, B.E. (1998):** The neutral cysteine proteinase of *Entamoeba histolytica* degrades IgG and prevents its binding . J. Infect. Dis., 177:508-511.
- Troll, H.; Marti, H. and Weiss, N. (1997):** Simple differential detection of *Entamoeba histolytica* and *Entamoeba dispar* in fresh stool specimens by sodium acetate-acetic acid-formalin concentration and PCR. J. Clin. Microbiol., 35:1701-1705.

- Urdaneta, H.; De Donato, M.; García, A. and Mora, L. (2008):** Epidemiologic and molecular study of *Entamoeba histolytica* and *Entamoeba dispar* strains in patients with diarrhea in Cumana, Sucre state, Venezuela. Invest Clin.;49(2):225-37.
- Van Guilder, H.D.; Vrana, K.E. and Freeman, W.M. (2008):** "Twenty-five years of quantitative PCR for gene expression analysis". Biotechniques 44: 619–626.
- Van hal , S.; Stark, D.J. Fotedar, R.; Marriott, D.; Ellis, J.T. and Harkness, J. (2007):** Amebiasis current status in Australia . Med .J. Aust., 186:412-416.
- Verweij, J. J.; Vermeer, J.; Brien, E.A.; Blotkamp, C.; Laeijendecker, D.;Van Lieshout, L. and Polderman, A.M. (2003):** *Entamoeba histolytica* infections in captive primates. Parasitol. Res., 90:100-103.
- Verweij, J. J.; Blange, R.A.; Templeton, K.; Schinkel, J.; Brien, E.A.; Rooyen, M.A.; Lieshout, L. V. and Polderman, A.M. (2004):** Simultaneous detection of *Entamoeba histolytica*, *Giardia lamblia*, and *Cryptosporidium parvum* in fecal samples by using multiplex real-time PCR. J. Clin. Microbiol., 42:1220-1223.
- Verweij, J. J.; Blotkamp, J.; Brien, E.A.; Aguirre, A. and Polderman, A.M. (2000):** Differentiation of *Entamoeba histolytica* and *Entamoeba dispar* cysts using polymerase chain reaction on DNA isolated from faeces with spin columns. Eur. J. Clin. Microbiol. Infect. Dis., 19:358-361.

- Vet, J.A. and Marras, S.A. (2005):** Design and optimization of molecular beacon real-time polymerase chain reaction assays. *Methods Mol Biol.*, 288:273-90.
- Visser, L. G.; Verweij, J.J.; Esbroeck, M.V.; Edeling, W.M.; Clerinx, J. and Polderman, A.M. (2006):** Diagnostic methods for differentiation of *Entamoeba histolytica* and *Entamoeba dispar* in carriers: performance and clinical implications in a non-endemic setting. *Int. J. Med. Microbiol.*, 296:397-403.
- Walsh, J.A. (1996):** Problems in recognition and diagnosis of amoebiasis. Estimates of global magnitude and mortality. *Rev Infect Dis.*, 8: 228-232.
- WHO. (1991):** Basic laboratory methods in medical parasitology. Reagents and solutions and their preparation., 100-105.
- WHO. (1997):** A consultation with experts on amoebiasis: Mexico City, Mexico 28-29. *Epidemiol Bull.*, 18(1):13-14.
- Williams, J.F. (2001):** Optimization strategies for the polymerase chain reaction. *Biotechniques.*, 7(7):762-9.
- Yassin, M.M.; Shubair, M.E.; Al-Hindy, A.I. and Jadallah, S. (1999):** Prevalence of intestinal parasites among school children in Gaza City. *J. Egypt. Soc. Parasitol.*, 29(2):365-373
- Zaher, S.M. (1993):** Laboratory diagnosis of infection with *C. parvum*. M. Sc. Thesis, Faculty of Medicine, Zagazig University.
- Zaman, S., Khoo, J.; Ng, S.W.; Ahmed, R.; Khan, M.A.; Hussain, R. and Zaman, V. (2000):** Direct amplification of *Entamoeba histolytica* DNA from amoebic liver abscess pus using polymerase chain reaction. *Parasitol. Res.*, 86:724-728.

- Zengzhu, G.; Bracha, R.; Nuchamowitz, Y.; Cheng, W.; and Mirelman, D. (1999):** Analysis by enzyme-linked immunosorbent assay and PCR of human liver abscess aspirates from patients in China for *Entamoeba histolytica*. J. Clin. Microbiol., 37:3034-3036.
- Zindrou, S.; Orozco, E.; Linder, E.; Tellez, A.; and Bjorkman, A. (2001):** Specific detection of *Entamoeba histolytica* DNA by hemolysin gene targeted PCR. Acta. Trop., 78:117-125.
- Zurauskas, J.P. and Mc Bride, W.J. (2001):** Case of amoebic liver abscess: Prolonged latency or acquired in Australia? Intern. Med. J., 31:565-566.