




References

References

Alabdulkareem, A.S.; Abahussein, A.A and Okoro, A.
(1998):

Severe alopecia areata treatment with systemic corticosteroids.

Int. J. Dermatol., 37(8): 622.

Arndt, K.A. (1986):

Definition of alopecia areata.

Manual of dermatologic therapeutic, 2: 17.

Arrazola, J.; Sendagorta, E. and Harto, A. (1987):

Treatment of alopecia areata with topical nitrogen mustard.

Int. J. Dermatol., 123: 165.

Ashworth, J.; Tuyp, E. and Macke, R.M. (1989):

Allergic and irritant contact dermatitis compared in the treatment of alopecia totalis and universalis.

Br. J. Dermatol., 120: 397.

Bachrach, V.H. (1998):

Sulfasalazine: a historical perspective.

Am. J. Gastroentol., 3: 487-93.

- Behrens-Williams, S.C.; Leiter, U.; Schiener, R.; Weidmann, M.; Peter, R.U. and Kerscher, M. (2001):
The PUVA-turban as a new option of applying a dilute psoralen solution selectively to the scalp of patients with alopecia areata.
J. Am. Acad. Dermatol., 44(2): 248.
- Bernardo, O.; Tang, L.; Lui, H.; Shapiro, J. (2003):
Topical nitrogen mustard in the treatment of alopecia areata: a bilateral comparison study.
J. Am. Acad. Dermatol., 49(2):291-4.
- Berth-Jones, J.; McBurney, A. and Hutchinson, P.E. (1994):
Diphencyprone is not detectable in serum or urine following topical application.
Acta. Derm. Venereol., 74(4): 312.
- Bissonnette, R.; Shapiro, J.; Zeng, H.; Mclean, D.I. and Lui, H. (2000):
Topical photodynamic therapy with 5-aminolaevulinic acid does not induce hair regrowth in patients with extensive alopecia areata.
Br. J. Dermatol., 143: 1032.
- Bodemer, C.; Peuchmaur, M.; Fraitaig, S.; Chatenoud, L.; Brousse, N. and De Prost, Y. (2000):
Role of cytotoxic T cells in chronic alopecia areata.
J. Invest. Dermatol. 114:112-116.

- Bolduc, C. and Shapiro, J. (2000):
DPCP for the treatment of alopecia areata.
Skin Therapy Lett., 5(5): 3.
- Box, S.A.; Pullar, T. (1997):
Sulphasalazine in the treatment of rheumatoid
arthritis.
Br. J. Rheumatol., 36: 382-6.
- Buckley, D.A. and Vivier, A.W. (2001):
The therapeutic use of topical contact sensitizers in
benign dermatosis.
Br. J. Dermatol., 145(3): 385.
- Buhl, A. (1991):
Minoxidil's action in hair follicles.
J. Invest. Dermatol., 96: 73.
- Bystryn, J.C. and Tamesis, J. (1991):
Immunologic aspects of hair loss.
J. Invest. Dermatol., 96(5): 88.
- Chua, S.H.; Goh, C.L. and Ang, C.B. (1996):
Topical squaric acid dibutylester therapy for alopecia
areata.
Ann. Acad. Med. Singapore, 25(6): 842.

- Claudy, A. and Gagnaire, D. (1983):
PUVA treatment of alopecia areata.
Arch. Dermatol., 119: 975.
- Colombe, B.W. and Lou, C.D.; Price, V.H. (1999):
Alopecia areata: HLA associations with patchy
alopecia areata versus alopecia universalis.
J. Investig. Dermatol Symp. Proc., 4(3, Dec): 216-19.
- Colon, E.A.; Popkin, M.K.; Callies, A.L.; Dessert, N.J. and
Hordinsky, M.K. (1991):
Lifetime prevalence of psychiatric disorders in
patients with alopecia areata.
Comprehensive Psychiatry, 32(3): 245.
- Cotellessa, C.; Peris, K.; Caracciolo, E.; Mordenti, C. and
Chimenti, S. (2001):
The use of topical diphenylcyclopropenone for the
treatment of extensive alopecia areata.
J. Am. Acad. Dermatol., 44(1): 73.
- Deleve, L.D. (1996):
Dinitrochlorobenzene is genotoxic by sister chromatid
exchange in human skin fibroblasts.
Mutat. Res., 371: 105.
- De Villez, R.L. (1990):
The therapeutic use of topical minoxidil.
Dermatol Clin., 8(2): 367.

- Deprost, Y.; Paquez, F. and Touraine, R. (1982):
Dinitrochlorobenzene treatment of alopecia areata.
Arch. Dermatol., 118: 542.
- Drake, L.A.; Ceilley, R.I. and Cornelison, R.L. (1992):
Guidelines of care for alopecia areata.
J. Am. Acad. Dermatol., 26: 247.
- Duhra, P. and Foulds, I.S. (1999):
Persistent vitiligo induced by diphencyprone.
Br. J. Dermatol., 123: 415-416.
- Ead, R.D. (1981):
Oral zinc sulfate in alopecia areata.
Br. J. Dermatol., 104: 483.
- Ebling, F.J.G.; Dawler, R. and Rook, A. (1986):
The hair. In: Rook, A.; Wilkinson, D.S.; Ebling, F.J.G.; Champion, R.H. and Burton, J.L. Textbook of Dermatology. 4th edition. Blackwell Scientific Publications, Oxford, London and Edinburgh.
- Elieff, D.; Sundby, S.; Kennedy, W.W. and Hordinsky, M. (1991):
Decreased sweat gland number and function in patients with alopecia areata.
Br. J. dermatol., 125(2): 130.

- Ellis, C.N.; Brown, M.F. and Voorhees, J.J. (2002):
Sulfasalazine for alopecia areata.
J. Am. Acad. Dermatol., 46(4): 541.
- Fathmann, C.G.; Codta, G.L. and Seroogy, C.M (2000):
Gene therapy for autoimmune disease.
Clin. Immunol., 95: 39-43.
- Fiedler, V.C. (1987):
Topical minoxidil solution (1% and 5%) in the
treatment of alopecia areata.
J. Am. Acad. Dermatol., 16: 737.
- Fiedler, V.C. (1992):
Alopecia areata. A review of therapy, efficacy,
safety, and mechanism.
Arch. Dermatol., 128(11): 1519-29 [Medline].
- Fiedler, V.C. and Alaiti, S. (1996):
Treatment of alopecia areata.
Dermatol. Clin., 14(4): 733-7 [Medline].
- Fiedler, V.C. and Buys, C.M. (1987):
Evaluation of anthralin in the treatment of alopecia
areata.
Arch. Dermatol., 123: 1491.

Fransway, A. and Muller, S. (1988):

Topical minoxidil 3% compared with placebo for the treatment of chronic severe alopecia areata.

Cutis, 41: 434.

Freyschmidt-Paul, P.; Seitel, S.; Zoller, M.; Konig, A.; Ziegler, A.; Sundberg, J.P.; Happle, R. and Hoffmann, R. (2000):

Treatment with an anti-CD44v10-specific antibody inhibits the onset of alopecia areata in C3H/HeJmice.

J. Invest. Dermatol., 115(4): 653.

Freyschmidt. Paul, P.; Hoffmann, R.; Levine, E.; Sundberg, J.P.; Happle, R. and McElwee, K.J. (2001 a):

Current and potential agents for treatment of alopecia areata.

Curr. Pharm., 7(3): 213.

Freyschmidt-Paul, P.; Ziegler, A.; McElwee, K.J.; Happle, R.; Kissling, S.; Sundberg, J.P. and Hoffmann, R. (2001 b):

Treatment of alopecia areata in C3H/HeJmice with topical immunosuppressant FK506 (Tacrolimus).

Eur. J. Dermatol., 11(5): 405.

- Friedli, A. and Harms, M. (2002):
Alopecia areata.
Ther. Umsch., 59(5): 233-7.
- Friedli, A.; Labarthe, M.P.; Engelhardt, E.; Feldmann, R.;
Salomon, D. and Saurat, J.H. (1998):
Pulse methyl prednisolone therapy for severe alopecia
areata.
J. Am. Acad. Dermatol., 39: 597.
- Friedmann, P.S. (1981):
Unsuccessful treatment of alopecia areata with
dapsone.
Br. J. Dermatol., 104: 597.
- Galadari, I.; Rubaie, S.; Alkaabi, J.; Galadari, H. (2003):
Diphenylcyclo-propenone (diphencyprone, DPCP) in
the treatment of chronic severe alopecia areata (AA).
Allerg. Immunol., (Paris) 35(10): 397-401.
- Galbraith, G.M.P; Thiers, B.H. and Fundenberg, H.H.
(1984):
An open label trial of immunomodulation therapy
with inosiplex (Isoprinosine) in patients with alopecia
totalis and cell mediated immunodeficiency.
J. Am. Acad. Dermatol., 11: 224.

Garcia, G. and Weiner, H.L (1999):

Manipulation of Th responses by oral tolerance.

Current topics in microbiology and immunology, 238:
123-45.

Giannetti, A. and Orecchia, G. (1983):

Clinical experience on the treatment of alopecia
areata with squaric acid dibutylester.

Dermatologica, 167: 280.

Gordon, P.M.; Aldrige, R.D.; McVittie, E. and Hunter, J.A.
(1996):

Topical diphencyprone for alopecia areata: evaluation
of 48 cases after 30 months follow-up.

Br. J. Dermatol., 134(5): 869.

Gundogan, C.; Greve, B.; Raulin, C. (2004):

Treatment of alopecia areata with the 308-nm xenon
chloride excimer laser: case report of two successful
treatments with the excimer laser.

Lasers. Surg. Med., 34(2):86-90.

Gupta, A.K.; Ellis, C.N.; Cooper, K.D.; Nickoloff, B.J.;
Ho, V.C.; Chan, L.S.; Hamilton, T.A.; Tellner, D.C.;
Griffiths, C.F. and Voorhees, J.J. (1990):

Oral cyclosporine for treatment of alopecia areata. A
clinical and immunohistochemical analysis.

J. Am. Acad. Dermatol., 22: 242.

- Gupta, A.K.; Grober, J.S.; Hamilton, T.A.; Ellis, C.N.; Siegel, M.T.; Voorhees, J.J.; et. al. (1995):
Sulfasalazine therapy for psoriatic arthritis: a double-blind, placebo controlled trial.
J. Rheumatol., 22(5): 894-8.
- Happle, R. (1991):
Topical immunotherapy in alopecia areata.
J. Invest. Dermatol., 96(Suppl): 71S.
- Happle, R.; Kalveran, K.; Buchner, U.; Echternacht-Hap, K.; Goggelman, W. and Summer, K.H. (1980):
Contact allergy a therapeutic tool for alopecia areata: application of squaric acid dibutylester.
Dermatologica, 161: 289-97.
- Happle, R.; Housen, B.M. and Wiesner-Menzel, L. (1983):
DPCP in the treatment of alopecia areata.
Acta. Derm. Venerol., 63: 49.
- Happle, R.; Perret, C.H.; Menzel, L.W. and MacDonald, D.M. (1984):
Treatment of AA with DPCP induces hair growth and affects the composition of peribulbar infiltration. 1st ed. New York, Butter Worths Co., 279.

Hartland, C. and Saihan, E. (1989):

Regression of cutaneous metastatic melanoma with
topical diphencyprone and oral cimetidine.

Lancet, 2: 455.

Hatzis, J.; Gougiotou, K.; Tosca, A.; et al. (1988):

Vitiligo as a reaction to topical treatment with
diphencyprone.

Dermatologica, 177: 146-148.

Hay, I.C.; Jamieson, M. and Ormerod, A.D. (1998):

Randomized trial of armothrapy. Successful
treatment for alopecia areata.

Arch. Dermatol., 135(5): 602.

Hoffmann, R. and Happle, R. (1996):

Topical immunotherapy in alopecia areata. What,
how, and why?.

Dermatol. Clin., 14(4) 739-44 [Medline].

Hoffmann, R.; Eicheler, W.; Huth, A.; Wenzel, E. and
Happle, R. (1996):

Cytokines and growth factors influence hair growth in
vitro. Possible implications for the pathogenesis and
treatment of alopecia areata.

Arch. Dermatol., 288(3): 153.

- Hull, S.M. and Cunliffe, W.J. (1992):
Treatment of alopecia areata with diphenyl
cycloprone.
Am. J. Acad. Dermatol., 26: 276.
- Hull, S.M. and Norris, J.F. (1988):
Diphencyprone in the treatment of long-standing
alopecia areata.
Br. J. Dermatol., 119: 367.
- Hutter, C.D. and Laing, P. (1996):
Multiple sclerosis: Sunlight, diet, immunology and
etiology.
Med. Hypotheses, 46(2): 67.
- Jansen, G.T. (1983):
Use of topical fluorouracil.
Arch. Dermatol., 119: 784.
- Julius, S. (1985):
Immunology of hair loss.
Dermatologic immunology and Allergy, 45: 645.
- Kalam, A.; Tahseen, M.D.; Islam, S.F.; Rahmatullah, M.D.
and Faruqi, N.A. (1991):
Dinitrochlorobenzene therapy in alopecia areata.
J. Indian Med. Assoc., 89(1): 9.

- Kaplan, A.L. and Olsen, E.A. (2004):
Topical 5-fluorouracil is ineffective in the treatment
of extensive alopecia areata.
J. Am. Acad. Dermatol., 50(6): 941-3.
- Kawvar, A.B. and Stiller, M.J. (1994):
Cyclosporine in dermatology: Pharmacology and
clinical use.
Int. J. Dermatol., 33: 86.
- Kern, F.; Hoffman, W.H. and Hambrich, G.W. (1973):
Alopecia areata: Treatment and immunological studies
of A.A.
Arch. Dermatol., 107: 407.
- Khoury, E.L.; Price, V.H.; Abdel-Salam, M.M.; Stern, M.
and Greenspan, J.S. (1992):
Topical minoxidil in alopecia areata.
J. Invest. Dermatol., 99(1): 40.
- Levien, C. and Maibach, H. (2002):
Exploration of alternative and natural drugs in
dermatology.
Arch. Dermatol., 138(2): 251.
- Li, L. and Hoffman, R.M. (1995):
The feasibility of targeted selective gene therapy of
the hair follicle.
Nat. Med., 1: 705-6.

Lowy, M.; Ledoux-cobusier, M.; Achten, G. and Wybran, J. (1985):

Clinical and immunologic response to isoprinosine in alopecia areata and alopecia universalis: Association with auto-antibodies.

J. Am. Acad. Dermatol., 12:78.

Lutz, G. (1994):

Effect of cyclosporine-A on hair.

Skin Pharmacol., 7(1-2): 101.

Lutz, G. and Kreysel, H.W. (1990):

Selective changes in lymphocytic differentiation antigens in the peripheral blood of patients with alopecia areata treatment with oral zinc.

Z. Hautkr., 65(2): 132.

MacDonald, N. (1999):

Alopecia areata: identification and current treatment approaches.

Dermatol. Nurs., 11(5): 356-9, 363-6 [Medline].

Madani, S. and Shapiro, J. (2000):

Alopecia areata update.

J. Am. Acad. Dermatol., 42(4): 549-66; quiz 567-70[Medline].

Maestroni, G.J. (2001):

The immunotherapeutic potential of melatonin.

Expert. Opin. Invest. Drugs, 10: 467.

Magee, K.L.; Hsu, S.M. and Tucker, S.B. (1990):

Trial of intralesional interferon-alfa in the treatment of alopecia areata.

Arch. Dermatol., 126(6): 760.

Mancuso, G.; Balducci, A.; Casadio, C.; Farina, P.; Staffa, M.; Valenti, L.; Milani, M. (2003):

Efficacy of betamethasone valerate foam formulation in comparison with betamethasone dipropionate lotion in the treatment of mild-to-moderate alopecia areata: a multicenter, prospective, randomized, controlled, investigator-blinded trial.

Int. J. Dermatol., 42(7): 572-5.

McElwee, K.J.; Rushton, D.H.; Trachy, R.; Oliver, R.F.(1997):

Topical FK506: a potent immunotherapy for alopecia areata? Studies using the Dundee Experimental bald rat model.

Br. J. Dermatol. 137: 491-97.

Micali, G.; Cicero, R.L.; Nasca, M.R. and Sapuppo, A.
(1996):

Treatment of alopecia areata with squaric acid
dibutylester.

Int. J. Dermatol., 35(1): 52.

Milgraum, S.S.; Mitchell, A.J. and Bacon, G.E. (1987):

Alopecia areata, endocrine function, and
autoantibodies in patients 16-years of age or younger.

J. Am. Acad. Dermatol., 17: 57.

Mitchell, A. and Douglas, M. (1985):

Topical photochemotherapy for alopecia areata.

J. Am. Acad. Dermatol., 12: 644.

Mitchell, A. and Krull, E. (1984):

Alopecia areata: Pathogenesis and treatment.

J. Am. Acad. Dermatol., 11: 763.

Monfrecola, G.; D'Anna, F. and Delfino, M. (1987):

Topical hematoporphyrin plus UVA for treatment of
alopecia areata.

Photodermatology, 4: 305.

Montes, L.F. (1977):

Topical halcinonide in alopecia areata and in alopecia
totalis.

J. Cutan. Pathol., 4: 47.

- Muller, S.A. and Winkelmann, A.K. (1963):
Alopecia areata: an evaluation of 736 patients.
Arch. Dermatol., 88: 290.
- Namazi, MR. (2003):
The potential efficacy of thalidomide in the treatment
of recalcitrant alopecia areata.
Med. Hypotheses., 60(4): 513-4.
- Nasca, M.R.; Micali, G.; Pulvirenti, N. and Cicero, R.L.
(1998):
Transient leukoderma appearing in an untreated area
following contact immunotherapy for alopecia areata.
Eur. J. Dermatol., 8(2): 125.
- Nelson, D.A. and Spielvogel, R.L. (1985):
Alopecia areata.
Int. J. Dermatol., 24(1): 26.
- Orecchia, G. (1989):
Thymopentin in treatment of alopecia areata.
Dermatologica, 178: 231.
- Orecchia, G. (1994):
The "Castling" phenomenon: a possible explanation
[Letter].
Eur. J. Dermatol., 4: 161.

- Orecchia, G.; Malagoli, R. and Santagostino, L. (1994):
Treatment of severe alopecia areata with squaric acid
dibutylester in pediatric patients.
Pediatr. Dermatol., 11(1): 65.
- Papadopoulos, A.J.; Schwartz, R.A. and Janniger, C.K.
(2000):
Alopecia areata. Pathogenesis, diagnosis, and therapy.
Am. J. Clin. Dermatol., 1(2, Mar-Apr): 101-5.
- Paquet, P.; arrese, J. and Pierard, G.E. (1992):
Oral cyclosporine and alopecia areata.
Dermatology, 185(4): 314.
- Pericin, M. and Trueb, R.M. (1998):
Topical immunotherapy of severe alopecia areata with
diphenylcyclopropenone: evaluation of 68 cases.
Dermatology, 196(4): 418.
- Perret, C.; Steiljen, P. and Happle, R. (1990):
Alopecia areata: Pathogenesis and topical
immunotherapy.
Int. J. Dermatol., 29: 83-8.
- Porter, D. and Burton, J.L. (1971):
A comparison of intralesional triamcinolone acetonide
in alopecia areata.
Br. J. Dermatol., 85: 272.

Potapenko, A.Y. and Kyagova, A.A. (1998):

The application of antioxidants in investigations and optimization of photochemotherapy.

Membr. Cell. Biol., 12(2): 269.

Price, V.H. (1987):

Topical minoxidil in extensive alopecia areata.

Dermatologica, 2: 36.

Price, V.H. (1991):

Alopecia areata: clinical aspects.

J. Invest. Dermatol., 96: 68.

Price, V.H. (1999):

Treatment of hair loss.

N. Engl. J. Med., 34: 964.

Price, V.H and Colombe, B.W (1996):

Heritable factors distinguish two types of alopecia areata.

Dermatol. Clin., 14: 679-89.

Price, V.H. and Khoury, E. (1991):

Progress in Dermatology.

Bull. Dermatol. Found., 25: 1.

Putt, S.C.; Weinstein, L. and Szindolet, M.T. (1994):

A case study: massage, relaxation, and reward for treatment of alopecia areata.

Psychol. Rep., 74: 1315.

Ranchoff, R. and Bergfeld, W. (1989):

Extensive alopecia areata: results of treatment with 3% topical minoxidil.

Cleve. Clin. J. Med., 56: 149.

Rappersberger, K.; Maingassner, J.G.; Fialla, R.; Födinger, D.; Sterniczky, B.; Rauch, S.; Putz, E.; Stütz, A. and Wolff, J. (1996):

Clearing of psoriasis by a novel immunosuppressive macrolide.

J. Invest. Dermatol., 106: 701-10.

Rokhsar, C.K.; Shupack, J.L.; Vafai, J.J.; Washenik, K. (1998):

Efficacy of topical sensitizers in the treatment of alopecia areata.

J. Am. Acad. Dermatol., 39 : 751-61 [Medline].

Ronald, J.S.; Provot, T.T. and Farmer, E.R. (1988):

Diseases of cutaneous appendages.

Current therapy in Dermatology, Toronto, Philadelphia, BC Decher Inc: 95.

- Ruzicka, T.; Assmann, T.; Lebwohl, M. (2003):
Potential future dermatological indications for
tacrolimus ointment.
Eur. J. Dermatol., 13(4): 331-42.
- Safavi, K.H.; Muller, S.A.; Suman, V.J.; et. al. (1995):
Incidence of alopecia areata in Olmsted County,
Minnesota.
Mayo. Clin. Proc., 70(7): 628-33 [Medline].
- Sahin, S.; Yalcin, B. and Karaduman, A. (1998):
PUVA treatment for alopecia areata.
Dermatology, 197(3): 245.
- Schreiber, S.L. and Crabtree, G.R. (1992):
The mechanism of action of cyclosporin A and FK506.
Immunol. Today, 13: 136-42.
- Schwartz, R.A. and Janniger, C.K. (1997):
Alopecia areata.
Cutis, 59: 238.
- Seiter, S.; Ugurel, S.; Tilgen, W. and Reinhold, U. (2001):
High-dose pulse corticosteroid therapy in the
treatment of severe alopecia areata.
Dermatology, 202(3): 230.

Shapiro, J. (1993 a):

Topical immunotherapy in the treatment of chronic severe alopecia areata.

Dermatol. Clin., 11: 611.

Shapiro, J. and Price, V.H. (1998):

Hair regrowth. Therapeutic agents.

Dermatol. Clin., 16(2): 341-56 [Medline].

Shapiro, J. and Madani, S.H. (1999):

Alopecia areata: diagnosis and management.

Int. J. Dermatol., 38(1): 19.

Shapiro, J.; Tan, J. and Ho, V. (1993 b):

Treatment of chronic severe alopecia areata with diphenylcyclopropenone and 5% minoxidil: a clinical and immunopathologic evaluation.

J. Am. Acad. Dermatol., 29: 729.

Shapiro, J.; Sundberg, J.P.; Bissonette, R.; McElwee, K.J.; McLean D.I.; Carroll, J.M.; Oliver, R.F.; Tang, L.; Lui, H. (1999):

Alopecia areata-like hair loss in C3H/HeJ mice and DEBR rats can be reversed using topical diphencyprone.

J. Invest. Dermatol. Symp. Proc., 4(3, Dec.): 239.

Sharma, U.K. (1996):

Pulsed administration of corticosteroids in the treatment of alopecia areata.

Int. J. Dermatol., 35(2): 133.

Sharma, U.K and Muralidhar, S. (1998):

Treatment of widespread alopecia areata in young patients with monthly oral corticosteroid pulse. Pediatr.

Dermatol., 15(4): 313.

Sharma, U.K.; Dawn, G. and Kumar, B. (1996 a):

Profile of alopecia areata in northern India.

Intern. J. Dermatol., 35: 22.

Sharma, U.K.; Kumar, B. and Dawn, G. (1996 b):

A clinical study of childhood alopecia areata in Chandigarh. India.

Pediatr. Dermatol., 13: 372.

Sharma, U.K. and Gupta, S. (1999):

Twice weekly 5 mg dexamethasone oral pulse in the treatment of extensive alopecia areata.

J. Dermatol., 26(9): 562.

Sharma, U.K.; Sialy, R.; Kumar, B. and Gupta, S. (1999):

Evaluation of thyroid function in north Indian with alopecia areata: response to intravenous injection of 100 micrograms thyrotropin releasing hormone (TRH).

J. Dermatol., 26: 339.

Sharquie, K.E.; Al-Obaidi, H.K. (2002):

Onion juice (*Allium cepa* L), a new topical treatment for alopecia areata.

J. Dermatol., 29(6): 343-6.

Shellow, V.W.; Edwards, J.E.; Koo, J.Y. (1992):

Profile of alopecia areata: a questionnaire analysis of patient and family.

Int. J. Dermatol., 31(3): 186-9 [Medline].

Shelly, W.B. and Shelly, E.D. (1987):

Alopecia areata.

Advanced Dermatologic Therapy, 9: 60.

Shevliakov, L.V. (1978):

Experience in treating alopecia areata.

Vestn. Dermatol. Venerol., 4: 70.

- Simonetti, O.; Lucarini, G.; Bernardini, M.L.; Simoncini, C.; Biagini, G.; Offidani, A. (2004):
Expression of vascular endothelial growth factor, apoptosis inhibitors (survivin and p16) and CCL27 in alopecia areata before and after diphencyprone treatment: an immunohistochemical study.
Br. J. Dermatol., 150(5): 940-8.
- Simpson, N.B. (1991):
Alopecia areata: In: Rook, A. and Dawber, R. editord.
Diseases of the hair and scalp. London: Blackwell Scientific Publications: 296.
- Smedegard, G. and Bjork, J. (1995):
Sulphasalazine: mechanism of action in Rheumatoid arthritis.
Br. Soc. Rheumatol., 34s: 7-15.
- Sonnichsen, N.; Reinnicke, C.; Herrmann, C. and Glatzel, E. (1984):
Zinc therapy of alopecia areata.
Dermatol. Monatssche., 170(7): 437.
- Summer, K. and Goggelmann, W. (1980):
1-chloro-2,4-dinitrochlorobenzene depletes glutathione in rat skin and is mutagenic.
Mutat. Res., 77: 91-3.

- Swarnson, A.; Mitchall, A.J. and Bruly, M.S. (1981):
Topical treatment of alopecia areata.
Arch. Dermatol., 117: 384.
- Tang, L.; Cao, L.; Sundberg, J.P.; Lui, H.; Shapiro, J.
(2004):
Restoration of hair growth in mice with an alopecia
areata»like disease using topical anthralin.
Exp. Dermatol., 13(1): 5-10.
- Taylor, C.R. and Hawk, J.L. (1995):
PUVA treatment of alopecia areata, totalis and
universalis: audit of 10 years' experience at St John's
Institute of Dermatology.
Br. J. Dermatol., 133(6): 914-8 [Medline].
- Thiers, B. (1991):
Isoprinosine treatment of alopecia areata.
J. Invest. Dermatol., 96: 725.
- Tosti, A.; DePadova, M.P.; Minghetti, G.; Veronesi, S.
(1986):
Therapies versus placebo in the treatment of patchy
alopecia areata.
J. Am. Acad. Dermatol., 15(2 Pt 1): 209-10.

- Tosti, A.; Mannuzi, P. and Gasponi, A. (1988):
Thymopentin in the treatment of severe alopecia areata.
Dermatologica, 177: 170.
- Tosti, A.; Morelli, R.; Bardazzi, F. and Peluso, A.M. (1994):
Prevalence of nail abnormalities in children with alopecia areata.
Pediatr. Dermatol., 11: 112.
- Tosti, A.; Guidetti, M.S.; Bardazzi, F. and Misciali, G. (1996):
Long-term results of topical immunotherapy in children with alopecia totalis or alopecia universalis.
J. Am. Acad. Dermatol., 35: 199.
- Tosti, A.; Piraccini, B.M.; Pazzaglia, M.; Vincenzi, C. (2003):
Clobetasol propionate 0.05% under occlusion in the treatment of alopecia totalis/universalis.
J. Am. Acad. Dermatol., 49(1): 96-8.
- Tsai, Y.M.; Chen, W.; Hsu, M.L.; Lin, T.K. (2002):
High-dose steroid pulse therapy for the treatment of severe alopecia areata.
J. Formos. Med. Assoc., 101(3): 223-6.

Unger, W. and Schemmer, R. (1978):

Corticosteroids in the treatment of alopecia totalis.

Arch. Dermatol., 114: 1486.

van Baar, H.M.; Van der Vleuten, C.J. and Van de Kerkhof, P.C. (1995):

Dapsone versus topical immunotherapy in alopecia areata.

Br. J. Dermatol., 133(2): 270.

van der Steen, P. and Happle, R. (1992):

Castling phenomenon.

Eur. Dermatol., 2: 151-3.

van der Steen, P.H.M.; van Baar, H.M.J.; Perret, C.M. and Happle, R. (1991):

Treatment of alopecia areata with diphenyl cyclopropenone.

J. Am. Acad. Dermatol., 24: 253-7.

van der Steen, P.H.; Boezeman, J.B. and Happle, R. (1992):

Topical immunotherapy for alopecia areata: re-evaluation of 139 cases after an additional follow-up period of 19 months.

Dermatology, 184(3): 198.

- van der Velden, E.M.; Drost, B.H.; Ijsselmuiden, O.E.;
Baruchin, A.M. and Hulsebosch, H.J. (1998):
Dermatography as a new treatment for alopecia areata
of the eyebrows.
Int. J. Dermatol., 37(8): 617.
- Weise, K.; Kretzschmar, L.; John, S.M. and Hamm, H.
(1996):
Topical immunotherapy in alopecia areata: anamnestic
and clinical criteria of prognostic significance.
Dermatology, 192(2): 129.
- Werth, V.P.; White, W.L.; Sanchez, M.R. and Franks, A.G.
(1992):
Incidence of alopecia areata in lupus erythematosus.
Arch. Dermatol., 128: 368.
- Whiting, D. (1987):
The treatment of alopecia areata.
Cutis., 40: 248.
- Whitmont, K.J. and Cooper, A.J. (2003):
PUVA treatment of alopecia areata totalis and
universalis: a retrospective study. Australas.
J. Dermatol., 44(2): 106-9.

- Wilkerson, M.; Connor, T.H.; Henkin, J.; Wilkin, J.K. and Matney, T.S. (1987):
Assessment of diphenylcyclopropenone for photochemically induced mutagenicity in Ames assay.
J. Am. Acad. Dermatol., 17: 606-11.
- Wilkinson, D.C.; Ebling, F.J.G.; Dawber, R. and Rook, A. (1986):
The hair.
Textbook of Dermatology, 3: 1985.
- Winter, R.; Kern, F. and Blizzard, R. (1976):
Prednisone therapy for alopecia areata.
Arch. Dermatol., 112: 1549.
- Wiseman, M.C.; Shapiro, J.; MacDonald, N. and Lui, H. (2001):
Predictive model for immunotherapy of alopecia areata with diphencyprone.
Arch. Dermatol., 137(8): 1063.
- Wolowa, F. and Jablonska, S. (1976):
Zinc in the treatment of alopecia areata. Biology and disease of hair. Baltimore, University, Park Press, 305.

- Yamazaki, M.; Miura, Y.; Tsuboi, R.; Ogawa, H. (2003):
Linear polarized infrared irradiation using Super
Lizer is an effective treatment for multiple-type
alopecia areata.
Int. J. Dermatol., 42(9): 738-40.
- Yoshino, T.; Asada, H.; Ando, Y.; Fujii, H.; Yamaguchi,
Y.; Yashikawa, K. and Itami, S. (2001):
Impaired responses of peripheral blood mononuclear
cells to T-cell stimulants in alopecia areata patients
with poor response to topical immunotherapy.
Br. J. Dermatol., 145(3): 415.
- Zoller, M.; Freyschmidt-Paul, P.; Vitacolonna, M.;
McElwee, K.J.; Hummel, S.; Hoffmann, R. (2004):
Chronic delayed-type hypersensitivity reaction as a
means to treat alopecia areata.
Clin. Exp. Immunol., 135(3): 398-408.