

References

Abdul Kareem, A.S.; Abahussein, A. A. and Okoro, A.(1998): Severe alopecia areata treatment with systemic corticosteroids.

Int. J. Dermatol., 37(8): 622.

Abel, E.A.; (1995): Narrow Band in Pityriasis rosea .

Dermatol Clinics., 13:841.

Aboul-Enein, H.Y.; Kladna, A.; Kruk, I., et al., (2003): Effect of psoralens on Fenton-like reaction generating reactive oxygen species.

Biopolymers., 72: 59.

Al-Abossi, M.M. and Ajam, Z. A. (1995): Oral photochemotherapy in vitiligo: Follow-up, patient compliance.

Int. J. Dermatol., 34: 206.

Anderson , R.R. and Parrish, J.A.(1998): Optical properties of human skin.

In :Regan . J.D.; Parnish, J.A. (eds). The Science of photo medicine. New York : Plenum press. P.147.Quoted from Hawk (1982).

Anderson, T.F. and Voorrhees, J.J. (1980): Psoralen photochemotherapy of cutaneous disorders.

Annu. Rev. Pharmacol. Toxicol., 20:235.

Anthony, F.A.; Laboda, H.M. and Costlow, M.E. (1997): Psoralen-fatty acid adducts activate melanocytes protein kinase C: a proposed mechanism for melanogenesis induces by 8-methoxypsoralen and ultraviolet A by light.

Photodermatol. Photoimmunol. Photomed., 13: 9.

Armstrong, R.B. (1992): Photobiology of ultraviolet radiation in : Photochemotherapy in Dermatology, edited by Abel E.A.; 2nd edition, Igaku-Shoin, Medical Publishers

Inc., New York, Tokyo, PP17.

Arndt, K.A. (1986): Definition of alopecia areata.

Manual of dermatologic therapeutic., 2:17.

Ashton, R.E.; Andre, P.; Lowe, N.J.; Whitefield, M. (1983): Historical and current perspectives.,

J. Am. Acad. Dermatol., 9: 173.

Baron, E.D. and Stevens, S.R. (2003): Photo therapy for cutaneous T- cell lymphoma.

Dermatol ther., 16:303.

Behrens, S.; Grundmann-Kollmann, M.; Schiener, R., et al., (2000): Combination phototherapy of psoriasis with narrow band UVB irradiation and topical tazarotene gel.

J. Am. Acad. Dermatol., 42 (3) : 493.

Behrens-Williams, S.C.; Letter, U.; Schiener, R., et al., (2001): The PUVA-turban as a new option of applying a dilute psoralen solution selectively to the scalp of patients with alopecia areata.

J. Am. Acad. Dermatol., 44(2): 248.

- Berg ,M. and Ros, A.M.(1994)** : Treatment of psoriasis with psoralens and ultraviolet A double-blind comparison of 8-MOP and 5-MOP.,
Photodermatol photoimmunol. Photomed., 10(5) : 217.
- Bertolino (2000):** Intralesional Steroid injection as a treatment of alopecia areata.
J. am. Acad. Dermatol., 44: 105.
- Bilsland, D.; Dawe, R.; Diffey, B.L., et al., (1997):** An appraisal of narrow-band (TL-01) UVB phototherapy.
Br. J. Dermatol., 137:327.
- Black, R.J. and Gaven , E.T. (2006):** Photocarcinogenecity of NB-UVB.
Br. J. Dermatol; 154:551.
- Bolduc, C.H. (2000):** Corlicosteroid in Alopecia areata.
Medicine Journal., 2:11.
- Bolduc, C .H. (2001) :** Alopecia Areata Association.
Arch. Dermatol., 88:290.
- Bolduc, C.H.; Lassus, A.; Eskelinen, A. et al (1999):** Treatment of alopecia areata with three different PUVA modalities.
Photodermatology., 74:118.
- Bolduc, C.H.; Lori, H.; Jerry, S. et al., (2001):** Efficacy of narrow band UVB in the treatment of AA.
www.ehrs.org/conferenceabstracts/2001tokyo/researchabstracts

Bolduc, C.H.; Muller, S.A. and winkelmann (2002): Alopecia areata clinic aspects and therapy.

Lancet., 11: 751.

Brajac, I.; Tkalcic, M.; Dragojevic, D.M. and Gruber, F. (2003): Roles of stress, stress perception, and trait – anxiety in the onset and course of alopecia areata.

Journal of Dermatology, 30: 871.

Brands, S.; Brakman, M.; Bos, J.D. and de Rio, M.A. (1999): No additional effect of calcipotriol ointment on low-dose UVB phototherapy in psoriasis.

J. Am. Acad. Dermatol., 41(6):991.

Brickl, R.; Schmidt, J. and Koss, F.W. (1984): Clinical pharmacology of oral psoralen drugs.

Photodermatology., 1:174.

Buckley, D.A. and Vivier, A.W. (2001): The therapeutic use of topical contact sensitizers in benign dermatosis.

Br. J. Dermatol., 145(3): 385.

Buhl, A. (1991): Minoxidil's action in hair follicles.

J. Am. Acad. Dermatol., 231:120.

Calazavara-Pinton, P.G.; Zone, C.; Candiago, E. and Facehetti, F. (2000): Blisters on psoriatic lesions treated with TL-01 lamps.

J. Dermatol., 200:115

Carraro, C. and Pathak, M.A. (1988): Studies on the nature of in vivo and in vitro photosensitization reactions by psoralens and porphyrins.

J. Invest. Dermatol., 90:267.

- Carrozza, P.; Hausermann, P.; Nestle, F.O. et al., (2000):** Clinical efficacy of narrow – band UVB (311-nm) combined with dithranol in psoriasis.
Dermatology. 200:35.
- Carson, D.A. and Lois, A. (1995):** Cancer progression and p53.
Lancet, 346:1009.
- Chua, S.H.; Goh, C.L. and Ang, C.B. (1996):** Topical squaric acid dibutylester therapy for alopecia areata.
Ann. Acad. Med. Singapore., 25(6): 842.
- Claudy, A. and Gagnaire, D. (1983):** PUVA treatment of AA.
Arch. Dermatol., 119: 975.
- Dark, C.; Dawe, R.C.; Evans, A. T. et al., (2000):** Narrow-band TL-01 phototherapy for patch-stage mycosis fungoides.
American Medical Association.,136:421.
- Der-Petrossian, M.; Seeber, A.; Honigsmann, H. and Tanew, A. (2000):** Half-side comparison study on the efficacy of 8-methoxypsoralen bath-PUVA versus narrow-band ultraviolet B phototherapy in patients with severe chronic atopic dermatitis.
Br. J. Dermatol., 142(1):39.
- Diawara, M.M.; Williams, D.E.; Oganessian, A. and Spitsbergen, J. (2000):** Dietary psoralens induce hepatotoxicity in C57 mice.
J. Nat. Toxins., 9(2): 1790.

Diffey, B.L. and Roelandts, R. (1986): Status of ultraviolet A dosimetry in methoxsalen plus ultraviolet A therapy.

J. Am. Acad. Dermatol., 15:1209.

Dogra, S. and Kanwar, A.J. (2004): Narrow band UVB phototherapy in dermatology.

Indian Dermatol Venereol Leprol., 70:205.

Dogra, S. and Parsad, D. (2003): Combination of narrow band UVB and topical calcipotriene in vitiligo.

Arch. Dermatol., 139: 393.

Dogra, S.; Parsad, D. and Handa, S. (2004): Narrowband ultraviolet B in airborne contact dermatitis : a ray of hope!

Br. J. Dermatol., 150:373.

Drake, L.A.; Ceilley, R.I. and Cornelison, R.L. (1992): Guididelines of care for alopecia areata.

J. Am. Acad. Dermatol., 26: 247.

Drake, L.A.; Ceilley, R.R.; Doner, W. et al., (1994): Guidelines of care for phototherapy and photochemotherapy.

J. Am. Acad. Dermatol., 31 :643.

Ead, R.D. (1981): Oral zinc sulfate in alopecia areata.

Br. J. Dermatol., 104 : 483.

Eisen, D. and Ellis, C.N. (1990): Effect of topical tacrolimus in some immunodisorder skin diseases.

J. Am. Acad. Dermatol., 23:1259.

- EI-Ghorr, A.A. and Norval, M. (1997):*** Biological effects of narrow band (311 nm TL-01) UVB irradiation.
J. Photochem. Photobiol. B., 38(2-3):99.
- Eller, M.S.; Yaar, M. and Gilchrest, B.A. (1994):*** DNA damage and melanogenesis.
Nature., 372:413.
- Ellis, C.N.; Brown, M.F. and Voorhees, J.J. (2002):*** Sulfasalazine for alopecia areata.
J. Am. Acad. Dermatol., 46:541.
- Fahmy, I.R. and Abu-Shady, H. (1947):*** Ammi Majus Linn: Pharmacological study and isolation of a crystalline constituent ammoidin. J. Pharm. Pharmac., 20:281.
Couted from Krull, E.A. and Kerker. (1986): Mechanism of action of psoralen.
Br. J. Dermatol., 120 : 70.
- Ferguson, J. (2004):*** NB-UVB is better than PUVA from Sneelman et al. Randomized half-side comparison of NB-UVB and trimethlypsoralen bath plus UVA treatments of psoriasis.
Acta. Dermatol. Venereologica., 84: 132.
- Fiedler, V.C. (1987):*** Topical minoxidil solution (1% and 5%) in the treatment of alopecia areata.
J. Am. Acad. Dermatol., 16: 737.
- Fiedler, V.C. (1992):*** Alopecia areata: A review of therapy, efficacy, safety and mechanism.
Arch. Dermatol., 128: 1519.

Fiedler, V.C. and Buys, C.M. (1987): Evaluation of anthralin in the treatment of alopecia areata.

Arch. Dermatol., 123: 1491.

Fisher, T.W.; Burmeister, G.; Schmidt, H.W. and Elsner, P. (2004): Efficacy of melatonin in treatment of alopecia areata.

Br J. Dermatol. Feb., 150 (2): 341.

Flindt-Hansen, H.; McFadden, V. and Eeg-Larsen, T. (1990): Effect of a new narrow-band UVB lamp on photocarcinogenesis in mice.

Acta. Derma. Venereol. (Stockh)., 71 : 245.

Fransway, A. and Muller, S. (1988): Topical minoxidil 3% compared with placebo for the treatment of chronic severe alopecia areata.

Cutis., 41: 434.

Friedli, A.; Labarthe, M.P.; Engelhardt, E. et al., (1998): Pulse methyl prednisolone therapy for alopecia areata.

J. Am. Acad. Dermatol., 39 : 597.

Gilhar, A.; Ullmann, Y.; Berkutski, T. et al., (1998): Autoimmune hair loss (alopecia areata) transferred by T lymphocytes to human scalp explants on SCID mice.

J. Clin. Invest., 101 : 62.

Gonzalez, E. (1995): PUVA for psoriasis.

Dermatol. Clin.,13:851.

Gordon, P.M.; Aldrige, R.D.; McVittie, E. and Hunter, J.A. (1996): Topical diphencyprone for alopecia areata: evaluation of 48 cases after 30 months follow-up.

Br. J. Dermatol., 134(5): 869.

Grewe, M.; Gyufka, K.; Budnik, A. et al., (1997): Interleukin receptors type I and II are differentially regulated in human keratinocytes by ultraviolet B radiation.

J. Invest. Dermatol., 107:865.

Guenther, L. (2000): Tazarotene combination treatment in psoriasis.

J. Am. Acad. Dermatol., 43:36.

Gupta, A.K. ; Ellis, C.N.; Cooper, K.D. et al., (1990): Oral cyclosporin for treatment of alopecia areata. A clinical and immunohistochemical analysis.

J. Am. Acad. Dermatol., 22:242.

Habif, T.P. (2004): Hair diseases. In Clinical Dermatology: A Color Guide to Diagnosis and Therapy.

Philadelphia: Mosby., 4th ed., pp. 834.

Halpern, S.M.; Anstey, A.V.; Dawe, R.S. et al (2000): Guidelines for topical PUVA: A report of workshop of the British Photodermatology Group.

Br. J. Dermatol., 142: 22.

Handerson, C.A. and Ichyshyn, A. (1995): Vitiligo complicating diphencyprone sensitization therapy for alopecia universalis.

Br. J. Dermatol., 133: 496.

Happle, R.; Perret, C.H.; Menzel, L.W. and MacDonald, D.M. (1984): Treatment of AA with DPC induces hair growth and affects the composition of peribulbar infiltration.

New York, Butter Worths Co.1st ed., 279.

Haratake, A.; Uchida, Y.; Schmuth, M., et al., (1997): UVB-induced alternations in permeability barrier function: Roles for epidermal hyperproliferation and thymocyte-mediated response.

J. Invest. Dermatol., 108:769.

Hawk, J.L.M. (1982): Responses of normal skin to ultraviolet radiation. In: Regan.

J.D.; Parrish, J.A. (eds.). The Science of Photomedicine. New York : Plenum Press.219.

Hay, I.C.; Jamieson, M. and Ormerod, A.D. (1999): Randomized trial of armotherapy. Successful treatment for alopecia areata.

Arch. Dermatol., 135(5): 602.

Healy, E. and Rogers, S. (1993): PUVA treatment for alopecia areata does it work ? A retrospective review of 102 cases.

Br. J. Dermatol., 129: 42.

Holder, R. and Young, M. (2000): New emerging therapies for vitiligo.

Dermatologic Clinics., 1:79.

Hönigsmann, H. (1994): Phototherapy and photochemo-therapy. Seminars in Dermatology, Vol. 9, 1 : 84.

- Hönigsmann, H.; Fitzpatrick, T.B. ; Pathak, M.A. et al., (1992):** Oral photo-chemotherapy with psoralens and UVA (PUVA): Principles and practice. In Fitzpatrick, T.B.; Eisen, A.Z.; Wolff, K. et al., (eds):
Dermatology in General Medicine (3rd ed). New York: McGraw-Hill., 153:1558.
- Hönigsmann, H.; Jaschke, E.; Nitsche, V. et al., (1982):** Serum levels of 8-MOP in two different drug preparations. Correlation with photosensitivity and UVA dose requirements for photochemotherapy.
J. Invest. Dermatol., 79 : 233.
- Hultsch, T.; Muller, K.D.; Meingassner, J.g. et al., (1998):** Ascomycin macrolactam dervative SDZ ASM 981 inhibits the release of granule associated mediators and of newly sytokines in RBL 2H3 mast cells in an immunophilin dependent manner.
Arch. Dermatol. Res., 290 : 501.
- Ikeda, T. (1965):** A new classification of alopecia areata. Dermatologica., 131: 421.
Quoted from Krull, E.A. and Mitchell, A.J. (1984): Alopecia areata: pathogenesis and treatment.
J. Am. Acad. Dermatol., 11: 763.
- Jensen, G.T. (1983):** Use of topical fluorouracil.
Arch. Dermatol., 119:784.
- Jimbow, K.; Chen, H.; Parks, J.S. et al., (1995):** Vitiligo leukodema: Abnormal retention of TRP-1 due to protein-protein interaction with cainexin. melanogenesis chaperone, leading to melanocytes apoptosis.
J. Invest. Dermatol., 104:664.

Jimbow, K.; Kaidbey, K.H.; Pathak, M.A. et al., (1974): Pigmentation stimulated by UV-B, UV-A and psoralens.

J. Invest. Dermatol., 62:548.

Julius, S. (1985) : Immunology of hair loss.

Dermatologic immunology and Allergy.,45: 645.

Jury, E.S.; Mchenry, E.D. ; Elever et al (2006): Narrow band in children.

Clinical and Experimental J. Dermatol; 31 : 196

Kaidbey, K.H. (1985): An action spectrum for 8-MOP sensitized inhibition of DNA synthesis in vivo.

J. Invest. Dermatol., 85:98.

Kaley , J. Whitmont and Alan, J. Cooper (2003): PUVA treatment of extensive alopecia areata

Australasian J. of Dermatol., 44, 106.

Kaplan, A.L. and Oslen, E.A. (2004): Fluorouracil in alopecia areata.

J. Am. Acad. Dermatol., 50 (6) : 941

Karashima, K. ; Weyer, U. and Radloff, H. (1992) : Immunological role of Tacrolimus 1-(FK 506).

J. Am. Acad. Dermatol., 27: 625.

Kavanagh, U.; Crosby, .J. and Norval, M. (1995): Urocanic acid isomers in human skin: analysis of site variation.

Br. J. Dermatol., 133(5) : 728

Keahey, T.M.; Lauker, R.M.; Laodbey, K.H. et al., (1994): Studies on the mechanism of clinical tolerance in solar urticaria.

Br. J. Dermatol., 110 : 327.

Kemp, E.H.; Waterman, E.A.; Hawes, B.E. et al., (2002): The melanin-concentrating hormone receptor 1, a novel target of autoantibody responses in vitiligo.

J. Clin. Invest., 109 (7) : 973.

Kerker, B.J.; Huang, C.P. and Morison, W.L. (1990): photochemotherapy of generalized granuloma annulare.

Arch. Dermatol., 359-361 .

Khaitan, B. ; Mittal, R. R. and Verma, K. (2004) : Studies-Extensive Alopecia areata treated with betamethasone oral mini pulse therapy: An open uncontrolled study.

Indian Journal of Dermatology, Venereology and Leprology. Vol.70, Num.6, P.349.

Khandpur, S. and Reddy, B.S. (2001): An association twenty-nail dystrophy with vitiligo.

J. Dermatol., 28(1) : 38.

Kirby, B. and Rogers, S. (1997): Treatment of PUVA itch with capsaicin.

Br. J. Dermatol., 137(1): 1 52.

Kochevar, I.E.; Pathak, M.A. and Parrisli, J.A. (1999): Photophysics, photochemistry and photobiology.

In: Fitzpatrick's Dermatology in General Medicine, 5th edition, chapter, 16 : 220.

Krutmann, J. (1995): UVA-1 induced immunomodulation.

In: Krutman,J., Llmets, C.A. (eds.). Photoimmunology. Oxford: Blackwell, p246.

Krutmann, J. and Morita, A. (1999): Mechanisms of ultraviolet UVB and UVA phototherapy.

Clinical and Experimental Photodermatology., 4(1): 70.

Kurz, M.; Gu, K. and Lohse, P.A. (2000): Psoralen photocrosslinked mRNA-puromycin conjugates: a novel template for the rapid preparation of mRNA-protein fusions.

Nucleic Acids. Res., 28 (18) : 83

Kwok, V.R., Anstey, A.V. and Hawk, J.L. (2002): Psoralen photochemotherapy (PUVA) is only moderately effective in widespread vitiligo: a 10-year retrospective study.

Clin. Exp. Dermatol., 27(2) : 104.

Langner, A.; Wotska, H.; Kowalski, J., et al. (1976): Photochemotherapy (PUVA) and psoriasis: comparison of 8-MOP and 8-MOP/5-MOP.

Int. J. Dermatol., 16: 688.

Lassus, A. and Kianto, (1980): PUVA treatment for AA.

Dermatologica., 161: 298.

Lehmann, B. ; Knuschke, P. and Meurer, M. (2000) : UVB-induced conversion of 7-dehydrocholesterol to 1 alpha, 25-dihydroxyvitamin D3 (calcitriol) in the human keratinocyte line Ha Ca T.

Photochem. Photobiol., 72 (6): 803.

Levien, C. and Maibach, H. (2002): Exploration of alternative and natural drugs in dermatology.

Arch. Dermatol., 138 (2) : 251.

Logan, R.A. and Hawk, J.L.M. (1985): Spontaneous photoonycholysis.

Br. J. Dermatol.; 113 : 605.

Lutz, G. (1994): Effect of cyclosporin-A on hair.

Skin Pharmacol., 7: 101.

Lutz, G. and Kreysel, H.W. (1990): Selective changes in lymphocytic differentiation antigens in the peripheral blood of patients with alopecia areata treatment with oral zinc.

J. Am. Acad. Dermatol., 65(2) : 132.

MacDonald Hull, S.P.; Gandour, M. and Carlassere, F. et al., (2003): Guidelines for the management of alopecia areata.

Br. J. Dermatol., 149: 692.

Madani, S. and Shapiro, J. (2000) : Alopecia areata update.

J. Am. Acad. Dermatol., 42: 549.

Mahmoud, F.; Abul , H.; Haines, D.; Al-Salah, C., et al. (2002): Decreased total numbers of peripheral blood lymphocytes with elevated percentages of CD4 + CD45 RO+ and CD4 + CD25 + of T + helper cells in non-vitiligo.

J. Dermatol., 29 (3) : 69.

Mansour, L.M.; Nasser, S.Q.; El-Sheikh, M.M and Abou Ammo, D. (1997): Immunological and ultrastructural studies of alopecia areata.

Egypt. J. Dermatol. & Venereol., 17 (1) : 20.

Markham, T.; Rogers, S. and Collins, P. (2003): Narrowband UV-B (TL-01) phototherapy vs oral 8-methoxypsoralen psoralen-UV-A for the treatment of chronic plaque psoriasis.

Arch .Dermatol.; 139 :325.

Marzano, C.; Baccichetti, F.; Carlassare, F. et al., (2000): DNA damage induced by 4,6,8,9- tertiary 1-2H-furo(2,3-h) quinolin-2-one, a new furocoumarine analog: biological consequences.

Photochem. Photobiol., 71(3) : 263.

Mehta, P.K.; Mamdani, B.; Shansky, R.M. et al., (1975): Severe hypertension. Treatment with minoxidil.

J. Am. Acad. Dermatol., 233: 249.

Meleland, J.; Fisher, C.; Farr, P. M. et al., (1991): Relationship between plasma psoralen concentration and psoralen-UVA erythema.

Br. J. Dermatol. 124 : 585.

Meola, T.; Soter, N.A.; Ostreicher, R. et al., (1993): The safety of UVB phototherapy in patients with HIV infection.

J. Am. Acad. Dermatol., 29:216.

Messenger, A.G. and Rundergren, J. (2004): Minoxidil: Mechanism of action on hair Growth.

Br. J. Dermatol. 150 (2) : 186.

Micali, G.; Cicero, R.L.; Nasca, M.R. and Sapuppo, A. (1996): Treatment of alopecia areata with squaric acid dibutylester.

Int. J. Dermatol., 35(1): 52.

Milgraum, S.S.; Mitchell, A.J. and Bacon, G.E. (1987): Alopecia areata, endocrine function, and autoantibodies in patients 16-years of age or younger.

J. Am. Acad. Dermatol., 17: 57.

Miller, R.A. (1982): Psoralens and UVA induced stellate hyperpigmented freckling.

Arch. Dermatol., 18 : 619.

Misery, L. (2000): The neuro-immuno-cutaneous system and ultraviolet radiation.

Photodermatol. Photoimmunol., 16(2):78.

Mitchell, A. and Douglas, M. (1985) : Topical photochemotherapy for alopecia areata

J. Am. Acad. Dermatol., 12: 644.

Mitchell, A. and Krull, E. (1984): Alopecia areata: Pathogenesis and treatment.

J. Am. Acad. Dermatol., 11: 763.

Momtaz, T.K. and Fitzpatrick, T.B. (1998): The benefits and risks of long-term PUVA photochemotherapy.

Dermatologic Clinics, 16 (2) : 227.

Moretti, J.G. (1995): The cause and treatment of vitiligo. Current opinion in dermatology.

Br. J . Dermatol., 105 : 271.

Moretti. S.; Spallanzani, A.; Hautmann G., et al. (2002): Vitiligo and epidermal microenvironment: possible involvement of keratinocyte-derived cytokines.

Arch. Dermatol., 138 (2) 273.

Morison, W.L. (1989): Effects of ultraviolet radiation on the immune system.

Photochem. Photobiol., 50:515. Quoted from Glimour et al. (1993).

Morison, W.L. (1999): Phototherapy and photochemotherapy and update.

Semin. Cutan. Med. Surg., 18(4):297.

Morison, W.L. and Pike, R.A. (1984): Spectral power distributions of radiation sources used on phototherapy and photochemotherapy.

J. Am. Acad. Dermatol; 10 : 64.

Mosher, D.B.; Fitzpatrick, T.B.; Ortonne, J.P. and Gori, Y. (1999): Hypomelanoses and hypermelanoses.

Br. J. Dermatol., 120:555.

Murphy, G.M.; wright, J.; Nicholls, D.S.H., et al. (1989): sunbedinduced pseudoporphyria.

Br. J. Dermatol., 120: 555.

Nasca, M.R.; Micali, G.; Pulvirenti, N. and Cicero, R.L. (1998): Transient leukoderma appearing in an untreated area following contact immunotherapy for alopecia areata.

Eur. J. Dermatol, 8(2): 125.

Nestle, F.O. (2000): Concepts for immune intervention in psoriasis vulgaris.

Schweiz. Rundsch. Med. Prax., 89 (37) : 468.

Nijsten, T.E.C. and Stern, S. (2003): The increased risk of skin cancer is persistent after discontinuation of psoralen plus ultraviolet A: a cohort study.

J. Invest. Dermatol., 121: 2528.

Njoo, M.D.; Bos, J.D.; Westerhof, R.M.; and Westerhof, W. (2000): Treatment of generalized vitiligo in children with narrow-band (TL-01) UVB radiation therapy.

J. Am. Acad. Dermatol., 42 (5 Pt 1) : 245.

Njoo, M.D. and Westerhof, W. (2001): Vitiligo. Pathogenesis and treatment.

Am. J. Clin. Dermatol., 2(3) : 167.

Norris, P.G.; Hawk, J.L.M.; Baker, G. et al., (1994): British Photodermatology Group guidelines for PUVA.

Br. J. Dermatol., 133: 1010.

Olsen, E.A.; Caron, S.C. and Turney, E.A. (1992): Systemic steroids with or without 2% topical minoxidil in the treatment of alopecia areata.

Arch. Dermatol., 128 (11) : 1467.

Olsen, E.A.; Karaduman, A.; Reinhold, U. et al., (2002): A randomized clinical trial of 5% topical minoxidil versus 2% topical minoxidil and placebo in the treatment of androgenetic alopecia in men.

J. Am. Acad. Dermatol., 47 : 377 .

Orecchia, G.; Malagoli, R. and Santagostino, L. (1994): Treatment of severe alopecia areata with squaric acid dibutylester in pediatric patients.

Pediatr. Dermatol., 11(1): 65.

Pagnoni, A. and Kligman, A.M. (1997): Ultraviolet photography to identify early photodamage in young children.

Br. J. Dermatol., 137 : 321.

Papadopoulos, J.L.; Vafai, J.J.; and Suman, V.J. (2001): Alopecia areata clinical aspects.

Cleve. Clin. J. Med., 56:149.

Park, Y.K.; Park, H.Y.; Bang, D.S. et al., (1986): Subcorneal pustular dermatosis treated with phototherapy.

Int. J. Dermatol., 25:124.

Patel, G.K.; Gould, D.J.; Hawk, J.L. and McGregor, J.M. (1998): A complex photodermatosis: solar urticaria progressing to polymorphic light eruption.

Clin. Exp. Dermatol., 23 (2): 77.

Pathak, M.A. and Fitzpatrick, T.B. (1992): The evolution of photochemotherapy with psoralens and UVA (PUVA): 2000 BC to 1992 AD.

J. Photochem. Photobiol., 14: 3.

Pathak, M.A. and Fitzpatrick, T.B. (1994): Preventive treatment of sunburn, dermatoheliosis, and skin cancer with sunprotective agents.

In: Dermatology in General Medicine (ed. 4). Fitzpatrick, T.B.; Eisen A.K.; Wolff, K., et al. (eds.) New York, McGraw Hill, pp. 1689.

Pathak, M.A. and Joshi, P.C. (1984): Production of active oxygen species by psoralens and ultraviolet radiation (320-400nm).

Biochem. Biophys. Acta., 798 : 11 5.

Perret, C.M.; Steijlen, P.M. and Happle, R. (1990): Pathogenesis and topical immunotherapy.

Int. J. Dermatol., 29 (2) : 83.

Pfeifer, S.L. ; Warburton, F ; Hawk, J.L. et al., (1997) : Melanoma and non-melanoma skin cancer following psoralen photochemotherapy.

Br. J. Dermatol. ; 137 (suppl 50) : 22.

Pirkhammer, D.; Seeber, A.; Honigsmann H. and Tanew, A. (2000): Narrow-band ultraviolet B (TL-01) phototherapy is an effective and safe treatment option for patients with severe seborrhoeic dermatitis.

Br. J. Dermatol., 143(5) : 964.

Porter, D. and Burton, J.L. (1977): A comparison of intralesional triamcinolone acetonide in alopecia areata.

Br. J. Dermatol., 85:272.

Price, V.H. (1987): Topical minoxidil in extensive alopecia areata.

Dermatologica., 2:36.

Price, V.H. (1991): Alopecia areata : clinical Aspects .

J. Invest. Dermatol., 96 (5) : 65.

Price, V.H.; Willy, A. and Chen, B.K. (2005): Topical tacrolimus in alopecia areata.

J. Am. Acad. Dermatol. 52 (1) : 138.

Putt, S.C.; Weinstein, L. and Szintolct, M.I. (1994): A case study: massage, relaxation, and reward for treatment of alopecia areata.

Psychol. Rep., 74: 1315.

Quevedo, W.C., Jr.; Holstein. T. J.; Dychman, J., et al. (2000): Inhibition of UVR-induced tanning and immunosuppression by topical applications of vitamins C and E to the skin of hairless (nu/nu) mice.

Pigment Cell Res, 13 (2) : 89.

Reolandts, R. (1991): The history of photochemotherapy Photodermatol.

Photoimmunol. Photomed., 8: 184.

Reolandts, R. (2002): The history of phototherapy: something new under the sun ?

J. Am. Acad. Dermatol. 46 : 926.

Riekki, R.; Jukkola, A.; Hoyhtya, M., et al. (2000): Modulation of skin collagen metabolism by irradiation.

Br. J. Dermatol., 142 : 874.

Rogers, S.; Marks, J. and Sinister, S. et al., (1981): Itch following photochemotherapy for psoriasis.

Acta. Derm. Venereol. (Stockh), 61:178.

Rokhsar, C.K.; Shupack, J.L.; Vafai, J.J. and Washenik, K. (1998): Efficacy of topical sensitizers in the treatment of alopecia areata.

J. Am. Acad. Dermatol., 39: 751.

Ronald, J.S.; Provot, T.T. and Farmer, E.R. (1988): Diseases of cutaneous appendages.

Current therapy in Dermatology.

Rosen, C.; Scki, T.; Farinelli, W., et al., (1987): A comparison of the melanocyte response to narrow band UVB and UVA exposure in vivo.

J. Invest. Dermatol., 88:774.

Rosenberg, E.W. and Drake, L. (1976): In discussion of Dunaway DA Alopecia areata.

Arch Dermatol., 112 : 256.

Safavi, K.H.; Muller, S.A. and Suman, V.J. (1995): Incidence of alopecia areata in Olmstead country.

Minnesota, 1975 through 1989. Mayo Clin. Proc., 70: 628.

Sahin, E.E. (1995) : Alopecia areata in childhood seminars.

Dermatol., 14 (1) : 9-14.

Sahin, S.; Yalcin, B. and Karaduman, A. (1998): PUVA treatment for alopecia areata.

Dermatology, 197(3) : 245

Sakuntabhai, A.; Mathews, J.N.S. and Farr, P.M. (1994): improved prediction of the minimal phototoxic dose in PUVA therapy.

Br. J. Dermatol., 130:605.

Schallreuter, K.U.; Darby, C.R. and Gibson, J.R. (2002): Effect of UVA on non – melanoma skin Cancer.

J. Am. Acad. Dermatol., 2 : 445.

Schwartz, R.A. and Janniger, C.K. (1997): Alopecia areata.

Cutis., 59: 238.

Seiter, S.; Colige, A.; Piquemal-Vivenot, P. et al., (2000): A full-UV spectrum absorbing daily use cream protects human skin against biological changes occurring in photoaging.

Photodermatol. Photomed., 16(4): 147.

Seiter, S.; Ugurel, S.; Tilgen, W. and Reinhold, U. (2001): High-dose pulse corticosteroid therapy in the treatment of severe alopecia areata.

Dermatology, 202(3): 230.

Shapiro, J. (1993) : Alopecia areata: Update on therapy.

Dermatol. Clin., 11 (1): 35.

Shapiro, J. and Madani, S.H. (1999): Alopecia areata: diagnosis and management.

Int. J. Dermatol., 38(1): 19.

Shapiro, J.; Tan, J. and Ho, V. (1993): Treatment of chronic severe alopecia areata with diphenylcyclopropenone and 5% minoxidil: a clinical and immunopathologic evaluation.

J. Am. Acad. Dermatol., 29: 729.

Sharma, V.K. (1996): Pulsed administration corticosteroids in the treatment of alopecia areata.

Int. J. Dermatol., 35: 133.

Sharma, U.K and Gupta, S. (1999): Twice weekly 5 mg dexamethasone oral pulse in the treatment of extensive alopecia areata.

J. Dermatol., 35:22.

Sharma, U.K and Muralidhar, S. (1998): Treatment of widespread alopecia areata in young patients with monthly oral corticosteroid pulse.

Pediatr. Dermatol., 15(4): 313.

Sharma, U.K.; Sialy, R.; Kumar, B. and Gupta, S. (1999): Evaluation of thyroid function in north Indian with alopecia areata : response to intravenous injection of 100 micrograms thyrotropin releasing hormone (TRH).

J. Dermatol., 26: 339.

Sharquie, K.E. and Al-Obaidi, H.K. (2002): Onion juice (*allium cepa* L), a new topical treatment for alopecia areata.

J. Dermatol., 29 (6): 343-6.

Shelly, W.B. and Shelly, E.D. (1987): Alopecia areata. *Advanced Dermatologic Therapy*., 9: 60.

Shephard, S.E. and Panizzon, R.G. (1999): Carcinogenic risk of bath PUVA in comparison to oral PUVA therapy.

Dermatology., 199 (2) : 106.

Simpson, N.B. (1999): Alopecia areata: In : Rook, A and Dawber, R. eds Diseases of the hair and scalp.

London: Black well Scientific Publications., 296 : 178.

Skinner, R.B. (1991): Psoralens. In: Systemic Drugs for Skin Diseases, edited by Wolverton, S.E. and Wilkin. J.K., W.B. Saunders Co., Philadelphia, p 219.

Snellman, E.; Ando, Y.; Greaves, M. et al (2004): Randomized half-side comparison of narrow band UVB and trimethylpsoralen bath plus UVA treatments for psoriasis. Acta. Dermatol. Venereologica., 84:132

Soter, N.A. (1990): Acute effects of ultraviolet radiation on the skin. Semin. Dermatol., 9:11. Ister. R.S.; Fitzpatrick, I.B; Parrish. J.A. et al. (1991): Oral psoralen photochemotherapy. In: Roenigk, K.,H. and Maibach, H.I. (eds.). Psoriasis. New York. Marcel Dekker, p 603.

Springer, K.; Sillevs, J.H.; Gooptuc. et al. (2003): Common hair loss disorder. American Family Physician., 68(1): 93-102.

Stern, R.S. (2001): The PUVA follow-up study. J. Am. Acad. Dermatol., 44:755.

Stern, R.S.; Fitzpatrick, T.B.; Parrish, J.A., et al. (1991): Oral psoralen photochemotherapy. In: Roenigk, K.H. and Maibach, H.I. (eds.). Psoriasis, New York, Marcel dekker, p 603.

Stern, R.S . ; Gooptuc and , Stoughton (2002): PUVA and genital Neoplasm , retrospective study. J . Am. Acad Dermatol.,38:860-1.

Stern, R.S. and Laird, N. (1994): The Carcinogenic risk of treatment for sever psoriasis :
Photochemotherapy follow-up study.

Cander., 73:2759

Stern, R.S.; Parrish, J.A. and Fitzpatrick, T.B. (1985): Ocular findings in patients
treated with PUVA.

J. Invest. Dermatol., 85:269.

Strauss, G.H.; Bridges, B.A.; Greaves, M. et al. (1980): Inhibition of delayed
hypersensitivity reaction in skin (DNCB test) by 8-methoxypsoralen
photoehemotherapy.

Lancet, 2:556.

Summer, K. and Goggelmann, W. (1980): 1-chloro -2,4 -dinitrochlorobenzene depletes
glutathione in rat skin and is mutagenic.

Mutat. Res., 77: 91.

Tanew , A. Ortel, B.; RaPerschberger,K., et al. (1988) :5-methoxypsoralen (Bergapten)
for photochemo-therapy : bioavailability, phototoxicity and clinical efficacy in
psoriasis of a new drug preparation.

J. Am Acad. Dermatol., 18:333

Taylor, C. and Hawk, J (1995): PUVA treatment of AA, partialis, totalis and universalis:
Audit of 10 years experiences at st john's Institute of Dermatology.

Br. J. Dermatol., 155: 914.

Thialo, G.; Frank, B. and Selfanie Boms.(2005): Indication of narrow band UVB beyond
psoriasis.

J. Am. Acad. Dermatol., 4:660.

Tosti, A.; Guerra, L. and Bardazzi, F. (1989): Pigmentary disorders during (topical immunotherapy.

Dermatologica., 21:196-7.

Tosti, A.; Morelli, R.; Bardazzi, f. and Peluso, A.M.(1994): Prevalence of nail abnormalities in children with alopecia areata.

Pediatr. Dermatol., 11: 112.

Tsoukas, M.M.; Lin, G.; Lee, M., et al. (1997): Predictive dosimetry of threshold phototoxicity in photodynamic therapy on normal skin. Red wavelengths produce more extensive damage than blue at equal threshold west. Dermatol., 108:501.

Vallant, V.P.; Gillcaudeau, P.; Batta, L. et al, (1994): PUVA bath therapy strongly suppresses immunological and epidermal activation in psoriasis: a possible cellular basis by remitting therapy.

J. Exp. Med.; ISO, 11:283 [abstract].

Van Der Schaar, W.W. and Sillevs Smitt, J.H. (1984): An evaluation of PUVA therapy for alopecia areata.

Dermatologica., 168: 250.

Vole-Platzer B. and Hönigsmann H. (1995): Photoimmunology of PUVA and UV-B therapy.

(J. Krutmann CA. Elmetts Eds) Oxford: Blackwell Science Ltd, 265.

Whiting, D. (1987): The treatment of alopecia areata.

Cutis., 40: 248.

Whitmont, K. and Copper, A (2003): PUVA treatment of AA. Totalis and univesalis: A retrospective study.

Australas. J. Dermatol., 0.44 (2): 106.

Wilkerson, M.; Connor, T.H.; Henkin, J., et al.(1987): Assessment of diphenylcyclopropenone for photochemically induced mutagenicity in Ames assay.

J. Am. Acad. Dermatol., 17: 606.

Winter , R.; Kern, F. and Blizzard, R. (1976): Prednisone therapy for alopecia areata. J. Dermatol., 112: 1549.

Wiseman, M.C.; Shapiro, J.; MacDonald, N. and Lui, H. (2001): Fiedictive model for immunotherapy of alopecia areata with diphencyprone.

Arch. Dermatol., 137(8): 1063.

Wolf, H.C. and Anderason, M.P. (1981): Distribution of 8-MOP and its metabolites in rat organs after a single oral administration.

J. Invest. Dermatol., 76:252.

Wolf, J.M. and Lashner, B.A. (2002) : Inflammatory bowel disease: sorting out the treatment options.

Cleve Clin J Med.;69:621.

Wolowa F. and Jablonska, S. (1976): Zine in the treatment of alopecia areata. Biology and disease of hair.

Baltimore, University, Park Press, 305.

Wolverton , S.E. and Wilkin , J.K. (1991) : Psoralens. In: Systemic drugs for skin diseases (eds.), 1st edit, W.B. Saunders Company.

Philadelphia, London., 219

Yeo, U.C.; Shin, J.H.: Yang, J.M., et al (2000): psoralen ultraviolet A induced erythema sensitivity correlates with the concentrations of psoralen in suction blister fluid. Br. J. Dermatol., 142(4):733.

Yoshino, T.; Asada, H.; Ando, Y., et al (2001): Impaired responses of peripheral blood mononuclear cells to T-cell stimulants in alopecia areata patients with poor response to topical immunotherapy.

Br. J. Dermatol., 145(3): 415.

Young , A.R. (1990): Cumulative effects of ultraviolet radiation on the skin: cancer and photoaging.

Semin. Dermatol., 9:25.

Zappacosta, A.R. (1980): Reversal of baldness in patient receiving minoxidil for hypertension.

N. Eng. J. Med; 303: 1480-1.