

REFERENCES

REFERENCES

- A dinolfi, L.E.; Utili R., Gaeta, P., Perna, P. and Ruggiero, G. (1987)
presence of endotoxemia and its relationship to liver dysfunction
in patients with typhoid fever.
Infect.;15:359-62.
- Alex, C.; Sonnen, W.,Leonard, J.A. and Grad, W. (1980):
Vibrios and spiral organisms.
Clin . Lab. Meth. And Dig. 8th ed. Vol2.; 1770-80
- Anderson, E.S.; Ward, L.R., De Saxe, M.J., Old, D.C., Barker, R.
and Duguid, J.p. (1978):
Correlation of phage-type, biotype and source in strains of
Salmonella typhimurium.
J. of . Hygiene,; 81 : 203-217.
- Antony, T.J.; Patwari, A. K., Anand, V. K., Pillai, P. K, Aneja, S. and
Sharma, D. (1993):
Duodenal string test in typhoid fever.
J. Infect. Dis.; 30 (5): 643-7.
- Bailey and Scott's. (1994):
Enterobacteriaceae
Diag. Microbiol.;9 thed, 28:362-85.
- Baquar , N.; Threlfall, E.J., Rowe, B. and Stanely, J.(1993):
Molecular subtyping within a single Salmonella typhimurium
Phage type, DT 204c, with a PCR- generated probe for IS200.
FEMS. Microbiol. Lett.; 112(2):217-21.
- Baquar, N.; Burnens, A. and Stanely, J.(1994):
Comparative evaluation of molecular typing of strains from a
national epidemic due to Salmonella brandenburg by r RNA gene
and IS200 probes and pulsed- field gel electrophoresis.
J. Clin. Microbiol.; 32 (8): 1876-80.
- Barker, R.M. and Old, D.C. (1989):
The usefulness of biotyping in studing the epidemiology and
phylogeny of salmonella.
J. Med. Micro biol.; 29 : 81-88.

- Barrett, T. J.; Snyder, J. D., Blake, P.A and Feeley, J.C. (1982):
Enzyme- linked immunosorbent assay for detection of Salmonella
typhi flagellin in urine from typhoid patients.
J. Clin. Microbiol.; 15:236-37.
- Bena ventel.; Gottuzzo, E., Guerra. J., Grados, O., Guerra, H. and Bracc,
N. (1994):
Diagnosis of typhoid fever using a string capsule device.
Trans. Roy. Soc. Trop. Med. Hyg.; 78*: 404-6.
- Beros, M.F.; Gonzalez, C., Mc Intosh, M. A. and Cabello, F.C. (1989):
Immune respons to iron-deprivation induced proteins of S. typhi
in typhoid fever.
Infect. Im mun.; 57 (4): 1271-5.
- Beuzon, C.R. and Casadesus, J. (1997) Apr:
Conserved structure of IS200 elements in Salmonella.
Nucleic. Acids. Res.; 25(7): 1355-61.
- Beuzon, C.R. and Casadesus, J. (1997) Nov:
Cloning with Mud-P22 hybrid prophages: mapping of IS200
elements on the chromosome of Salmonella typhimurium LT2.
Mol. Gen Genet.; 256 (5): 586-88.
- Beuzon, C.R.; Schiaffino, A., Leori, G., Cappuccin elli, P., Rubino, S.
and Casadesus, J. (1997) May:
Identification of Salmonella obrtuovis by PCR amplification of a
Sero var- specific IS200 element.
Appl. Environ. Microbial.; 63 (5): 2082-85.
- Brown, D.J.; Munro, D.s. and Plastt, D.J. (1986):
Recognition of the cryptic plasmid, PSLT, by restriction
fingerprintong and a study of its incidence in Scottish Salmonella
Isolates.
J. Hyg .(Lond); 97(2): 193-97
- Brown, D.; Threlfall, E.J. and Rowe, B. (1991):
Instability of multiple drug resistance plasmids in Salmonella
typhimurium isolated from poultry.
Epidemiol. Infect.; 106(2) : 247-57.

- Buchmeies, N.A. and Heffron, F. (1989):
Intracellular survival of wild type *Salmonella typhimurium* and
macrophages- sensitive mutants in diverse population of
Macrophages Infect. Immun.; 57(1) : 1-7.
- Buezko, G.B. and Mclean, J. (1994):
Typhoid fever associated with adult respiratory distress syndrome.
Chest.; 105 (6) : 1873-4.
- Burnens, A.P.; Stanley, J., Sechter, J. and Nicolet, J.(1996):
Evolutionary origin of a monophasic *Salmonella* serovar, 9,
12:1,v:- revealed by IS200 profiles and restriction fragment
polymorphisms of the *fli B* gene.
J. Clin. Microbial.; 34(7): 1641-45.
- Burnens, A P.; Stanley, J., Sack, R., Hunziker, P., Brodard, I. and Nicolet,
J. (1997):
The Flagellin N- Methylase gene *fli B* and an adjacent serovar-
specific IS200 element in *Salmonella typhimurium*.
Microbiology.; 143 (pt5): 1539-47.
- Butler, T.; Bell W.R. Levin, J., Linh, N.N. and Arnold, K. (1978):-
Typhoid fever : studies of blood coagulation, bacteremia and endo
toxemia.
Arch. Intern . Med.; 138: 407-10.
- Calva, E.; Ordonez, L.G., Fernandez-Mora, M., Santana, F.J.,
Bobadilla, M. and Puente, J.L.(1997):
Distinctive IS200 insertion between *gyr* and *rcs C* genes in
Salmonella typhi.
J. Clin. Microbiol.; 35 (12): 3048-53
- Cano, R.J.; Torres, M.j., Klem, R.E., Palomares, J.C. and Casadesus,
J.(1992)
Detection of *Salmonella* by DNA hybridization with a fluorescent
alkaline phasphatase substrate.

- J. App. Bacteriol.; 72(5): 393-99.
- Casadesus, J. and Roth, J.R. (1989):
Absence of insertion among spontaneous mutants of *Salmonella typhimurium*.
Mol. Gen. Genet.; 216(2-3): 210-16.
- Casnueva, V.; Cidx, Cavicchioli, Oelk, M., Cofre, J. and Chiang, M.T. (1992):
Serum adenosine deaminase in the early diagnosis of typhoid fever.
Pediat. Infect. Dis. J.; 11(10): 828-30.
- Chandler, M. and Fayet, O. (1993):
Translational frame shifting in the control of transposition in Bacteria.
Mol. Microbiol.; 7:497-505.
- Chang, H.R., Loo, L.H., Jeyaseelan, K., Earnest, L. and Stackebrandt, E. (1997):
Phylo genetic relationships of *Salmonella typhi* and *Salmonella typhimurium* based on gene sequence analysis.
Inter. J. Syst. Bact., 47(4):1253-4.
- Chart, H., Rowe, B. and Cheesbrough, J.S. (1997):
Serological response of patients infected with *Salmonella typhi*
J. Clin. Path.; 50(11): 944-6
- Chaudhry, R., Laxmi, B.V., Nisar, N., Ray, K. and kumar, D. (1997):
Standardisation of poly merase chain reaction for the detection of *Salmonella typhi* in typhoid fever.
J. Clin Path., 50(5): 437-9.
- Choo, K.E.; Davis, T.M. Tsamil, A. and Ong K.H. (1997):
Longevity of antibody responses to a *Salmonella typhi* specific outer membrane protein: interpretation of a dot enzyme immunosorbent assay in an area of high typhoid fever endemicity.
American. J, Trop. Med. And Hyg.; 57(6): 656-9.
- Choo, K.E.; Oppenheimer, S.J., I smail, A.B. and Ong, K.H. (1994)
Rapid diagnosis of typhoid fever by clot enzyme immunoassay in an endemic area.
Clin. Infect. Dis., 19(10): 172-76.

- Duguid, J.P.; Anderson, E.S. and Compell, J. (1966):
Fimbriae and adhesive properties in Salmonella
J. Path. and. Bact.; 92: 107-38.
- Duguid, J.P. and Compell, J. (1969):
Antigens of type-1 fimbriae of Salmonella and other
enterobacteria.
J. Med. Microbiol.; 2: 535-53.
- Ernst, R.k.; Dombroski, D.M. and Merick, J.M.(1990):
Anaerobiosis, type 1 fimbriae and growth phase are factors that
affect in vasion of Hep-2 cells by Sal. typhi.
Infect. Immum.; 58(6) 2014-6.
- Escamilla, J.; Ugarte, H.F. and Kilpatric, M.E.(1986):
Evaluation of blood clot cultures for isolation of S. typhi, S.
paratyphi A. and Brucella melitensis.
J. Clin. Microbiol.; 24(3): 388-90.
- Esteban, E., Snipes, K., Hird, D., Kasten, R. and Kinde, H. (1996):
Ribotyping for characterization of Salmonella serotypes.
J. Clin. Microbiol., 31: 244-37
- Ewing, W.H. (1986):
Edwards and Ewing's, Identification of Enterobacteriaceae, 4 th.
Elsevier, New york, pp 1-536.
- Ezquerria, E.; Burnens, A., Jones, C. and Stanley, J.(1993):
Genotypic typing and phylogenetic analysis of Salmonella
Paratyphi B and S. java with IS200.
J. Gen. Microbiol.; 139 (pt 10): 2409-14.

Fang, F.C., Libby, S.T., Buchmeier, N.A., Loewen, P.C., Switala, J.,
Harwood, J. and Guiney, D.C. (1992):
Salmonella virulence.
Proc. Natl. Acad. Sci. USA. 89:11978-82.

Ferrecio, C., Morris, J.G. and Valdivieso, C. (1988):
Efficacy of ciprofloxacin in the treatment of chronic typhoid
Carriers.
J. infect. Dis.; 157 : 1235-9.

Fica, A.; Fernandez — Beros, M.E., Aron-Hott, L., Rivas, A., D'ottone
K., Chumpitaz, J., Guevara, J.m., Rodriguez, M. and Cabello, F. (1997):
Antibiotic-resistant Salmonella typhi from two outbreaks: few
ribotypes and IS200 types harbor Inc HI 1 Plasmids.
Microb. Drug. Resist.; 3(4) : 339-43.

Frankel, G.; (1994):
Detection of Salmonella typhi by PCR (letter).
J. Clin. Microbiol.; 32(5):1415.

Frankel, G.; Newton, S.M.C., Coolink, G.K. and Stocker, B.A.D.
(1989):
Intragenic recombination in a flagellin gene: characterization of the
HI-j gene of Sal. ty.
EMBO. J.; 8:13149-52.

Fricker, C.R. (1987):
The isolation of Salmonella and Compylobacters.
J. App. Bact.; 63: 99-116.

Galas, D.J. and Chandler, M. (1989):
Bacterial insertion sequences.
In mobile DNA, pp. 109-162
Edited by D.E. Berg and M.M. Howe Washington, D.C.
American Society for Microbiology.

Garcia-Vescoui, E., Soncini, F.C. and Graismen, E.A. (1996):
Enviromental regulation of Salmonella virulence
Cell.; 82 : 165-174.

Gibert, I.; Barbe, j. and Casadesus, J. (1990).
Distribution of insertion sequence IS200 in Salmonella and
Shigella.
J. Gen. Microbiol.; 136(pt 12): 2555-560.

Gilman, R.H.; Terminel, M., Liveine, M.M., Mengoza, P.H and Hornick,
R.B. (1975):
Relative efficacy of blood, urine, rectal swab, bone marrow, and
rose spot cultures for recovery of Sal. typhi in typoid fever.
Lancet.; 31 : 1211-3.

Gotuzzo, E.; Frisancho, O., Sanchez, J., Liendo, G., Carillo, C. and
Black, R.E. (1991):
Association between the acquired immunodeficiency syndrome
and infection with Salmonella typhi or Salmonella paratyphi in
endemic typoid area.
Arch. Inter. Med.; 151 : 381-82.

Grossman, N. and Leivel. (1984):
Complement activation via the alternative pathway by purified Sal.
lipopolysaccharide is affected its structure but not its O-antigen.
J. Immunol.; 131 : 376-82.

Heising, P.; Kratz, B., Halle, E. Groser, Y. Altwegg, M., Rabsch, W. and
Faber, J.P. (1995):
Identification of DNA gyrase A mutations in ciprofloxacin
resistant isolates of Salmonella typhimurium from men and cattle
in Germany.
Microb. Drug. Resist.; 1(3): 211-18.

- Hermans, P.W.; Saha, S.K., Van Leeuwen, W.J., Verbrugh, A.h., Van Belkum, A. and Goessens, W.H. (1996):
Molecular typing of *Salmonella typhi* strains from Dahaka (Bangladesh) and development of DNA probes identifying plasmid-encoded multidrug-resistant isolates.
J. Clin. Microbiol.; 34(6): 1373-79.
- Hilton, A.C; Banks, J.G. and Penn, C.W. (1997):
Optimization of RAPD for fingerprinting *Salmonella*.
Lett. App. Microbiol.; 24(4): 243-48.
- Hirsch, H.J.; Starliner, P. and Brachet, P. (1972):
Two kinds of insertions in bacterial genes.
Mol. Gen. Genet.; 119: 191-206.
- Hoffman, S.L. (1984):
Typhoid fever.
In : Strickland GT, ed. *Hunter's tropical medicine*. 6 the ed.
Philadephia, PA: WB Saunders.: 282-97.
- Hoffman, S.L.; Edman, D.C., Punjabi, N.H., Lesmana, M. and Harahap, J. (1987):-
Bone marrow aspirate culture superior to sterpto kinase clot culture and 8 ml 1:10 blood to broth ratio blood culture for diagnosis of typhoid fever.
Am. J. Trop. Med. Hyg.; 35(4): 836-39.
- Hoffman, S.L.; Flaniga, T.P., klaucke, D., Punjabi, N.H., Sutomo, A. and Moechtar, M.A(1986):
The Widal slide agglutination test, a valuable rapid diagnostic test in typhoid fever patients at the infection diseases hospital of Jarkata.
Am. J. Epidemiol.; 123(5): 869-75.
- Hoffman, S.L.; Punjobi, N.H., Rockill, R.c., Sutomo, A., Rivai, A.R. and Pulungish, S.P. (1984):
Duodenal string capsule culture compared with bone marrow, blood, rectal swab cultures for diagnosis of typoid and paratyphoid fever.
J. Infect. Dis.; 149(20): 157-61.

- Hook, E.W. (1995):
Salmonella species (including typhoid fever).
 In Mandell, G.L., Douglas, R.G., Bennett, J.E. Principles and
 Practice of infectious diseases.
 New York: Churchill Livigstone.; 1700-16.
- Hugenholtz, P.; Pitulle, C., Hershberger, K.L. and Pace, N.R. (1998):
 Novel division level bocterial diversity in a yellowstone hot spring.
 J.Bact.; 180(2): 366-76.
- Jesudson, M.V.; Sridharan, G., Mukundom, S. and John, T.J.(1994):
 Vi-specific latex agglutination for early rapid detection of
 Salmonella serotype in blood cultures
 Diag. Microbial. Dis.; 18(2): 75-8.
- Joan, G., Fungand Richard, G. and tilton. (1985):
 Detection of bacterial antigens by counter immunoelectrophoresis,
 Co. agglutination and latex agglutination. Chapter 90. Edwin H.;
 Lennette, E. Akbert Balous, William, J. and Hasler, JR.
 Manual CL. Microbiol.; 883-9.
- Joan, D. and John, B. (1984):
 Reexamination of the protective role of the capsular poly
 saccharide (Vi-antigens) of Salmonella typhi.
 J. Infect. Dis.; 150(3): 436-47.
- Johnson, S.; Wilkinson, R. and Davidson, R.N. (1994):
 Acute tropical infections and lung.
 John.; Jul; 49(7): 714-8.
- Kang, G.; Sridharan, G., Jesudasen, M.V and John, T.J. (1992):
 Evaluation of modified passive haemagglutination assay for Vi
 antibody estimation in Salmonella typhi infections.
 J. Clin. Path.; 45(8): 740-1.
- Kapperud, G.; Lassen J., Dommarsnes, k., Kirstiansen, B.E., Caugant,
 D.A., Ask, E. and Jahkola, M. (1989):
 Comparison of epidemiological marker methods for identification
 of Salmonella typhimurium isolates from outbreak caused by
 contaminated chocolate.
 J. Clin Microbiol.; 27(9): 2019-24

- Katouli, M.; Wollin, R., Gunnarsson, A., Kuhn, I. and Mollby, R. (1994):
Biochemical phenotypes of *Salmonella* Livingstone isolated from
Humans, animals and feedstuffs in Swedn.
Acta. Vet. Scand.; 35(1): 27-36
- Kauffmann, F. (1966):
The bacteriology of Enterbacteriaceae.
Munksgaard, Copenhagen, pp1 – 400.
- Koay, A.S., Jegathesan, M., Rohani, M.Y and Cheong, Y.M. (1997):
Pulsed-field gel electrophoresis as an epidemiologic tool in the
investigation of laboratory acquired *Salmonella* typhi infection.
Southeast. Asian. J. Trop. Med. Public. Health.; 28 (1): 82-84.; 28
(1): 82-84.; 28 (1): 82-84.
- Lam, S. and Roth, J.R. (1983) Oct:
Salmonella-specific insertion sequence.
Cell.; 34(3): 951-60.
- Lam, S. and Roth, J.R.; (1983) Dec:
Genetic mapping of IS200 copies in *Salmonella* typhimurium
strain LT2.
Genetics.; 105(4): 801-11.
- Lam, S. and Roth, J.R., (1986):
Structural and functional studies of insertion element.
J. Mol Biol.; 186 (2): 157-67.
- Leclerc, G.J.; Tartera, C. and Metcall, E.S. (1998):
Enviromental regulation of *Salmonella* typhi invasion-defective
mutants.
Jnf. Immun.; 66(2): 682-91.
- Le Minor L. (1993):
Antigenic formulas of the *Salmonella* serovars 6 th rev. WHO
Collaborating Centre for Reference and Research on *Salmonella*.
Institut posteur, Paris, pp. 1-145.
- Liebisch, B. and Schwarz, S. (1996):
Molecular typing of *Salmonella enteriea* subsp. *Enterica* serovar
Enteritidis isolates
J.Med. Microbiol.; 44(1):52-59.

- Lim, P.L.; Tam, F.C., Cheong, Y.M. and Jegathesan, M. (1998):
One step 2-minuts test to detect typhoid specific antibodies based
on particle seperation in tubes.
J. Clin. Microbiol .; 36(8): 2271-78.
- Limdborg, A.A. (1980):
Bacterial virulence factors with particular refernce to Salmonella
bacteria.
Scand. J. Infect. Dis.; 24 : 86-92.
- Lin, A.W.; Usera, M.A., Barrelt, T.J. and Golds by, R.A. (1996):
Application of random amplified polymorphic DNA analysis to
differentiate strains of Salmonella enteritidis.
J.Clin. Microbial.; 34(4): 870-76.
- Liu, S.L.; Kzaki, T., Miura, H., Matsui, K. and yobnnchi, E. (1988):
Intact motility as a Salmonella typhi invasion-related factor.
Infect. Immun.; 56(8): 1967-73.
- Liu, S.L and Sanderson, K.E. (1995):
Reasrangments in the genome of the bacterium Salmonella typhi
Proc-Natl-Acad-Sci-U-S-A 14 ; 92 (4) : 1018-22
- Luby, S.P.; Faizan, M.K.; Fisher-Hoch, S.p. and Mintz, E.D. (1997):
Risk factors for typhoid fever.
Epidem. And Infect. 119(2);129-138
- Lucho, V.J.; and Fould, J. (1990):
Heterogenicity of Lipopoly saccharide phenotype among Sal. typhi
strains.
J.Infect. Dis.; 162: 763-4.
- Lucho, V.J.; Joiner, K.A., Foulds, J., Frank, M.M. and Leive, L. (1987):
C3b generation is affected by the structure of the O antigen
lipopoly saccharide in lipopoly saccharide from Salmonella.
J. Immunol.; 139(4): 1253-9.
- Mak, Y.M. and Ho, K.K. (1992):
An improved method for the isolation of chromosomal
DNA from various bacteria and cyanobacteria
Nucleic. Acid. Res.; 20:4101

- Mandell, G.L.; Douglas, R.G. and Bennrtt. J.E. (1985):
Principles and practice of infectious diseases.
Churuchill, Living stone.: 1700-16.
- Maniatis, T.; Fritsch, E.F. and Sambrook, J. (1982):
Molecular cloning: a laboratory manual. Cold Spring Harbor
Laboratry, Cold Spring Harbor, NY.
- Martin, M.G.; Gonzalez, M.A., Moro, J. and Mendoza, M.C. (1997):
Genetic typing methods applied to the differentiation of clonal
lines among *Salmonella enterica* serogroup G strains causing
human Salmonellosis.
FEMS. Jmmunol. Med. Microbiol. ; 19(3): 215-21.
- Matsutani, S.; Ohtsubo, H., Maeda. Y. and Ohtsubo, E. (1987):
Isolation and characterization of IS elements repeated in the
bacterial chromosome.
J.Mol. Biol.; 196(3): 445-55
- Mikhail, J.A.; Sanborn, W.R. and Sipple, J.E. (1983):

Rapid economical diagnosis of enteric fever by blood clot culture
Coagulation procedure.
J. Clin. Microbiol.; 17(3). 564 – 5.
- Millemann, Y.; Lesage, M.C., Chaslus Dancla, E. and Lafont J.P. (1995):
Value of plasmid profiling ribotyping and detection of IS200 for
tracing avian isolates of *Salmonella typhimurium* and *S. enteritidis*
J. Clin. Microboil.; 33(1): 173-79.
- Mohammed, K.; Coovadia, Y.M., Connoly, C. and Sturm, A.W. (1998):
The early diagnosis of typhoid fever prior to the Widel test and
bacteriological culture results Acta.
Tropica.; 69 (2) : 165.73.
- Moshitch, S.; Doll, L., and Rubinfeld, B.Z (1992):
Mono and biphasic *Salmonella typhi*: genetic homogeneity and
distinguishing characteristics.
Mol. Microbiol.; 6(18):2589-97.
- Mukherjee, C.; Malik, A.H., Khan and Malik, A. (1993):
Rapid diagnosis of typhoid fever by co-agglutiantion in an Indian
hospital.

- J.Med. Microboil.; 11(39): 74-77.
- Murase, T.; Oskitu, T., Suzuki, R., Morozumi, H., Matsushima, A., Nakamura, A. and Yamai, S. (1995):
Evaluation of DNA fingerprinting by PFGE as an epidemiologic tool for Salmonella infections.
Microboil. Jmmunol.; 39(9): 673-76.
- Murray, J.F.; Matthay, M.A., Luce, J.M. and Flick, M.R. (1988):
An expanded defintion of adult respiratory distress syndrome.
Am. Rev. Resp. Dis.; 138: 220 – 33.
- Naas, T.; Blot, M., Fitch, W.M and Arber, W. (1995):
Dynamics of IS-related genetic rearrangements in resting E.coli K12.
Mol. Biol. Envol.; 12: 198-207.
- Nadakumair, K.s.; Palanive, L.V and Muthakkavuppan. (1993):
Dignosis of typhoid fever detection of S.typhi antibodies by inhibition ELISA.
Clin. Ex. Immunol; 94(2): 317-21.
- Nair, S.; Poh, C.L., Lim, Y.S., Tay, L. and Goh, K.T. (1994):
Genome fingerprinting of Salmonella typhi by pulsed field gel electrophoresis for subtyping common phage types.
Epidemid. Infect.; 11(3): 391-402.
- Nardiello, S.; Pizzello, T., Russo, M. and Galanti, B. (1984):
Serodiagnosis of typhoid fever by enzme linked immunosorbent assay determination of anti Salmonella typhi lipopolysaccharide antibodies.
J. Clin. Microbiol.; 20(4):718-21.
- Nastasia, A. and Mammina, C. (1995):
Epidemiological evaluation by PCR ribotyping of sporadic and outbreak-associated strains of Salmonella enterica serotype typhimurium.
Res. Microbiol.; 146(!): 99-106.
- Nastasia, A.; Mammina, C. and Villafrate, M.R. (1991):
rDNA fingerprinting as a tool in epidemiological analysis of Salmonella typhi infections.
Epidemiol. Infect.; 107(3): 565-67.

- O'Brien, A.D. (1982):
Innate resistance of mice to *Sal. typhi* infection.
Infect. Immun.; 38: 948-52.
- O'Reilly, C.; Black, G.W., Laffey, R. and Mc Connell, D.J. (1990):
Molecular analysis of an IS200 insertion in the gpt gene of
Salmonella typhimurium LT2.
J.Bacteriol.; 172(11):6599-6601.
- Ochman, H.; Anne, S., Gerber, F. and Daniel, L.H.(1988):
Genetic applications of an inverse polymerase chain reaction.
Genetics.; 120 : 621-23.
- Odaret, M.; Derche, P. and Simonet, M. (1996):
Molecular typing of *Yersinia pseudotuberculosis* by using an IS200
like element.
J. Clin. Microbiol.; 34(9): 2231-35.
- Ohnishi, K. and Kimura, K. (1998):
Serum thrombomodulin level in patients with typhoid fever.
Trans. of The. Royal. Soci. Of Trop. Med. and Hyg.
92(3): 320 -21.
- Old, D.C. (1992):
Nomenclature of *Salmonella*.
J. Med. Microbiol.; 37: 361-63.
- Olsen, J.E.; Skov, M.N., Angen, O., Threlfall, E.J. and Bisgaard, M.
(1997):
Genomic relationships between selected phage of *Salmonella*
enterica subsp. Serotype typhimurium defined by ribotyping,
IS200 typing and PFGE.
Microbiol.; 143(pt4): 1471-79.
- Olsen, J.E.; Skov, M.N., Brown, D.J., Christensen, J.P. and Bisgaard,
M. (1996) Jun:
Virulence and genotype stability of *Salmonella enterica* serovar
Berta during natural out break.
Epidemiol. Infect.; 116(3): 267-74.
- Olsen, J.E.; Skov, M.N., Christensen, J.P. and Bisgaard, M. (1996):
Dec:
Genomic lineage of *Salmonella enterica* serotype Gallinarum.

- J. Med. Microbiol.; 45(6): 413-18.
- Orskov, F. and Orskov, I. (1978):
 Serotyping of Enterobacteriaceae with special emphasis on K
 antigen determination.
 Meth Microbiol.; 1: 1-77.
- Orskov, F. and Orskov, J. (1983):
 Summary of a workshop on the clone concept in the epidemiology,
 taxonomy and evolution of the Enterobacteriaceae and other
 bacteria.
 J. Inf. Dis.; 148: 346-57.
- Orskov, I.; Orskov, F. Jann, B. and Jann, K. (1977):
 Serology, chemistry and genetics of O and K antigens of E. coli.
 Bact. Rev.; 41 : 667-710.
- Pang, T. and Puthucherry, S.D. (1983):
 Significance and value of the Widal test in the diagnosis of typhoid
 fever in an endemic area.
 J. Clin. Path.; 36:471-5.
- Parker, M.T.; Collier, L.H., Smith, G.R. and Easmo, C.S.F. (1990):
 Tooley and Wilson's
 Principles of Bact, virol, Immun, 8 th ed., vol 3 Edward. Arnold,
 London.; 424-6.
- Pedler, S.J. and Orr, K.E. (1990):
 Examination of faeces for bacterial pathogens.
 J. Clin. Path.; 43: 410-15.
- Pillai, P.K. and Parkash, K. (1993):
 Drug resistance and phage type of Salmonella typhi in India.
 India J. Med. Res; 97: 154 – 8.
- Platt, and Brown, D.J. (1988):
 Old and new techniques together resolve a problem of infection by
 Salmonella typhimurium.
 Epidemiol. Infect.; 99(1) : 137- 42
- Platt, D.J.; Chesham, I.S., Brown, D.J., Kraft, C.A. and Taggart, J.(1986):
 Restriction enzyme fingerprinting of enterobacterial plasmid. A
 simple strategy with wide application.

J. Hyg. (lond).; 97 (2): 205-10.

Popoff, M.Y. and Le Minor, L. (1992).

Antigenic formulas of Salmonella serovars, WHO Collaborating center for Reference and Research on Sal.

Institut Pasteur, Paris, PP 1-145

J. Biol. Chem.; 266(29): 19510-18.

Qadri A. (1997):

Identification of specific recognition molecules on murine mononuclear phagocytes and B lymphocytes for Vi capsule polysaccharide;

modulation of MHC Class II expression on stimulation the polysaccharide.

Immunology. 92 (1): 146-52.

Qadri, A.; Ghosh, S., Prakash, K., Kumar, R. and Talwar, G.P. (1990):

Sandwich enzyme immunoassay for detection of Salmonella typhi.

J. immuno. assay.; 11(12): 251-70.

Rahman, H.; Singh, V.B., Sharma, V.D. and Harne, S.D. (1991):

Co-agglutination test for rapid detection of Salmonella enteroroxin.

Int. J. Microbiol.; 275(3): 303-11.

Ralph, A. and Giannella, A. (1989):

The Mannual. Clinic. Microbiol.; (21): 317-25.

Ramadass, P.; Meerarain S., Venkatesha, M.D., Senthilkumar, A. and Nachimuthu, K. (1997):

Characterization of leptospiral serovars by randomly amplified Polymorphic DNA fingerprinting.

Int. J. Syst. Bacteriol.; 47(2): 575-76.

Rasaily, R.; Dutta, P., Saha, M.R., Mitra, U., Bhattacharya, S.K. and Manna, B. (1993):

Value of a single Widal test in the diagnosis of typhoid fever.

Indian. J. Med. Res.; 97: 104 -7

- Reynolds, A.E.; Felton, J. and Wright, A. (1981):
Insertion of DNA activates the cryptic *bgI* operon in *E. coli* K12.
Nature; 293 : 625 – 29.
- Riley P.A.; Threlfall, E.J., Cheasty, T., Wooldridge, K.G, Williams, P.H. and Phillips, I. (1993):
Occurrence of *Flme* plasmids in multiply antimicrobial resistant *Escherichia coli* isolated from urinary tract infection.
Epidemiol. Infect.; 110 (3): 459-68.
- Rowe, B., word, L.R. And Threlfall, E.J. (1997):
Multi drug – resistant *Salmonella typhi*; a world wide epidemic.
Clin. Infect. Dis.; 24. Suppl 1: S106-9.
- Roy, S.K.; Speelman, P., Butter, T., Nath, S. Rahman. H. and Stoll, B.J. (1985):
Diarrhea associated with typhoid fever.
J. Infect. Dis.; 151 (6): 1138-42.
- Rubin, F.A; Mc Whirter, P.D., Burr, D., Punjabi N.H., Lesmana, M., Sukri, and Hoffman, S.L.(1990):
Rapid diagnosis of typhoid fever through identification of *S. typhi* within 18 hours of specimen acquisition by culture of the mononuclear cell-platelet fraction of blood.
J. Clin. Microbiol.; 28: 825-27.
- Saedler, H.; Reif, HJ., Hu, S. and Davidson, N. (1974):
IS is a genetic element for turn off and turn on of genetic activity in *E. coli*.
Mol. Gen. Genet.; 132: 365-89.
- Sanderson K.E.; Sciore, P., Lui, S.L. and Hessel, A. (1993):
Location of IS200 on the genomic cleavage map of *Sal. typhimurim* LT2.
J.Bacteriol.; 175: 7624-28.
- Sawyer, S.A.; Dykhuizen, D.e., Dubose, R.F and Green, L. (1987):
Distribution and abundance of insertion sequences among natural isolates of *E. coli*.

Genetics.; 115-63.

- Saxena, M.; Gupta, J.K. and Vadehra, D.V. (1990):
Phospholipase, a production by isolates of *Sal* Species.
Indian. J. Med. Res.; 91(A): 177-81.
- Schiaffino, A.; Beuzn, C.R., uzgau, S., Leori, G., Coppuccinelli, P.,
Casadesus, J. and Rubino, S. (1996):
Strain typing with IS200 finger prints in *Salmonella abortusovis*
Appl. Environ. Microbiol.; 62(7): 2375-80.
- Schroder, S.A. (1968):
Interpretation of serologic test for typhoid.
JAMA.; 206 : 839-40.
- Schwarz, S. and Liebisch, B. (1994):
Use of ribotyping, IS200 typing and plasmid analysis for the
identification of *Salmonella enterica* subsp. *enterica* serovar
typhimurium vaccine strain Zoosaloral H and its differentiation
from wild type strains of the same serovar.
Int. J. Med. Microbiol. Viral. Parasitol. Inf Dis.; 28(4): 442-50.
- Senewirtane, B. and Senewritane, K. (1977):
Reassessment of the Widal test in the diagnosis of typhoid.
Gastroenterology.; 73: 233-36.
- Sharma, M., Datta, U., Roy, P., Verma, S. and Sehgal, S. (1997):
Low sensitivity of counter immunoelectrophoresis for
serodiagnosis of typhoid fever.
J. Med. Microbiol.; 46(12): 1039 – 42.
- Sharma, A.M. and Sharma, O.P.(1992):
Pulmonary mainfestation of typhoid fever.
Chest.; 101 (4): 1144-46.
- Shukla, S., Patel, B. and Chitnis, D.S. (1997):
100 years of Widal test and its reappraisal in an endemic area.
Indian. J. Med. Research. 105: 53-7.
- Song, J.H. (1994):
PCR diagnosis of typhoid fever.
J. Clin. Microbiol.; 32(8):2033.

- Song, J.H.; Cho H. Park, M.Y., Na, D.S., Moon, H.B. and Pai, C.H.(1993):
Detection of *S. typhi* in the blood of patient with typhoid fever by polymerase chain reaction.
J. Clin. Microbiol.; 31(6): 1439-43.
- Soria, G.; Barbe, J. and Gibert, H. (1994):
Molecular fingerprinting of *Sal. typhimurium* by IS200 – typing as a tool for epidemiological and evolutionary studies.
Microbiologia.; 10(1-2): 57-68.
- Stanley, J.; Baquar, N. and Burnens, A. (1995):
Molecular subtyping scheme for *Salmonella panama*.
J. Clin. Microbiol.; 33 (5): 1206-11.
- Stanley, J.; Baquar, N. and Threlfall, E.J. (1993):
Genotypes and phylogenetic relationships of *Salmonella typhimurium* are defined by molecular finger printing of IS200 and 16 *Srrn* loci.
J. Gen. Microbiol.; 139(pt 6): 133-40.
- Stanley, J.; Chowdry, N., Powell, N. and Threlfall, E.J. (1992):
Chromosomal genotypes of *Sal. berta*.
FEMS. Microbiol. Lett. 74 (2-3):247-52.
- Stanley, J.; Jones, C.S. and Threlfall, E.J. (1991):
Evolutionary lines among *Salmonella enteritidis* phage types are identified by insertion sequence IS200 distribution.
FEMS. Microbiol. Lett.; 66(1): 83-89.
- Stanley, J.; Powell, N., Jones, G. and Burnens, A.P(1994):
A framework for IS200, 16 *Sr* RNA gene and plasmid profile analysis in *Salmonella* serogroup DI.
J. Med. Microbiol.; 41 (2): 112-19.
- Stanley, J. and Saunders, U. (1996):
DNA insertion sequence and the molecular epidemiology of *Salmonella* and *Mycobacterium*.
J. Clin. Microbiol.; 45 : 236-51.
- Stone, G.G.; Chengappa, M.M., Oberst, R.D., Gabbert, N.H., Mc Vey, S., Hennessy, K.J., Muenzenberger, M. and Staats, J. (1993):
Application of polymerase chain reaction for the correlation of *Salmonella* serovars recovered from greyhound feces with their diet.

J. Vet. Diagn. Invest.; 5(3): 378-85.

Tacket, Co., Sztein, M.B., Losonsky, G.A., Wesserman, S.S., Nataro, J.P., Edelman, R., Pickard, D., Dougan, G., Chatfield, S.N. and Levine, M.M. (1997):

Safety of live oral *Salmonella typhi* vaccine strains.
Infect. and Immun.; 65(2): 452-6.

Takaski, L.; Grossman, N. and Leive, L. (1983):

Salmonella activate complement differentially via the alternative Pathway on the structure of their lipopolysaccharide O-antigen.
J. Immunol.; 130: 1867-72.

Tassios, P.T.; Markogiannakis, A., Vatopulos, A.C., Kotsanikau, E. Velenakis, E.N., Kourea kremastinou, J. and legakis, N.J. (1997):

Molecular epidemiology of antibiotic resistance of *Sal. enteritidis* during a 7 years period in Greece.
J. Clin. Microbiol.; 35(6): 1316-21.

Tenover, F.C. (1985):

Plasmid fingerprinting. A tool for bacterial strain identification and surveillance of nosocomial and community – acquired infections.
Clin. Lab. Med.; 5(3): 413-36.

Therlfall, E.J.; Torre, E., Ward, L.R., Djalos perez, A., Rowe, B. and Gibert, I. (1994):

Insertion sequence IS200 fingerprinting of *S.typhi*: an assessment of Epidemiological applicability.
Epidemiol. Infect.; 112(2): 1153-61.

Therlfall, E.J.; Torre, E., Ward, L.R., Rowe, B. and Gibert, I., (1993):

Insertion sequence IS200 can differentiate drug-resistant and drug - sensitive *Sal.typhi* of Vi-phage types E1 and M1.
J. Med. Microbiol.; (6): 454-58.

Tompkins, L.S.; Troup, N., Labigne-Russel, A. and Coben, M.L. (1986):

Cloned, random chromosomal sequences as probes to identify *Salmonella* species.
J. Infect. Dis.; 154(1): 156-162.

Tong, M.J.; Youel, D.B, and Cotton, G.L. (1972):

Acute pneumonias in tropical infection.
Am. J.Trop. Med Hyg.; 21: 50-57.

Torre, E.; Threlfall, E.J., Hampton, M.D., Ward, L.R., Gibert, I. and Rowe, B. (1993):

Characterization of *Salmonella virchow* phage types by plasmid profile and IS200 distribution.

J. Appl. Bacteriol.; 75(5): 435-40.

Tsang, R.S. and Chan, P.Y. (1981):

Serological diagnosis of typhoid fever by counter immunoelectrophoresis.

Br. Med. J.; 282: 150-7.

Tsang, R.S.; Chan, P.Y., Lam, S.k., Brooy, J. T. and Rowley, D. (1981):

Antibody response to lipopolysaccharide and protein. Antigens of *S.typhi* during typhoid infection measurement serum antibodies by radioimmunoassay.

Clin. Exp. Immunol.; 46: 508-14.

Van Lith, L.A. and Arats, H.J. (1994):

Polymerase chain reaction identification of *Salmonella* serotypes.

Lett. Appl. Microbiol.; 19(4): 273-76.

Waston, K.C. (1978):

Laboratory diagnosis of recovery *S.typhi* from

Weir

f, ES (1997):

Human intestinal epithelial

cell - derived interleukin - 6.

Wil:

Isolation of *Bact. typhosum* by means of bismuth sulphite medium in water and milk borne epidemics.

J. Hyg.; 38.; 507-19.

- Xi, W.; Ni, X., Iu, T., Zhang, J. Xu, J., Huang, M. and Wa, C. (1993):
Early diagnosis of typhoid fever using monoclonal Ab.
Chinese. Med. Sci. 8(2): 123-24.
- Zhang, X.L.; Morris, C. and Hackett, J. (1997):
Molecular cloning, nucleotide sequence, and function
of a site specific recombinase encoded in the major
" pathogenicity island " of *Salmonella typhi*
Gene.; 202(1-2): 139 -46