

References

REFERENCES

- Achterman C and Kalamchi A (1979):** Congenital deficiency of the fibula. J Bone Joint Surg; 61B: 133-137.
- Aldegheri R (2000):** Distraction osteogenesis for lengthening of the tibia in patients who have limb-length discrepancy or short stature. J Bone Joint Surg Am.; 82-A(8): 1195.
- Amstutz HC (1972):** Natural history and treatment of congenital absence of the fibula. In proceedings of the American Academy of Orthopaedic Surgeons. J Bone and Joint Surg; 54-A: 1349.
- Anderson M; Green, WT and Messner MB (1963):** Growth and prediction of growth in the lower extremities J bone Surg Am 45: 1-14. Apoll A les pseudoarthrosis congenitalis de jambe. Rev Chir Orthop; 56: 38-120, 1969.
- Birch JG; Walsh SH; Small JM; Morton A; Kock KD; Smith C; Cummings D and Buchanan R (1999):** Syme amputation for the treatment of fibular deficiency. An evaluation of long-term physical and psychological functional status. J. Bone and Joint Surg; 81-A: 1511-1518.

<i>References</i>

- Boakes JL; Stevens PM and Moseley RF (1991):** Treatment of genu valgus deformity in congenital absence of the fibula. *J Pediatr Orthop*; 11: 721-724.
- Bohne WH and Root L (1977):** Hypoplasia of the fibula. *Clin Orthop*; 125: 107-12.
- Caskey PM and Lester EL (2002):** Association of fibular hemimelia and clubfoot. *J Pediatr Orthop.*; 22(4): 522-5.
- Catagni MA (1992):** Management of fibular hemimelia using the Ilizarov method. In *Instructional course lectures, American Academy of Orthopaedic surgeons. Vol. 41, pp.431-434. Park Ridge, Illinois, American Academy of Orthopaedic surgeons.*
- Catagni MA; Bolano L and Cattaneo R (1991):** Management of fibular hemimelia using the Ilizarov method. *Orthop Clin of North America*; Vol. 22, No, 4: 715-722.
- Cheng JC, Cheung KW and Ng BK (1998):** Severe progressive deformities after limb lengthening in type-II fibular hemimelia. *J Bone Joint Surg Br.*; 80(5): 772-6.

<i>References</i>

Chio IH; Kumar J and Bowen JR (1990): Amputation or limb lengthening for partial or total absence of the fibula. J Bone Joint Surg; 72A: 1391-1399.

Choi IH, Lipton GE, Mackenzie W, Bowen JR and Kumar SJ (2000): Wedge-shaped distal tibial epiphysis in the pathogenesis of equinovalgus deformity of the foot and ankle in tibial lengthening for fibular hemimelia. J Pediatr Orthop.; 20(4): 428-36.

Choi IH; Kumar SJ and Bowen JR (1990): Amputation or limb-lengthening for partial or total absence of the fibula. J. Bone and Joint Surg; 72-A: 1391-1399.

Coventry MB and Johnson EW (1952): Congenital absence of the fibula. J Bone Joint Surg; 34A: 941-955.

Dalmonte A and Donzelli D (1988): Comparison of different methods of limb lengthening. J Pediatr Orthop; 8: 62-70.

Dutoit M; Pigault P and Padovani JP (1990): The fate of children undergoing bone lengthening in congenital hypoplasia of the legs. Rev Chir Orthop Reparatrice Appar Mot; 76: 1-7.

<i>References</i>

- Epps CH and Schneider PL (1989):** Treatment of Hemimelias of the lower extremity. J Bone Joint Surg; 71A: 273-277.
- Exner GU (2003):** Bending osteotomy through the distal tibial physis in fibular hemimelia for stable reduction of the hindfoot. J Pediatr Orthop B.; 12(1): 27-32.
- Gainé WJ and McCreath SW (1996):** Syme's amputation revisited: a review of 46 cases. J Bone Joint Surg Br.; 78(3): 461-7.
- Gardner E (1971):** Osteogenesis of the human embryo and fetus. In Bourne G: The Biochemistry and Physiology of bone New York, Academic Press.
- Grogan DP, Holt GR and Ogden JA (1994):** Talocalcaneal coalition in patients who have fibular hemimelia or proximal femoral focal deficiency. A comparison of the radiographic and pathological findings. J Bone Joint Surg Am.; 76(9): 1363-70.
- Guidera KJ; Brinker MR and Kouseff BG (1993):** Overgrowth management in Klippel-trenaunay-Weber and proteus syndromes. J Pediatr. Orthop; 13: 459-466.

<i>References</i>

Maffulli N and Fixsen JA (1991): Fibula hypoplasia with absent lateral rays of the foot. J Bone Joint Surg; 73B: 1002-1004.

Maudie D, Hamdy RC, Fassier F, Morin B and Duhaime M (1997): Management of fibular hemimelia: amputation or limb lengthening. J Bone Joint Surg Br.; 79(1): 58-65.

McCarthy JJ, Glancy GL, Chang FM and Eilert RE (2000): Fibular hemimelia: comparison of outcome measurements after amputation and lengthening. J Bone Joint Surg Am.; 82-A(12): 1732-5.

Michael C (2004): Fibular Hemimelia. EMedicine Specialists, Orthopedic Surgery, Knee.

Miller LS and Bell DF (1992): Management of congenital fibular deficiency by Ilizarov technique. J Pediatr Orthop; 12: 651-7.

Naudie D; Hamdy RC; Fassier F; Morin B and Duhaime M (1997): Management of fibular hemimelia: amputation or limb lengthening. J. Bone and Joint Surg; 79-B (1): 58-65.

<i>References</i>

- Ogden JA (1980):** Chondro-osseous development and growth. In Urist MR: fundamental and clinical bone physiology. Philadelphia JB Lippincott.
- Ogden JA and Conlogue GJ (1981):** Spontaneous Rickets in the wild arctic Fox: *Alopex lagopus*. *Skeletal Radiol.* 7:43-54.
- Ogden JA and Grogan DP (1987):** Prenatal development and growth of the musculoskeletal system. In Albright. JA. Brand RA (eds): The scientific basic of Orthopedics. New York Appleton and Lange.
- Ogden JA and Southwick WO (1977):** Endocrine dysfunction and slipped capital femoral epiphysis *Yale J Biol. Med.* 50: 1-16.
- Ogden JA; Conlogue GJ and Bronson ML (1979):** Radiology of postnatal skeletal development. III. The Clavicle *skeletal Radiol.* 4: 196-203.
- Oppenheim WL (1991):** Fibular deficiency and indications for Syme's amputation. *Prosthet Orthop Int.* 41: 435-6.
- Paley (2004):** Lengthening reconstruction surgery of the foot and ankle for fibular hemimelia. Asami, Turkey.
- Peterson HA (1990):** The evolution of femoral lengthening concept and techniques, in de pablos J, Canadell J (ed.) bone lengthening Current Trends and Controversies. Pamplona, Spain, Universal de Navarra. pp 15-18.

<i>References</i>

- Raymond TM and Stuart L W:** Lovell and Winter's Pediatric Orthopaedics. In: The Limb-Deficient Child. Chapter 30; p. 1218-1233.
- Roux MO and Carlloz H (1999):** Clinical examination and investigation of the cruciate ligaments in children with fibular hemimelia. J Pediatr Orthop.; 19(2): 247-51.
- Shapiro F; Haltrop M and Climcher M (1977):** Organization and cellular biology of the perichondrial Ossification Groove of Ranvier. J Bone Joint Surg; 59A: 703-723.
- Sharma M, MacKenzie WG and Bowen JR (1996):** Severe tibial growth retardation in total fibular hemimelia after limb lengthening. J Pediatr Orthop.; 16(4): 438-44.
- Simmons ED Jr, Ginsburg GM and Hall JE (1996):** Brown's procedure for congenital absence of the tibia revisited. J Pediatr Orthop.; 16(1): 85-9.
- Stanitski DF and Stanitski CL (2003):** Fibular hemimelia: a new classification system. J Pediatr Orthop; 23(1): 30-4.
- Stevens PM and Arms D (2000):** Postaxial hypoplasia of the lower extremity. J Pediatr Orthop.; 20(2): 166-72.

References

Wagner H (1978): Operative lengthening of the femur. Clin. Orthop. 136: 125-142.

Wolfgang GL (1984): Complex congenital anomalies of the lower extremities femoral bifurcation, tibial hemimelia and Diastasis of the ankle. Case report and review of the literature. J Bone Surg; 66A: 4534-458.