



References

- Anderson, R.H.; Becker, A.E. and Tynan, M. (1996):** Description of VSD by non-invasive index for combined systolic and diastolic ventricular function. J. Cardiol. 13: 267.
- Andro, A.; Pellet, ph. D.; Wiliam, G. and Tolar, B.S. (2004):** The Tei index methodology and disease state value Echocardiography, A Jour. Of CV ultrasound allind tech. vol. 21 No. 7.669.
- Bargeron, L.M.; Elliot, L.P.; Soto, B.; Beam, P.R. and Curry, G.C. (1999) :** Axial cineangiography in congenital heart disease. Section 1. concept, technical and anatomical considerations. Circulation 56: 1075- 1083.
- Barzoe, (2004):** Doppler derived myocardial performance index in healthy children in the shiraz (I. JNS) Vol. 24, No.2 85-89.
- Beker, E.J.; Ayton, V. and Smith, M.A. (1999):** Magnetic resonance imaging at a high field strength of VSD in infants. Br. Heart. J, 62: 305.
- Blackstone, E.H.; Kirklin, J.W.; Bradley, E.L.; Dushane, J.W. and Appelbaum, A. (2000):** Optimal age and results in repair of large ventricular septal defects. Journal of thoracic and cardiovascular surgery 72: 661-679.



References

- Brenner, L.D.; Caputo, G.R.; Mostbeck, G.; Steiman, D.; Dulce, M.; Cheitlin, M.D.; O'Sullivan, M. and Higgins, C.B. (1998):** Quantification of left to right atrial shunts with velocity-encoded cine nuclear magnetic resonance imaging. J Am Coll Cardiol. 20: 1246-1250.
- Bridges, N.D.; Perry, S.B.; Keane, J.F.; Goldstein, S.A.N. Nandell, V.; Mayer, J.E.; Jr, Jonas, R.A.; Castaneda, A.R. and Lock, J.E. (2000):** Preoperative transcatheter closure of congenital muscular ventricular septal defects N Eng J Med 1, 324: 1312.
- Bruch, C.; Schemender, A. and Mortin, D. (2000):** Tei index in patients with mild to moderate congenital heart failure. Euro. Heart Journ, 21, 1888- 95.
- Bryd, B.F. ; Schiller, N.B.; Botvinick, E.H. and Higgins, C.B. (1995):** Normal cardiac dimensions by magnetic resonance imaging. Am J Cardiol; 55: 1440-2.
- Chan, K. (2000):** Usefulness of the myocardial performance index for assessing left ventricular function in congenital heart disease. Am J Cardiol; 86: 654-658.
- Cook, D.R. (2000) :** Left ventricular wall motion velocities in healthy children measured by pulsed wave Doppler tissue echocardiography, rate. J Am Soc Echocardiogr; 13: 1002-11.
- Davis, P. (2000) :** Left ventricular function in endurance trained children by tissue Doppler imaging. Med Sci Sports Exerc; 36: 1507-1513.



References

- Docherty, J.R. and McGrath, J.C. (2002):** Value of the Doppler index of myocardial performance in the early phase of acute myocardial infarction. *J Am Soc Echocardiogr*; 13: 723-730.
- Dujardin, K.S.; Tel, C. and Yeo, T.C. (1998) :** Prognostic value of a doppler index combining systolic and diastolic performance in idiopathic- dilated cardiomyopathy. *Am J Cardiol*; 82: 658-664.
- Dushane, J.W.; Krongrand, E.; Ritter, D.G. and Me-Goon, D.C. (1999):** The fate of raised pulmonary vascular resistance after surgery in ventricular septal defect. In: Rowe RD, Kidd BSL, eds. *The child with congenital heart disease after surgery*. Mount kisco, New York: Futura.
- Eidem and Edwards (2001):** New index of combined systolic and diastolic myocardial performance: A simple and reproducible measure of cardiac function a study in normals and dilated cardiomyopathy. *J car diol*; 26:357-366.
- Eidem and Tei (1999):** Pulsed wave Doppler tissue echocardiography assessment of the long axis function of the right and left ventricles during the early neonatal period. *Hart*: 90: 175-180.
- Eidem, B.W.; Tei, C. and O'Leary, P.W. (1998):** Serial evaluation for myocardial performance in fetuses and neonates using a new Doppler index. *Pediatr Int*; 41: 722-727.
- Eillis, J.H.; Moodie, D.S.; Strerba, and R. Gill, C.C. (1998):** Ventricular septal defect in the adult: Natural and unnatural history. *Am Heart J*. 114: 115-120.



References

- Falkensammer and Paul (2001):** Doppler assessment of cardiac function at 11-14 weeks' gestation in fetuses with normal and increased nuchal translucency. *Ultrasound Obstet Gynecol*; 24: 390-398.
- Feigenbaum, H. (1998):** Echocardiography: Echocardiographic evaluation of cardiac chambers (5th edition) Lea & Febiger philadelphia- Baltimore- Hong Kong- London Sydney- Tokyo, p. 134-172.
- Felner, J.M.; Blusmentein, B.A. and Schlant, R.C. (2000):** Sources of variability in echocardiographic measurements. *Am. J. Cardiol.* 45: 995.
- Force, T.L.; Folland, E.D. and Abischer, U. (1991):** Echocardiographic assessment of ventricular function. In Marcus, M.L.; Skorton, D. J.; Schelbert, H.R. and Wolf, G.L. (eds): *Cardiac imaging: A comparison to Braunwold's heart disease*, Philadelphia, W.B. Saunders company, P. 347-401.
- Freedom, R.M.; Culham, J.M.G. and Moes, C.A.F. (1999):** Ventricular septal defect, Chapter 15, *Angiocardiography of Congenital Heart Disease*.
- Fyler, D.C. M.D. (1998):** Ventricular septal defect in Nadas pediatric cardiology, Edited by Donald C fyler, Md, Hanley and Belfus, Inc, Philadelphia, Mosby, Year Book, Inc, Snt Louis. Baltimore Boston, Chicage, London, Philadelfia, Syndney, Toronto, Ch. 28, p. 435-4557.



References

- Gardier, R.W. (2001):** Quantitative assessment of myocardial tissue velocities in normal children with Doppler tissue imaging. Am J Cardiol; 77: 1254-7.
- Gerad, P.; Aurgimma, M.; Pamelan, D.; William, H. (1997):** Quantitive evaluation of left ventricular structure, wall stress, and systolic function. Practical clinical echocardiography (Otto) 2nd ed., p. 1-24.
- Goresky, G. (2001):** Assessment of global left ventricular function in normal children and in children with dilated cardiomyopathy. J Am Soc Echocardiogr; 12: 1058-1064.
- Graham, T.P.; Bender, H.W. and Spatch, M.S. (1996):** Ventricular septal defect in Adams FH, Emmanouildes GC, Reimenschneider TA (eds) Moss Heart disease in infants, children and adolescent, Williams and Wilkins, Baltimore, London, H.G. P 189;
- Graham, T.P.; Jr, and Gutgesell, H.P. (1999):** Ventricular septal defects, Heart diseases in infants, Children and adolescent. Vol. 1, First edition, P. 724: 745; 1996.
- Grossman, W. (1990):** Shunt detection and measurement Cardiac catheterization and Angiography, third Edition, P. 155-169.
- Hardin, J.T.; Muskett, A.D.; Canter, C.E.; Martin, T.C. and Spray, T.L. (1992):** Primary surgical closure of large ventricular resptal defects in small infants, Ann Thorac Surg 53: 397-401.



References

- Harrison, D.A.; Liu, P.; Walters, J.E. et al., (2002) :** Cardiopulmonary function in adult patients late after Fontan repair. J Am Coll Cardiol; 26: 1016- 21. Full Text Bibliographic Links.
- Helbing, W.A.; Bosch, H.G.; Maliepaardc, et al., (1997):** Comparison of echocardiographic methods with magnetic resonance imaging for assessment of right ventricular function in children. Am. J. Cardiol; 76: 589-94.
- Heuvel, A.C. (1998):** Treatment of ventricular septal defects. Congenital Cardiovascular defects. 741: 746;
- Higgins, C.B. (1999) :** Magnetic resonance imaging in patients with congenital heart disease. Circulation (70): 851.
- Hirooka, K.;Yasumura, Y. and Tsujita, Y. (2001) :** An enhanced method for left ventricular volume and ejection fraction by triggered harmonic contrast echocardiography. Int J Cardiovasc imaging; 17: 253-61.
- Hoenecke, H.G.; Golderg, S.J. and Shan, D.J. (1992):** Effect of beam directional alterations on left ventricular shortening fraction. Am. J. Cardiol., 50: 1120.
- Hoffman, J.I.E. (1999):** Congenital heart disease. Incidence and inheritance pediatric. in. North Am. (37): 25.
- Hole, T. (2003):** Impaired left ventricular diastolic function in children with chronic renal failure. Kidney int; 65: 1461-1466.
- Jacksno, E.B. (2001):** Measuring left ventricular myocardial performance index in fetuses. Am J Cardiol; 91: 919-921.



References

Jensen, S.E. (2000) : Assessment of myocardial velocities in healthy children using tissue Doppler imaging. *Ultrasound Med Biol*; 26: 229-37.

Kalbunde, R. (2004): Abnormal left ventricular mass and aortic distensibility in pediatric dialysis patients. *Pediatr Nephrol*; 20: 64-68.

Kapusta, L.; Thijssen, J.M.; Cuypers, M.H. et al., (2000): Assessment of myocardial velocities in healthy children using tissue Doppler imaging. *Ultrasound Med Biol*, 26: 229-37. full text bibliographic links.

Keith, J.D. (1998): Ventricular septal defect in Heart disease in infancy and childhood. Edited by Keith JD, Rowe RD and Vlad P., 3rd (ed). Macmillan publishing Co. New York Collier Macmillan Canada, Ltd. Toronto Bailliere Tindal London. P. 320.

Kersting- Sommerhoff, B.A.; Diethelm, L.; Teitel, D.F.; Sommerhoff, C.P.; Higgins, S.S.; Higashino, S.S. and Higgins, C.B. (1999): Magnetic resonance imaging of congenital heart disease: sensitivity and specificity using receiver operating characteristic curve analysis. *Am Heart J*. 118: 155-161.

Khambadkone, S.; Li, J. de Leval, M.R. et al., (2003) : Basal pulmonary vascular resistance and nitric oxide responsiveness late after fontan- type operation. *Circulation*; 107: 3204-8. ovid full text bibliographic links.



References

- Kidd, I.; Driscoll, D.J. and Gersony, W.M. (1998):** Second natural history study of congenital heart defects: results of treatment of patients with ventricular septal defects. *Circulation* 87 (suppl 1): 138-151.
- Klein, J. (2001):** Prognostic value of global myocardial performance indices in acute myocardial infarction. Comparison to measures of systolic and diastolic left ventricular function. *Chest*; 124: 1645-1651.
- Kurt, Amplatz, James, H.; Moller Wilfrido, R. and Castaneda – Zuniga, (1998):** Ventricular septal defect in Radiology of congenital heart disease. Vol. 1 Ch. 9. P. 219-252.
- Lerman, J. (1999):** Quantitative assessment of left ventricular function: establishing normal values of the myocardial performance index in the infant Echocardiography; 18: 9-13.
- Levin, A.E.; Spach, M.S. and Canent, R.V. Jr, (1997):** Ventricular pressure flow dynamics in VSD. *Circulation*, 35: 430.
- Little, W.C. (2001):** Assessment of normal and abnormal cardiac function. In : Braunwald E, Heart disease: A textbook of cardiovascular medicine. Vol. 1. 6th ed. Philadelphia: WB. Saunders; 479-502.
- Lock, J.E.; Kean, J.F. and Fellow, K.E. (1998):** The use of catheter intervention procedures for congenital heart disease. *J Am Coll. Cardiol.* (7): 1420.



References

- Macleod, S.M. (2000):** Value of a Doppler derived index combining systolic and diastolic time intervals in predicting outcome in primary pulmonary hypertension. *Am J Cardiol*; 81: 1157-1161.
- Mancini, G.; Costello, D.; Bhargava, G.; Lew, W.; LeWinter, M. and Karliner, J. (1999):** The isovolumic index: a new noninvasive approach to the assessment of left ventricular function in man *Am J Cardiol*; 50: 1401-1408.
- Merchandise, B.; Schroder, E. and Bosl, A. (1998):** Early detection of doxorubicin cardiotoxicity. *Am. Heart J.* 11:892-898.
- Michele, M.D. (2003):** Usefulness of tissue Doppler echocardiography for evaluating ventricular function in children with heart disease. *Am J Cardiol*; 91: 570-4.
- Mikel, D. and Smith, M.D. (1997):** Left ventricular diastolic function-clinical by Doppler echocardiography. *The practical of clinical echocardiography (Otto) 2nd ed., P. 49-74.*
- Moller, J.E.; Sondergaard, E. and Poulsen, S.H. (2001) :** The Doppler echocardiographic myocardial performance index predicts left-ventricular dilation and cardiac death after myocardial infarction. *Cardiology*; 95: 105-111.
- Mori, K.; Hayabuchi, Y.; Kuroda, Y.; et al., (2000) :** Left ventricular wall motion velocities in healthy children measured by pulsed wave Doppler tissue echocardiography: normal values and relation to age and heart rate. *J Am Soc Echocardiogr*; 13: 1002-11. [ovid full text bibliographic links.](#)



References

- Nasr, M.A.; Hashish, H. and Dardir, M.D. (1999):** Role of Lung biopsy in the assessment of operability in patients with left to right shunt and border line pulmonary hypertension. Egypt Heart J. 39: 129. The Egyptian Society of Pediatric Cardiologists (Newsletter).
- Neutze, J.M.; Ishikawa, T.; Clrkson, P.M.; Calder, A.L.; Barratt-Boyes, B.G and kerr, A.R. (2000):** Assessment and follow- up of patients with ventricular septal defect and elevated pulmonary vascular resistance. Am J Cardiol; 63: 327; 1989.
- Nugent, E.W.; Plauth, W.H.; Edwards, J.E.; Schlant, R.C. and Williams, W.H. (1998):** Congenital Heart Disease in the heart arteries and veins. Edited by Hurst, J.W., Logue, R.B., Rackley, C.E., Schlan, R.C., Sonnenblick, E.H., Wallace, A.G., and Wengey, N.K. Sixth eidition. Mggraw- HILL Book Col. P.
- Park, J.K.; Dell, R.B.; Ellis, K. and Gersony, W.M. (1998):** Surgical management of the aorta and ventricular septal defect. J Am Coll Cardiol, 20: 176.
- Park, M.K. (1996):** Textbook of pediatric cardiology. 3rd edition St. Louis, Mosby- Year book. P. 250-259.
- Park, M.K. (1996):** Ventricular septal defect, pediatric cardiology, 3rd edition, page 135-142.
- Payne, K.; Mattheyse, F.J.; Libenberg, D. and Dawes, T. (2001):** Effect of preload alternations on a new Doppler echocardiographic index of combined systolic and diastolic performance. J Am Soc Echocardiogr; 135: 1065-1072.



References

- Pearson, A.; Gudipati, C.V. and Labovitz, A.J. (1997):** Systolic and diastolic flow abnormalities in patients with hypertensive hypertrophic cardiomyopathy. J. Am. Coll Cardiol 12: 989-995.
- Perloff, J.K. (1997):** Ventricular septal defect in the clinical recognition of congenital heart disease. Edited by perloff J.K., Second edition W.B. Saunders Company. Philadelphia. London, Toronto. P. 396.
- Poulsen, S.H.; Jensen, S.E. and Nielsen, J.C. (2000):** Serial changes and prognostic implications of a Doppler- derived index of combined left ventricular systolic and diastolic myocardial performance in acute myocardial infarction. Am J Cardiol; 85: 19-25.
- Razai and Weiss (2001):** Importance of the index of myocardial performance in evaluation of left ventricular function. Echocardiography; 19: 273-278.
- Rezai, K.; Weiss, R. and Stanford, W. (1999) :** Relative accuracy of three scintigraphic methods for determination of right ventricular ejection fraction : a correlative study with ultra fast computed tomography. J nucl med; 32: 429-35
- Salder, W. et al., (1998):** frontal section through the heart of a 6-mm embryo showing the primary interventricular foramen and the entrance of the antrum the primitive left ventricle .



References

- Senzaki, H.;** Masutani, S.; Kobayashi, J. et al., (2002) : Ventricular afterload and ventricular work in Fontan circulation : comparison with normal two- ventricle circulation and single- ventricle circulation with Blalock- Taussig shunts. *Circulation*; 105: 2885-92. Ovid Full Tex Bibliographic Links.
- Silverman, N.H. (2000):** Quantitative methods to enhance morphological information using M- mode, Doppler, and cross sectional ultrasound. In Silverman, N.H. (ed): *Pediatric Echocardiography*. Baltimore, Williams and Wilkins, pp. 35-108.
- Silverman, N.H.; Ports, T.A. and Snider, A.R. (1999):** Determination of left ventricular volume in children. Echocardiographic and angiographic comparisons. *Circulation*, 62: 584.
- Sivasubramanian, S. (1996):** Ventricular septal defect. Madras, India, October1.
- Smith, V.E.; White, W.B.; Mecran, M.K. and Karimeddin, M.K. (1996):** Improved left ventricular filling accompanies reduced left ventricular mass during therapy of essential hypertension. *J. Am. Cardiol.*, 81: 1449-1454.
- Snider, A.R.; Serwer, G.A. and Ritter, S.B. (1996):** Defects in cardiac septation in: *Echocardiography in pediatric Heart Disease*. (sec.ed.), Editros Snider A R.; Serwer, G.A; Ritter S B. Year Book Medical publishers, Inc. Chicago, London, Boca Raton, Littleton.
- Soto, B. and Pacifico, A.D.(2000):** Ventricular septal defect. *Echo cardiography in Congenital Heart Malformations*, 327-349.



References

Stoylen, A. and Skjaerpe, T. (2003): Systolic long axis function of the ventricle: global and regional information. Scand cardiovasc J 2003, 37: 253-8. ovid full text bibliographic links.

Tei and Ling (2002): Fetal tissue Doppler echocardiography: detection rates of cardiac structures and quantitative assessment of the fetal heart. Ultrasound Obstet Gynecol; 21: 26-32.

Tei and Nishimura (2002): Noninvasive Doppler- derived myocardial performance index: Correlation with simultaneous measurement of cardiac catheterization measurements. J Am Soc Echocardiogr; 10: 169-178.

Thomas, R.; Kimball, M.D.; Stephen, R.; Daniel, M.D.; Richard, A.; Meyer, M.D.; David, W.; Hannon, M.D.; Philip Khoury, M.S.; and Schwartz, D.C. (1991): Relation of symptoms to contractility and defect size in infants with ventricular septal defect. Am J Cardiol. (67): 1097- 1102.

Tsevdos, E.J. (2001): Late ventricular geometry and performance changes of functional single ventricle assessed by magnetic resonance imaging. J Am Coll Cardiol; 28: 212-21.

Tutschek, B. (2003): Characterization of left ventricular diastolic function by tissue Doppler imaging and clinical status in children with hypertrophic cardiomyopathy. Circulation; 109: 1756-62.

Wahr, D.W. and Schiller, N.B. (1997): Left ventricular volumes determined by two- dimensional echocardiography in normal adult population. J. Am. Coll. Cardiol., 1: 863-868.



References

Weissler, A. (2000): Myocardial performance index in pediatric patients after cardiac transplantation. J Am Soc Echocardiogr; 17: 439-422.

Weissler, A.; Harnis, W. and Shcoenfeld, C. (1998): Systolic time intervals in beart failure in man. Circulation; 37: 149-159.