

REFERENCES

REFERENCES

- Aach, R.D.; Stevens, C.E.; Hallinger, B.F. et al. (1981):*
Hepatitis C virus infection in post-transfusion hepatitis. An analysis with first and second generation assays. The New England Journal of Med., 325: 1325-9.
- Abdel-Wahab, M.; Hashem, A.; Kamal, M. et al. (1993):*
Frequency of anti HCV among school children in Menofeya. International Conference on Schistosomiasis, Cairo, Egypt, P. 91 (Abst.).
- Abdel-Wahab, M.F.; Zakaria, S.; Kamel, M. et al. (1994):*
High seroprevalence of hepatitis C infection among risk groups in Egypt. Am. J. Trop. Med. Hyg., 51 (5): 563-67.
- Abu El-Fotouh, A. (1995):*
Hepatitis C virus genotyping in Egyptian patients with chronic liver disease. M.Sc. Thesis Tropical medicine, 1995, Cairo University.
- Adachi, Y.; Harii, K.; Takahashi, Y.; Tanihata, M.; Ohba, Y. and Yamamoto, T. (1980):*
Serum glutathione S-transferase activity in liver disease. Clin. Chem. Acta., 106: 243-55.
- Ahmed, L.; Salama, H.; Hassan, E. et al. (1995):*
The diagnostic value of HCV IgM antibody detection in patients with acute and chronic HCV infection. The journal of Egyptian medical association, 78 (16): 153-63.
- Akahane, Y.; Kojima, M.; Sugai, Y. et al. (1994):*
Hepatitis C virus infection in spouses of patients with type C chronic liver disease. Ann. Intern. Med., 120: 748.
- Akira, H.; Naboru, S.; Hiromu, Y.; Naok, Y.; Takuji, T.; Hideki, M. (1993):*
Immunoblot analysis of placental form of glutathione S-transferase in protein extracted from paraffin-embedded human glioma tissue. Cancer Res. Clin. Oncol., 119: 493-6.

- Alberti, A. and Realdi, G. (1991):**
Parenterally acquired NANB (Type C) hepatitis. In McIntyre, N.; Benhamou, J.P.; Bircher, J. eds. Oxford Textbook of Clinical Hepatology, Oxford University Press, 605-17.
- Al-Nasser, M.N. (1992):**
Intrafamilial spread of hepatitis C virus a major mode of spread in the Saudi Arabia population. Ann. Trop. Paed., 12-211.
- Alter, M.J.; Hadler, S.C.; Francis, D.P. et al. (1985):**
Epidemiology of NANB hepatitis in USA. Prog. Clin. Biol. Res., 182: 71.
- Alter, H.J.; Purcell, R.H.; Shih, J.W. et al. (1989):**
Detection of antibody to HCV in prospectively followed transfusion recipients with acute and chronic NANB hepatitis. The New England Journal of Medicine, 321: 1492-500.
- Alter, M.J.; Hadler, S.C.; Judson, F.N. et al. (1990):**
Risk factors for acute NANB hepatitis in United State., JAMA, 264-2231.
- Alter, M.J. (1991):**
Inapparent transmission of HCV. Foot Prints in the Sand. Hepatol., 14 (2): 389-91.
- Alter, M.J.; Margolis, H.S.; Krawczynski, K. et al. (1992):**
The natural history of hepatitis C in the united state. N. En. J.J. Med., 327-1899.
- Ban, N.; Takahashi, Y.; Takayama, T. et al. (1996):**
Transection of glutathione S-transferase pi antisense complementary DNA increases the sensitivity of a colon cancer cell line to adriamycin, cisplatin, melaphalan and etoposide. Cancer Res., 56 (15): 3577-82.
- Barabara, J.A. and Garson, J.A. (1993):**
PCR and transfusion microbiology. Vox Sanguinis; 64: 73-81.
- Bassily, S.; Hyams, K.C.; El-Masry, N.A. et al. (1992):**
HCV and hepatosplenic schistosomiasis. Scand. J. Infect. Dis., 24: 687-88.

- Bassily, S.; Dunm, M. and Farid, Z. (1993):**
Chronic hepatitis B in patients with schistosomiasis. J. Trop. Med. Hyg., 86: 67-71.
- Beach, M.J.; Meeks, E.L.; Mimms, L.T. et al. (1992):**
Temporal relationships of HCV RNA and antibody responses following experimental infection of chimpanzees. J. Med. Virol., 36: 226-37.
- Beckett, G.J. and Hayes, J.D. (1987):**
Plasma glutathione S-transferase measurements and liver disease in man. J. Clin. Biochem., 2: 1-24.
- Beckett, G.J.; Hayes, J.D. (1993):**
Glutathione S-transferase" Biochemical applications. Adv. Clin. Chem., 30: 281-380.
- Beckett, G.J.; Hussey, A.J.; Howie, A.F.; et al. (1989):**
Measurements of glutathione S-transferase B₁ in plasma after birth asphyxia: an early indication of hepatocellular damage. Clin. Chem., 35 (6): 995-99.
- Beckett, G.J.; Foster, G.R.; Hussey, A.T. et al. (1989):**
Plasma glutathione S-transferase and proteinase more sensitive than alanine aminotransferase as markers of paracetamol induced liver damage. Clin. Chem., 35 (11): 2186-9.
- Beckett, G.J.; Donovan, J.W.; Hussey, A.J. et al. (1990):**
Intravenous N-acetylcysteine, hepatotoxicity and plasma glutathione S-transferase in patients with paracetamol over dosage. Hum. Exp. Toxicol., 9 (3): 183-6.
- Bernvil, S.S.; Andrew, V.J. and Sasich, F. (1994):**
Second generation anti-HCV screening in Saudi Arabia donor population. Vox. Sang., 66: 33.
- Blacker, K.L.; Olson, E.; Vessey, D.A. and Boyer, T.D. (1991):**
Characterization of glutathione S-transferase in cultured human keratinocytes. J. Invest. Dermatol., 97 (3): 442-6.
- Blight, K.; Lesniewski, R.; La, Brooy, J.T. et al. (1994):**
Detection and distribution of HCV specific antigens in naturally infected liver. Hepatol., 20: 553.

- Board, P.G. and Mannervik, B. (1991):**
The contribution of the C-terminal sequence to the catalytic activity of glutathione S-transferases a human alpha class glutathione S-transferase. *Bioch. J.*, 275 (1): 171-4.
- Boyer, T.D. (1989):**
The glutathione S-transferase an update. *Hepatology*, 9 (3): 486-96.
- Brettler, D.B.; Mannucci, P.M.; Gringeri, A. et al. (1992):**
The low risk of HCV transmission among sexual partners of HCV infected haemophiliac males: An international multicenter study. *Blood*, Jul. 1990 (2): 540-43.
- Brillanti, M.; Masci, C.; Ricci, P. et al. (1992):**
Significance of IgM antibody to HCV in patients with chronic hepatitis C. *Hepatology*, 15: 998-1001.
- Brillanti, S.; Foli, M.; Giani, S. et al. (1993):**
Persistent hepatitis C viraemia without liver disease. *Lancet*, 341: 464-65.
- Brown, B.R. and Gandolfi, A.J. (1987):**
Adverse effects of volatile anesthetics. *Br. J. Anaesth.*, 59: 14-23.
- Buffet, G.; Charnaux, N.; Laurent-Puig, S. et al. (1994):**
Enhanced detection of antibodies to HCV by use of a third generation recombinant immunoblot assay. *J. Med. Virol.*; 43: 259-61.
- Bukh, J.; Purcell, R.H. and Miller, R.H. (1993):**
At least 12 genotypes of hepatitis C virus predicted by sequence analysis of the putative E1 gene of isolates collected worldwide. *Proc. Natl. Acad. Sci.*, 90: 8234-48.
- Buller, A.; Clapper, M.L. and Tew, K.D. (1987):**
Glutathione S-transferase in nitrogen mustard-resistant and sensitive cell lines. *Mol. Pharmacol.*, 31: 575-8.

- Caccuri, A.M.; Aceta, A.; Piemonte, F.; Di-Ilio, C.; Rosato, N. and Federici, G. (1990):**
Interaction of hemin with placental glutathione S-transferase.
Eur. J. Biochem., 189 (3): 493-7.
- Campbell, E.; Takahashi, Y.; Abramovitz, M.; Peretz, M. and Listowsky, I.A. (1990):**
Distinct testis and brain Mu class glutathione S-transferase.
Molecular cloning and characterization of a form present even
in individuals lacking hepatic type Mu isoenzymes. J. Biol.
Chem., 265 (16): 9188-93.
- Camps, J.; Cordoba, J. and Esteban, J.I. (1993):**
Pathophysiology of HC. In progress in Hepatology by J.P.
Miguet and D. Dhumeaux, Published by Libbey Eurotext,
Paris, 63-68. Newsletter of Egyptian Society of Hepatology,
Vol. 1, No. 3.
- Canada, A.T.; Roberson, K.M.; Vessela, R.L.; Trump, D.L.; Robertson, C.N. and Fine, R.L. (1996):**
Glutathione S-transferase in benign and malignant prostate.
Biochem. Pharmacol., 51(1): 87-90.
- Caramelo, C.; Alberola, M.L.; Bermejilla, T. et al. (1994):**
Evidence against transmission of HCV through hemodialysis
ultrafiltrate and peritoneal fluid. Nephron; 66: 470-73.
- Cha, T.A.; Beall, E.; Irvine, J. et al. (1992):**
At least 5 related but distinct HCV genotypes exist. Proc. Natl.
Acad. Sci., 89: 7144.
- Chen, D.S.; Kuo, G.C.; Sung, J.L. et al. (1990):**
HCV infection in an area hyperendemic for HBV and chronic
liver disease. The Taiwian experience. J. Infect. Dis., 162:
817-22.
- Chen, P.J.; Wang, J.T.; Hwang, L.H. et al. (1992):**
Transient immunoglobulin M antibody response to HCV capsid
antigen in posttransfusion hepatitis C: Putative serological
marker for acute viral infection. Proc. Natl. Acad. Sci. USA., 1,
89 (13): 5971-75.

- Chen, P.J.; Hwang, L.H.; Chang, L.Y. et al. (1993):**
Immunoglobulin M antibodies against HCV antigens: significance and application. *Gastroenterol.*, 28: Suppl. 5: 67-70.
- Chibla, J.; Ohba, H.; Matsura, Y. et al. (1991):**
Serodiagnosis of HCV infection with an HCV core protein molecularly expressed by a recombinant. *Baculovirus*, *Proc. Natl. Acad. Sci.*, 88: 4641-5.
- Choo, Q.L.; Kuo, G.; Weiner, A.J. et al. (1989):**
Isolation of cDNA clone derived from a blood borne NANB viral hepatitis genome. *Science*, 244: 359-62.
- Choo, Q.L.; Weiner, A.J.; Overby, L.R. et al. (1990):**
Hepatitis C virus: The major causative agent of viral NANB hepatitis. *Br. Med. Bull.*, 46: 423-41.
- Choo, Q.; Richman, K.; Han, J. et al. (1991):**
Genetic organization of hepatitis C virus. *Proc. Natl. Acad. Sci. USA*, 88: 2451-5.
- Choo, Q.L.; Richman, K.H.; Han, J.H. et al. (1991):**
Genetic organization and diversity of the HCV. *Proc. Natl. Acad. Sci. USA.*, 38: 2451-55.
- Clemenens, J.M.; Tuskur, S.; Chau, K. et al. (1992):**
IgM antibody response in acute hepatitis C viral infection. *Blood*, 79: 169-72.
- Darwish, M.A.; Raouf, T.A.; Rushdy, P. et al. (1993):**
Risk factor associates with a high seroprevalence of HCV infection in Egyptian blood donors. *Am. J. Trop. Med. Hyg.*, 49: 440.
- Deakin, M.; Elder, J.; Hendrickse, C.; et al. (1996):**
GSTT, genotypes and susceptibility to cancer carcinogenesis. 17 (4): 881-4.
- De-Luca, M.; Ascienne, A.; Vacca, C. et al. (1992):**
Are health care workers really at risk of HCV infection? *Lancet*, 339: 1364.

- Denis, F.; Aussel, L.; Ranger, S. et al. (1994):**
Prevalence of antibodies to HCV among patients with leprosy in several African countries and the Yemen. *J. Med. Virol.*, 43-1.
- Dienstag, J.L. (1983):**
NANB hepatitis II: Experimental transmission, putative virus agents markers and prevention. *Gastroenterol.*; 85: 743-68.
- Dow, B.C.; Fellett, E.A.C.; Jordon, T. et al. (1994):**
Testing of blood donation for hepatitis C virus. *Lancet*, 343: 477-178.
- Driven, H.A.; Dictus, E.L.; Boaders, N.L. et al. (1995):**
The role of human glutathione S-transferase isoenzymes in the formation of glutathione conjugates of the alkylating cytostatic drug thiotepa. *Cancer Res.*, 55 (8): 1701-6.
- Dusheiko, G.M. (1992):**
Hepatitis C virus. Recent advances in gastroenterology. Pounder R. ed. Churchill Livingstone, 195-216.
- Dusheiko, G.M. and Rizzetto, M. (1992):**
The management of chronic hepatitis C. Synopses in viral hepatitis, Adelphi communications, LTD: 1-20.
- Dusheiko, G.; Schmilovitz, W.H.; Braun, D. et al. (1994):**
HCV genotypes: An investigation of type specific difference in geographic origin and disease. *Hepatology*, 19 (1): 13-18.
- Ebeling, F.; Naukkarinen, R.; Leikola, J. et al. (1990):**
RIBA for hepatitis C virus antibody as predictor of infectivity (Letter) *Lancet*, 335: 982-3.
- El-Khashab, M.; Mourad, A.; Soliman, H. et al. (1995):**
Prevalence of HCV infection among military personnel with acute hepatitis. 13th annual congress of Egyptian Society of hepatology. Cairo, 19-21 April.
- Ellis, L.A.; Brown, D.; Conrad, J.D. et al. (1990):**
Prevalence of hepatitis C in South Africa: Detection of anti-HCV in recent and stored serum. *J. Med. Virol.*, 32: 249-51.

- El-Gohary, A.; Hassan, A.; Nooman, Z. et al. (1995):**
HCV genotypes in acute and chronic hepatitis C infection in Suez. 2nd Annual meeting of the ESTIP, January, 1995, Cairo.
- El-Sahly, A.M.; Nour El-Din, M.A.B.; Abel-Rahman, H.N. et al. (1991):**
Prevalence of antibodies to HCV in Egyptian patients with schistosomiasis. The 9th National Congress of the Egyptian Society of Hepatology, 1991.
- El-Zayadi, A.; Selim, O.; Rafik, M. and El-Haddad, S. (1992):**
Prevalence of HCV among NANB related chronic liver disease in Egypt. Hepatol., 14 (2): 417.
- Engvall, E. and Berlmán, P. (1971):**
Enzyme linked immunosorbent assay (ELISA). Quantitative assay of IgG. Immunochemistry, 8: 781-4.
- Esteban, J.I.; Esteban, R.; Viladomir, L. et al. (1989):**
Hepatitis C virus antibodies among risk groups in Spain. Lancet, 2: 294-7.
- Fan, K.C.; Huang, Y.C.; Li, C.H. (1995):**
Radioimmunoassay for plasma glutathione S-transferase Pi and its clinical application in gastrointestinal cancer. Cancer, 76 (8): 1363-7.
- Farci, P.; Alter, H.J.; Wang, D. et al. (1991):**
A long term study of HCV replication in NANB hepatitis. N. Eng. J. Med., 325: 98-104.
- Feucht, H.H.; Polywka, S.; Zollner, B. and Laufs, R. et al. (1994):**
Greater amount of HCV RNA in tears compared to blood. Microbiol. Immunol., 38 (2): 157-8.
- Fleischner, G.M.; Kamisaka, K.; Gamaitan, Z. and Arias, I.M. (1976):**
Immunological studies of rat and human ligandin in glutathione metabolism and function. Edited by Arias, I.M. Raven Press, New York, 259-69.
- Fuchs, K.; Mortz, M.; Schreier, E.; Zachoval, R. et al. (1991):**
Characterization of nucleotide sequences from European HCV isolates. Gene, 103: 163-9.

- Gaffer, Y.A.; Fattoh, S.A.; Kamel, M. et al. (1991):**
The impact of endemic schistosomiasis with acute viral hepatitis. *Am. J. Trop. Med. Hyg.*, 45 (6): 743.
- Garloni, G.; Iacovacci, S.; Sargiacomo, M. et al. (1993):**
Susceptibility of human liver cell culture to HCV infection. *Arch. Virol: Suppl.*, 8: 31-9.
- Garson, J.A.; Tedder, R.S.; Briggs, M. et al. (1990):**
Detection of HCV sequences in blood donation by nested PCR and prediction of infectivity. *Lancet*; 335: 1419-22.
- Gerber, M.A. and Thung, S.N. (1992):**
Immunopathology of hepatitis C. Immunology and liver. Falk symposium No. 70, Basel, 1992, Abstract book: 19-20.
- Gerlich, W.H. and Thomssen, A. (1991):**
Terminology, structure and laboratory diagnosis of hepatitis virus. In McIntyre, N.; Benhamou, J.P.; Bircher, J.; Rizzetto, M.; Rodes, J., eds. *Oxford Textbook of Clinical Hepatology*, Oxford University Press, 537-65.
- Giralt, M.; Cervello, I.; Nogues, M.R. et al. (1996):**
Glutathione S-transferase and reactive oxygen species of human scalp sebaceous glands in male pattern baldness. *J. Inves. Dermatol.*; 107 (2): 154-8.
- Girardi, E.; Zoccarelli, M.; Tossini, G. et al. (1990):**
HCV infection in travenous drug users, prevalence and risk factors. *Scand. J. Infect. Dis.*, 22: 751.
- Guesden, J.L.; Ternynck, T. and Avrameas, S. (1979):**
The use of avidin-biotin interaction in immunoenzymatic techniques. *J. Histochem. Cytochem.*, 27: 1131-9.
- Habig, W.H.; Kanisaka, K.; Ketley, J.N. et al. (1976):**
The human hepatic glutathione S-transferase in glutathione metabolism and function. Raven Press, New York; 223-31.
- Habig, W.H.; Pabst, M.J. and Fleischer, G.M. (1974):**
The identity glutathione S-transferase B with ligandin a major binding protein of liver. *Proc. Natl. Acad. Sci. USA*, 71: 3879-82.

- Habig, W.H.; Pabst, M.J. and Jacoby, W.B. (1974):**
Glutathione S-transferase. A first enzymatic step in mercapturic acid formation. *J. Biol. Chem.*, 249: 7130-9.
- Harrison, D.J. and Hayes, P.C. (1990):**
Immunological classification of glutathione S-transferase in human renal and liver diseases. In: Hayes, J.D.; Hayes, P.G.; Mantle, T.J. and Pickett, C.B. *Glutathione S-transferase and drug resistance*. London: Taylor and Francis, 431-41.
- Hasebe, C. and Sekiya, C. (1994):**
Diagnosis of type C hepatitis using PCR technology. *Rinsho-Byori*, 42 (3): 223-9.
- Hassan, N.F. (1993):**
Prevalence of HCV antibodies in patient groups in Egypt. *Trans. Soc. Trop. Med. Hyg.*, 87: 638.
- Hayes, J.D. and Mantle, T.J. (1986):**
Use of immunoblot techniques to discriminate between the glutathione S-transferase and study to their distribution in extrahepatic tissues. Evidence of three immunologically distinct groups of transferase in rat. *Bioch. J.*, 23: 779-88.
- Hayes, J.D.; Gilligan, D.; Chaman, B.J. and Beckett, C.J. (1983):**
Purification of human hepatic glutathione S-transferase and the development of a radioimmunoassay for their measurement in plasma. *Clin. Chem. Acta*, 134: 147-121.
- Hayes, P.C.; Husser, A.J.; Keating, J. (1988):**
Glutathione S-transferase levels in autoimmune chronic active hepatitis: a more sensitive index of hepatocellular damage than aspartate aminotransferase. *Clin. Chem. Acta.*, 172: 211-16.
- Hayes, P.C.; Harrison, D.J.; Bouchier, L.A.; McLellan, L.I. and Hayes, J.D. (1989):**
Cytosolic and microsomal glutathione S-transferase isoenzyme in normal liver and intestinal epithelium. *J. G. Gut*, 30: 854-59.
- Hayes, J.D.; Kerr, L.A. and Cronshaw, A.D. (1989):**
Evidence that glutathione S-transferase B₁B₁ and B₂B₂ are the products of separate genes and that their expression in human liver is subject to inter-individual variation. *Biochem. J.*, 264: 431-45.

- Hayes, P.C.; Hayes, J.D.; Hussay, A.J. et al. (1990):**
Changes in plasma glutathione S-transferase B1 concentration after alcohol ingestion in man, a measure of hepatocellular sensitivity to chronic alcohol excess. Clin. Chem. Enzymol. Commun., 2: 189-94.
- Hayes, P.C.; Bouchier, I.A. and Beckett, G.J. (1991):**
Glutathione S-transferase in human in health and disease. Gut, 32: 813-18.
- Hayes, P.C.; May, L.; Hayes, J.D. and Harrison, D.J. (1991):**
Glutathione S-transferase in human liver cancer. Gut, 32 (12):1546-9.
- Hibbs, R.G.; Carwin, A.L.; Hassan, N.F. et al. (1993):**
The epidemiology of antibody to hepatitis C in Egypt. J. Infect. Dis., 168: 789.
- Hirmamatsu, N.; Hayashi, N.; Haruna, Y. et al. (1992):**
Expression of HCV-related antigen and liver damage in patients with chronic hepatitis C. Hepatology, 16: 544.
- Hollinger, F.B.; Mosley, J.W.; Szmuress, W. et al. (1982):**
NANB hepatitis following blood transfusion. In: International symposium. Szumness, W.; Alter, H.J.; Maynard, J.Z. eds. Franklin Institutes Press, Philadelphia, 361-376.
- Holt, D.E.; Hawie, A.F.; Beckett, G.J. et al. (1995):**
Measurement of fetal plasma levels of glutathione S-transferase B1 as an indicator of decay to the liver caused by hypoxia in utero. Fetal Diagn. Ther., 10 (1): 11-16.
- Houghton, M.; Weiner, A.; Han, J. et al. (1991):**
Molecular biology of the hepatitis C virus: Implication for diagnosis, development and control of viral disease. Hepatology, 14: 381-88.
- Howie, A.F.; Patrick, A.W.; Fisher, B.M. et al. (1989):**
Plasma hepatic glutathione S-transferase concentrations after insulin-induced hypoglycemia in normal subjects and diabetic patients. Diabetic Med., 6 (3): 224-7.

- Howie, A.F.; Spencer, E. and Beckett, G.J. (1992):**
Aspartate aminotransferase, alanine aminotransferase and glutathione S-transferase in plasma during and after sedation by low dose isoflurane or midazolam. Clin. Chem., 38 (4): 476-9.
- Hussey, A.J.; Aldridge, L.M.; Paul, D.; Ruy, D.C.; Beckett, G.J. and Allan, L.G. (1988):**
Plasma glutathione S-transferase concentration as a measured of hepatocellular integrity following using general anaesthesia with halothane, enflurane or isoflurane. Br. J. Anaesth., 60: 130-5.
- Iersel, M.L.; Plomen; Struik, I.; Van Amersfoort, C. et al. (1996):**
Inhibition of glutathione S-transferase activity in human melanoma cells by alpha, beta unsaturated carbonyl derivatives. Chem. Biol. Interact., 102 (2): 117-32.
- Jakobason, P.J.; Mancini, J.A.; Ford-Hutchinson, A.W. (1996):**
Identification and characterization of a novel human microsomal glutathione S-transferase with leukotriene C4 synthase activity and significant sequence identity to 5-lipoxygenase-activity protein and leukotriene C4 synthase. J. Bio. Chem., 271 (36): 22203-10.
- Jakoby, W.B. and Habig, W.H. (1980):**
Glutathione S-transferase. In Jakoby, W.B. ed. Enzymatic basis of detoxification, Vol. 2 New York. Academic Press, 83-94.
- Jeffrey, B.; Wheatly, M.; Kathleen, K. et al. (1994):**
Examination of glutathione S-transferase isoenzyme profiles in human liver using high performance affinity chromatography. Journal of Chromatography. A., 663: 53-63.
- Jeffers, I.J.; Cheinquer, H.; Hunt, W. et al. (1992):**
Correlation of ALT level and severity of hepatitis histology with a new quantitative HCV-RNA method in patients with chronic hepatitis C. Hepatol., 16: 551.
- Jeffers, L.J.; Perez, G.O.; De Medina, M.D. et al. (1990):**
HCV infection in two urban hemodialysis units. Kid. International, 38: 320.

- Jendrassik, L. et al. (1938):**
Measurement of serum bilirubin. *Biochem. Z.*, 297-81.
- Ji, Y.; Akerboom, T.P.; Sies, H. (1996):**
Microsomal formation of S-nitroglutathione from organic nitrites; possible role of membrane bound glutathione S-transferase. *Biochem. J.*, 313(2): 377-80.
- Jochen, A.B.B. (1992):**
Occupationally-acquired hepatitis C virus infection. *Lancet*, 339: 304-305.
- Juronen, E.; Tasa, G.; Uskula, M. et al. (1996):**
Purification, characterization and tissue distribution of human class theta glutathione S-transferase T1-1. *Biochem. Mol. Biol. Int.*, 39 (1): 21-9.
- Kai-Chun, M.; Ying-Chun, Hai, L. (1995):**
Radioimmunoassay for plasma glutathione S-transferase pi and its clinical application in gastrointestinal cancer. *Cancer*, 76 (8): 1363-7.
- Kamel, M.A.; Ghaffar, Y.A.; Wassef, M.A. et al. (1992):**
High HCV prevalence in Egyptian blood donors. *Lancet*, 340:327.
- Kamisaka, K.; Habig, W.H.; Ketley, J.N. et al. (1975):**
Multiple forms of human glutathione S-transferase and their affinity for bilirubin. *Eur. J. Biochem.*, 60: 153-61.
- Kao, J.H.; Chen, P.J.; Loi, M.Y. and Chen, D.S. (1993):**
Superinfection heterogenous HCV in a patient with chronic type C hepatitis. *Gastroenterology*, 105: 583-87.
- Kapolowitz, N.; Percy, R.O. and Jantti, N.S. (1973):**
Role of hepatic anion-binding protein in bromulphthalen conjugation. *Exp. Med.*, 138: 483-7.
- Kaprell, H.P.; Camps, J.; Simpson, B. et al. (1993):**
Characterization of a new assay for the specific detection of HCV core IgM antibodies. Conference on viral hepatitis. Venice, Italy, No. 93.

- Katayama, T.; Mazda, T.; Kikuchi, S. et al. (1992):**
Improved serodiagnosis of NANB hepatitis by an assay detecting antibody to hepatitis C virus core antigen. *Hepatology*, 15: 391-94.
- Kato, N.; Hijikata, M.; Ootsuyama, Y. et al. (1990):**
Molecular cloning of the human HCV genome from Japanese patients with NANB hepatitis. *Proc. Natl. Acad. Sci. USA*, 87:9524-28.
- Kato, N.; Ootsuyama, Y.; Tanaka, T. et al. (1992):**
Marked sequence diversity in the putative envelope protein of hepatitis C virus. *Virus-Res.*, 22 (2): 107-23.
- Kato, N.; Yokosaka, O.; Hasada, K. et al. (1993):**
Quantification of HCV competitive reverse transcription-polymerase chain reaction: Increase of the virus in advanced liver disease. *Hepatology*, 18: 16-20.
- Kaziol, D.E.; Holland, P.V.; Alling, D.W. et al. (1986):**
Antibody to hepatitis B core antigen as a paradoxical marker for NANB hepatitis agents in donated blood. *Ann. Int. Med.*, 104: 488-495.
- Keefe, E.B. and Benner, K.G. (1993):**
Hepatitis A through E: distinguishing the etiologic agents. *Modern Medicine of the middle east*, 10: 48-62.
- Keir, M. (1992):**
Identification of molecular species of porcine luteal glutathione S-transferase and its hormonal regulation by hypophyseal gonadotropins. *Hokkaido Igaku-Zasshi*, 67 (5): 595-605.
- Kelen, G.D.; Green, G.B.; Purcell, R.H. et al. (1992):**
Hepatitis B and hepatitis C in emergency department patients. *N. Engl. J. Med.*, 320 (21): 1399-404.
- Kiyosawa, K.; Sodeyama, T.; Tanaka, E. et al. (1991):**
Intrafamilial transmission of hepatitis C virus in Japan. *J. Med. Virol.*, 33: 114-116.

- Kiyosawa, K.; Sodeyama, T.; Tanaka, E. et al. (1991):**
HCV in hospital empolyers with needlestick injuries. *Ann. Intern. Med.*, 115: 367.
- Koziel, M.J.; Dudley, D.; Choo, Q.L. (1993):**
Recognition of HCV proteins by intrahepatic cytotoxic T-lymphocytes in persons with chronic hepatitis. *Hepatology*, 18, No. 4, P 12. AASLD Abstract, 111A: 218.
- Krawczynski, K.; Beach, M.; Di Bisceglie, E.A. et al. (1990):**
Hepatocellular HCV related antigen and antibody viral specificity and relation to natural history of infection. *Hepatology*, 12: A905.
- Kuhnl, P.; Sedl, S.; Stangel, W. et al. (1989):**
Antibodies to hepatitis C virus in German blood donors. *Lancet*, 2: 324.
- Kuo, G.; Choo, Q.L.; Alter, H.J. et al. (1989):**
An assay for circulating antibodies to a major etiologic virus of human NANB hepatitis. *Science*, 244: 362-64.
- Lee, W.P.; Lee, C.L.; Lin, H.C. (1996):**
Glutathione S-transferase and glutathione peroxidase are essential in the early stage of adriamycin resistance before glycoprotein over expression in HOB1 lymphoma cells. *Cancer, Chemother. Pharmacol.*, 38 (1): 45-51.
- Lee, S.D.; Trai, Y.T.; Hwang, S.J. et al. (1991):**
A prospective study of post transfusion NANB (type C) hepatitis following cardiovascular surgery in Taiwan, *J. Med. Virol.*, 33: 188.
- Lefskowitch, J.H.; Schiff, E.R.; Davis, G.L. et al. (1993):**
Pathological diagnosis of chronic hepatitis C: A multicenter comparative study with chronic hepatitis B. *Gastroenterology*, 104: 595-603.
- Lin, H.H.; Koo, J.H.; Hsu, H.Y. et al. (1994):**
Possible role of higher maternal viraemia in perinatal transmission of hepatitis C virus. *J. Infect. Dis.*, 169: 638.

- Lizuka, T.; Sato, T. and Sugimoto, M. (1991):**
Glutathione S-transferase in alcoholic fatty liver. *Arukoni. Kenkyuto, Yakubutsu, Ison.* 26 (3): 428-46.
- Loguercio, C.; Caporaso, N.; Tuccillo, C. et al. (1998):**
Alpha glutathione transferases in HCV-related chronic hepatitis: a new predictive index of response to interferon therapy? *J. Hepatol.*, 28 (3): 390-5.
- Lok, A.S.; Ma, O.C.K.; Chan, T.M. et al. (1991):**
Overestimation of the prevalence of antibody to HCV in retrospective studies on stored sera. *Hepatol.*, 14: 756-62.
- Lowry, O.H. Rosebrough, N.J.; Farr, A.L. and Randall, R.J. (1951):**
Protein measurement with Folin phenol reagent. *J. Biol. Chem.*, 193: 265-75.
- Lucey, M. and Traber, P.G. (1991):**
Detection of hepatitis C virus RNA by PCR. *Hepatol.*, 13: 193-5.
- Mannervik, B. (1985):**
Identification of three calsses of cytosolic glutathione S-transferase common to several mammalian species. *Proc. Natl. Acad. Sci., USA*, 82: 7202-6.
- Maple, P.A.; Mckee, T.; Desselberger, U. et al. (1994):**
HCV infections in transplant patients: serological and virological investigations. *J. Med. Virol.*, 44: 43.
- McGill, D.B.; Rakalu, J.; Zinsmeister, A.R. and Ott, B.T. (1990):**
A21 year experience with major hemorrhage after percutaneous liver biopsy. *Gastroenterology*, 99: 1396-400.
- McHatchinson, J.G.; Person, J.L.; Govindarjan, S. et al. (1992):**
Improved detection of HCV antibodies in high risk population. *Hepatol.*, 15: 19-25.
- Meikle, I.; Hayes, J.D. and Walker, S.W. (1992):**
Expression of an abundant alpha class glutathione S-transferase in bovine and human adrenal cortex tissues. *J. Endocrinol.*, 132 (1): 83-92.

- Mellbye, M.; Biggaro, R.J.; Watzin, P. et al. (1990):**
Sexual transmission of hepatitis C virus: Cohort Study (1981-1989) among European homosexual men. *Br. Med. J.*, 301:210.
- Mera, N.; Ohmori, Itahashi, K. et al. (1994):**
Immunochemical evidence for the occurrence of Mu class glutathione S-transferase in human fetal liver. *J. Biochem. Tokyo*, 116 (2): 315-20.
- Meyer, D.J.; Coles, B.; Pemble, S.E. et al. (1991):**
Theta a new class of glutathione S-transferase purified from rat and man. *Biochem. J.*, 274 (2): 409-14.
- Michel, V.; Olivieri, C.; Isabelle, B. et al. (1995):**
Plasma alpha glutathione S-transferase assessed as a marker of liver damage in patients with chronic hepatitis. *C. Clin. Chem.*, 41 (12): 1716-19.
- Mitra, A.; Govindwar, S.; Joseph, P. and Kulkarni, A. (1991):**
Inhibition of human term placental and fetal liver glutathione S-transferase purified from rat and man. *Biochem. J.*, 274 (2): 409-14.
- Morgenstern, R.; DePierre, J.W. and Journuall, N. (1985):**
Microsomal glutathione S-transferase primary structure. *J. Biol. Chem.*, 13976-83.
- Mori, S.; Kato, N.; Yagyn, A. et al. (1992):**
A new type of HCV in patients in Thailand. *Biochemical, Biophysical research. Communications*, 183: 334-42.
- Mulder, T.P.; Verpaget, H.W.; Sier, C.F. et al. (1995):**
Glutathione S-transferase Pi in colorectal tumour is predictive for overall survival cancer. *Res.*, 55 (12): 2696-702.
- Mulder, P.J.; Wilbert, H.M.; Peters, Daniel, A. and Jan, B. (1996):**
Sandwich ELISA for glutathione S-transferase alpha1: plasma concentration in control and patients with gastrointestinal disorders. *Cancer, Res.*, 1996.
- Muratori, L.; Giostra, F.; Cataleta, M. et al. (1994):**
Testing for HCV sequences in peripheral blood mononuclear cells of patients with chronic hepatitis C in the absence of serum hepatitis C virus RNA. *Liver*, 14: 124-8.

- Murray, J.M.; Rowlands, B.J.; Trinick, T.R. (1992):**
Indocyanine green clearance and hepatic function during and after prolonged anaesthesia. Comparison of halothane with isoflurane. *Br. J. Anaesth.*, 68 (2): 168-71.
- Mursonen, M.; Wolf, H.; Autrup, H. (1996):**
Inhibition of glutathione S-transferase activity in human melanoma cell. *Chem. Biol. Interact.*, 102 (2): 116-32.
- Nagayama, R.; Miyake, K.; Tsuda, F. and Okamoto, H. (1994):**
IgM antibody to HCV core peptide (CP14) for monitoring activity of liver disease in patients with acute or chronic HCV. *J. Med. Virol.*, 42 (3): 311-17.
- Nakatsuji, Y.; Matsumoto, A.; Tanaka, E. et al. (1992):**
Detection of chronic HCV infection by four diagnostic system: First and second generation ELISA, second generation RIBA and nested PCR analysis. *Hepatol.*, 16: 300-5.
- Nare, B.; Smith, J.M. and Prichard, R.K. (1992):**
Mechanism of inactivation of schistosoma mansoni and mammalian glutathione S-transferase activity by the antischistosomal drug otipraz. *Biochem. Pharmacol.*, 43 (6): 1345-51.
- Nelson, D.R.; Lim, H.L.; Oliver, D. et al. (1995):**
Alpha glutathione S-transferase as a marker of hepatocellular damage in chronic hepatitis C-virus infection. *Am. J. Clin. Pathol.*, 104 (2): 193-8.
- Niitsu, Y.; Takahashi, Y. and Sairo, T. (1989):**
Serum glutathione S-transferase as a tumour marker for gastrointestinal malignancies. *Cancer*, 63: 317-23.
- Novati, R.; Thiero, V.; Maisonneuve, P. et al. (1992):**
Mother to child transmission of HCV detected by nested polymerase chain reaction. *J. Infect. Dis.*, 165: 720-23.
- Ohna, T.; Mizokami, M.; Suzuki, K. et al. (1993):**
Four groups and eight genotypes of hepatitis C virus. *Hepat.*, 18 (4). Pt. 2 AASLD Abstract 83 A.

- Ohno, T.; Mizokami, M.; Tibbs, C.J. et al. (1994):**
New genotype of HCV in South Africa. J. Med. Virol., 42: 409-13.
- Ogata, N.; Alter, H.J.; Miller, R.H. and Purcell, R.H. (1991):**
Nucleotide sequence and mutation rate of the H strain of hepatitis C virus. Natl. Acad. USA, 88: 3392-6.
- Okamoto, M.; Baba, M.; Kodama, E. et al. (1993):**
Detection of HCV genome in human serum by multi-targeted PCR. J. Med. Virol., 41 (1): 6-10.
- Okamoto, H.; Kurai, K.; Okada, S. et al. (1992):**
Full length sequence of HCV genome having poor homology to reported isolates: comparative study of four distinct genotypes. Virology, 188 (1): 331-41.
- Okamoto, H.; Sugiyama, Y.; Okada, S. et al. (1992):**
Typing HCV by PCR with type-specific primers: Application to clinical surveys and tracing infectious sources. J. Gen. Virol., 73 (3): 673-9.
- Overby, L.R. (1992):**
Hepatitis C : looking at a virus that hasn't been seen, viral hepatitis management: Standard for the future Cannes, France, PP, 11-12.
- Patel, A.; Sherlock, S.; Dusheiko, G. et al. (1991):**
Clinical course and histological correlations in post-transfusion hepatitis C: The Royal free hospital Experience. Eur. J. Gastroenterol. Hepatol., 3: 491-95.
- Pawlotsky, J.M.; Ben Yahia, M.; Andre, C.; Voisin, M.C.; Intrator, L. et al. (1994):**
Immunological disorders in C virus chronic active hepatitis. Hepatology, 19: 841-8.
- Pekka, T. and Kimmo, K. (1994):**
Ultrasensitive time- resolved immunofluorometric assay of glutathione S-transferase, alpha in serum. Clin. Chem., 40 (2): 184-9.

- Plamen, J.H.; Van Ommen, B. and Van Bladeren, P.J. (1991):**
Irreversible inhibition of human glutathione S-transferase isoenzymes by tetrachloro-1,4-benzoquinone and its glutathione conjugate. *Biochem. Pharmacol.*, 41 (11): 1665-9.
- Polesky, H.F. (1990):**
Would another test (anti-HCV) have helped? Editorial comment. *Am. J. Clin. Pathol.*, 93 (1): 155.
- Pontisso, P.; Morsica, G.; Cavalletto, D. et al. (1992):**
HCV-RNA and liver disease in chronic asymptomatic carriers of anti-HCV. *Hepatology*, 16: 584 (Abstr.).
- Poterucha, J.J.; Rakela, J.; Lumeng, L. et al. (1992):**
Diagnosis of chronic hepatitis C after liver transplantation by the detection of viral sequences with PCR. *Hepatology*, 15: 42-45.
- Quiroga, J.A.; Campillo, M.L.; Catillo, I. Et al. (1991):**
IgM antibody to HCV in acute and chronic hepatitis C. *Hepatology*, 14: 38-43.
- Ramzy, M.F.; Ramsis, R.; Mahrous, A. et al. (1995):**
HCV liver disease in transplant recipient. 13th Annual congress of Egyptian Society of Hepatology. Cairo, 19-21, April.
- Rollins, D.E. and Buckpitt, A.R. (1979):**
Live cytosol catalyzed conjugation of reduced glutathione with a reactive metabolite of acetaminophen. *Toxicol. Appl. Pharmacol.*, 47: 331-9.
- Saeed, A.A.; Al-Admawi, A.M.; Al-Rasheed, A. et al. (1991):**
Hepatitis C virus infection in Egyptian voluntary blood donors in Riyadh. *Lancet*, 338: 459-60.
- Saleh, M.; Pereiral, L. and Tibbs, C. (1994):**
High prevalence of hepatitis C virus in normal libyan population. *Transactions of the Royal society of tropical medicine and hygiene*, 88: 292-94.

- Soma, Y.; Satoh, K. and Sato, K. (1986):**
Purification and subunit structural and immunological characterization of fine glutathione S-transferases in human liver and the acidic form as a hepatic tumour marker. *Biochem. Biophys. Acta.*, 869: 247-58.
- Sanchez-Quijano, A.; Rey, C.; Agudo, I. et al. (1990):**
Hepatitis C virus infection in sexually promiscuous groups. *Eur. J. Clin. Microbiol. Inf. Dis.*, 9: 610-12.
- Sankery, T.; Romeo, J.; Ulrich, P. et al. (1992):**
Detection of HCV in seronegative donors with raised ALT. *Lancet*, 340: 249-50.
- Senjo, M.; Ishibashi, T. and Imai, Y. (1985):**
Purification and characterization of cytosolic liver protein facilitating heme transport into apocytochrome C5 from mitochondria. *J. Biol. Chem.*, 260: 9191-6.
- Shapiro, C.N. (1994):**
Transmission of hepatitis virus. *Ann. Intern. Med.*, 120(1):82.
- Sherlock, S. and Dooley, J. (1993):**
Diseases of the liver and biliary system. Ninth edition. Blackwell Scientific Publications.
- Sherman, M.; Bass, N.M.; Campbell, J.A. and Kirsch, R.E. (1983):**
Radioimmunoassay of human ligandin. *Hepatology*, 3: 162-9.
- Sherman, M.; Campbell, J.A.; Timuss, S.A. et al. (1983):**
Glutathione S-transferase in human hepatocellular carcinoma. *Hepatology*, 3(2): 170-6.
- Simmonds, P.; Alberti, A.; Alter, H.J. et al. (1994):**
A proposed system for the Nomenclature of hepatitis C viral genotypes. *Hepatology*, 19: 1321-25.
- Simmonds, P.; Holmes, E.C.; Cha, T.A. et al. (1993):**
Classification of HCV into six major genotypes and a series of subtypes by phylogenetic analysis of NS5 region. *J. Gen. Virol.*, 74 (11): 2391-9.

- Simmonds, P.; Modmish, F.; McGullough, P. et al. (1992):**
Contamination of immune-assay control with hepatitis C virus.
Lancet, 334-1607.
- Simmonds, P.; Zhango, L.Q.; Waston, H.G. et al. (1990):**
Hepatitis C quantification and sequencing in blood products,
Haemophiliacs and drug abusers. Lancet, 336: 1469-72.
- Singh, S.V.; Leal, T. and Awasthi, Y.C. (1980):**
Inhibition of human glutathione S-transferase by bile acids.
Toxicol. Appl. Pharmacol., 5: 248-54.
- Skidmore, S.J.; Collingham, K.E. and Drake, M. (1994):**
Sexual transmission of HCV. J. Med. Virol., 42: 247-48.
- Soni, P.N.; Khakoo, S.I. and Dusheiko, G.M. (1994):**
Viral hepatitis. Recent advances in Gastroenterology, Pounder
Red. Churchill Livingstone, 235-66.
- Stegers, E.A.P.; Mulder, T.P.J.; Biseeling, J.G.A. et al. (1995):**
Glutathione S-transferase as a marker in hepatocellular damage
and HELLP syndrome. Department of Obstetrics/ Gynaecology
and Gastroenterology / Hepatology. University Hospital St.
Radboud Lancet, 345: 1571-2.
- Stevens, C.E.; Aach, R.D.; Hullinger, F.B. et al. (1984):**
Hepatitis B virus antibody in blood donors and the occurrence
of NANB hepatitis in transfusion recipients: An analysis of the
transfusion transmitted viruses. Ann. Intern. Med., 101:733-8.
- Stevens, C.E. (1990):**
Perinatal and sexual transmission of HCV in Hollinger R.B.,
Lemon S.M.; Margolis eds. Viral hepatitis and liver diseases.
Baltimor, 1990; Williams and Wilkins. Co.
- Study Group, UK Haemophilia Center (1989):**
Review of ovirucidols for inactivation of viruses in blood
derivatives. Lancet, 1: 814.
- Suginoto, M. (1995):**
Glutathione S-transferases (GSTs). Nippon-Rinsho, 53 (5):
1253-9.

- Sugimoto, M.; Shimada, N.; Aikawa, K.; Sugiyama, Y. (1993):**
Relationship between content of hepatic glutathione S-transferases and the kinetics of indocyanine green elimination in various liver diseases. *Biopharm- Drug-Dispos.*, 14 (7): 567-78.
- Sugitani, M.; Inchanspe, G.; Shindo, M. and Prince, A. (1992):**
Sensitivity of serological assay to identify blood donors with hepatitis C Viraemia. *Lancet*, 339: 1018-19.
- Szasz, G. and Persijn, J.P. (1974):**
Measurement of Gamma-Glutamyl Transferase. *Clin. Chem. Clin. Biochem.*, 12: 228.
- Takada, N.; Takase, S. and Takada, A. (1992):**
Clinical features of acute NANB non-C and non-E Probably type F hepatitis. *Hepato.*, 16: 597.
- Takahashi, M.; Yamada, G.; Miyamoto, R. et al. (1993):**
Natural course of chronic hepatitis C. *The American J. of Gastroenterology*; 88, 2: 240-43.
- Takamizawa, A.; Mori, C.; Fuke, I. et al. (1991):**
Structure and organization of HCV genome isolated from human carriers. *J. Virol.*, 65: 1105-13.
- Tamai, K.; Satoh, K.; Tsuchida, S. et al. (1990):**
Specific inactivation of glutathione S-transferase in class Pi by SH-modifiers. *Bioch. Biophys. Res. Commun.*, 167 (1): 331-8.
- Tanaka, E.; Kiyosawa, K.; Sodeyama, T. (1991):**
Significance of antibody to hepatitis C virus in Japanese patients with viral hepatitis, relationship between anti-HCV antibody and the prognosis of non-A, non-B post-transfusion hepatitis. *Med. Virol. J.*, 33: 117-22.
- Tedder, R.S.; Gilson, R.J.C.; Briggs, M. et al. (1991):**
Hepatitis C virus: Evidence for sexual transmission. *Bn. Med. J.*, 302: 1299-302.
- Thaler, M.I.; Landers, D.W.; Wara, D. et al. (1990):**
Transmission of HCV detected by polymerase chain reaction. *Hepatology*, 12: A829.

- Tholer, M.M.; Park, C.K.; Lunders, D.V. et al. (1991):**
Vertical transmission of HCV. *Lancet*, 338: 17-18.
- Thorburn, D.; Bird, G.L.; Spence, E. et al. (1996):**
Alpha glutathione S-transferase levels in chronic hepatitis C infection and the effect of alpha interferone therapy. *Clin. Chem. Act.*, 253(1-2): 171-80.
- Tiainen, P. and Karhi, K.K. (1994):**
Ultrasensitive time-resolved immuofluorometric assay of glutathione transferase alpha in serum. *Clin. Chem.*, 40 (2): 184-9.
- Tiainen, P.; Lindgren, L. and Rosenberg, P.H. (1995):**
Disturbance of hepatocellular integrity associated with propofol anaesthesia in surgical patients. *Acta. Anaesthesiol. Scand.*, 39 (6): 840-4.
- Tibbs, C.J.; Polmer, S.J.; Cocker, R. et al. (1991):**
Prevalence of hepatitis C in tropical communities, the importance of confirmatory assay. *J. Med.*, 34: 143.
- Tori, J.; Libre, J.M.; Carbonell, M. et al. (1990):**
Sexual transmission of hepatitis C virus and its relation with HBV and HCV. *B.M.J.*, 301: 1130-33.
- Trepo, C. (1992):**
Diagnostic marker of viral hepatitis B and C viral hepatitis management: standards for the future. *Cancers*, 1992, Abstract book. Macclesfield, U.K. Adelphi Communications, LTD, 21-22.
- Trull, A.K.; Facey, S.P.; Rees, F. et al. (1994):**
Serum alpha-glutathione S-transferase a sensitive marker of hepatocellular damage associated with acute liver allograft rejection. *Transplantation*, 58 (12): 1345-51.
- Vander Jagt, D.L.; Hunsaker, L.; Carcia, K.B. and Roger, R.E. (1985):**
Isolation and characterization of the multiple glutathione S-transferase from human liver. *J.B. Chem.*, 260: 11603-10.

- Van der Poel, C.L.; Cuypers, H.T.; Lelie, P.N. et al. (1991):*
Confirmation of HCV infection by new four-antigen
recombinant immunoblot assay. *Lancet*, 337: 317-19.
- Van der Poel, C.L.; Cuypers, H.T. and Reesink, H.W. (1994):*
Hepatitis C virus six years on. *Lancet*, 344: 1475-79.
- Van der Poel, C.L.; Reesink, H.W.; Lelie, P.N. et al. (1990):*
Anti-HCV and transaminase testing of blood donors. *Lancet*,
336: 187-8.
- Vaubourdolle, M.; Chazouilleres, O.; Briaud, I. et al. (1995):*
Plasma alpha-glutathione S-transferase assessed as a marker of
liver damage in patients with chronic hepatitis C. *Clin. Chem.*,
41 (12 pt. 1): 1716-9.
- Vitale, C.; Tricerri, A.; Marongello, M. et al. (1993):*
Epidemiology of HCV infection in dialysis units, First versus
second generation assays. *Nephrol.*, 64: 315.
- Valler, A.; Bidwell, D.E. and Engvall, E. (1974):*
A microplate method of enzyme linked immunosorbent assay
and its application to malaria. *Bull. Wld. Org.*, 51: 209-11.
- Wang, Y.L.; Lee, S.D.; Hwang, S.J. et al. (1994):*
Incidence of post-transfusion hepatitis before and after
screening for hepatitis C virus antibody. *Vox Sang*, 67: 187-90.
- Weiland, O.; Zhang, Y.Y. and Widell, A. (1993):*
Serum HCV RNA levels in patients with chronic hepatitis C
given a second course of alpha IFN treatment after relapse
following initial treatment. *Scand. J. Infect. Dis.*, 25: 25-30.
- Weiner, A.J.; Han, J.; Alan, J. et al. (1990):*
HCV testing in low risk population. *Lancet*, 336: 695.
- Weiner, A.J.; Kuo, G.; Bradley, D.W. et al. (1990):*
Detection of HCV sequences in NANB hepatitis. *Lancet*, 335:
1-3.

- Wejstal, R.; Hermodsson, S. et al. (1990):**
Mother to infant transmission of HCV infection. *J. Med. Virol.*, 30: 178-180.
- Wheatly, J.B.; Kelley, M.K.; Montali, J.A. et al. (1994):**
Examination of glutathione S-transferase isoenzyme profiles in human liver using high performance affinity chromatography. *J. Chromatogr. A.*, 663 (1): 53-63.
- Wilkinson, J.H. (1976):**
Measurement of AST in: Principles and practice for diagnosis enzymology. Year Book Medical Publishers.
- William and Robert (1996):**
Identification and characterization of human microsomal glutathione S-transferase. *J. Biol. Chem.*, 271 (36): 22203-10.
- Wolff, C.; Schluter, K.; Prohaska, W. and Kleesiek, K. (1992):**
Improved detection of HCV RNA by RT-PCR. *European J. Clin. Chem. and Clin. Biochem.*, 30: 717-27.
- Yanagi, M.; Kaneko, S.; Unoura, M. et al. (1991):**
HCV in fulminant hepatic failure. *N. Engl. J. Med.*, 324:1895.
- Yano, M.; Yastsuhashi, H.; Inow, O. et al. (1991):**
Epidemiology of HCV in Japan: Role in chronic liver diseases and hepatocellular carcinoma. *J. Gastroenterol. and Hepatol. Suppl.*, 1: 31-35.
- Yoshioka, K.; Aiyama, T.; Okumura, A. et al. (1996):**
Hypervariable region sequence in cryoglobulin-associated hepatitis C virus in sera of patients with chronic hepatitis C: relationship to antibody response against hypervariable region genome. *Hepatology*, 24 (6): 1346-50.
- Yoshioka, K.; Kakumu, S.; Wakita, T. et al. (1992):**
Detection of HCV by PCR and response to interferon alpha therapy: Relationship to genotypes of HCV. *Hepatology*, 16: 293-99.

Zakaria, S.; Mabrouk, M.A.; Watts, D. et al. (1994):
Acute sporadic NANB hepatitis in Egyptian military recruits
hospitalized at Almaza fever hospital in Cairo, Egypt. Military
Med. J., 39 (1): 54.

Zuckerman, A.J. (1990):
Hepatitis C virus a giant leap formed. Hepatol., 11 (2): 320-22.