

REFERENCES

REFERENCES

- Als, F.C.; Edwolhy, S.M. and Bloch, D.A. (1986):*
The American Rheumatism Association 1987 Revised Criteria for the classification of Rheumatoid Arthritis. *Arthriti Rheum*, 315-24.
- Als, O.S.; Rills, B.J.; Gofredsen, A. Christiansen, C, Deflens, L.J. (1988):*
Biochemical markers of bone turnover in rheumatoid arthritis. *Acta Med. Scand*, 219:209-13.
- Bab, I., Passi-Even, L.; Gazit, D. Sekels, E.; Ashoton, B., Play Lan-Ramu, N.; Ziv, I. and Ulmansk, (1988):*
Osteogenesis in vivo diffusion chamber cultures of human maceow cells. *Bone Miner.* 4:373.
- Bady, J.J.; Cleeren, A. Pot, M. and Borkowskia (1987):*
Serum Osteocalcin in tumor. Associated hypercalcemia, *J. Bone Miner Res* 2 (1) 486 A.
- Barham, D. and Trinder P. Bartel, H. and Bohmer M., (1972):*
Clin. Chem. Acta 37: 193.
- Bataille, R. Delmas, P. and Sany, J. (1987):*
Serum BCP in multiple meloma. *Cancer*, 59:32934.
- Bayram, M. Ylimaz, N. Erbagci, A.B.; Kilincer, M.S. (1999):*
Diagnostic value of biochemical markers of bone turnover and postmenopausal osteoporosis *Clin-Chem-lab-Med.* Feb; 37 (2): 137-43.
- Biuret. (1977):* Determination of total serum protein.

Black, A., Schokecht, P.A., Ralston, S.L. and Shapses, S.A. (1999):
Diurnal Variation with age a difference in the biochemical markers
of bone turnover. *Analyst*, Feb. 7 (2) 140-5.

Black, D. Duncan, A. Robins, S.P. (1988):
Quantitative analysis of the pyridinium cross links of collagen in using
ion-pair reversed-phase high performance liquid chromatography.
Anal Biochem, 167-203.

Bonde, M. Ovist P. Fledelusus, C. Riis; B.J.; Christanses, C. (1995):
Application of an enzyme immunoassay for a new marker of bone
resorption cross-link: follow-up on hormone replacement therapy and
osteoporosis risk assesment, *J. clin. Endocrinol Metab.* 80:864-868.

*Borsalino, G.; Borzoni, R. Caravaggiof, Rocchi, R. Pedrazzoni, M.
palummeri, E. and Passevie, M. (1985):*
Changes in serum osteocalcin after orthopedic surgery procedures, *J*
Clin. Med. Bologana, 66:393-400.

Brincat, M. and Studd, J.W.W. (1988):
Menopause a multi-system disease obstet. *Gynecol.* 2:289-316.

*Brown, J.P.; Delmas, P.D.; Malaval, L.; Edouard, C.; Chapuy, M.C. and
Meunier, P.J. (1984):*
Serum bone Gla Protien: a specific marker for bone formation in
postmenopausal osteoporosis. *Lancet*, I. 109-3.

Buritit, W.J.; Brady, T.G. and Oral, Off, J.J. (1994):
Immunochemical characterization of cirulating parathyroid hormone
related protien in patient with cancer, *N. Engl. J. Med*, 322:1106-
1112.

Cann, C.E.; Genant, H.K.; Kolb, F.O.; and Ettinger, B. (1985):
Quantitive computed tomography for prediction of vertabral fracture
risk. *Bone* 6:1.

Charbon, S.A.; Delnas, P.D.; Malaval, L. Chavassieux, P.M. Arlot, M. Chapuy, Meunier, P.J.; Serum bone Gla-protein in renal osteodystrophy (1986):

Comparison with bone histomorphometry. J. Clin-Endocrinol Metab, 63:892-7.

Christiansen, C. and Juel, Riis, B. (1990):

Risk assessment using biochemical analysis. In smith, R. (ed) osteoporosis, PP. 158-61. (London: Royal Collage of physicians).

Christiansen, C.; Chistainsen, M.S.; Mc Nair, P. et. Al. (1990):

Prevention of early post menopausal bone loss controlled 2-year study in 315 normal females, European, J. Clin. Invest, 10:273-279.

Clark & Collip (1925): Determination of Ca in urine by periceptation by oxalate and titration.

Coleman, R.E.; Whitaker, K.B.; Moss D.R.; Mashiter, G.; Fogelaman, I. and Rubens, R.D. (1988):

Biochemical predication of responses of bone metastases to treatment. Brit, J. Cancer, 58:205-10.

Colle, M.; Ruffe, A. Rudas, E. and Chebbo, M. (1987):

Differences in osteocalcin levels according to age and sex Arch Fr Pediat, 44:839-41.

Consensus conference, Osteoporosis (1984):

J. Am. Med. Assoc, 252, 799-802.

Cooper, C.; Campion, G. and Melton, L. (1992):

J 3rd. hip Frcture in the edarly: A world wide projection Osteoporo. Int. Nov; 2 (6): 285-9.

Dela Piedra, C. Toural, V. and Rapado, A. (1997):

Osteocalcin and urinary hydroxyproline creatinine ratio in the differential diagnosis of primary hyperparathyroidism and hypercalcemia of malignancy. Scand J Clin lab invest, 47:587-92.

Delmas, P.D. (1988):

Biochemical markers of bone turnover in osteoporosis in: Riggs, B.L. Melton 12 J. eds. Osteoporosis New York Raven press, 297-316.

Delmas P.D. (1991):

Biochemical markers of bone turnover: Methology and clinical use in osteoporosis.

Am. J Med. 91 (Supl 58) 595-636.

Delmas, P.D.; (1995):

Biochemical markers for bone turnover in Riggs, B.L.; Melton, L.J.; Editors. Osteoporosis: etiology, diagnosis, and managmnt. Philadelphia: lippincot Revan, 319-33.

Delmas, P.D.; Gienyts, E.Bertholin, A. Garners, P. Marchand, F. (1993):

Immunossay of Pyridinoline excretion in normal adults and paget's disease. J Bone Miner Res. 8:643-648.

Delmas, P.D.; Selnner, D. Wahner, H.W. Mann,, K.G and Riggs, B.L. (1983):

Increase in serum bone gammcarboxy glutamie acid protein with aging in women. J Clin Invest 71:13.16-21.

Delmas, P.D.; Slennr, D. Whaner, H.W.; Mann, K.G. and Riggs, B.L. (1989):

Increase in serum bone carboxyglut amic acid protien with aging in women, J. Clin Invest., B 16-1321.

Demiaux, B. Arolt, M.E.; Chapuy, M.C.; Meunier, P.L.; Delmas, P.D. (1992):

Serum Osteocalcin is increased in patients with osteomalacia: Correlation with biochemical and histomorphometric findings. J Clin Endocrinol Metab, 74:1146-51.

Dounas P.T., Watson W.A. and Biggs H.G. (1970):

Determination of S. albumin

Duda, R.J.; O'Brien, J.F.; Katzmann, J.M.; Man, K.J and Riggs, B.L. (1988):

Concurrent assays of circulating bone Gla-Protein and bone alkaline phosphatase: Effects of sex, age and metabolic bone disease. J Clin Endocrinol Metab, 66:951-7.

Eastell, R. Collwell, A., et al., (1997):

Biochemical markers of bone resorption compared with estimates of bone resorption from radiotracer kinetic studies in osteoporosis. J Bone Miner Res, 12:59-65.

Ebeling (1996):

Bone turnover markers and bone density across the menopausal transition 340:278-9.

Ehenstam, E.A.; Liunghall, S. Hallgren, R. (1986):

Serum osteocalcin in rheumatoid arthritits and other inflammatory arthritis Relation between inflammatory activity and the effect of glucocortoids and remission inducing drugs. Ann Rheum, Dis 45:484-90.

Epstein, S.M.C. clintck, R, Bryce, G. Poser, J. Johnston, J.R.C and it is (1984):

Difference in serum bone Gla-protein with age and sex. Lancet, 1:307-10.

Epstein, S. (1988):

Serum and urinary markers of bone remodeling assessment of bone turnover. *Endocrinol. Rev.* 9:437-49.

Eyre D. (1992):

New biochemical markers of bone resorption *J Clin. Endocrinol. Metab.* 24:47.

Eyre, D.R. Koob, T.J. Van New K.P. (1984):

Quantitation of hydroxypyridinun cross-links in collagen by high performance liquid chromatography. *Anal Biochem.* 137:380-388.

Fermo, I., Arch Ilonic, Casari, E. Paroni, R. (1997):

Urine pyridinium cross-link determination by Beckman cross link kit. *Clin Chem.* 43:2186-7.

Fujita, T., Satomura, A., Hidaka, M., Ohswal, Endo, M., Ohi, H. (2000):

Acute alteration in bone mineral density and biochemical markers for bone metabolism in nephrotic patients receiving high-dose glucocorticoids and one-cycle etidronate therapy. *Calcif. Tissue Int.*, 3,66:3, 195-9.

Fujimoto, D. Suzuki, M. Uchiyama, S. Miyamotoes Inoue, T. (1993):

Analysis of pyridinoline cross-linking compound of collagen fibers in human urine. *J Biochem (Tokyo)* 99:1144-1146.

Garnero, P. Gimaux, M. Demiaux, B. Preaudat sugin, P.Delmas, P.D. (1992):

Measurement of serum osteocalcin with a human-specific radioimmunoassay. *J. Bone Miner Res.* 7:1389-1397.

Garnero, P. Gineyts, E. Riou, J.P. Delmas, P.D. (1994):
Assessment of bone resorption with a new marker of collagen degradation in patients with metabolic bone disease. J Clin Endo Metab, 79:780-5.

Garnnero, P. and Delmas, P.D. (1993):
Assessment of the serum levels of bone alkaline phosphatase with a new radioimmunoassay in patients with metabolic bone disease J Clin Endocrinol Metab. 77:1046-1053.

Geant, H.K.; Cann, C.E.; Ettinger, R. and Gordan, G.S. (1982):
Quantitative computed tomography of vertebral spongiosa. Ann Of intern. Med. 97:699-705.

Geusens, P. (1993):
Effects of anabolic steroids on established vertebral osteoporosis, Rev, Esp Rheumatol. Yol. 20. Supplement 1, July, PP. 75.

Gevers, G. Devos Pand De Roo, M. (1986):
Increased levels of osteocalcin in rheumatoid arthritis. British J Rheumatol, 25:260.

Godscha L.K., and Downs, R.W. (1988):
Effect of short term glucocorticoids on serum osteocalcin in healthy young men. J Bone Min Res, 3:113-5.

Gorden, S. (1976):
In clinical management of osteoporosis, Gordan, S. and Vaughan, C. (eds), H.M.M Publisher, England, Chap. 2, p.19.

Granero, P., Gineyts, E., Arbault, P. christiansen, C. Delmas P.D. (1995):
Different effects of his phosphonate and estrogen therapy on free and peptide-bound bone cross-links excretion J Bone Miner Res. 10:641-649.

Granero, P., Grimaux M., Seguin, Delmas, P.D. (1994):

Characterization of immunoreactive forms of human Osteocalcin generated in vivo and vitro. J Bone in vivo and vitro. J Bone Miner Res, 9:255-264.

Granero, P. Sornay-Rendu. E. Delmas, P.D. (1993):

Assesment of age-related changes of bone tunover in cocmal women with new bio-chemical markers J.Bone Miner Resg (Suppl 1) is 389.

Gruber, H.E.; Ivery, J.L. Baylink, D.J.; Mathews, M. Nelp, W.B.; nison, K. and Chesunt, C.H.III (1984):

Long-term calcitonin therapy in post menopausal osteoporosis Metabolism, 33:295-303.

Gundherg, C.M.; Wilson, M.S.; Gallop, P.M. and Parfill, A.M. (1985):

The determination of osteocalcin in human serum comparison of with a wella-characterized assay. Clin Chem, 31:1720-3.

Hansen M.A., Overgard K., Ris B.J. and Christiansen C. (1991):

Role of peak bone mass and bone loss in post menopausal osteoporosis. Br. Med. J 303:961-964.

Hansen, D.A. Weis, M.A.E.; Bollen, A.M. Maslan, S.H.; Signer, F.R.; Eyre, D.R. (1992):

A specific Immunoassay for monitoring human bone resorption: quantitation of type I collagen cross- linked- N-telopeptides in mice. J Bone Miner Res. 7:1251-1258.

Harris, S.T.; Gertz, B.T.; Genant, K.K. et al., (1993):

The effect of short term treatment with alendronate on vertebral density and biochemical markers of bone remodelling in early post menopausal women. J Clin Endocrinol Metab, 76:1399-1406.

- Harvey, R.D.; Mchardy, K.C.; Reid I.W. (1991):*
Measurement of bone collagen desradation in hyperthyroidism and during thyroxin replacement therapy uring pyriidinium crosslnks as a specific urinary markers. J Clin Endocrinal Meathb, 72:1189-1194.
- Hasling, C.; Eriksen, E.F.; Melkko, J.; Ristelli, L.; Mosekilde, L. and Risteli, I. (1991):*
Effects of a combined estrogen gestagen regiomen on serum levels of the carboxyterminal propeptide of human type I procollagen in osteoporosis. J. Bone, Min. Res, 6:1295-1300.
- Hassager, C.; Collwell, A.; Assiri, A.M.A. Eastell, R. Russell R.G.G; Christiansen, C. (1992):*
Effect of menopause and Hormone replacement therapy on urinary exctetion of pyridinium cross-links: a longitudinal and cross-sectional study. Clin Endocrinol (Oxf) 37:5-50.
- Hauschka, P.V.; Lian, J.B. Cole-DEC. and Gumberg C.M. (1989):*
Osteocalcin and matrix Gla protien: Vitamin K-dependant proteins in bone physio Rev, 69 (3): 990-1047.
- Health, H. and Sizemor, G.M. (1997):*
Calcitonin in normal man. J. Clinical Invest. 60:1135.
- Heaney R.P., Recker R.R. and Seville P.D. (1978):*
Menopausal changes in bone remodeling. J Lab. Clin. Med. 192:964-970.
- Henry R.J., Cannon D.C. and Winkelman J.W. (1974):*
Clinical chemistry, principles and techniques Harper & Row 2nd Ed.
- Hillyard, C.J.; Stevenson, J.C.; and Maclantyre, L (1978):*
Relative deficiency of plasma calcitonin in normal women, lancet, 1:961-962.

Hodsman, A.B.; Fraher, L.J.; Watson, P.H.; Obtye, T.; Sti, L.W. Adachi, J.D.; Taves, D.H. Drost D: (1997):

A randomized control trial to compare the efficacy of cyclical parathyroid hormone versus cyclical parathyroid hormone and sequential calitonin to improve bone mass in post menopausal women with osteoporosis J Clin. Endocrinol. Metab Feb; 82 (2): 620-8.

Hoshino, H.; Kushida, K.; Takahashi, M.; Denda, M.; Yamazaki, K. Yamanashi, A, Invo, T. (1997):

Short-term effect of parathyroidectomy on biochemical markers in primary and secondary hyperthyroidism. Miner Electrolyte Metab, 23-93-9.

Hough, S. (1998):

Fast and slow bone losers relevance to the management of osteoporosis Drugs. Aging, 12 suppl. 1:1-7.

Hyldstrup, L.; Clemmensen, L.; Jensen, B.A. and Transble, I (1998):

Non invasive evaluation of bone formation Measurement of serum alkaline phosphatase, Whole body retention of diphosphonate and serum osteocalcin in metabolic bone disorders and thyroid diseases S.C and J Clin Lab Invest, 48:611-9.

Jackson, J.A.; and Kleerekoper, M. (1990):

Osteoporosis in men. Diagnosis, Pathophysiology, and prevention, Medicine 69:137.

Jahansen, J.S.; Giwer C. man A. Hartuell, D. Nielsen, C.T. Price, P.A.; Christiansen C.V. and Shakkebaek, N.E. (1988):

Serum bone Gla protein as a marker of bone growth in children and adolescents correlation with age hight serum insulin like growth factor. and serum testosterone. J Clin Endocrinol Metab, 67:273-8.

Jahansen, J.S.; Thomsen K.; and Christiansen C. (1987):

Plasma bone Gla protein concentration in healthy adults. Dependence on sex, age and glomerulus filtration. Scand J Clin Labs Invest, 47:345-5.

*

Jaworski, Z.F.; (1981):

Physiology and pathology of bone remodeling orth. Clin North America, 12,485.

Johnston, C.; Slemenda, C.W.; Mettison 111 L.J.; (1991):

Clinical use of bone densitometry. N Engl. J Med. 324:1105-1109.

Johnston, C. and Epstein, S. (1981):

Clinical biochemical, radiographic and economic features of osteoporosis. Orth. Clin. North America 12:559.

Kerner, S.A.; Scott R.A.; Pike, J.W. (1989):

Sequence elements in the human osteoclicin gene for basal activation and inducible response to hormone D3. Proc Nat Acad Sci U.S.A, 86:4455-59.

Kind and King (1954):

Colorometric determination of S.AL.P

Knais, J.A. (1990):

Skeletal disorders. In: Recent advance in orthopedics, Mckibbin B. (ed), p. 247.

Krolner, B.; Dunbar, O. and Buhr, A. (1982):

Physical exercise and age related bone loss. Clin. Orth, 103:17.

Kruse, K.; Kracht, U., Wohlfart, K.; Kruse, U. (1989):

Biochemical markers of bone turn over, inact serum parathyyroid hormone and renal calcium excretion in patients with pseudohypoparathyroidism and hypoparathyroidism before and during vitamin D. treatment. Eur, J. Pediat 148:532-539.

Lane, J.M. and Vigorita, V.M. (1984):

Osteoporosis, orth. Clin North America, Vol. 15.No. 4:159.

Lewenhaupt, A.; Ekman, P.; Eneroth P. and Kalner. A. (1988):

On reference values for tumor markers in protstic carcinoma. S.C. and J Urol Nephrol suppl, 110:119-23.

Lian, J.B.; (1985):

Factors contributing to interactivity calcification in: Rubin, R.P.; Weiss, G.B. and Jutney, J.W.J.R. (eds): Calcium in biological system. New York: Renum, pp 633-44.

Liel, Y., Edwards, J., Shary J., Speicer, K.M., Gordonl and bell, N.H.: (1988):

The effects of race and body habitus on bone mineral density of the radius, hip and spine in premenopausal women. J Clin. Endocrinol Metab. 66:1246.

Lindsay, R.; (1987):

Estrogen therapy in prevention and management of osteoporosis. An J. Obstet Gyneco 115:1347; 1987.

Lips, P.; courprop. Meunier P.J.; (1978):

Mean wall thickness of trabecular bone packets in the human iliac crest. Changes with age calcif. Res. 26-13-17.

Lipton, A.; Demers L, Dniloffy, curley, E.; Hamitton, C.; Harvey, H.; et al, (1993):

Increased urinary excretion of pyridinium cross-links in cancer patients. Clin. Chem 39:614-8.

Lukert, B.P.; Higgins, J.C.; Stoskopf, M.M.; (1986):

Serum osteocalcin is increased in patients with hyperthyroidism and decreased in patients receiving glucorticoids. J Clin. Endocrinol Metab. 62:1056-8.

Lutchmen, E.C.; Hardwick, T. N.; Biener, R.; Choudury, H. A.; Traut, J. R. Shapses, S. A. (1998):

Longitudinal study of urinary hydroxypyridinium cross links and growth in healthy infants higher values with breast feeding and after daytime sleep. Exp- Clin Endocrinol- Diabetes 106 (1): 51-6

Magaro, M. A.; homoote, L.; Mirocey, Zoli, A., Gonvino, G.; (1989):

Bone GLA protien (BGP) Levels and bone tunover in heumatiod arthritis. Br J Rheumatol 28: 207-11.

Marhoffer, W., Scharz, H.; Stracke, H.; Uikman, J Schrmidic, K.; (1991):
Serum osteocalcin concent rations in male (Abstract) Pediatr Res, 18:98 A.

Matti J. Valimaki, Rittal Tahtela, James D, James M and Blewrene Rigg 1994:

Bone resorption in healthy and osteoporotic post menopausal women.

Melick R.A., Farrugia W. and Quelch K.J. (1985):

Plasma osteocalcin in men. Aust in. J Med. 15:410-16.

Melton 111 L.T., Atkinson E.J., O'Fallon W.M., Wahnes, H.W., Riggs, B.L. (1993):

Long-term fractures prediction by bone mineral assessed in different skeletal sites. J Bone Miner Res. 8:1227-1233.

Melton, J.J.; (1988):

Epidemiology of fractures. In Riggs, B. L. and Melton, L. J. (ed), osteoporosis, Etiology, Diagnosis and Management, New York, Raven Press, p. 133-54.

Melton, L. J.; Thamer M. Ray N. F.; Chan J. K.; Chesnut, C. H.; Enhom T. A. et al (1997)

Fractures attributable to osteoporosis : Report for the national osteoporosis Foundation. J Bone Miner Res, 12: 16-23

Merer: (1987):

Marked decline in trabecular bone mineral content in healthy men with age: lack of association with sex steroid levels, J. Am. Geriatr. Soc. 35: 189.

Merle, B. and Delmas, P. D (1990):

Normal carboxylation of circulating osteoclast bone Gla-protein in paget is disease of bone. Bone Miner 11:237-45

Michnovicz, J. J.; Hershscof, R.j.; Naganuma H.; Bradlow, H.L. and Fishman, J. Increased 2 (1986)

Hydroxylation of estradiol a possible mechanism for the anti- estrogen effect of cigarette smoking, N. Engl. J. Med. 315- 1305.

Mizunashi, k. Furukaway, Miur, R.; Yumitas, H.E. and Yoshinak (1988):

Hormone on the serum osteocalcin in idiopathic Invest, 82:861-65.

Mole, P. A. Walkinshaw, M.H.; Robins, S. P. Paterson, C. R. (1992):

Can urinary pyridinium cross-links and urinary oestrogens predict bone mass and rate of bone loss after the menopause? Invest. 22:767-771.

Nielsen, H. K.; Charles, P and Mosekild, L. (1988):

The effect of single oral doses of prednisone on the circadian rhythm of serum osteocalcin in normal subjects.

Nilas, L. and Christiansen, C (1987):

Bone mass and its relationship to the menopause J Clin Endocrinol Metab 65 : 697- 702.

Norden, B.E.C.; (1960):

Osteomalacia osteoporosis and calcium deficiency, Clin, Orthop, 17, 253-57.

Nottelovitz, M. (1986):

Post menopausal osteoporosis, A practical approach to its prevention, Acta obstet Gynecol. S. C and SPP. 134 : 67.

Orme, S. M. and Belebetz, P. E. (1994):

Is a low skin fold thickness an indicator of osteoporosis, Clin: Endocrinol, O. X. F., September 41 (3) 283-7

Parfitt, A. M. (1987):

Bone remodeling relation ship to the amount and structure of bone and the pathogenesis and prevention of fracture in : Riggs, B1, Melton 111 LJ, eds, Osteoporosis, m New York Raver Press, 45-94.

Parfitt A. M.; Mathews, C.H.E.; Villaneieva, A. R.; Kleerekoper, M.; Freme, B.; Rao, D. S. (1983):

Relationship between surface, volume, and thickness of trabecular bone in aging in osteoporosis. J Clin. Invest, 72: 1396.

Pastoureau, P. and Delmas P.D. (1990):

Serum bone Gla-protein BGP in human with BGP- based radio immunoassay. Clin. Chem. 36:1920-1924.

Pidethcha, P.; Intramane, S.; Patrakarn, K.; Leelahakal. P.; Fungsapa, S. (1999):

Reference value for urinary deoxypyridinoline as a specific marker for measuring bone resorption in Thais, J.; Med. Assoc- Thai Nov; 82(N): 1136-41.

Pietschmann, P., Machold, K.P. and Woloszczu K.W. (1995):

Serum osteocalcin concentrations in patients with rheumatoid arthritis. *Annals of Rheumatic Dis.* 48:654.

Podenphant, I.; Johansen, J. S.; Thomsen, K.; Ril SBJ, leth, A, chirstiansen, C. (1987):

Bone turnover in spinal osteoporsis. *Jbone Miner Res* 2: 497-503.

Posen, S.; Cornish C. and Kleene, Koper, M. (1987):

Alkaline phospatase and metabolic bone disorders. In Alvioli, L. V. and Krane, S. M. (eds): *Metabolic bone disease*. New York Academic, pp 141 – 810.

Poser, J.W.; Esch Fs, ling Nc, Price P.A. (1980):

Isolation and sequence of the vitamin, k-dependent portein from human bone. *J Biol Chem* 255:8685-91.

Power, M. j and Fottrell, P.F (1994):

Osteocalcin: Diagnostic methods and clinical applications. *Crit Rev Clin lab Sci*, 28:297-335.

Pratt, D. A.; Daniloff, Y.; Duncan, A, Robins, S. P. (1992):

Automated analysis of the pyridinium cross links of collagen in tissue and urine using solid phase extraction and reversed- phase high performance liquid chromatography. *Anal Biochem*, 207 : 168-75.

Price P. (1983).

Osteocalcin. In: *Bone and minreal research Annual I*. Amsterdam: Excerpt amdica 157- 190.

Price P. A.; Parthemore, J. B. and Deftos, L. J. (1988):

New biochemical marker for bone metabolism: Measurement by radioimmunoassay of bone Gla protein in the plasma of normal subjects and patients with bone disease *J Clin Invest*, 66: 876-83.

Price, P. a.; Williamson, M. K.; Lothringer, J.W.; origin of the vitamin K (1981):

Dependent bone protein found in the plasma and its clearance by kidney and bone. J Biol Chem, 256: 12760 -6.

Quick, A.J. (1942):

The hemorrhagic diseases and the Physiology of Homeostasis. Thomas , ILL onois.

Raymakers, J. A.; Kapelle, J. W.; Van - Berestijn, E. C & Duursma, S. A. (1992):

Assessment of osteoporotic spine deformity, A new method. Skeletal Radiol. 19 (2) : 91-7

Rec. GSCC, (1972) :

Detetermination of ALP J. Clin. Chem Clin . Biochem. 10:128

Reeve, J., (1990):

The use of biochemical and isotopic studies in the investigation of bone disorders. In Stervenson, J. C. (ed) New techniques in Metabolic bone Disease, pp.. 92-109 (london- Weight).

Reeve, J.; Meunier, P.J.; Parson, J. A.; et al., (1980):

Anabolic effect of human parathyroid hormone fragment on trabecular bone in involutinal osteoporosis : a multicentre trial. Br Med, J. 1340 - 1366.

Rico, H.; Corbranse, J. A.; Cabello, J. Game, Z Catresana F and Hernande E. R. (1987):

Low serum osteocalcin in acute alcohol intoxication : a direed toxic effect of alchol on osteoblasts. J Bone Min Res, 2:221- 25.

Riggs, B.L. (1991):

Over view of osteoporosis. West. J Med. 154-77.

Riggs, B. L. & Melton, L. J: (1986):

Involutional osteoporosis N. Eng. J. Med. 314 : 1676.

Riggs, B. L.; Melton 111 L. J. (1992):

The prevention and treatment of osteoporosis. N Eng I Med 377 : 620 -7.

Riggs, B.L., Wahner, H.W., Melton, J., Richelson, L.S., Judd, H.L. and Offord K.P. (1986):

Rates of bone loss in the appendicunai and axial skeletons of women. J Clin. Invest. 77:1487.

Risteli, J.; Elo, max, I.; Niemi, S., Novamo, A.; Ristelil (1993):

Radioimmunoassay for the pyridinoline cross-linked carboxy terminal telopeptied of type I collagen : a new serum marker of bone collagen degradation. Clin Chem, 39: 653-40.

Robins, C.P.; Black, D.; Paterson C. R.; Reid, D. M. and Duncan, A. (1991):

Evaluation of urinary hydroxypyridinium cross-link measurements as resorption markers in metabolic diseases. Eur. J.; Clin. Invest, 2:310- 315.

Robins, S.p.; Duncan, A. and Riggs, B. L. (1990):

Direct measurement of free hydroxypyridinium cross-link of collagen in urine as new markers of bone resorption in osteoporosis. In : Christiansen, C.; Overgard, K.; eds. Osteoporosis, Copenhagen: Osteoporosis, 465-468.

Robins, S. P., Duncan, A.; Wilson N.; Evans, B. J.; (1996):

Standardization of pyridinium cross links, pyridinoline and deoxy pyrdinoline, for use of biochemical markers of collagen degradation, Cin Chem. 42: 1621 -6.

Robins, S. P.; Woitge, Hesley, R.; J.U.J.; Seyedins, S.; Seibel, M.J. (1994):

Direct enzyme-linked immunoassay for measuring bone resorption, J.; Bone Miner, Res. 9:1643- 1649.

Ross, P. D. and Knowlton, W. (1998):

Rapid bone loss is associated with increase levels of biochemical markers. J. Bone Miner. Res.; 13 (2) : 297 – 302.

Ross, R.K., Paanini, Hill A., Mack T.M. et al (1981):

Menopausal estrogen therapy and protection from ischaemic heart disease. Lancet; 1:858-860.

Sakhaee, M.; Studd, J.W.; Fofelment. L.; Dooley, M.; Montgomery, J. and Murphy, B. (1985):

Skeletal effects of oral estrogen compared with subcutaneous estrogen and testosterone, in postmenopausal women, B. R. J Med 297: 331.

Salnder, R.B., Slenenda, C.W., Laporte, R.E., Cawly, J.A., Schram, M., Barresi, M.L. and Kriska, A.M. (1985):

Post menopausal bone milk consumption in childhood and adolescence Am. J Clin. Nutr. 42,2704.

Sandles, R. B.; Slemenda, C. W., Laporte, R. E., Cauely, J. A., Schram, M.M., Barresi M.L and Kriske A. M., (1985).

Post menopausal bone milk consumption in childhood and adolescence. A M. J. Clin Nutr., 42, 270 –4.

Seible, M. J. G.; Wotige H.; Scheidt, Nave C, Leidig- Brackner, G, Duncan, A.; Nicol, P.; et al (1994):

Urinary hydropyridinium crosslinks of collagen in population based screening for vertebral osteoporosis: results of apilot study. J Bone Miner Res, 9: 1433 –40.

Seible, M. J.; Gartenberg, F., Ratcliffe, A.; Robins, S. P.; Sillberg, J.; Bilezikian, J. P.; (1992):

Urinary hydroxy pyridinium cross links of collagen as specific indices of bone resorption in primary hyperparathyroidism, J.; Clin Endocrinol, Metab, 74: 481-6.

Seyedin S.M.; Kung, V.T.; Danilo, F.F.Y.N. (1993):

Immunoassay for urinary pyridinoline: The new marker of bone resorption, J. Bone, Miner, Res 8:635-641.

Seyedin, S. M. and Rosen D.M. (1990):

Matrix Proteins of the skeleton. Opin cell Biol, 2:914-19.

Singer, F. R.; and Mills, B. G. (1984):

Paget disease of bone etiologic and therapeutic aspects. In : peck, W.W.; (ed) : Bone and Mineral research, New York: Elsevier, 2:394-421.

Singer, F.R.; (1992):

Paget's Disease of the bone Cecil Textbook of Medicine, 19 Edition , pp. 1931-33. Smith, D.M. Osteoporosis, osteomalacia diagnostic index. J. Lab. Clin Med 87:882, 1976.

Slovik, R. M.; Gunberg, C. M.; Neer, R. M. and Lain J. B. (1948):

Clinical evaluation of bone turnover by serum osteoclin measurements in hospital setting, J. Clin Endocrinal Metab, 59:228 -30.

Speancer and price (1977):

Determination of albumin by Bromocresol Green method

Stepan, J.J., Postichalj, Presl, J.; Pacovsky (1987):

Bone loss and biochemical indices of bone remodeling in surgically post menopausal women. Bone 7:279- 284.

Takahsai, M.; Kushida, K., Hoshinolt, Inoue T., (1997):

Discrimination ability of pyridinoline cross links related markers for bone resorption in post menopause and osteoporosis. *Endocrinol. Res*, 23: 105-17

Tanka, K. A.; H.; Shiota, H.; Kuwada, M.; Trea, K., and Katayama, K. (1988):

Plasma and bone osteocalcin levels in rats with type II collagen induced arthritis. *Arthritis Rheum*, 31:1414-20.

Tarallo, P. Henny, J. Fouriner, B.; and Siestg (1991):

Plasma osteocalcin: biological variation and reference limits. *Scand, J.; Clin lab invest*, so: 649-55.

Taylor, A.K.; Link hart, S. G.; Mohan, S.; and Baylink, D. J. (1988):

Development of radioimmunoassay specific for human osteocalcin *Metabolism* 872-77

Thomas, K. A., Cook, S.D.; Bennett, J.T.; Whitecloud, T. S.; and Rice, J. C. (1991):

Femoral neck and lumber spine densities in a normal population 3-20 years of age. *J Pediatr. Orthop.* 11:48.

Trinder, P.(1969): *Ann. Clin, Biochem* 6,24

Tsuji, T.; Hughes, F.J.; Mc Culloch, C. A. G. and Meleher, A. H (1990):

Effects of doner age on osteogenic cells of rat bone maceow in vitro. *Mech. Age. Dev.* 51:121.

Uebelhart, D.; Schlemmen, A.; Johansen. J.S.; Gineyts, E.; Christiansen, c.; Delmas, P.D. (1991) :

Effect of hormone replacement therapy on the urinary excretion of pyridinium cross-links, *J Clin Endocrinol Metab*, 72:367-373.

REFERENCES

Uebelhart, D.; Ginenyts, E.; Delmas, P. D. (1990):

Urinary excretion of pyridinum cross-links: anew marker of bone resorption in metabolic disease, Bone Miner, 8:87:96.

Yasumura, A. S. (1987):

Management in: Novak's textbook of Gynecology. 11th edition Library of Congress. Cataloging, P.428.

Yilmaz, N., Bayram, M.; Erabgci, A. B. Kilincer, M. S.; (1999):

Diagnostic value of biochemical markers of bone turnover and postmenopausal osteoporosis. Clin- Chem- lab, Med Feb, 37 (2) : 137-43.

Yilmaz, N., Ozaslan, J. (2000):

Biochemical bone turnover markers in patients with ankylosing spondylitis. Clin. Rheumatol. 1,19:2, 92-8.

Yoneda, M. L.; and Frazer, D. (1988):

Contraversies concerning the safety of oestrogen replacement therapy. Am. J. Gynecol, 156: 1313.