

REFERENCES

Adpick NS et al (1985):

Major childhood tumour resection using nomovolaemic hemodilution anaesthesia and Hetastarch. *Journal of Paediatric surgery*, 20 (4): 372-375.

American Society of Anaesthesiologists (1995):

Directory of Members. park Ridge, IL.

Amindson B and Haljamae (1980):

Skeletal microcirculating and cellular metabolic effects of whole blood, ringer's solution and dextran-70 infusion in haemorrhagic shock. *Cir. Shock*, 7: 111.

Anderson and Swedlow DB (1986):

Invasive assessment of the failing circulation from cardiovascular problems in pediatric critical care. *Clinics in Critical Care Medicine*. New York, Churchill Livingstone, p. 157-162.

Aperia A and Zetterstrom R (1982): Renal control of fluid homeostasis in the newborn infant. Clin Perinatal 9: 523.

Ardailou R, Sraer J, Chansel D et al (1987):

The effect of angiotensin II on isolated glomeruli and cultured glomerular cells. Kidney Int 31 (Supl. 20): S74.

Awazu M et al (1990):

From paediatric textbook of fluid and electrolytes. Edited by Faurel Carven, 418.

Barker SJ, Tremper KOK (1985):

Transcutaneous oxygen tension. A physiological variable for monitoring oxygenation. J. Clin. Monit. 1: 130.

Behrman SW, Fabian Tc, Kudsk KA et al. (1991):

Microcirculatory flow changes after initial resuscitation of haemorrhagic shock with 7.5% to hypertonic saline and 6% dextran 70. J. trauma 31: 589.

Bennett EJ (1975):

Fluid balance in the newborn. *Anaesthesiology* 43: 210.

Bennett, ED (1986):

Colloid osmotic pressure of plasma substitutes. *Proc. Confer., Manchester, Birmingham and London*, 23-25.

Berry FA (1983):

The renal system. In Gregory GA, editor: *Paediatric anaesthesia*. New York. Churchill Livingstone.

Berry FA (1990):

Practical aspects of fluid and electrolyte therapy, in *Anaesthetic Management of Difficult and Routine Pediatric Patients*. Berry FA, editor. New York, Churchill Livingstone, p 89-120.

Blajehman MA and Perrault RA (1988):

"Blood component therapy in anaesthetic practice" *Cond. Anaesth. Soc.J.*, p. 282.

Boldt J et al (1993):

Influence of different volume therapy on platelet function in patients undergoing cardiopulmonary bypass. *Anaesth. Analg.*

Bradbury MW (1977):

Physiology of body fluids and electrolytes, from physiology for the anaesthesiologist edited by Nishan G and Karamanin, p. 345:363.

Brenner BNM, Beeuwkes R III (1978):

The renal circulations. *Hosp Pract.* 13: 35.

Brinkmeyer L, Safar P and Motoyama EK (1983):

Superiority of colloid over electrolyte solution for severe normovolaemic hemodilution in treatment of hemorrhage. *Disaster Med* 1: 171.

Brooks TD and Gravenstein N (1986):

Pulse oximetry for early detection of hypoxemia in anaesthetized infants. *J Clin Monit* 1: 135.

Campbell D and Spence AA (1985):

Intravenous therapy, Anaesthetic resuscitation and intensive care, 6th ed., Churchill Livingstone, Edinburgh, London and New York, 75-89.

Cervera AL and Moss G (1988):

Crystalloid requirements and distribution when resuscitating with RBC, and non-colloid solutions during haemorrhage. *Circ. Shock*, 5: 357-364.

Charles F, Simmons and Leukouni Lchikawa (1990):

External balance of water and electrolytes, from paediatric textbook of fluid and electrolytes. Edited by Laurel Careven, 23.

Charles F, Simmons et al. (1990):

External balance of water and electrolytes, from pediatric textbook of fluid and electrolytes edited by faurel craven, p. 21.

Cote CJ (1990):

NPO after midnight for children, *Anaesthesiology* 72: 589.

Cohen IT and Etsuro K. Motoyama (1996):

Intraoperative and postoperative management. from Smith's anaesthesia for infants and children. Edited by Etsuro KO and Peter J., 316.

Cross JS, Griber DP, Burchard KW et al. (1988):

Hypertonic saline fluid therapy following surgery. *J trauma*, 29: 817.

Davis JO and Freeman RH (1986):

Mechanisms regulating renin release. *Physiol. Rev.* 56:1.

Davis P and Derbishire (1990):

Postoperative fluid and electrolyte requirements for infants and children. *Textbook of Anaesthesia*, p. 748.

Demling RH (1980):

Response of the pulmonary microcirculation to fluid loading after haemorrhagic shock and resuscitation. *Surgery*, 87: 552-559.

Dobb GJ (1980):

Body water, electrolytes and parenteral fluid therapy. In: Wylin and Churchill-Davidson's ed.' *A practice of anaesthesia*, 5th ed., London, 568-586.

Donald SP and Mali Mathru (1977):

Acid-base, fluids and electrolytes. *Clinical anaesthesia*, edited by Paul G Barash et al., Lippincott-Raven Publishers, Philadelphia, 164.

Dong W, Beldose SW, Chadwick HS et al (1986):

Electrical correlates of brain injury resulting from severe hypotension and hemodilution in monkeys. *Anaesthesiology* 65: 612.

Edwards BR (1981):

Postnatal development of urinary concentrating ability in rates. Paris, Masson, P 233.

Eichhorn JH (1986):

Standards for patient monitoring during anaesthesia at Harvard Medical School. *JAMA* 256: 1017.

Ellis D, Avner et al. (1990):

Normal neonates and the maturational development of homeostatic mechanisms, from pediatric textbook of fluid and electrolytes. Edited by Laurel Carven, p. 117.

Ellis D, Avner ED (1985):

Fluid and electrolyte disorders in paediatric patients. In Puschett JB, editor: Disorders of fluid and electrolyte balance. New York, Churchill Livingstone, p. 217.

Etsuro KM (1996):

Respiratory physiology in infants and children, from Smith's Anaesthesia for Infants and children, Edited by Peter J. Davis, p 43-47

Filston (1992):

Fluid and electrolyte management in paediatric surgical patients. Surg. Clinics of Nor. America, 72: 1189-1205.

Fontana, Leila Welborn, Paul D, Morgan et al. (1995):

Oxygen consumption and cardiovascular function in children during profound intraoperative normovolaemic hemodilution. Anaesth. Analg, 80: 219-225.

Frederic A Berry (1992):

Perioperative fluid management for paediatric patients. ASA refresher courses, p. 422.

Frederic A Berry (1997):

Fluid and electrolyte therapy in paediatrics. ASA refresher course, 166: 1-7.

Furman EB, Roman DG, Lemmer LAS et al. (1985):
Specific therapy in water, electrolyte and blood
volume replacement during paediatric surgery.
Anaesthesiology 42: 187.

Gala GJ, Lilly MP, Thomas SE et al (1991):
Interaction of sodium and volume in fluid
resuscitation after haemorrhage. J Trauma 31: 545.

Gardner WS (1993):
Intraoperative fluid management. Surgical clinic of
North America. 73: 229-241.

Gelman S, Fowlerk Smith L (1984):
Regional blood flow during isoflurane and halothane
anaesthesia. Anaesth. Analg. 63: 557.

Gertz KH, Mangos JA, Braum G et al. (1966):
Pressure in the glomerular capillaries of the rat
kidney in relation to arterial blood pressure.
Pflugers Arch 288: 369.

Giesecke AH (1989):
Perioperative fluid therapy. In: Miller, R.D. ed.,
Anaesthesia, 3rd ed. Churchill Livingstone Inc., New
York.

Goodnough LT, Price TH et al (1989):

Increased preoperative collection of autologous blood with recombinant erythropoietin therapy. N. Eng.J. Med. 321: 1163.

Graham JFM (1979):

Preoperative starvation and plasma glucose concentrations in children undergoing outpatient anaesthesia. Br. J. Anaesth. 51: 161-164.

Graham TP (1983):

Invasive evaluation of the child with congenital heart disease. In Gold Smith HS (ed). Practice of surgery. Philadelphia, harper and Row, 1-16.

Haberkem M and Dangel P (1991):

Normovolaemic hemodilution and intraoperative autotransfusion in children. Eur. J. Pediatr. Surg. 1: 30.

Haljamae H (1985):

Rationale for the use of collids in treatment of shock and hypovolaemia. Acta Anaesth. Scand. 29: 48-54.

Halvorsen L, Gunther RA, Dubick MA et al (1991):

Dose response characteristics of hypertonic saline dextran solutions. J Trauma 31: 285.

Hammermann et al. (1993):

Volume replacement with hydroxyethyl starch solution in children. British J. of Anaesth., 70: 661-665..

Herbacznska-Cedrok (1973):

Contribution of intrarenal generation of prostaglandins in autoregulation of renal blood flow in the dog. Circ. Res. 33: 428.

Herman ML and Nearman HS (1991):

Toxic effects of colloids in the intensive care unit. Crit. Care. Clin. Jul; 7 (3); 313-23.

Hindman BJ, Dexter F, Cutkomp J et al (1992): Brain blood flow and metabolism do not decrease at stable brain temperature during cardiopulmonary bypass in rabbits. Anaesthesiology, 77: 342.

Hodge D and Fleisher G (1985):

Pediatric catheters flow rates. *Am. J Emerg Med* 3: 403.

Holliday MA and Segar WE (1977):

The maintenance need for water in parenteral fluid therapy. *Pediatrics*, 19: 823-832.

James D (1990):

Blood transfusion and notes on related aspects of blood clotting. *Scientific foundation of anaesthesia*. 4th ed, Churchill Livingstone, London, Chapter 7.

James MS, David EC and Robert J (1996):

Anesthesia equipment and monitoring, from *Smith's Anesthesia for Infants and children*, edited by Etsuro K and Peter J, p. 248-250.

Jose PA, Staincova V, Felix et al. (1975):

Autoregulation of renal blood flow in puppy. *Am. J. Physiol.* 229: 983.

Kagan BM, Staincova V, Felix NS et al. (1972):

Body composition of premature infants. *Am. J. clin./ Nutr.* 25: 1153.

Kallos T and Smith TC (1974):

Replacement of intraoperative blood loss.
Anaesthesiology 41: 239.

Kandel G and Aberman A (1983):

Mixed venous oxygen saturation; its role in the
assessment of circulatory ill patient. Arch Intern
Med; 143: 1400-2.

Kemmotsu O, Ueda M, Otsuka H et al (1991):

Blood pressure measurement by arterial tonometry
in controlled hypotension. Anesth. Analg. 73: 54.

Kerner JA (1983):

Caloric requirement from manual of paediatric
parenteral nutrition, New York, John Wiley and
Stone, 63-69.

Kimble KJ, Darnall and Yeldermon (1981):

An automated oscillometric technique for estimating
mean arterial pressure in critically ill newborns.
Anaesthesiology, 54: 423.

Lawless S, Orr R (1989):

Axillary arterial monitoring of pediatric patients.
Pediatrics 84: 273.

Leila G Welborn (1991):

Fluid management for paediatric patients, ASA
refresher course, p. 216.

Liebman J, Plonsey R et al (1996):

Anaesthesia equipment and monitoring, from
Smith's Anaesthesia for infant and children, edited
by Etsuro K and Peter J. 252.

Lowe HJ (1981):

Monitoring metabolism during closed circuit
anaesthesia, from the quantitative practice of
anaesthesia. Edited by Lowe H, p. 149.

Luscher TF, Boulanger CM, dohiy et al. (1992):

Endothelium-derived contracting factors. Hyper-
tension 19: 117.

Maier M, Starlinger M, Wagner M et al. (1981):

The effect of hemorrhagic hypoperfusion on urinary
kallikrein excretion. Circ Res. 48: 386.

Manninen PH et al (1985):

The effect of high dose mannitol on serum and urine electrolytes and osmolarity in neurological patients. *Canad. Anaesth. Soc. J*, 82-92.

Mapes A and Jones BR (1994):

Enlarging paediatric IV catheters. *Paediatr Emerg. Care* 10: 18.

Marsden PA, Brenner BM (1991):

Nitric oxide and endothelins, novel autocrine/paracrine regulators of the circulation. *Semin Nephrol* 11: 169.

Mazze RI (1986):

Renal physiology and the effects of anesthesia. In Miller RD, editor; *Anaesthesia*. Churchill Livingstone.

McCrory WW (1988):

Developmental nephrology. Cambridge, MA, Harvard University Press.

McKlveen RE, Cooke JE, Loftus JR (1990):

Increased fluid flow through the streamline intravenous catheter. *Anaesth. Analg* 70: 328.

Miller BR, Tharp JA, Jssac WB (1990):

Gastric residual volume in infants and children following a 3-hours fast, *J Clin. Anaesth* 2: 301.

Miller RD and Brzica SM (1990):

"Blood, Blood component and colloid therapy" In Miller, RD, 3rd ed. *anaesthesia*.

Moller JT, Johannessen NW, Espersen K et al (1993):

Randomized evaluation of pulse oximetry in 802 patients. *Anaesthesiology* 78: 445.

Mollison P.L. (1988):

Blood transfusion in clinical Medicine. Blackwell Scientific Publications, Oxford.

Morray JP, Caplan RA, Geiduschek JM et al (1993):

A comparison of pediatric and adult anaesthesia closed malpractice claims. *Anaesthesiology*, 78: 461.

The effect of carbon monoxide inhalation on pulse oximetry and transcutaneous PO₂. *Anaesthesiology*, 66: 677.

Payne K, Ireland P (1984):

Plasma glucose level in the perioperative period in children. *Anaesthesia* 39: 868-872.

Perloff WH (1990):

Physiology of the heart and circulation. *Clinics in Critical care medicine*; New York, Churchill Livingstone.

Porter SS and Goldberg RJ (1986):

Intraoperative allergic reactions to hydroxy-ethyl starch. *Can. Anaesth. Soc. J.*, 33 (1): 394-398.

Ring J., Seifert J, Messmer KO and Brendel W (1976):

Abnaphylactoid reactions due to hydroxy-ethyl starch infusion. *Eur. J. Surg. Res.*, 8: 389-399.

Risberg et al (1992):

Pulmonary microvascular leakage after micro-embolization and hemodilution. *Surgery*, 29: 409-415.

Rosen KR and Rosen DA (1986):

Comparative flow rates for small bore peripheral intravenous catheters. *Pediatric Emerg Care* 2: 153.

Rothen HU, Lauber R, Mosimann M (1992):

An evaluation of the rapid infusion system. *Anaesthesia* 47: 597.

Rudolph AM, Yemann MA, Teramo et al. (1978):

Studies on the circulation of the previable human fetus. *Pediatr Res* 5: 452.

Saldeen T (1983):

Clotting microembolism and inhibition of fibrinolysis in adult respiratory distress syndrome. *Surg. Clin. North Am.*, 63: 285-304.

Salem MR (1987):

Hypercapnia, hypocapnia, hypoxemia. *Semin Anaesth.* 6: 202.

Salem MR (1996):

Blood conservation. In: *Smith anaesthesia for infant and children*, edited by Etsuro and Peter, p. 347-381.

Schaller RT, Schaller J, Morgan A et al. (1983):
Haemodilution anaesthesia. *Am. J. Surg.* 146: 79.

Schlondorff D, Wever H, Trizna W (1978):
Vasopressin responsiveness of renal adenylate cyclase in newborn rats. *Am J Physiol* 234: F16.

Schmitt HJ, Schuetz WH, Proesihel et al (1996):
Accuracy of pulse oximetry in children with cyanotic congenital heart disease. *J. Cardiol. Thoracic Vasc. Anaesth.* 7: 61.

Schnermann J, Brigges JP, Weber DC (1984):
Tubulo-glomerular feedback, prostaglandins, and antigens in the autoregulation of glomerular filtration rate. *Kidney Int.* 25: 53.

Schreiner MS, Triebwasser A, Keon TP (1990):
Ingestion of liquids compared with perioperative fasting in paediatric outpatients, *Anaesthesiology*, 72: 593.

Schreiner MS et al. (1992):
Should children drink before discharge from Day Surgery. *Anaesthesiology*, vol 76 (4): 528-533.

Schultz DM, Giordano DA, Schultz DM (1962):

Weights of organs of fetuses and infants. Arch Pathol
74: 244.

Schuster DP and Lefrakss (1988):

Shock. In: Civetta J.M.; Taylor R.W., Kirby R.R.
(eds): Critical Care. Philadelphia, J.B. Lippincott, p.
903.

Schwartz GL, Brion LP, Spitzer A (1987):

The use of plasma creatinine concentration for
estimating glomerular filtration in infants, children
and adolescents. Pediatr Clin North Am. 34: 571.

Schwentker EP (1978):

Posterior fusion of the spine for scoliosis. Surg.
rounds 1: 12.

Seiber FE, Smith DS, Traysman et al (1987):

Glucose, A reevaluation of its intraoperative use.
Anaesthesiology, 67: 72.

Selkurt EE, Hall PW, Spencer MP (1949):

Influences of graded arterial pressure decrement on
renal clearance of creatinine. Am. J. Physiol. 159:
369.

Sheehan AM, Palange K, Karosr Js et al (1992):
Significantly improved peripheral intravenous catheter performance in neonates. *J Perinatol* 12:369.

Shermine D, Demetrius E and Alan B (1996):
Regulation of fluids and electrolytes in infant and children from Smith's anaesthesia for infant and children, edited by Etsuro and Peter, 105-137.

Shires GT (1987):
Pathophysiology of fluid replacement in hypovolaemic shock. *Ann. Clin. Res.* , 9: 144-150.

Shires T et al. (1964):
Fluid therapy in hemorrhagic shock. *Arch. Surg* 88: 688.

Shirley Graves (1990):
Fluid therapy for paediatric patients, ASA refresher course, p. 263.

Shoemaker WC et al (1981):
Fluid therapy in emergency resuscitation, clinical evaluation of colloid and crystalloid regimes. *Crit. Care Med.*, 9: 367-368.

Singh S and Schaeffer (1983):

Cardiorespiratory effects of volume overload with colloids in dogs Crit. Care Med., 11: 585-590.

Small A and Homer LD (1989):

An explanation of impaired solute mixing in extracellular fluid after haemorrhagic hypotension. Am. J Physiol, 236: 440-446.

Smith HW (1943):

Lectures on the kidney. Lawrence, KS, University Extension Division of University of Kansas,. p. 97.

Steele JM, Berger EY and Dunning MF (1990):

Total body water in man. Am. J. Physiol. 162: 313.

Stehling L and Zauder HL (1991):

Acute normovolemic hemodilution. Transfusion 31: 857.

Sten GE (1988):

Energy expenditure and fluid and electrolytes requirements in anaesthetized infants and children, anaesthesiology, vol. 69, No. 3, p 377.

Steward DJ (1995):

manual of Padiatric Anaesthesia, 4th ed. Churchill Livingstone.

Strauss RG (1981):

Review of the effects of hydroxy-ethyl starch on the blood coagulation system. Transfusion, 21: 299.

Thompson W, Shoemaker W, Ayres S (1989):

Textbook of critical care. Philadelphia, WB Saunders: 492-6, 1146-48.

Thompson WL (1988):

Hydroxy-ethyl starch, in blood substitutes and plasma expanders. new York, Alan R. Liss, Inc., p. 283.

Toshimasa Yashioka et al. (1990):

Body fluid compartments, from Pediatric Textbook of Fluid and Electrolytes, edited by Laurel Craven, P. 14.

Tucker and Blentz (1977):

An analysis of the determinants of nephron filtration rate. Am. J. Physiol. 232: 447.

Tullis PK and Waxman (1985):

"Artificial plasma expanders". In: *Advances in Anaesthesia*. New York, Alan R., Liss, Inc. vol. II, p. 293-316.

Van Workens E, Trouborst A, Van Lanschot (1992):

Profound haemodilution. *Anaesth. Analg.*, 75: 818-21.

Versmold HT, Kitterman JA et al (1981):

Aortic blood pressure during the first hours of life in infants. *Pediatrics* 67: 607.

Walker WF, Johnston J (1971):

Metabolic basis of surgical care. edited by William Heinemann, London, p. 210-223..

Weill MH, Henning RJ and Puri VK (1990):

Colloid osmotic pressure. Clinical Significance. *Crit. Care Med.*, 7: 113.

Welbron LG, Willis AM, Nisselson CL (1986):

Perioperative blood glucose concentration in paediatric outpatients, *Anaesthesiology* 65: 543-547.

Widman FK (1991):

Standards for blood banks and transfusion services,
ed 14. Arlington. American Association of Blood
Banks.

Winters RW (1982):

Principles of paediatric fluid therapy, 2 ed., Boston,
Little, Brown & Co.

Witte MK, Hill JH and blumer JL (1987):

Shock in paediatric patients. *Adv. Paediatr.* 34: 139.

Yacobi A, Stoll RG and Sun CY (1982):

Pharmacokinetics of hydroxy-ethyl starch in normal
subjects. *J Clin. Pharmacol.*, 22: 206-212.

Yelderman M, New W (1983):

Evaluation of pulse oximetry. *Anaesthesiology.* 59:
349.

Ynonne McMahan et al. (1990):

Management of patients immediately after surgery
from paediatric textbook of fluid and electrolytes.
Edited by Laure Carven, 459.

Zarins CK et al (1978):

Lymph and pulmonary response to isobaric reduction in plasma oncotic pressure in baboons. Circ. Res., 43: 925-930.

Zauder LH (1991):

Intraoperative and postoperative blood salvage devices. In Stehling LC, editor: Perioperative autologous transfusion. American Association of Blood Bank, p. 153-172.

Zimmerman and Defendini R (1977):

Hypothalamic pathways containing oxytocin, vasopressin and associated neurophysins. International Congress on the Neurohypophysis, Basel.