
REFERENCES

- Alo, K.M.; Yland, M.J.; Kramer, D.L.; Charnov, J.H.; Redko, V. (1998):* Computer assisted and patient interactive programming of dual octrode spinal cord stimulation in the treatment of chronic pain. *Neuromodulation*; 1: 30-39.
- American Cancer Society (1979):* Report on social economic and psychological needs of patients in Califomia. Major findings and implications. San Francisco, American Cancer Society, California Division.
- Anderson, G.B. (1991):* Sensitivity, specificity, and predictive value. In: Frymoyer JW ed. *The adult spine: principles and practices*. New York: Raven Press, 277-87.
- Angarola, R.T.; Mendelsohn, C.B. (1998):* Pain and health policy. In: *The management of pain*. Ashburn MA, Rice LJ (Eds.), Churchill Livingstone, New York, PP. 199-204.
- Arndr, S.; Lindblom, U.; Meyerson, B.A.; Molander, C. (1990):* Prolonged relief of neuralgia after regional anesthetic blocks. A call for further experimental and systematic clinical studies. *Pain*, 43: 287-97.
- Arner, S. (1991):* Intravenous phentolaminc test: diagnostic and prognostic use in reflex sympathetic dystrophy. *Pain*, 46: 17-22.
- Arnold, J.M.; Teasel!, R.W.; MacLeod, A.P.; Brown, J.E.; Carruthers, S.C. (1993):* Increased venous alpha-adrenoceptor responsiveness **in** patients with reflex sympathetic dystrophy. *Ann. Intern. Med.*, 118: 619-21.
- Augustinsson, L.E. (1995):* Spinal cord stimulation for chronic pain. *Neurosurg. Clin. North Am.*, 6: 157.
- Bahar et at (1997):* Spinal anaesthesia with midazolam in the rat. *Can. J. Anaesth.*, 44: 208-215.

-
-
- Baker, D.M.; Lamerton, A.J. (1994):* Operative lumbar sympathectomy for sever lower limb ischaemia still a valuable treatment option. *Ann. R. Coll. Sttrg. Engl.*, 76: 50-53.
- Barolat, G.; Ketcik, B. and He, J. (1998):* Long-term outcome of spinal cord stimulation for chronic pain management. *Neuromodulation*, 1: 19-30.
- Barolet, (1994):* Current status of epidural spinal cord stimulation. *Neurosurg.*, 5: 98.
- Benedetti, C. (1987):* Intraspinal analgesia, a historical overview, *acta Anesthesiol Scant*, 31 Supplementum, 85: 17-24.
- Bengtsson, M.; Lofstrom, J.B. (1990):* Nerve block in pancreatic pain. *Acta. Chir. Searid.*, 156: 285-91.
- Bhalla, A.P. (1996):* Glossopharyngeal neuralgia: a diagnostic conundrum-management by the multimodality approach (Abstract). //th World Congress of Anaesthesiologists; 1996 Apt 14-19; Sydney, Australia. Sydney: World Federation of Societies of Anaesthesiologists, 313.
- Blackie, J.D.; Lees, A.J. (1990):* Botulinum toxin treatment in spasmodic torticollis. *J. Neurol. Neurosurg. Psychiatry*, 53: 640-643.
- Bogduk, N. (1996):* Reply to: assessing a new procedure: thoracic radiofrequency dorsal root ganglion lesions. *Clin. J. Pain*; 12: 76-78.
- Bogduk, 1V.; Marsland, A. (1986):* On the concept of third occipital headache. *J. Neurol. Neurosurg. Psychiatry.*, 49: 775-80.
- Bonica, J.J. (1990):* The management of pain. 2nd ed. Rhea & Febiger. Philadelphia, P. 1982.
- Bonica, J.J. and Buckley, F.P. (1990):* Regional analgesia with local anaesthetics. In: Bonica JJ (eds): The management of pain, 2" ed. Philadelphia, Lea & Febiger, P. 1883.

-
-
- Borodic, G.E.; Ferrante, R.J.; Pearce, L.B.; Alderson, K. (1994):* Pharmacology and histology of the therapeutic application of Botulinum toxin. In: *Therapy with Botulinum Toxin*, ed. by Jankovic J, Hallett M. New York: Marcel Dekker, Inc., 119-157.
- Bovim, G.; Berg, R.; Dale, L.G. (1992):* Cervicogenic headache: anesthetic blockades of cervical nerves (C2-05) and facet joint (C2-C3). *Pain*, 49: 315-20.
- Bowen-Wright, R.M. (1990):* Precautions against injection into the spinal artery during celiac plexus block. *Anaesthesia*, 45: 247-48.
- Bowersox, S.S.; Gadbois, T.; Singh, T.; PeRus, M. et al (1996):* Selective N-type neuronal voltage sensitive calcium channel blocker, SNX-111 produces spinal antinociception in rat models of acute persistent and neuropathic pain. *J. Pharm. and Exp. Ther.*, 279: 1243-1249.
- Brown, D.L.; Bulley, C.K.; Quiel, E.L. (1987):* Neurolytic celiac plexus block for pancreatic cancer pain. *Anesth. Analg.*, 66: 869-73.
- Brown, D.L.; Rorie, D.K. (1994):* Altered reactivity of isolated segmental lumbar arteries of dogs following exposure to ethanol and phenol. *Pain*, 56: 139-43.
- Busch, E.; Wilson, P.R. (1989):* Atlanto-occipital and atlanto-axial injections in the treatment of headache and neck pain (Abstract). *Reg. Anesth.*, 14(52): 45.
- Campero, M.; Verdugo, R.J.; Ochoa, J.L. (1993):* Vasomotor innervation of the skin of the hand: a contribution to the study of human anatomy. *J. Anal.*, 182: 361-68.
- Casale, R.; Glynn, C.J.; Buonocore, M. (1990):* Autonomic variations after stellate ganglion block: are they evidence of an autonomic afference? *Funct. Neurol.*, 5: 245-46.
- Cherny, IV.L et al. (1994):* Opioid responsiveness of cancer pain caused by neuropathic or nociceptive mechanisms. A combined analysis of controlled single dose studies. *Neurology*, 44: 857-61.

-
-
- Cheshire, W.P.; Abashian, S.W.; Mann, .I.D. 0994):* Botulinum toxin in treatment of myofascial pain syndrome. *Pain*, 59: 65-69.
- Choe, H.; Choi, ES.; Kim, EH.; Ko, S.H.; Choi, H.G.; Han, Y.J.; Song, H.S. (1997):* Epidural morphine plus ketamine for upper abdominal surgery-improved analgesia from preincisional versus postincisional administration. *Anesthesia & Analgesia*, 84(3): 560-563.
- Choi, Y.K.; Chou, S. (1995):* Recuts syndrome. Another cause of abdominal pain. *Reg. Anesth.*, 20: 347-51.
- Christie, J.M.; Martinez, C.R. (1995):* Computerized axial tomography to define the distribution of solution after stellate ganglion nerve block. *J. Clin. Anesth.*, 7: 306-11.
- Claeys, L. (1999):* Spinal cord stimulation and chronic critical limb ischemia. *Neuromodulation*; 2: 1-3.
- Clarke, A.J.; Sitnpson, K.H.; Ellis, F.R. (1990):* Continuous brachial plexus block in the management of intractable cancer pain in the arm. *Palliar Med.*, 4: 123-25.
- Coffey, R.J. (1996):* Neurosurgical management of intractable pain. P. 3428. In: Yomans JR (ed): *Neurological Surgery*, 4th Ed. WB Saunders; Philadelphia.
- Coldwell, D.M.; Loper, K.A. (1989):* Regional anaesthesia for hepatic arterial embolization. *Radiology*, 172: 21039-40.
- Cosman, E.r.; Nashold, B.S.; Ovehnan-Levitt, J. (1984):* Theoretical aspects of radiofrequency lesions in the dorsal root entry zone. *Neurosurgery*; 15: 945-50.
- Cousins, M.J. (1989):* Selective spinal analgesia. *Lancet*, 1: 1141-1142.
- Cui, J.G.; Meyerson, B.A.; Solleni, A. and Linderorth, B. (1998):* Effects of spinal cord stimulation on tactile hypersensitivity in mononeuropathic rats is potentiated by GABA-B and adenosine receptor activation. *Neurosci. Lett*; 247: 183-186.
-
-

-
-
- et/4 J.C.; Soiletei, A.; Linderoih, B. and Meyerson, B.A. (1997)!*
Adenosine receptor activation suppresses tactile hypersensitivity and potentiates effect of spinal cord in mononeuropathic rats. *Neurosis Lett.*, 223: 173-176.
- Dangoisse, M.; Collins, S.; Glynn, C.J. (1994):* Haemothorax after attempted intercostals catheterization, *Anaesthesia.* 49; 961-63.
- Davies, D.D. (1993):* Incidence of major complications of neurolytic celiac plexus block. *J.R. Soc. Med.*, 8: 264-66.
- De Leon-Casasola, O.A.; Kent, E.; Lema, M.J. (1993):* Neurolytic superior hypogastric plexus block for chronic pelvic pain associated with cancer. *Pain*; 54: 145-51.
- De Lissovoy, G. et al (1997):* Cost effectiveness of long-term intrathecal morphine therapy for pain associated with failed back surgery syndrome. *Clin. Ther.*, 19(1): 96-112.
- DeConno, F. et al (1994):* Pain measurement in cancer patients. A comparison of six methods. *Pain*, 57: 161-66.
- Deer, T. (1997):* Interventional pain management American Society of Anesthesiologists, annual refresher course lectures, 176: 1-7.
- Deer, T. (1998):* The role of neuromodulation by spinal cord stimulation in chronic pain syndromes; current concepts. *Techniques in Regional Anesthesia and Pain Management.* Vol. 2, No. 3: 161-167.
- Deer, T.; Winkelmuller, W.; Erdine, S.; Bedder, M.; Burchiel, K. (1999):* Intrathecal therapy for cancer and non malignant pain; *Neuromodulation*, Vol. 2, No. 2; 55-66.
- DeLeon-Casasala, O.A. (1997):* Superior hypogastric block and ganglion impar neurolysis for pain associated with cancer. *Tech. Reg. Anesth. Pain Manage*, 1: 27.
- Denemijn, P.L.; Fields, H.L.; Allen, R.R.; McKay, W.R.; Rowbotham, M.C. (1994):* The interpretation of pain relief and sensory changes following sympathetic blockade. *Brain*; 117: 1475-87.
-
-

Dijkema, H.E.; Weil, E.H.H.; Mijs, P.T.; Janknegt, R.A. (1993):

Weuromoclutation of sacral nerves for incontinence and voiding dysfunctions. *Eur. Urol.*, 24: 72-76.

Dowman, R. (1991): Spinal and supraspinal correlates of nociception in man. *Pain*, 45: 269-281.

Dubuisson, D. (1994): Root surgery. In: Wall PD, Melzack R. eds. *Textbook of Pain*. Edinburgh: Churchill Livingstone, Third edition, 1055-65.

Duncan, L.; Wildsmith, J.A. (1995): Liposomal local anaesthetics. *Br. J. Anaesth.*, 75: 260-61.

DuPen, S.L.; Kharasch, E.D.; Williams, A. et al. (1992): Chronic epidural bupivacaine-opioid infusion in intractable cancer pain. *Pain*, 49: 293.

Durrani, Z.; Winnie, A.P.; Ikuta, P. (1988): Interpleural catheter analgesia for pancreatic pain. *Anesth. Analg.*, 67: 479-81.

Eisenach, J.; DuPen, S.L.; Dubois, M. et al. (1995): Epidural clonidine analgesia for intractable cancer pain. *Pain*, 61: 391.

Eisenberg, E.; Carr, D.B.; Chalmers, T.C. (1995): Neurolytic celiac plexus block for treatment of cancer pain: a meta-analysis. *Anesth. Analg.*, 80: 290-95.

Elias, M.; Chakerian, M.U. (1994): Repeated stellate ganglion blockade using a catheter for pediatric herpes zoster ophthalmicus. *Anesthesiology*, 80: 950-52.

Elliot Krames (1999): The Egyptian international congress on pain.

Emery, P.; Bowman, S.; Wedderburn, L.; Grahame, R. (1989): Suprascapular nerve block for chronic shoulder pain in rheumatoid arthritis. *BMJ*, 299: 1079-80.

Erdine, S.; Yucel, A. (1995): Complications of drug delivery systems. *Pain Reviews*, 2: 227-242.

Erdine, S.; Yucel, A. (1996): Intrathecal morphine by implanted manual pump for cancer pain. *Pain Digest*, 6: 161-165.

-
-
- Erickson, S.J.; Hogan, Q.H. (1993):** CT-guided injection of the stellate ganglion: description of technique and efficacy of sympathetic blockade. *Radiology*; 188: 707-709.
- Esses, S.L; Moro, J.K. (1993):** The value "of facet joint blocks in patient selection for lumbar fusion. *Spine*, 1: 185-90.
- Felder, C.; Whitworth, T.; Brookof, D.; Alo, K.; Viand, M.J. (1998):** Sacral nerve root stimulation for the treatment of intractable pain due to interstitial cystitis. Abstracts of American Pain Society Meeting.
- Ferrandiz, M.; Aliaga, L.; Catala, E.; Villar-Landeira, J.M. (1994):** Thoracic paravertebral block in chronic postoperative pain. *Reg. Anesth.*, 19: 221-22.
- Ferraz, M.B. et al (1990):** Reliability of pain scales in the assessment of literate & illiterate patients with rheumatoid arthritis. *J. Rheumatol.*, 17: 1022-24.
- Fields, H.L. (1996):** Treatment of trigeminal neuralgia. *N. Engl. J. Med.*, 334: 1125-26.
- Foley, K.M. (1987):** Assessment of pain in patients with cancer in cancer pain ed. M. Swerdlow and V. Ventafridda, 37-44.
- Fredman, B.; Ben Nun, B.; Zohar, E. et al (1999):** Epidural steroids for treating "Failed back surgery syndrome". *Anesth. Analg.*, 88: 367-72.
- Fujita, Y. (1993):** CT-guided neurolytic splanchnic nerve block with alcohol. *Pain*, 55: 363-66.
- Gamer, L.; Coats, R.R. (1994):** Ipsilateral stellate ganglion block effective for treating shoulder pain after thoracotomy. *Anesth. Analg.*, 78: 1195-96.
- Ganong, W.F. (1995):** Review of medical physiology. Appleton Lange California, 103-119.
- Gordh, T.; Karlsten, R.; Kristenson, J. (1995):** Intervention with spinal NMDA, adenosine, and No systems for pain modulation. *Ann. Med.*, 27: 229-234.
-
-

-
-
- Gordon, A.; Zechmeister, K.; Collin, J. (1994): The role of sympathectomy in current surgical practice. Eur. J Vase Strg., 8: 129-37.*
- Gracely, R.H.; Lynch, S.A.; Bennett, G.J. (1992): Painful neuropathy: altered central processing maintained dynamically by peripheral input. Pain, 51: 175-94.*
- Grant, G.J.; Vermeulen, K.; Langerman, L.; Zakowski, M.; Turndof, H. (1994): Prolonged analgesia with liposomal bupivacaine in a mouse model. Reg. Anesth., 19: 264-69.*
- Greenwold, H.P. (1991): Interethnic differences in pain perception. Pain, 44: 157-63.*
- Guyatt, G.; Walter, S.; Norman, G. (1987): Measuring changeover time: assessing the usefulness of evaluative instruments. J. Chron. Dis., 40: 171-78.*
- Guyton, A.C. and Hall, J.E. (1996): Pain pathway-textbook of medical physiology. W.B. Saunders, Philadelphia, 609-620.*
- Haddox, J.D. (1996): A call for clarity. In: Campbell JN ed. Pain, 1996 an updated review. Seattle: IASP Press, 97-99.*
- Hamann, W.; Hall, S. (1992): Acute effect and recovery of primary afferent nerve fibres after graded radiofrequency lesions in anaesthetized rats (Abstract). Br. J. Anaesth., 68: 443.*
- Hanna, M.H.; Peat, S.J. (1989): Ketanserin in reflex sympathetic dystrophy. A double-blind placebo controlled cross-over trial. Pain; 38: 145-50.*
- Hannington-Kiff J.G. (1991): Doses failed natural opioid modulation in regional sympathetic ganglia cause reflex sympathetic dystrophy? Lancet; 338: 1125-27.*
- Hansson, C.; Holm, J.; Lillieborg, S.; Syren, A. (1993): Repeated treatment with lidocaine/prilocaine cream (EMLA) as a topical anaesthetic for the cleansing of venous leg ulcers. A controlled study. Acta. Derm. Venereol., 73: 231-33.*
-
-

-
-
- Hassenbush, S.J. et al (1995):* Long-term intraspinal infusions of opioids in the treatment of neuropathic pain. *J. Pain Sympt. Management*, 10: 527.
- Hatangdi, V.S.; Boas, R.A. (1985):* Lumbar sympathectomy: a single needle technique. *Br. J. Anaesth.*, 57: 285-89.
- Hogan, Q.; Doston, R.; Erickson, S.; Kettler, R.; Hogan, K. (1994):* Local anesthetic myotoxicity: a case and review. *Anesthesiology*, 80: 942-47.
- Hogan, Q.H. (1998):* Neural blockade for diagnosis and treatment of painful conditions. In: Ashburn MA, Rice LJ (eds): *The Management of pain*, New York, Churchill Livingstone, P. 275.
- Hogan, Q.H. and Abram, S.E. (1998):* Diagnostic and prognostic neural blockade. In: Cousins MJ, Bridenbaugh PO (ed): *Neural Blockade in Clinical Anesthesia and Management of Pain*, 3'd ed. Philadelphia, Lippincott-Raven, P. 837.
- Hogan, Q.H.; Erickson, S.J.; Haddox, J.D.; Abram, S.E. (1992):* The spread of solutions during stellate ganglion block. *Reg. Anesth.*, 17: 78-83.
- Hoogbergen, M.M.; Kauer, J.M. (1992):* An unusual ulnar nerve-median nerve communicating branch. *J. Anal.*, 181: 513-16.
- Hord, A.H.; Rooks, M.D.; Stephens, B.O.; Rogers, H.G.; Fleming, L.L. (1992):* Intravenous regional bretylium and lidocaine for treatment of reflex sympathetic dystrophy: a randomized, double-blind study. *Anesth. Analg.*, 74: 818-21.
- Houraly, P.; Vangertruyden, G.; Verduyck, F.; Trimpeneers, F.; Hendrickx, J. (1995):* Endoscopic extraperitoneal lumbar sympathectomy. *St. Endosc.*, 9: 530-33.
- Hsaue, M. (1993):* Pain and the nerve root. An inter-disciplinary approach. *Spine*, 18: 2053-58.
- Igarashi, T.; Hirabayashi, Y.; Saitoh, Fukuda, H. (1998):* Thoracic and lumbar extradural structure examined by extraduroscope. *Br. J. Anaesth.*, 81: 121-125.
-
-

-
-
- Janig, W. (1996):* The puzzle of reflex sympathetic dystrophy. In: Janig W, Startton-Hicks Meds. Reflex sympathetic dystrophy: a reappraisal. (Progress in pain research and management Vol. 6) Seattle: IASP Press, 1: 24.
- Jensen, M.B.; Turner, J.A. et al (1995):* What is the maximum number of levels needed to pain intensity measurements? Pain, 58: 387-92.
- Jensen, M.P. (1986):* The measurement of clinical pain intensity. A comparison of six methods. Pain, 27: 117-26.
- Johansson, A.; Hao, J.; Sjund, B. (1990):* Local corticosteroid application blocks transmission in normal nociceptive c-fibres. Acta. Anaesthesiol. Scand., 34: 335-338.
- Johansson, A.; Sjolund, B. (1996):* Nerve blocks with local anesthetics and corticosteroids in chronic pain: a clinical follow-up study. J. Pain Symptom. Manage., 11: 181-87.
- Jones, M.W. (1993):* Commissural myelotomy for relief of intractable pain. In Arbut E (ed): Management of cancer related pain. Futura, Mount Kisco, NY.
- Justins, D.M.; Rubin, A.P. (1993):* Pain and autonomic blocks. In: Wildsmith JAW, Armitage EN eds. Principles and practice of regional anaesthesia, second edition. Edinburgh: Churchill Livingstone, 233-47.
- Kapral, S.; Krafft, P.; Gosch, M.; Fleischmann, D.; Weinstabl, C. (1995):* Ultrasound imaging for stellate ganglion block: direct visualization of puncture site and local anesthetic spread. A— or study. Reg. Anesth., 20: 323-28.
- Katz, N. (1994):* Role of invasive procedures in chronic pain management. Semin. Neurol., 14: 225-36.
- Katz, IV. and Ferrante, F.M. (1993):* Nociception. Vade Boncouer T.R., editors: Postoperative pain management, New York, Churchill-Livingstone.

-
-
- Kawmata, M. et al (1996):* Comparison between celiac plexus block and morphine treatment on quality of life in patients with pancreatic cancer pain. *Pain*; 64(3): 597-602.
- Kim, D.H.; Kline, D. (1995):* Surgical outcome for intra- and extrapelvic femoral nerve lesions. *J. Neurosurg.*, 83: 783-90.
- Kissin, L (1996):* Preemptive analgesia. Why its effect is not always obvious. *Anesthesiology*, 84: 1015-19.
- Konig, H.M.; Koster, H.G.; Niemeijer, R.P. (1991):* Ischaemic spinal cord lesion following percutaneous radiofrequency spinal rhizotomy. *Pain*; 45: 161-66.
- Korsten, H.H.; Hellebrekers, L.J.; Grouls, R.J. et al (1990):* Long-lasting epidural sensory blockade by n-butyl p-aminobenzoate in the dog: neurotoxic or local anesthetic effect? *Anesthesiology*, 73: 491-98.
- Krames, E. et al. (1993):* Intrathecal infusional analgesia for intractable pain; patient management Guidelines. *J. Pain Symptom Manage*, 9: 36-46.
- Krames, E.; Olson, K., Clinic Realities and Economic Considerations (1997):* Patient selection in intrathecal therapy. *Pain Symptom Manage*, 14: S3-S13.
- Krames, E.S. (1993):* Intrathecal infusional therapies for intractable pain: patient management guidelines. *J. Pain Sympt. Manage*, 8: 36.
- Krames, E.S.; Schuchard, M. (1995):* Implantable intraspinal infusional analgesia: management guidelines, *pain Reviews*, 2: 243-267.
- Kramis, R.C.; Robbers, W.J.; Gillette, R.G. (1990):* Post-sympathectomy neuralgia: hypotheses on peripheral and central neuronal mechanisms. *Pain*, 64: 19.
- Lamacraft, G. (1998):* Neural blockade in chronic and cancer pain. *Neural blockade*; 131-153.

-
-
- Lamer, T.J. (1994):* Treatment of cancer-related pain: when orally administered medications fail. *Mayo. Clin. Proc.*, 69: 473-80.
- Lamotte, R.H.; Shaine, C.N.; Simone, D.A. and Tsai, E.P. (1991):* Neurogenic hyperalgesia: Psychophysical studies of underlying mechanisms. *Journal of Neurophysiology*, 66: 190-211.
- Lee, E.M. (1997):* Pathogenesis and mechanisms of phantom pain. *Current Review of Pain*; 1: 310-319.
- Lee, R.B.; Stone, K.; Magelssen, D.; Belts, R.P.; Benson, W.L. (1986):* Presacral neurectomy for chronic pelvic pain. *Obstet. Gynecol.*, 68: 517-21.
- Levin, A.B.; Katz, J.; Benson, R.C. (1980):* Treatment of pain of diffuse metastatic cancer by stereotactic chemical hypophysectomy: long term results and observations on mechanism of action. *Neurosurgery*, 6: 258.
- Lillemoe, K.D.; Cameron, J.L.; Kaufman, H.S.; Yeo, C.J.; Pitt, I.L.A.; Sauter, P.K. (1993):* Chemical splanchnicectomy in patients with unresectable pancreatic cancer. A prospective randomized trial. *Ann. Surg.*, 217: 447-57.
- Lin, C.C.; Mo, L.R.; Lin, Y.W.; Yau, M.P. (1994):* Bilateral thoracoscopic lower sympathetic splanchnicectomy for upper abdominal cancer pain. *Eur. J. Surg. Suppl.*, (572): 59-62.
- Linker, IV.J.; Camm, A.J.; Ward, D.E. (1991):* Dynamics of ventricular repolarisation in the congenital long QT syndromes. *Br. Heart J.*, 66: 230-37.
- Lipfert, P.; Seitz, R.J.; Arndt, J.O. (1987):* Ultralong-lasting block: triethyldodecyl ammonium bromide is probably a neurotoxin rather than a local anaesthetic. *Anesthesiology*, 67: 896-904.
- Loeser, J.D.; Sweet, W.H.; Tew, J.M.; van Loveren, H. (1990):* Neurosurgical operations involving peripheral nerves. In: Bonica JJ, Loeser JD, Cllaproan CR, Fordyce WE eds. *The management---b)---pi'* in. Philadelphia, PA: Lea and Febiger, 2044-66.

-
-
- Lofstrom, J.B.; Cousins, M.J. (1988):* Symphetic neural blockade of upper and lower extremity. In: Cousins MJ, Bridenbaugh PO eds. *Nettral blockade in clinical anesthesia and management of pain*. Second edition. Philadelphia, PA: Lippincott, 461-502.
- Lord, S.M.; Barnsley, L.; Bogduk, IV. (1995):* Percutaneous radiofrequency neurotomy in the treatment cervical zygapophysial joint pain: a caution. *Neurosurgery*; 36: 732-39.
- Lord, S.M.; Barnsley, L.; Wallis, B.J.; Bogduk, IV. (1994):* Third occipital nerve headache: a prevalence study. *J. Neurol. Neurosurg. Psychiatry.*, 57: 1187-90.
- Lord, S.M.; Barnsley, L.; Wallis, B.J.; Bogduk, IV. (1997):* A randomized, double-blind, controlled trial of percutaneous radiofrequency neurotomy for the treatment of chronic cervical zygapophysial joint pain. 8th World Congress on Pain; 1996 Aug. 17-22; Vancouver, Canada. Seattle: International Association for the Study of Pain, 382.
- Low, P.A.; Opfer-Gehriking, T.L.; Dyck, P.J.; Litchy, W.J.; O'Brien, P.C. (1995):* Double-blind, placebo controlled study of the application of capsaicin cream in chronic distal painful polyneuropathy. *Pain*, 62: 163-68.
- Marks, R.C.; Houston, T.; Thulbourne, T. (1992):* Facet joint injection and facet nerve block: a randomized comparison in 86 patients with chronic low back pain. *Pain*; 49: 325-28.
- Marmheimer, C.; Eliasson, T.; Augustinsson, L.E.; Blondstrand, C.; Emanuelsson, H.; Larrson, S. et al (1998):* Electrical stimulation versus coronary artery bypass surgery in sever angina pectoris: The ESBY study. *Circulation*, 97: 1157-1163.
- Maurri, S.; Barontini, F. (1990):* Responsiveness of idiopathic spasmodic torticollis to Botulinum A toxin injection: A critical evolution of five cases. *Clin. Neurol. Neurosurg.*, 92: 165-168.

-
-
- Mauskop, A. (1993):* Trigeminal neuralgia (tic douloureux). *J. Pain Symptom Manage.*, 8: 148-54.
- McMahon, S.B. (1991):* Mechanisms of sympathetic pain. *Br. Med. Bull.*, 47: 584-6(10).
- McMahon, S.M. and Koltzenurg, M. (1990):* The changing role of primary afferent neurons in pain. *Pain*, 43: 269-72.
- Mercadante, S. (1993):* Celiac plexus block versus analgesics in pancreatic cancer pain. *Pain*, 52: 187-92.
- Mercadante, S. (1995):* Octreotide in the treatment of diarrhoea induced by celiac plexus block. *Pain*, 61: 345-46.
- Meyerson, B.A.; Cui, J.G.; Yakhnitsa, V. et al (1997):* Modulation of spinal pain mechanisms, by spinal cord stimulation and the potential role of adjuvant pharmacotherapy. *Stere-tact funct. Neurosurg.*, 68: 129-140.
- Michael, A.; Weaver, Thomas, M. Root, Marc, B. Hahn (1998):* Implantation therapy for pain management. In: Michael A., Ashburn, M.D. *The management of pain*. 26: 419-485.
- Miller (2000):* Ronald D. Miller, *Anesthesia*, Fifth edition, P. 2351.
- Miller (1998):* Ronald D. Miller, *Anesthesia*, Fourth edition, P. 2365.
- Molloy, A.R.; Kalski, R.; Hong, W.; Lawless, M.; Cousins, M.J. (1996):* Topical morphine in the management of chronic eye pain (Abstract). 11th World Congress of Anaesthesiologists; 1996 Apr 14-19; Sydney, Australia. Sydney: World Federation of Societies of Anaesthesiologists, 321.
- Moore, N.R.; Dixon, A.K.; Freer, C.E. (1989):* The anatomy of the first sacral nerve root sheath shown by computed tomography. *Br. J. Radiol.*, 62: 710-15.
- Murali, R. (1990):* Peripheral nerve injections and avulsions in the treatment of trigeminal neuralgia. In: Rovit RL, Murali R, Jannetta PJ eds. *Trigeminal neuralgia*. Baltimore, MD: Williams and Wilkins, 95: 108.

Murphy, 7".M. (1994): Chronic pain. In: Miller RD (ed): Anaesthesia, 4th ed. New York, Churchill Livingstone, P. 2345.

Myers, R.R.; Katz, J. (1988): Neuropathology of neurolytic and semidestructive agents. In: Cousins M.J., Bridenbaugh PO eds. Neural blockade in clinical anesthesia and management of pain, second edition. Philadelphia, PA: Lippincott, 1031-51.

Nachbur et aL (1994): Spinal cord stimulation for unreconstructable chronic limb ischemia. Eur. J. Vasc. Surg., 8: 383.

National Mstitutes of Health (NIH) (1997): Consensus Development Conference on Acupuncture. Bethesda, Md. NTH.

Neidre, A.; MacNab, L (1983): Anomalies of the lumbosacral nerve roots. Review of 16 cases and classification. Spine, 8: 294-99.

Nelson, D. (1996): Psychological selections criteria for implantable spinal cord stimulators. Pain Forum; 5(8): 93-103.

Neuromed Bull (1995): Neruomed Bull, 2, 5.

Nishikimi, 1V.; Sakurai, T.; Shionoya, S.; Oshima, M. (1992): Microcirculatory characteristics in patients with Buerger's disease. Angiology; 43: 312-19.

Nitescu, P.; Sjoberg, M.; Appलगren, L. et aL (1995): Complications of intrathecal opioids and bupivacaine in the treatment of "refractory" cancer pain. Clin. J. Pain, 11: 45.

Nitta, H.; Tajima, T.; Sugiyama, H.; Moriyama, A. (1993): Study on dermatomes by means of selective lumbar spinal nerve block. Spine, 18: 1782-86.

North, R. et aL (1991): Failed back surgery syndrome: 5-year follow-up after spinal cord stimulation implantation. Neurosurgery; 28: 692-99.

North, R.B. (1993): Neurosurgical procedures for chronic pain: general neurosurgical practice. Clin. Neurosurg., 40: 182.

-
-
- North, R.B. (1993):** Spinal cord stimulation for chronic, intractable pain. Adv. Neurol., 63: 289.
- North, R.B.; Boark, G.L. (1995):** Spinal cord stimulation for chronic pain. Neurosurg. Clin. North Am., 6: 45.
- North, R.B.; Han, M.; Zahurak, M.; Kidd, D.H. (1994):** Radiofrequency lumbar facet denervation: analysis of prognostic factors. Pain, 57: 77-83.
- North, R.B.; Kidd, D.H.; Campbell, J.N. et al. (1991):** Dorsal root ganglionectomy for failed back surgery syndrome: a five year follow-up study. J. Neurosurg., 74: 236.
- North, R.B.; Kidd, D.H.; Zahurak, M.; Piantadosi, S. (1996):** Specificity of diagnostic nerve blocks: a prospective, randomized study of sciatica due to lumbosacral disease. Pain, 65: 77-85.
- Oakley, J.C. and Weiner, R.L. (1999):** Spinal cord stimulation for complex regional pain syndrome: a prospective study of 19 patients at two centers. Neuromodulation; 2: 47-50.
- Omar Tawfik (1999):** 2nd annual conference of anaesthesia, intensive care and pain management, P. 12.
- Omar Tawfik (2001):** Conference on interventional pain therapy. Elein Elsokhna, 2-5 May, 2001.
- Paramore, L.O (1997):** The use of alternative therapies: Estimates from the 1994 Robert Wood Johnson Foundation National Access to Care Survey. J. Pain Symptom Manage., 13: 83.
- Parris, W.C.; Reddy, B.C.; White, H.W.; McGrath, D.M. (1991):** Stellate ganglion blocks in pediatric patients. Anesth. Analg., 72: 552-56.
- Pali, R.B. (1993):** Cancer pain. P. 285, Lippincott-Raven, Philadelphia.
- Penn, R.D.; Paice, J.A.; Kroin, J.S. (1990):** Intrathecal octreotide for cancer pain. Lancet, 738.

-
-
- Plancarte, R.; Amescua, C.; Patt, R.B.; Aldrete, J.A. (1990): Superior hypogastric block for pelvic pain. Anesthesiology; 73: 236-39.*
- Poewe, W.; Wissel, J. (1994): Experience with Botulinum toxin in cervical Dystonia. In: Therapy with Botulinum toxin, ed. by Jankovic J, Hallett M. New York: Marcel Dekker Inc., 267-278.*
- Porzelius, J. (1995): Memory for pain after nerve-block injections. CI in. J. Pain, 11: 112-20.*
- Prasanna, A.; Murthy, P.S. (1993): Sphenopalatine ganglion block and pain of cancer (Letter). J. Pain Symptom. Manage, 8: 125.*
- Prithivi Raj (1997): Botulinum toxin, current Review of pain, 1: 403-416.*
- Prithvi Raj (1996): Pain mechanisms: Pain Medicine a comprehensive review. Mosby-Year Book, PP. 12-34.*
- Purcell-Jones, G.; Pither, C.E.; Justins, D.M. (1989): Paravertebral somatic nerve block: a clinical, radiographic, computed tomographic study in chronic pain patients. Anesth. Analg., 68: 32-39.*
- Rains, C.P.; Bryson, H.M. (1995): Topical capsaicin. A review of its pharmacological properties and therapeutic potential in post-herpetic neuralgia, diabetic neuropathy and osteo-arthritis. Drugs Aging, 7: 317-28.*
- Raj, P.P. (1997): Botulinum toxin in the treatment of pain associated with musculoskeletal hyperactivity. Anaesthetic techniques in pain management in current review of pain; 1: 403-416.*
- Raj, P.P. and Evans, D. (1997): Accupuncture. In: Raj PP (ed): Pain Medicine: A comprehensive Review. St. Louis, CV Mosby, P. 309.*
- Raja, S.N.; Choi, Y.; Asano, Y.; Holmes, C.; Goldstein, D.S. (1995): Arteriovenous differences in plasma concentrations of catechols in rats with neuropathic pain. Anesthesiology, 83: 1000-1008.*
-
-

-
-
- Ramamurthy, S.; Hoffman, J. (1995): Intravenous regional guanethidine in the treatment of reflex sympathetic dystrophy/causalgia: a randomized, double-blind study. Guanethidine Study Group. Anesth. Analg., 81: 718-23.*
- Rawal, N. (1995): Spinal antinociception: Clinical aspects. Annuals of Medicine, 27: 263-268.*
- Reddy, P.D.; Zelicof, S.b.; Ruotolo, C.; Holder, J. (1995): Interdigital neuroma. Local cutaneous changes after corticosteroid injection. Clin. Orthop., 317: 185-87.*
- Rizor, R.F. (1994): Epiduroscopy: technique for examination of the lumbar epidural space. ASRA nineteenth annual meeting; Chicago.*
- Rocco, A.G. (1995): Radiofrequency lumbar sympatholysis. The evolution of a technique for managing sympathetically maintained pain. Reg. Anesth., 20: 3M2.*
- Rocco, A.G.; Palombi, D.; Raeke, D. (1995): Anatomy of the lumbar sympathetic chain. Reg. Anesth., 20: 13-19.*
- Rosenberg, P.H. (1996): Fiberoptic epidurosocopy-current research. 711 International Pain Clinic, Istanbul, Turkey, Abstract, P. 124.*
- Saberski, L.; Kitahata, K. (1995): Direct visualization of the lumbosacral epidural space through the sacral Hiatus. Anesth. Analg., 80: 839-40.*
- Saberski, L.R. (1996): Spinal endoscopy: current concepts. In: International Pain Management. Eds Waldman SD and Winnie, AP. WB Saunders Co., PP. 137-149.*
- Sanders, M.; Zuurmond, W. (1995): Safety of unilateral and bilateral percutaneous ceruical cordotomy in 80 terminally ill cancer patients. J. Clin. Oncol., 13: 1509.*
- Sato, S.; Yamashita, S.; Iwat, M.; Mizuyama, K.; Satsumae, L (1994): Continuous interscatene block for cancer pain. Reg. Anesth., 19: 73-75.*

-
-
- Scudding, J.W. (1994):* Peripheral neuropathies. In: Wall PD, Melzack R. Textbook of pain, third edition, Edinburgh: Churchill Livingstone, 667-83.
- Schott, G.D. (1994):* Visceral afferents: their contribution to "sympathetic dependent" pain. *Brain*, 117: 39-413.
- Schwarzer, A.C.; Aprill, C.N.; Derby, R.; Fortin, J.; Kine, G.; Bogduk, IV. (1994):* The false-positive rate of uncontrolled diagnostic blocks of the lumbar zygapophysial joints. *Pain*, 58: 195-200.
- Serlin, R.C. et al (1995):* When is cancer pain mild, moderate or severe? Grading the pain severity by its interference with function. *Pain*, 61: 277-84.
- Shetter, A. (1997):* Spinal cord stimulation in the treatment of chronic pain. *Current Review of Pain*; 1(3): 213-222.
- Sidall, P.J.; Cousins, M.J. (1998):* Introduction to pain mechanisms: Implications for neural blockade. In Cousins MJ, Bridenbaugh PO (eds): *Neural Blockade in Clinical Anaesthesia and Management of Pain*, 3rd ed. Philadelphia, Lippincott-Raven, P. 675.
- Siddall, P.J.; Cousins, M.J. (1995):* Recent advances in pain management. *Aust. NZJ Surg.*, 65: 674-85.
- Sinatra, R.S. (1998):* Acute pain management and acute pain services. In Cousins MJ, Bridenbaugh PO (ed): *Neural blockade in Clinical Anaesthesia and Management of Pain*, 3rd ed., Philadelphia, Lippincott-Raven, P. 793.
- Sjaastad, O.; Fredriksen, T.A.; Pfaffenrath, V. (1990):* Cervicogenic headache: diagnostic criteria. *Headache*, 30: 725-26.
- Sluiter, M.E. (1988):* The use of radiofrequency lesions for pain relief in failed back patients. *Int. Disabil. Stud.*, 10: 37-43.
- Smith, H.P.; McWhorter, J.M.; Challa, V.R. (1981):* Radiofrequency neurolysis in a clinical model: neuropathological correlation. *J. Neurosurg.*, 55: 246-53.

Staats, P.S.; Kost-Byerly, S. (1995): Celiac plexus blockade in a 7 year old child with neuroblastoma. *J. Pain Symptom. Manage.*, 10: 321-24.

Stanton-Hicks, M.; Abram, S.; Nolte, H. (1986): Sympathetic blocks. In: Raj PP ed. *Practical management of pain*. Chicago, IL: Year Book, 661—!-1.

Stanton-Hicks, M.; Raj, P.P.; Racz, G.B. (1996): Use of regional anesthetics for diagnosis of reflex sympathetic dystrophy and sympathetically maintained pain: a critical evaluation. In: Janig W. Startton* Hicks M eds. *Reflex sympathetic dystrophy: a reappraisal*. Seattle: IASP Press, 217-37.

Stanton-Hicks, M.; Salmon, J. (1997): Stimulation of the central and peripheral nervous system for the control of pain. *J. Clin. Neurophysiol.*, 14: 46-62.

Stechison, M.T.; Mullin, B.B. (1992): Surgical treatment of Greater occipital neuralgia: an appraisal of strategies. *Acta. Neurochirurgia*; 131: 236-240.

Stein, C. (1995): Morphine- a local anaesthetic. *Pain Clin. Updates*, 3: 1-4.

Stein, C.; Milian, M.J.; Shippenberg, T.S.; Peter, K.; Herz, A. (1989): Peripheral opioid receptors mediating antinociception in inflammation. Evidence for involvement of mu, delta, and kappa receptors. *J. Pharmacol. Exp. Ther.*, 248: 1269-75.

Stoelting, R.K. (1993): Pharmacology and physiology in anesthetic practice. J.B. Lippincott, Philadelphia, 70-100.

Stolker, R.J.; Vervest, A.C.; Groen, G.J. (1993): Percutaneous facet denervation in chronic spinal pain. *Acta. Neurochir. Wien*, 122: 82-90.

Stolker, R.J.; Vervest, A.C; Groen, G.J. (1994): The management of chronic spinal pain by blockades: a review. *Pain*; 58: 1-20.

Taha, J.M.; Tew, J.M. (1996): Comparison of surgical treatments for neuralgia: reevaluation of radiofrequency rhizotomy. *Neurosurgery*, 38: 865-71.

-
-
- Taha, J.M.; Tew, J.M.; Buncher, C.R. (1995): A prospective 15-year follow up of 154 consecutive patients with trigeminal neuralgia treated by percutaneous stereotactic radiofrequency thermal rhizotomy. J. Neurosurg., 83: 989-93.*
- Taha, J.W.; Tew, J.M. (1995): Long-term results of radiofrequency rhizotomy in the treatment of cluster headache. Headache, 35: 193-96.*
- Tavola, T.; Gana, G.; Conte, G. et al. (1992): Traditional Chinese acupuncture in tension type headache: A control study. Pain, 48: 325.*
- Tawfik, M.D. (1994): Mode of action of intraspinal opioids. Pain Reviews; 1: 275-294.*
- Timothy, R. (1999): Interventional Pain Medicine American Society of Anesthesiologists, annual refresher course lectures, 145: 1-7.*
- Timothy, R. (2000): Interventional Pain Medicine American Society of Anesthesiologists, annual refresher course lectures, 216: 1-7.*
- Torebjor, K.H.E.; Lundberg, L.R. and Lamotte, R.H. (1992): Central changes in processing of mechanoreceptive input in capsaicin induced secondary hyperralgesia in humans. Journal of Physiology, 448: 765-780.*
- Torebjork, E. et al (1995): Noradrenaline-evoked pain in neuralgia. Pain; 63: 11:20.*
- Turner, J.A.; Deyo, R.A.; Loeser, J.D.; Von Korff, M.; Fordyce, W.E. (1994): The importance of placebo effects in pain treatment and research. JAM, 271: 1609-14.*
- Valley, M.A.; Bourke, D.L.; Foster, D.C. (1994): The use of superior hypogastric plexus block to predict pain relief outcome of presacral neurectomy (Abstract). Reg. Anesth., 19: S16.*
- Van Kleef, M.; Baendse, G.A.; Dingemans, W.A. et al (1995): Effects of producing a radiofrequency lesion adjacent to the dorsal root ganglion in patients with thoracic segmental pain. Clin. J. Pain; 11: 325-32.*
-
-

-
-
- Van Kleef, M.; Spaans, F.; Dingemans, W.; Barendse, G.A.M.; Floor, E.; Sluijter, M.E. (1993):* Effects and side effects of a percutaneous thermal lesion of the dorsal root ganglion in patients with cervical pain syndrome. *Pain*; 52: 49-53.
- Van Rheda, van der Kloot, E.; Krukker, J.; Lemmens, ILA.; Greep, J.M. (1986):* The high thoracic sympathetic nerve system its anatomic variability. *J. Surg. Res.*, 40: 112-19.
- Vecchio, P.C.; Adebajo, A.O.; Hazleman, B.L. (1993):* Suprascapular nerve block for persistent rotator cuff lesions. *J. Rheumatol.*, 20: 453-55.
- Verdugo, R.J.; Ochoa, J.L. (1993):* Use and misuse of conventional electrodiagnosis, quantitative sensory testing, thermography, and nerve blocks in the evaluation of painful neuropathic syndromes. *Muscle Nerve*, 16: 1056-62.
- Verdugo, R.J.; Ochoa, J.L. (1994):* Placebo response in chronic, causalgiform, neuropathic pain patients: study and review. *Pain Rev.*, 1: 33-46.
- Verdugo, R.J.; Ochoa, J.L. (1994):* Sympathetically maintained pain. I. Phentolamine block questions the concept. *Neurology*; 44: 1003-10.
- Vervest, A.C.M.; Stolker, R.J. (1991):* The treatment of cervical pain syndromes with radiofrequency procedures. *Pain Clin.*, 4: 103-12.
- Vulin, K.N.; Overgaauw, D.M.; Jessurum, G.; Tenvaarwerk, L.; Kropmans, T. et al (199):* The effects of spinal cord stimulation on quality of life in patients with therapeutically chronic refractory angina pectoris. *Neuromodulation*; 2: 33-40.
- Waldman, S.D. (1993):* Sphenopalatine ganglion block-80 years later. *Reg. Anesth.*, 18: 274-76.
- Waldman, S.D.; Wilson, W.L.; Kreps, R.D. (1991):* Superior hypogastric plexus block using a single needle and computed tomography guidance: description of a modified technique. *Reg. Anesth.*, 16: 286-87.
-
-

-
-
- Walker, S.M.; Cousins, M.J. (1998):* General principles of cancer pain management. Gildenberg P & Tasker R (eds) textbook of stereotactic and functional neurosurgery. McGraw Hill, N.Y., 1353-1379.
- Wall, P.D.; Melzack, R. (eds) (1994):* Textbook of pain, 31d ed., PP. 787-788, Edinburgh, Churchill, Livingstone, 1994.
- Wceks, L.; Batty, A.; Wolff; T.; Firtell, J.; Scheker, L. (1994):* Regional anaesthesia and subsequent long-term pain. J. Hand Sttrg. (Br), 19: 342-46.
- Wcyland, A.; Weyland, W.; Carduck, H.P.; Hidebrandt, J. (1990):* Which paravertebral approach is optimal for lumbar sympathetic blocks? Results of computer tomographic simulation studies. Pain, 55: 489.
- Weinstabl, C.; Porges, P.; Plainer, B.; Werba, A.; Spiss, C.K.; Seitz, H. (1993):* Coeliac plexus block with bupivacaine reduces intestinal dysfunction in neurosurgical ICU patients. Anaesthesia, 48: 162-64.
- Wetzel, F.T.; LaRocca, S.H.; Adinolli, M. (1992):* The treatment of chronic extremity pain in failed lumbar surgery. The role of lumbar sympathectomy. Spine, 17: 1462-68.
- Wilder, R.T. (1996):* Rellex sympathetic dystrophy in children and adolescents: differences from adults. In: Janig W, Statton-Hicks M eds. Reflex sympathetic dystrophy: a reappraisal (Progress in pain research and management. Vol. 6) Seattle: IASP Press, 67-77.
- Wong, C.S.; Liaw, W.J.; Tung, C.S.; Su, Y.F.; St, H.O. (1996):* Ketamine potentiates analgesic effect of morphine in postoperative epidural pain control. Regional Anesthesia, 21(6): 534-541.
- Wong, G.Y.; Brown, D.L. (1997):* Celiac plexus block for cancer pain. Teach Reg. Anesth. Pain Manage., 1: 18.
- Woolf, C.J. (1993):* Preemptive analgesia-treating postoperative pain by preventing the establishment of central sensitization. Anaesthesia, 77: 362-379.
-
-

Woolf, C.J. (1995): Somatic pain-pathogenesis and prevention. Br. J. Anaesth., 75: 169-76.

Woolf, Cols; Bennett, G.J.; Doherty, M.; Dubner, R. (1998): Towards a mechanism-based classification of pain. Pain, 77: 227-229.

Wulf, H.; Mater, C; Scheie, H.A. (1991): Bupivacaine toxictry after stellate ganglion block. In response. Anesth. Analg., 73: 505-506.

Wulf, H.; Mater, C.; Schele, H.A.; Wabbel, W. (1991): Plasma concentrations of bupivacainc after stellate ganglion blockade. Anesth. Analg., 72: 546-48.

Yaksh, T.L.; Malmberg, A.B. (1994): Central pharmacology of nociceptive transmission. In: Wall, P, Melzack R, Eds. Textbook of Pain, 2" ed. Edinburgh, UK: Churchill Livingstone, 165-200.