

### REFERENCES

- Abraham, E.C.; Perry, R.E.; and Sealling, M. (1983):  
Application of affinity chromatography for  
separation and quantification of glycosy-  
lated haemoglobin.  
J. lab. Clin. Med., 102 (2) : 187.
- Allen, D.W.; Schroeder, W.A.; and Balog, J. (1958):  
Observations on the chromatographic hetero-  
genecity of normal adult and fetal human  
haemoglobin. A study of the effect of crys-  
tallization and chromatography on the hetero-  
gencity and isoleucine content.  
J. Am. Chem. Soc., 80 : 1628.
- Alperin, J.B.; Dow, P.A.; Stout, L.C. (1979) :  
Comparison of haemoglobin A<sub>1c</sub> in human and  
baboon blood.  
Acta. Haematol., 61 : 334-338.
- Bains, T. (1985) :  
Serum fructosamine - the pH factor.  
Br. Med. J., 291 : 1050.
- Baker, J.R.; O'Lonner, J.P.; Metcalf, P.A.; Lawson ,  
M.R.; Johnson, R.N. (1983) :  
Clinical usefulness of estimation of serum  
fructosamine concentration as a screening  
test for diabetes mellitus.  
Br. Med. J., 287 : 863-867.

- Baker, J.R.; Johnson, R.N.; Scott, D.J. (1984) :  
Serum Fructosamine Concentrations in type  
II - non insulin dependent diabetes mellitus.  
Br. Med. J., 288 : 1484 - 1486.
- Baker, J.R.; Metcalf, A.; Johnson, N.; Newman, D.;  
Rietz, P. (1985) :  
Use of protein-based standards in automated  
colorimetric determinations of fructosamine  
in serum.  
Clin. Chem., 31 (9) : 1550 - 1554.
- Bookchin, R.M.; and Gallop, B.M. (1968) :  
Structure of HbA<sub>1</sub> : Nature of the N-terminal  
B- chain blocking group.  
Biochem. Biophys. Res. Commun., 32 : 86-93.
- Boucher, B.J.; Burrin, J.M.; Gould, B.J. (1983) :  
A collaborative study of measurement of gly-  
cosylated haemoglobin by several methods in  
seven laboratories in the United Kingdom.  
Diabetologia 24 : 265 - 71.
- Bridges, K.R.; Schmidt, G.J.; Jensen, M.; Cerami, A.;  
Bunn, H.F. (1975) :  
The acetylation of haemoglobin by aspirin in  
vitro and in vivo.  
J. Clin. Invest., 56 : 201-207.
- Brownlee, M.; Cerami, A. (1981) :  
The biochemistry of the complications of  
diabetes mellitus.  
Ann. Rev. Biochem., 50 : 385-432.

Bunn, H.F. (1979) :

Evaluation of glycosylated haemoglobin in  
diabetic patients.

Diabetes 30 : 613-617.

Bunn, H.F. (1981) :

Non-enzymatic glycosylation of protein :  
relevance to diabetes.

Am. J. Med., 70 : 325 - 330.

Bunn, H.F.; Haney, D.N.; Kamin, S.; Gabbay, K.H.; and

Gallop, P.M. (1976) :

The biosynthesis of human haemoglobin A<sub>1</sub> .

J. Clin. Invest., 57 : 1652 - 59.

Bunn, H.F.; Gabbay; K.H.; and Gallop, P.H. (1978) :

The glycosylation of haemoglobin. Relevance  
to diabetes mellitus.

Sci., 200 : 21.

Butler, W.J.; Ostrander, L.D.; Carman, W.J. and

Lamphiear, D.E. (1985) :

Mortality from coronary heart disease in the  
Tecumseh study : Long-term effect of diabetes  
mellitus, glucose tolerance and other risk  
factors.

Am. J. Epidemiol., 121 : 541-7.

Carlson, L.A.; and Bottiger, L.E. (1972) :

Ischemic heart-disease in relation to fasting  
values of plasma triglycerides and cholesterol

Lancet 1 : 865 - 868.

- Chou, P.P.; Kerkay, J.; Gupta, M.K. (1981) :  
Development of a laser nephelometric  
method for the quantitation of human glyco-  
haemoglobins.  
Anal. Lett. 14 : 1971 - 1087.
- Cole, R.A.; Soeldner, J.S.; Dunn, P.J.; Bunn H.F.(1978):  
A rapid method for the determination of gly-  
cosylated haemoglobins using high pressure  
liquid chromatography.  
Metabolism 27 : 289 - 301.
- Day, J.F.; Thornburg, R.W.; Thorpe, S.R.; Baynes, J.W.  
(1979) :  
Nonenzymatic glycosylation of rat albumin.  
Studies in vitro and in vivo.  
J. Biol. Chem., 254 : 9394 - 9400.
- Dolhofer, R.; Widland, O.H. (1980) :  
Increased glycosylation of serum albumin in  
diabetes mellitus.  
Diabetes 29 : 417 - 422.
- Dolhofer, R.; Wieland, O.H. (1981) :  
Improvement of the thiobarbituric acid assay  
for serum glycosylated protein determination.  
Clin. Chim. Acta 112 : 107-204.
- Doumass, B.T.; and Biggs, H.G. (1972) :  
Standard methods of clinical chemistry  
Vol. 7, Academic Press, New York.

- Doumas, B.T.; Watson, W.A.; and Biggs, H.G. (1971) :  
 Albumin standards and the measurement of  
 serum albumin with bromeresol green.  
 Clin. Chim. Acta 31 : 87-96.
- Dozy, A.M.; and Huisman, T.H.J. (1969) :  
 Studies of heterogeneity of haemoglobins.  
 XIV. Chromatography of normal and abnormal  
 human haemoglobin types on M.sephadex.  
 J. Chromatogr., 40 : 62.
- Ducimetiene, P.; Eschwege, E.; Papoz L.; Richard, J.L.;  
 Claude, J.R.; and Rosselin G. (1980) :  
 Relation of plasma insulin levels to the  
 incidence of myocardial infarction and  
 coronary heart disease mortality in a middle-  
 aged population .  
 Diabetologia 19 : 205-210.
- Dupree, E.A. and Meyer, M.B. (1980) :  
 Role of risk factors in complications of  
 diabetes mellitus.  
 Am. J. Epidemiol., 112 : 100-112.
- Dunn, P.J.; Cole, R.A.; Soeldner, J.S. et al.  
 (1979) :  
 Temporal relationship of glycosylated haemo-  
 globin concentration to glucose control in  
 diabetics.  
 Diabetologia 17 : 213-220.

- Ellis, E.N.; Brouhard, B.H.; Travis, L.B.; et al.  
(1980) :  
Urinary albumin and N-acetyl-beta-D-glucosaminidase in children with diabetes mellitus type I.  
Diabetes care 6 : 251-5.
- Epstein, F.M. (1967) :  
Hyperglycaemia - A risk factor in coronary heart disease :  
Circulation 26 : 609-619.
- Epstein, F.H. (1975) :  
Glucose intolerance and the risk of coronary heart disease.  
Arch. Mal. Coeur., 68 : 51 - 56.
- Ettinger, P.D.; Oldewurtel, H.A.; Weisse, A.B. et al.  
(1968) :  
Diminished glucose tolerance and immunoactive insulin response in patients with non ischemic heart disease.  
Circulation 38 : 559 - 67.
- Fitzgerald, M.D.; Cauchi, M.N. (1980) :  
Glycosylated haemoglobins in patients with a haemoglobinopathy.  
Clin. Chem., 26 : 360-361.
- Fluckiger, K.; Winterthaler (1976) :  
In vitro synthesis of haemoglobin A<sub>1c</sub>  
FEDS letters 71 : 356-360.

- Fuller, J.H.; Keen, H.; Jarrett, R.J. (1979) :  
Haemostatic variables in diabetes and their  
relationship to diabetic complications.  
Br. Med. J., 2 : 964 - 966.
- Fuller, J.H.; McCartney, P.; Jarrett, R.J.; Keen, H.;  
Rose, G.; Shipley, M.J.; and Hamilton ,  
R.J.S. (1979) :  
Hyperglycaemia and coronary heart disease.  
The whitehall study.  
N. Chron. Dis., 32 : 721 - 728.
- Fuller, J.H.; Shipley, M.J. Rose G. et al. (1983) :  
Mortality from coronary heart disease and  
stroke in relation to degree of glycemia.  
The Whitehall Study.  
Br. Med. J., 287 : 867-870.
- Gabbay, K.H.; Hasty K.; Breslow, J.L.; Ellison, R.C.;  
Hunn H.F.; Gallop, P.M. (1977) :  
Glycosylated haemoglobins and long-term blood  
glucose control in diabetes mellitus.  
N. Clin. Endocrinol. Metab., 44: 859-864.
- Gallop, P.M.; Fluckiger, R.; Hanneken, A.; Mininsohn,  
M.M.; Gappay, K.H. (1981) :  
Chemical quantitation of haemoglobin glyco-  
sylation : Fluorimetric detection of  
formaldehyde released upon periodate oxi-  
dation of glycoglobin.  
Anal. Biochem., 117 : 427 - 432.

- Goldstein, D.E.; Ide, C. and Wilson, R. (1983) :  
Should glycosylated haemoglobin be used for  
monitoring diabetic control.  
Practical cardiology 9 (3) : 187.
- Gordon, T.; Castelli W.P.; Hjortland M.C. et al.  
(1977) :  
High density lipoprotein as a protective  
factor against coronary heart disease.  
Am. J. Med., 62 : 707 - 714.
- Graf, R.J.; Halter, J.B.; and Porto, D. (1978) :  
Glycosylated Hb in normal subjects and  
subjects with maturity onset diabetes.  
Diabetes 27 : 834-839.
- Graf, R.J.; Halter, J.B.; Halan, E.; et al. (1979):  
Nerve conduction abnormalities in untreated  
maturity onset diabetes : relation to levels  
of fasting plasma glucose and glycosylated  
haemoglobin.  
Ann. Intern. Med., 90 : 248 - 303.
- Graf, H.; Stummvoll, H.K.; Schernthaner, G.; Muller,  
M.M. (1980) :  
Glycosylated haemoglobin in renal failur.  
Diabetologia 19 : 555-6.
- Henien, M.Y.A. (1987) :  
In "Serum fructosamine levels in obesity".  
Thesis for M. Sc. Internal Medicine, Benha  
Faculty of Medicine, Zagazig University.

- Higgins, P.J.; Garlick, P.L. and Bunn, H.F. (1982) :  
Glycosylated haemoglobin in human and  
animal red cells. Role of glucose permeabi-  
lity.  
Diabetes 31 : 743.
- Hindle, E.J. and Glenise, R. (1985) :  
Serum fructosamine as a measure of blood  
glucose control in type I diabetes mellitus  
Br. Med. J., 291 : 676.
- Hindle, E.J.; Rostron, M. and Gatt, J.A. (1985) :  
The estimation of serum fructosamine : an  
alternative measurement to glycated haemo-  
globin.  
Ann. Clin. Biochem., 22 : 84-89.
- Hoch, M. (1955) :  
The steady state, a test for electrophoresis  
homogeneity.  
Biochem. J., 46 : 199.
- Hodge, J.E. (1955) :  
The Amadori rearrangement.  
Adv. Carbohydrate Chem., 10 : 169-210.
- Holmquist, W.R.; and Schroeder, W.A. (1966) :  
A new N-terminal blocking group involving  
a schiff base in haemoglobin A<sub>1c</sub>.  
Biochemistry 5 : 2489-2503.
- Hommons, G.T.; Junger, K.; McDonald, J.M.; Ladenson,  
J.H. (1982) :

Evaluation of three minicolumn procedures  
for measuring haemoglobin A<sub>1</sub>.  
Clin. Chem., 28 : 1775-8.

Horton, B.F. and Huisman, T.H.J. (1965) :  
Studies on the heterogeneity of haemoglobin :  
VII. Minor haemoglobin components in haema-  
tological diseases.  
Br. J. Haematol., 11 : 296-304.

Huisman, T.H.T.; and Meyerling, C.A. (1960) :  
Studies on the heterogeneity of haemoglobin  
I. The heterogeneity of different haemoblo-  
bins types on carboxymethyl cellulose and on  
Ambertile IRC- 50 chromatography. Quantita-  
tive aspects.  
Clin. Chem. Acta 5 : 103.

Hurst, J.W. (1986) :  
Chest pain due to myocardial infarction.  
In " The heart" McGraw-Hill Inc., New York .,  
P 112, 113

Hurst, J.W.; King III, S.B.; Walter, P.F.; Fresinger,  
G. C.; and Edwards, J.E. (1982) :  
Atherosclerotic coronary heart disease :  
Angina pectoris, Myocardial infarction, and  
other manifestations of Myocardial Ischemia,  
in Hurst, J. Willis :  
The heart. McGraw-Hill Inc., New York.,:1009.

International Collaborative Group (1979) :  
Summary and joint discussion.  
J. Chron. Dis., 32 : 829 - 837.

- Jarrett, R.J. ; Keen, H.; McCartrey, M. (1978) :  
 Glucose tolerance and blood pressure in two  
 population samples : Their relation to  
 diabetes mellitus and hypertension.  
 Int. J. Epidemiol., 7 : 15-24.
- Javid, J.; Pettis, P.K.; Koenig, R.J. and Cerami, A :  
 (1978) :  
 Immunologic characterization and quantifi-  
 cation of HbA<sub>1c</sub>.  
 Br. J. Haematol., 38 : 329-337.
- Johnson, R.N.; Metcalf, P.A.; Baker, J.R. (1983) :  
 Fructosamine, a new approach to the estima-  
 tion of serum glycosyl protein. An index of  
 diabetic control.  
 Clin. Chim. Acta 127 : 87-95.
- Jones, R.T.; and Schroeder, W.A. (1963) :  
 Chromatography of human haemoglobin. Factors  
 influencing chromatography and differentia-  
 tion of similar haemoglobins.  
 J. Chromatogr., 10 : 421.
- Jones, M.D.; Koler, R.D.; Jones R.T. (1978) :  
 Microcolumn method for the determination of  
 haemoglobin in minor fractions A<sub>1a+b</sub> and A<sub>1c</sub>  
 Haemoglobin 2 : 53-58.
- Jovanovic, L.; Peterson, C.M. (1981) :  
 The clinical utility of glycosylated haemo-  
 globin.  
 Am. J. Med., 70 : 331 - 338.

- Jovansvic, I.; Peterson, C.M.; Saxena, B.B.; Dawood, M.Y.; Saudek, C.D. (1980) :  
Feasibility of maintaining normal glucose profiles in insulin-dependent pregnant diabetic women.  
Am. J. Med., 68 : 105-112.
- Keen, H. and Jarrett, R.J. (1973) :  
Macroangiopathy - its prevalence in asymptomatic diabetes.  
Arch. Metab. Dis. (Suppl) 2 : 3-9.
- Keen, H.; Rose, G.; Pyke, D.A. et al. (1965):  
Blood sugar and arterial disease.  
Lancet 2 : 505-508.
- Kennedy, A.L. and Baynes, J.W. (1984) :  
Nonenzymatic glycosylation and the chronic complications of diabetes : an overview  
diabetologia 26 : 93-98.
- Kingsley, L.A.; Dofl, B.H.; and Orchard, T.J. (1983):  
Long term metabolic control and diabetic retinopathy in an adolescent IDD cohort.  
Diabetes 32 (Suppl. I) : 9 A.
- Koenig, R.; Peterson, M.D. (1976) :  
HbA<sub>1c</sub> as indicator of the degree of glucose intolerance in diabetes.  
Diabetes 25 : 230.
- Koenig, R.; Peterson, C.M.; Jones, R.L.; Saudek, C.; Mehrman, M.; Grami, A. (1976) :

- Mayer, T.K. and Freedman, Z.R. (1983) :  
Protein glycosylation in diabetes mellitus,  
a review of laboratory measurement and  
their clinical utility.  
Clin. Chim. Acta 127 : 147-184.
- McDonald, M.; Shapiro, R.; Bleichman, M.; Soloway, J.  
and Bunn, M.F. (1978) :  
Glycosylated minor components of human  
adult Hb.  
J. Biol. Chem., 253 : 2327-2332.
- McFarland, K.F.; Catalano, E.W.; Day, J.F.; Thrope,  
S.R.; Baynes, J.W. (1979) :  
Non enzymatic glycosylation of serum proteins  
in diabetes mellitus.  
Diabetes 28 : 1011 - 1014.
- Menard L.; Dempsey, M.E.; Blankstein, L.A.; Aleyassine,  
I.; Wacks, M.; Soeldner, S. (1980) :  
Quantitative determination of glycosylated  
haemoglobin A<sub>1</sub> by agar gel electrophoresis.  
Clin Chem., 26 : 1598-1602.
- Mortensen, H.B.; Nielsen, L.; Soegaard, U.; Svendsen,  
I.; Nerup, J. (1983) :  
Comparison of six assays for glycosylated  
haemoglobin determination.  
Scand. J. Clin. Lab. Invest., 43 : 357-62.
- National diabetes Data group (1979) :  
Classification and diagnosis of diabetes  
mellitus and other categories of glucose  
intolerance.  
Diabetes 28 : 1039-1057.

haemoglobin and home capillary blood glucose level in diabetics.

Diabetologia 19 : 31-34.

Pan, W.H.; Cedres, L.B.; Liu, K.; Dayer, A.; James A.; Schoenberger; J.A.; Shekelle, R.B.; Slamler, R.; Smith, D.; Collette, P.; and Stamler, J. (1986):

Relationship of clinical diabetes and asymptomatic hyperglycaemia to risk of coronary heart disease mortality in men and women.

Am. J. Epidemiol., 123 :604 -516.

Panzer S.; Kronik, G.; Lechner, R.; Betteheim P.;

Neuman, G.; and Dudezak R. (1982) :

Glycosylated haemoglobins : an index of red cell survival.

Blood 59 : 1348.

Paulsen, E.P.; and Koury, M., (1976) :

HbA<sub>1c</sub> levels in insulin dependent and independent diabetes mellitus.

Diabetes 25 : 890.

Peacock, J. (1984) :

Glycosylated haemoglobin, measurement and clinical use.

J. Clin. Pathol., 37 : 841-851.

Peacock, I.; Tattersall, R.B. (1982) :

Methods of self monitoring of diabetic control

Clin. Endocrinol Metab., 11 : 485 - 501.

- Pyorala, K.; Savolainen, E.; Lehtovirta, E.; Punsar, S.; and Siltanen, P. (1979) :  
Glucose tolerance and coronary heart disease:  
Helsinki Policemen study .  
J. Chron. Dis., 32 : 927-745.
- Rahbar, S.; Blumenfeld, O.; Ranney, H.M. (1969) :  
Studies of an unusual haemoglobin in patients  
with diabetes mellitus.  
Biochem. Biophys. Res. Commun., 36: 838-43.
- Reaven, G.M., and Olefsky, J.M. (1978) :  
Role of insulin resistance in the pathogenesis  
of hyperglycaemia. In : Katzen M.M., Mahler,  
R.J. (eds).  
Diabetes, Obesity, and vascular disease.  
Halsted Press. Washington, P 229-266.
- Reunanen, A.; Pyorala, K.; Aromaa A. et al. (1974):  
Glucose tolerance and coronary heart disease  
in middle aged finnish men : Social Insurance  
Institution's coronary Heart disease study.  
J. Chron. Dis. , 32 : 747-758.
- Rhinesmith, H.S.; Schroeder, W.A. and Martin, N. (1958):  
The N-terminal sequence of the B-chain of  
normal adult human haemoglobin.  
J. Am. Chem. Soc., 80 : 3358.
- Roberts, B. and Baker, R. (1986) :  
Serum fractosamine : A screening test for  
diabetes in pregnancy.  
Am. J. Obstet. Gynaecol., 154 : 1027-1030.

- Roberts, A.B.; Court, D.J.; Henly, P.; Baker, J.R.;  
James, A.G. and Ranayne, I.D. (1983) :  
Fructosamine in diabetic pregnancy.  
Lancet ii , 998-1000.
- Rodkey, F.L. (1965) :  
Direct spectrophotometric determination of  
albumin in human serum.  
Clin chem., 11 : 478-87. Apr. 65.
- Rosenthal, P.K.; Vazquez, D.A.; Seckinger, D.L. (1981):  
Determination of glycohaemoglobins by a rapid  
batch separation.  
Am. J. Clin. Pathol., 75 : 45-49.
- Ruderman, N. and Handenschild, C. (1984) :  
Diabetes as an atherogenic factor progress in  
cardiovascular diseases Vol XXVI, No. 5.  
(March/April) .
- Ryall, R.G.; Graham, J.J. (1978) :  
Routine measurement of glycosylated haemoglo-  
bin.  
Lancet ii : 740.
- Sandor, G. (1966) :  
Serum proteins in health and disease.  
London : Shapman and Hall.
- San-Gil, F.; Schier, G.M.; Moses, R.G.; and Gan, T.I.  
(1985) :  
Improved estimation of fructosamine, as a  
measure of glycated serum protein, with the  
technican RA 1000 analyses.  
Clin. Chem., 31 (12): 2005-2006.

- Schellekens, A.P.M.; Sanders, G.T.B.; Thornton, W.;  
Van Groenestein T. (1981) :  
Sources of variation in the column-chromatographic determination of glycohaemoglobin (HbA<sub>1</sub>).  
Clin. Chem., 27 : 94-99.
- Schmidt, M.I.; Duncan, B.B.; Gitelman, H.J.; et al. (1983) :  
High levels of urinary N-acetyl-beta-D-glucosaminidase (NAG) in children with diabetes : frequency and correlation with glucose control.  
Diabetes 32 (Suppl 1 ) : 30 A.
- Schneck, A.G.; and Schroeder, W.G. (1961) :  
The relation between the minor compartments of whole normal human adult Hb as isolated by chromatography and starch block electrophoresis.  
J. Am. Chem. Soc. 83 : 1472.
- Schroeder, W.A.; Cua, J.J.; Mastuda, G. and Renninger, W.D. (1963) :  
Haemoglobin F and acetyl-containing haemoglobine  
Biochem. Biophys. Acta 63 : 532.
- Schwartz, H.C.; King, K.C.; Schwarty, W.L. (1976) :  
Effect of pregnancy on HbA<sub>1c</sub> in normal, gestational diabetes and diabetic women.  
Diabetes 25 : 1118.

Yue, D.K.; Morris, K.; McLennan, S.; Turtle, J.R. (1980):  
 Glycosylation of plasma proteins and its  
 relation to glycosylated haemoglobin in  
 diabetes.  
 Diabetes 29 : 296-300.

Yue, D.K.; McLennan, Church, D.B.; Turtle, J.R. (1982):  
 The measurement of glycosylated haemoglobin in  
 man and animal by aminophenylboronic acid  
 affinity chromatography.  
 Diabetes. 31 : 701-5.