

REFERENCES

- Abbott W.M., Malomey R.D., McCabe C.C., Lee, C.E.
and Wirthlim L.S. (1982): Atrial
embolism a 44 years perspective.
Am. J. Surg. 1243:460-64.
- Aberg H. (1968): Atrial fibrillation. Acta Med.
Scand; 184:425-31.
- Abildskov J.A., Kay Millar, and Mary J.O. Burgees
(1971): Atrial fibrillation.
Am. J. of Cardiol. Vol.28, Sept., p.
263-267.
- Arthur Selzer (1982): Atrial fibrillation revisted
The New Engl. J. of Med. Vol.306 No.17,
Apr. 29, p. 1044-1045.
- Askey J.M. (1962): Embolism and atrial fibrillation.
Am. J. Cardiol. 9:491-95.
- Baily G.W.H., Brainff B.A., Hancock E.W., and
Cohon K.E. (1968): Relation of left
atrial pathology to atrial fibrillation
in mitral valve disease. Ann. Intern.
Med. 69:13-20.
- Barker P.S., Bohning A.L., and Wilson F.N.(1982):
Auricular fibrillation in Grave's disease.
Am. Heart J. 8:121-27.
- Bar-Sela' S., Ehrenfeld M., and Eliakim M.(1981):
Atrial embolism in thyrotoxicosis with
atrial fibrillation. Arch. Intern. Med.
141:1191-92.

- Bell, R., Barber P.V., Bary C.L., and Beton D.C.
(1977): Incidence of thyroid disease
in cases of hypertrophic cardiomyopathy.
Br. Heart J. 40:1306.
- Blinks J.R., Koch-Wesert I. (1961): Analysis of
the effects of changes in rate and rhythm
upon myocardial contractility. J.Pharmacol.
Exp. Ther. 134:373-389.
- Bootsma B.K., Hoelen A.J., and Strackee J. '(1971):
Analysis of P-R intervals in patients
with atrial fibrillation at rest and
during exercise. Circulation; 41:783-794.
- Burkhardt D., Staub J.J., Kranmmlim M., and Reader
E. (1978): The systolic time intervals
in thyroid dysfunction. Am. Heart J.95:187.
- Caldwell G., Kellett H.A., Gow S.M., Beckett
G.J., Sweeting V.M., Seth J., and Toft
A.D. (1985): A new strategy for thyroid
function testing. The Lancet, Sat. 18
May: 1117 - 1119.
- Chopra I.J. (1972): A radioimmunoassay for measurement
of thyroxine in unextracted serum. J.Clin.
Endocrinol and Metabol. 34:938.
- Chopra, I.J., Jerome M.H., William M.P., and John
T.N. (1983): Thyroid function in non-
thyroidal illnesses. Ann. Intern. Med. 98:
949-957.

- Cobler, J.L., Mark E., and Philip G. (1984):
Thyrotoxicosis in institutionalized elderly
patients with atrial fibrillation.
Arch. Intern. Med. Vol. 144, p. 1758-1760.
- Cotoi S., Carasea E., Georgescu C., and Lazer P.
(1978): Two variations of atrial fibrillation studied by monophasic action potential recording. Jpn. Heart J. Jul. 19(4). p. 479-84 (AB).
- Croxson M.S. and Ibbertson M.H. (1975):
Serum digoxine in patients with thyroid disease. Br. Med. J. 3:566.
- David J., Sahn, Anthary DeMaria, and Joseph Kisslo
(1978): M-mode echocardiography.
Circulation; 58:1072.
- De Groot W.J., Leonard J.J. and Nedvid J. (1970)
Hyperthyroidism as a high cardiac output state. Am. Heart J. 79:265.
- De Groot W.J., Gunn, A. and Felt J. (1975):
Cardiovascular changes in hyperthyroidism
Br. Med. J. 3:460.
- De Silva R.A., and Lown B. (1982): Cardioversion
for atrial fibrillation indications and complications. In: Kulbrtus H.E., Olsson S.B., Schlepper M., eds. Atrial fibrillation. Molndal, Sweden (AB) Massle, 231-39.

- Dewar H.A & Weightman D. (1983): A study of embolism in mitral valve disease and atrial fibrillation. Br. Heart J., 49:133-40.
- Dodge, H.T., Kirkham J.T., and King C.V. (1957): Ventricular dynamics and atrial fibrillation. Circulation; 15:335-347.
- Dunn M., Alexander J., De Silva R. and Hildner F. (1986): Antithrombotic therapy in atrial fibrillation. Chest; 89-2-(Suppl.), 685-74 (Review).
- Edmonds R.E., Greenspan K., and Fisch C. (1970). The role of inotropic variation in ventricular function during atrial fibrillation. J. Clin. Invest.; 49:738-746.
- Edward Lown (1967): Electrical reversion of cardiac arrhythmias. Br. Heart J.: 29:469.
- Elkhardt M.J., Harford T.C., and Kaiber C.T. (1981): Health hazards associated with alcohol consumption. J.A.M.A., 246:648-666.
- Ellis L.B. & Harken D.E. (1961): Atrial embolism in relation to mitral valvuloplasty. Am. Heart J. 62:611.
- Eric H.S., Robert P., Bernard D.K., Juan, G. Richard R.S., and Isaac W. (1982): Clinical use of oral verapamil in chronic and paroxysmal atrial fibrillation. Chest, 81: 3 p. 308-11.

- Eugene, C.R., Constance S., Brian C. (1985):
Alcohol-related acute atrial fibrillation . A case-control study:
a review of 40 patients. Arch. Intern.
Med. Vol. 145; 830-833.
- Ewy G.A., Ulfers L., Hager W.D., Rosenfeld
A.R., Roske W.R., and Goldman S.
(1980): Response of atrial fibrillation
to therapy. J. Electrocardiol. 13:119.
- Feely J. and Peden N. (1984): Use of adrenoceptor
blocking drugs in hyperthyroidism.
Drugs ; 27(5) P. 425-46. Review (AB).
- Felt V., Genkova V, and Nedvid (1982): Changes of
 T_3 and T_4 in patients with thyro-
cardiac disease and their relations to
systolic time intervals. Endocrinologie ;
80:325.
- Forfang K., Simonsen S., Andersen A., and Efskind
L. (1977): Atrial septal defect of
secundum type in the middle aged. Am.
Heart J., 94:44.
- Forfar J.C., Anthony D. Toft, and Leonard J.J.
(1982a): Thyrotoxic atrial fibrillation:
an underdiagnosed condition. Br. Med. J.
285:909.
- Forfar J.C. and Caldwell G.C. (1985): Hyperthyroid
heart disease. Clin. Endocrinol. and
Metabol.; 14(2) p. 491-508.

- Forfar J.C., Millar H.C., and Toft A.D.(1979):
Occult thyrotoxicosis: a correctable cause
of "idiopathic" atrial fibrillation.
Am.J. Cardiol. 44:9-12.
- Forfar J.C., Stewart A., Sewers J., and Toft A.D.
(1982 b): Cardiovascular responses in
hyperthyroidism before and during B-
adrenoceptor blockade: Evidence against
adrenergic hypersensitivity. Clin. Endo-
crinol. 16:441.
- Friedman M.J., Robert D., Okada, Gordon A.,
Ewy, and Dorothy J.H. (1982): Left ventri-
cular systolic and diastolic function
in hyperthyroidism. Am. Heart J.: 104:1303.
- Fuster V, Gersh B.J., Guiliani E.R., Tajik A.T.,
Brandenberg R.O., and Frye R.L. (1981):
The natural history of idiopathic dilated
cardiomyopathy. Am. J. Cardiol. 47:525-31.
- Goldberger E and Wheat M.W. (1982): Supraventricular
tachyarrhythmias. Treatment of cardiac
emergencies. 3rd ed. London, The C.V.Mosby
Company, 80-107.
- Greenfield J.C., Harley A. and Thompson H.K.(1968):
Pressure-flow studies in man during atrial
fibrillation. J.Clin. Invest. 47:2411-2421.
- Greenwood W.F., Albridge M.E., and Mchalvey A.D.(1963):
Effects of mitral commissurotomy on duration
of life, functional capacity, hemoptysis
and systemic embolism. Am.J. Cardiol.11:348.

- Gross M.M., Lewis E., and Hastey J. (1974): Acute alcohol withdrawal syndrome, in Kissin B, Beveleiter H. (eds): The Biology of Alcoholism. New York, Random House; Vol. 3 p. 191.
- Grossman W., Robin, Jonson L.W., Brooks M.P., and Dexter L. (1971): The enhanced myocardial contractility in thyrotoxicosis. Ann. Intern. Med., 74 : 869.
- Hecht H.M., Osher W.J. and Samuels A. (1951): Cardiovascular adjustments in subjects with organic heart disease before and after conversion of atrial fibrillation to normal sinus rhythm (AB). J. Clin. Invest., 30:647.
- Hinton R.C., Kistler J.P., Fallon J.T., Friedlich A.L., and Fisher C.M. (1977): Influence of etiology of atrial fibrillation on incidence of systemic embolism. Am. J. Cardiol, 40:509-13.
- Hoffenberg R. (1981): Hyperthyroidism, hypothyroidism and thyroid function testing. Intern. Med. Vol. 1, No.6: 256-266.
- Hurst J.W., Paulk E.A., Proctor H.D., and Schlant R.C. (1964): Management of patients of with atrial fibrillation. Am. J. Med. 837:728-41.

- Jerzy G., Richard B., and Singer B. (1981): Mortality in an insured population with atrial fibrillation. J.A.M.A. Vol. 245, No.13. 1540-1544.
- Joel M., Leonard N.H. Mark E.J., and John A.K.(1979): Relationship of atrial fibrillatory wave amplitude to left atrial size and etiology of heart disease. Am. Heart J. 97(2): 184-186.
- John Mancini G.B. and Ary and Goldberger (1982): Cardioversion of atrial fibrillation: consideration of embolization, anticoagulation, prophylactic pacemaker and long-term success. Am. Heart J., 617-21.
- Kannel W.B., Abbot R.D., Savage D.D. and NcMamara P.M. (1982): Epidemic features of chronic atrial fibrillation: The Framingham Study. N. Engl. J.M.; 306: 1018-22.
- Kedrov A.A. and Somova L.V. (1984): Clinical considerations in the treatment of atrial fibrillation (in thyrotoxicosis patients) Ter. Arkh. 56(9) p.62-6 (AB).
- Killip T. (1963): Synchronized D.C. precordial shock for arrhythmias: Safe new technique to establish normal rhythm may be sutilized on an elective or emergency basis. J.A.M.A. 186:1.

- Larsen P.R. (1972): Direct immunoassay of triiodo-thyronine in human serum. J. Clin. Invest. 5L: 1939.
- Lendrum B., Feinberg H., and Boyd E. (1960): Rhythm effects on contractility of the beating isovolumic left ventricle. Am. J. Physiol.;199:1115-1120.
- Lewis B.S., Ehrenfeld E.N., and Lewis N. (1979): Echocardiographic L.V. function in thyrotoxicosis. Am. Heart J. 97:460.
- Lown B (1967): Electrical reversion of cardiac arrhythmias. Br. Heart J. 29:469.
- Lown B., Perborth M.G., Kaidbey S., Abe T., and Harken D.E. (1963): Cardioversion of atrial fibrillation N. Engl. J. Med;269:325.
- Mauricio B.R., Pablo A.C., M.Susana H., Gerado J.N., julio P., Raul J.L., Julio O, and Marcelo V. (1976): Clinical efficacy of amiodarone as an antiarrhythmic agent. Am. J. Cardiol. 38:934-943.
- Merillon J.P., Phassa P.H., Chastre, J., and Wolf A. (1981): Left ventricular function and hyperthyroidism. Br. Heart J.,46:137.
- Merillon J,P., Motte J., and Pronsky D. (1979): Cardiovascular response in hyperthyroidism. Br. Med. J. 1:1502.

- Michael V.C., Ira C., Schulman, and Angelo S.(1981):
Effects of thyroid hormone on left
ventricular functions in patients treated
for thyrotoxicosis. Am. J. Cardiol. 48:33.
- Millar K., Eich R.H., and Burgess M.J. (1969):
Enhancement of cardiac output by irregular
rhythm (AB). Circulation; 40(3):146.
- Moe G.K. (1962): On the multiple wavelet hypothesis
of atrial fibrillation. Arch. Intern.
Pharmacodyn. 140:183-188.
- Moe G.K., and Abildskov J.A. (1955): Atrial fibrilla-
tion as a self-sustaining arrhythmia
independent of focal discharge. Am. Heart
J. 48: 59-70.
- Moe G.K., Rheinboldt W.C., and Abildskov J.A.(1964):
A computer model of atrial fibrillation.
Am. Heart J., 67:200-220.
- Morris J.J., Entman M.L., and Thompson H.K. (1963):
Cardiac output in atrial fibrillation and
sinus rhythm (AB). Circulation; 28:772.
- Morris J.J., Peter R.H., and McIntosh M.D. '(1966):
Electrical cardioversion of atrial fibrilla-
tion. Immediate and long term results and
selection of patients. Ann. Intern. Med.
65:216.

- Murphy B.E.P., and Pattee C.J. (1984): Determination of thyroxine utilizing the property of protein. binding. J. Clin. Endocrinol. 24:187.
- Nadeau R.A., Roberge F.A., and Billette J. (1970): Role of the sinus node impulses and atrial fibrillation. Circulation (Res.); 27:129-138.
- Nakazawa M.K. and Sakuragai K. (1980): Timing of cardioversion application in thyrotoxic A.F. Circulation: 62:296.
- Nakazawa H.K., Sakuragai K., Hamada N., Momotani N. and Ito K. (1982): Management of atrial fibrillation in the post-thyrotoxic state. Am. J. Med. 72(6):903-6 (AB).
- Nyström E., Peterson K., Lindstedt G., and Lundberg P. (1986): Screening for thyroid disease in women $\geq$ 50 years of age seeking hospital care: Influence of common non-thyroidal illness on serum free thyroxine as determined by analog radioimmunoassay. Clin. Chemistry. 32(4):603-608.
- Ollis C.A., Monroe D.S., and Tomlinson S. (1982): Characters of thyrotrophin-stimulated cyclic AMP production in cultured human thyroid cells. From Thyrotoxic tissue. J. Endocrinol.; 85:237.

- Olsson S.B. (1981): Nature of cardiac arrhythmias and electrolyte disturbances. Role of potassium in atrial fibrillation. Acta. Med. Scand. (Suppl.). 647:33-7.
- Peterson L.M., Fisher R.D., Reis R.L., and Morrow A.G. (1969): Cardiac operations in patients with left atrial thrombus. Ann. Thorac. Surg. 8:402.
- Philip O.E., Chia F.W., and Catalino De La Cruz (1978): Arrhythmias and the "Holiday Heart" alcohol associated cardiac rhythm disorders. Am. Heart J., 95(5):555-562.
- Popovici D and Nihai N.U. (1980): Abnormalities of oxidative phosphorylation due to excess or deficiency of thyroid hormones. Endocrinologie; 18:143.
- Popp R.L. (1972): Echocardiographic assessment of cardiac disease. Circulation; 54:538.
- Prinzmetal M., Rakita L, and Borduas J. (1955): The nature of spontaneous auricular fibrillation in man. With comments on the action of antiarrhythmic drugs. J.A.M.A. 157:1175-1182.
- Proskey A.T., Saksena F., and Uther J. (1977): Myocardial infarction associated with thyrotoxicosis. Chest; 72:109.

- Rich E.C., Siebold C., and Brian C. (1985):
Alcohol related acute AF. Arch. Intern.
Med. Vol. 145:830-833.
- Sano T., Suzuki F., and Sato S. (1967): Sinus node
impulses and atrial fibrillation.
Circulation (Res.); 21:507:513.
- Scherf D. (1955): Atrial arrhythmias, N. Engl. J.
Med.; 252: 928-933.
- Sobel B.E., and Braunwald E. (1977): Cardiac dys-
rhythmias. In: Thorn G.W., Adams R.D.,
Braunwald E., Isselbacher K.S., Petersdorf
A.G. eds. Harrison's Principles of
internal medicine. 8th ed. New York.²
Mc Graw-Hill, 11878-206.
- Sultan Ahmad, Gilbert E., Carl. J. and Philip O.
(1972): Systolic time intervals as a
measure of the left ventricular myocardium
in man. Circulation; 26:559.
- St. John, Sutton M.G., Tajik A.T., McGoon D.C. (1981):
Atrial septal defect in patients aged
60 years or older: operative results and
long term postoperative follow-up. Cir-
culation; 64:402.
- Stephen W.M., Gray E., Branch N.S., and William
J.M. (1980): Efficacy of intravenous
procainamide infusion in converting
atrial fibrillation to sinus rhythm in
relation to left atrial size.
Br. Heart J., 44:589-95.

- Storstein O., and Tventon M. (1955): Quinidine treatment of established auricular fibrillation. Acta. Med. Scand.; 153: 57-66.
- Sung R.J., Myerburg R.T., and Castallanos A.(1978): Electrophysiologic demonstration of concealed conduction in the human atrium Circulation; 58:940.
- Symons C.(1979): Thyroid heart disease. Br. Heart J. 41:257.
- Tajiri J., Hamaski S., Shimada T., Morita M., Migashi K., and Ohishi S. (1986): Masked thyroid dysfunction among elderly patients with atrial fibrillation. Jpn. Heart J., Mar. 27(2) p. 183-90 (AB).
- Thomas F.B., Mazzaferri E.L., and Skillman T.G. (1970): Apathic thyrotoxicosis: A distinctive and laboratory entity . Ann. Intern. Med.; 72:679-685.
- Verhoeven R.P., Visser T.J., Docter R., and Henneman C. (1977): Plasma T_4 , T_3 and $r T_3$ during B-adrenergic blockade in hyhperthyroidism. J.Clin. Endocrinol. and Metabol.; 44:1002.
- Ward D.E., Camm A.J., and Spurrell A.A.J. (1980): Clinical antiarrhythmic effects of amiodarone in patients with resistant paroxysmal tachycardias. Br. Heart J., 44:91-5.

- Waxman H.L., Myerburg R.J. Appel R., and Sung R. (1981): Verapamil for control of ventricular rate in paroxysmal supra-ventricular tachycardia and atrial fibrillation or flutter. Ann. Intern. Med.; 94:1-6.
- Wayne E.J. (1983): Wayne's clinical diagnostic index for thyrotoxicosis. Baily and Love's Short Practice of Surgery. ELBS Edition Reprinted.
- Weber M., Weissel P., and Maintiz M. (1984): Antiarrhythmic effect and side effects of amiodarone. Wine-klin. Wochenscher Sep.; 28. 96(18). p. 685-96 (AB).
- Wilkin T.J. (1983): Hyperthyroidism and the heart Br. Heart J., 286:1459.
- Williams R.S., Lefowitz R.J., and Williams L.(1979): Thyroid hormone regulation of alpha adrenergic receptor. Cardiovasc. Pharmacol. 1:181.
- Witherbee D.G., Brown M.G., and Holzman D.(1952): Ventricular rate response following exercise during auricular fibrillation and after conversion to normal sinus rhythm. Am. J. Med. Sci., 223:667-670.

- Wolf P.A., Dawber P.R., Thomas H.E., and Kannel W.B. (1978): Epidemiologic assessment of chronic atrial fibrillation and risk of stroke: The Framingham Study. Neurology; 28:973.