

REFERENCES

REFERENCES

- Abell M.R., M.D. and Littler E.R.M.D.(1975):** Benign metastasizing uterine leiomyoma, multiple lymph nodal metastases. *Cancer*, 3G: 2206.
- Adachi N., Yamaguchi, Miyakiy, et al., (1990):**Parathyroid hormone related protein is a possible autocrine growth inhibitor for lymphocytes. *Biochem. Biophys. Res. Commun.*166: 1088.
- Alnemri, E.S., Robertson V.M., Fernandes T.F., et al.,(1992):** Overexpressed full-length human bcl-2 extends the survival of Baculovirus infected SFq insect cells. *Proc. Natl Acad. Sci USA.*, 89:7295.
- Akkersdijk G.J., Flu P.K., Giard R.W.,et al. (1990):** Malignant leiomyomatosis peritonealis disseminata. *Am.J. obst.Gynecol*,163:591
- Anderson J.R.(ed) (1980):** Benign tumors of the uterine corpus, *Muir's text book of pathology 11th edn.* Edward Arnald, london, chap.7, pp. 942.
- Andersen J., Grine E.,Carla L.Y., et al.,(1993):** Expression of Connexin 43 in human myometrium and leiomyoma. *Am.J.obst.Gynecol.*, 169:1266.
- Andersen J., Barbieri R.L.(1995):** Abnormal gene expression in uterine leiomyomas J.Soc. Gynecol. Invest., 2:663.
- Andersen J.(1996):** Growth factors and cytokines in uterine leiomyomas. *Semin.Reprod. Endocrinol*,14:269.
- Arrends M.J., Morris R.G., and wyllie A.H.(1990):** Apoptosis is the role of the endonuclease.*Am.J.path*, 136: 593.
- Aterman K., Fraser G.M., and lea R.H. (1977):** disseminated peritoneal leiomyomatosis .*Virchow Arch.(path.Anat.)*, 374:13.
- Banooni F., Labes J. and Goodman P.A.(1971):** Uterine leiomyoma Containing metastatic breast Carcinoma.*Am.J.obst.Gynecol.*, 111:427.

REFERENCES

- Barnes H.M., and Richardson P.J.(1973):** Benign metastasizing fibroid leiomyoma. *J. obst. Gynecol. Br commonw*, 8: 369.
- Bissonnette R.P., Echeverri F., Mohoubi A., et al.(1992):** Apoptotic cell death induced by C-myc is inhibited by bcl-2. *Nature*, 359:552.
- Blackwood E.M. and Eisenman R.N.(1991):** Max: a helix-loop helix zipper protein that forms a sequence-specific DNA-binding Complex with myc. *Science*, 251:1211.
- Bodrug S.E., Aime-Sempec, sato T., et al.,(1995):** Biochemical and functional comparison of Mcl-1 and Bcl-2 proteins :evidence for a novel mechanism of regulating bcl-2 family protein function. *Cell death differ.*, 2:173.
- Boise L.H., Gonzalez-Garcia M., Postema C.E., et al., (1993):** Bcl-x, a bcl-2 related gene that functions as dominant regulator of apoptotic cell death. *Cell*, 74:597.
- Bortner C.D. (1995):** The role of DNA fragmentation in apoptosis. *Trends Cell Biol.*, 5:21.
- Bossert N.L., Nelson K.G., Ross K.A., et al.,(1990):** Epidermal growth factor binding and receptor distribution in the mouse reproductive tract during development. *Dev.Biol.*, 142:75.
- Bradford M.(1976):** A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.*, 72: 248.
- Brandon D.D., Bethea Cl., strawn E.Y., et al.,(1993):** Progesterone receptor messenger ribonucleic acid protein are overexpressed in human uterine leiomyomas. *Am.J. obst. Gynecol.*, 169: 78.
- Buchi K.A. and keller P.J (1980):** Estrogen receptors in normal and myomatous human uteri. *Gynecol. Obst.invest.*, 11:59.
- Buchi K.A.,and Keller P.J .(1983):** Cytoplasmic progestin receptors in myomal and myometrial tissues. *Acta. Obst. Gynecol. scand.*, 62:487.
-

REFERENCES

- Buttram V.C., and Reiter R.C.(1981):** uterine leiomyomata, eitiology, symptomatology and management. *Fert.steril.*, 36: 433.
- Buttram V.C. (1986):** uterine leiomyomata, eitiology, symptomatology, and mangement. *Prog.clin. Bioch.Res.*, 225: 275.
- Campos L., Rovault J.p., sabido O.,et al., (1993):** High expression of bcl-2protein in acute myeloid leukemia cells is associated with poor response to chemotherapy. *Blood*, 81: 3091.
- Casey M.L., Mibe M., Erk A., et al.,(1992):** Transforming growth factor - β stimulation of parathyroid hormone-related protein expression in human uterine cells in culture: m RNA level and protein secretron. *J.clin.Endocrinol. Metab.*, 74:950.
- Cheng E.H-Y., levine B., Biose L.H., et al., (1996):** Bax-independent inhibition of apoptosis by bclx_i. *Nature*, 379: 554.
- Chittenden T., Harrington E.A., O'Connor R., et al., (1995):** Induction of a poposis by the bcl-2 homologue Bak. *Nature*, 374:733.
- Chresta C., Masters J.,and Hickman J.(1996):** Hypersensitivity of human testicular tumors to etopside-induced apoptosis is associated with functional p53 and a high bax: bcl-2 ration. *Cancer Res.*, 56:1834.
- Clarke C.L.and sutherland R.L.(1990):** Progestin regulation of cellular proliferation. *Endoc. Rev*, 11: 266.
- Clarke A.R., Purdie C.A., Harrison D.J., et al.,(1993):**Thymocyte apoptosis lnuded by p53- dependent and independent pathways. *Nature*, 362:849.
- Clement B.P.(1988):** I.V. leiomyoma of the uterus. *path.annu.*, 23: 153.
- Cohen J.J., Duke R.C., fadok V.A., et al., (1992):**Apoptosis and prograded cell death in immunity. *Annu.Rev.Immunol.*, 10:267.
- Cohen J.J.(1993):** The cell biology of acute myocardial ischemia. *Hosp. proc.* 28;35.
-

REFERENCES

- Colombel M., Symmans F., Gil S., et al.,(1993):** Detection of the apoptosis-suppressing oncoprotein bcl-2 in hormone-refractory human prostate cancers. *Am.J.path.*, 143:390.
- Cramer S.F.,Meyer J.S., Kaner J.F., et al.,(1980):** Metastasizing leiomyomata of the uterus-S phase fraction, estrogen receptor, and ultrastructure. *Cancer*, 45: 932.
- Danila-Muster A., Sohleanu V.,and Simionescu L.(1963):** Le role pathologique del' hormone somatotrope dans la fibromatose uterine *Rev. fr. Endocrinol. Clin.*, 4:243.
- Derynk R.(1988):** Transforming growth factor- α .*Cell*, 54:593.
- Ducke R.C., Chervenak R., and Cohen J.J. (1983):** Endogenous endonuclease- induced DNA fragmentation.An early event in cell mediated cytolysis. *Proc. Natl.Acad. Sci.USA.*,80: 16361.
- Duke R.C. and Cohen J.J. (1986):** IL- 2 addiction withdrawal of growth factor activates a suicide programme in dependent T cells. *Lymphokine Res.*, 5:289.
- Duval E., and Wyllie A.H. (1986):** Death and the cell. *Immunol. Today.*, 7:115.
- Ellis R.E., Yuan J., and Horwitz H.R. (1991):** Mechanism and function of cell death. *Annu. Rev.cell Biol.*, 7:663.
- EvanG.I. Wyllie A., Gibert C.S., et al.,(1992):** Induction of apoptosis in fibroblasts by C-myc protein. *Cell.*, 63:119.
- Evan G.I. and littlewood T.D.(1998):** Amatter of life and cell death. *Science*, 281: 1317.
- Faber M., Gnrad S., Heinrichs W.L., et al., (1972):** Estradiol binding by frbrord tumors and normal myometrium. *obst. Gynecol.*, 40:479.
- Fady I., Sharara M.D., Lynnette K., et al.,(1995):** Growth hormone receptor messenger ribonucleic acid expression in leiomyoma and surrounding myometrium. *Am.J.obst Gynecol.*, 172:814.

REFERENCES

- Farrow S.N., White J.H.M., Martinou I., et al., (1995):** Cloning of a bcl-2 homologue by interaction with adenovirus EIB 19kD *Nature*, 374:731.
- Ferguson J.E., Gorman J.V., Bruns D.E., et al., (1992):** Abundant expression of parathyroid hormone- related protein in human amnion and in association with labour. *Proc. Natl. Acad. sci. USA.*, 89: 8384.
- Friedman A-J., Harrison, Atlas D., et al., (1989):** A randomized, placebo-controlled, double-blind study evaluating the efficacy of leuprolide acetate depot in the treatment of uterine leiomyoma. *Fert. Steril.*, 51:251.
- Fujii S., Okamura H., Nakashima N., et al., (1980):** Leiomyomatosis peritonealis disseminata. *obst. Gynecol.*, 55:79s.
- Garcia I., Martinou I., Tsujimoto Y., et al., (1992):** Prevention of programmed cell death of sympathetic neurons by the bcl-2 protooncogene. *Science*, 258:302.
- Garfield R.E., and Hayash R.H. (1980):** Presence of gap- Junctions in the myometrium of women during various stages of menstruation *Am. J. obst. Gynecol.*, 5:569.
- Garfield R.E., kannan M.S. and Daniel E.E. (1980):** Gap-Junction formation in myometrium: control by estrogens, progesterone and prostaglandins. *Am J. physiol.*, 238:C81.
- Garfield R.E. (1985):** Cell-to cell communication in smooth muscle. In :Grover Ak, Daniel E.E., eds. Calcium and Contractility-smooth muscle. *Clifton, Newjersey:Humana., P.P. 143.*
- Giudice L.C. (1992):** Insuline-like growth factors and ovarian follicular development. *Endocrine Rev.*, 13: 641.
- Golden T. and stout A.R. (1941):** Smooth muscle tumors of the gastrointestinal tract and retroperitoneal tissues. *Surg. Gynecol obst.*, 73: 784.
- Golden R.D., Hunt N.C., Clark J., et al. (1993):** Apoptotic bodies in a murine model of alcoholic liver disease reversibility of ethanol induced changes. *J. pathol.*, 171:73.

REFERENCES

- Goldzieher J.W., Moqueo M., Ricaud L., et al.,(1966):** induction of degenerative changes in uterine myomas by high-dose progestin therapy. *Am.J. obst. Gynecol.*, 96: 1078.
- Golstein p. ojeius D.M., and young J.D.(1997):** Controlling cell death. *Science*,275:1081.
- Goustin A.S., Leof E.B., Shipley G.D., et al., (1986):** Growth factors and Cancer. *Cancer Res.*, 46: 1015.
- Grella L., Arnold T.E., kuilekual K.H. et al.,(1994):**Intravenous leiomyomatosis.vas.surg., 20: 987.
- Guidice L.C., Irwin J.C., Dsupin B.A., et al., (1993):** insulin-like growth factor (IGF), IGF binding protein (IGFBP) and IGF receptor gene expression and IGFBP synthesis in human uterine leiomyomata. *Hum.Rep.*, 8: 1766.
- Hannigan F.w and Gomez L.G.(1979):** uterine leiomyosarcoma: a review of prognostic clinical and pathologic features. *Am.J. obst.Gynecol.*,134:559.
- Harrington. E.A., Bennett M.R., fanidi A., et al., (1994):** C-myc-induced apoptosis in fibroblasts is inhibited by specific cytokines. *EMBOJ.*, 13: 3286.
- Harrison M.L., Jones D.S.,and smith S.K.(1994):** Quantification of messenger ribonucleic acid for epidermal growth factor in human myometrium and leiomyomata using reverse transcriptase polymerase chain reaction. *J.clin. Endocrinol. Metab.*,78:1179.
- Heinonen P.K., Taina E., NerdrumT., et al.,(1984):** Intravascular leiomyomatosis. *Ann.chir.Gynecol.*, 73: 100.
- Henderson J.E., Kremer R., Rhim J.S., et al.,(1992):** Identification and functional characterization of adenylate cyclase linked receptors for parathyroid hormone- linked peptide on immortalized human keratinocytes. *Endocrin.*, 130:449.

-
- Hendrickson M.R., and kempson R.I.(1980):** Surgical pathology of the uterine Corpus.philadlphia:WB Saunders, chap.4, P.P.275.
- Hengartner M.(1998):** Death cycle and Swiss army knives. *Nature*, 391:441.
- Henkart P.A.(1994):**2-chlorodeoxyadenosine chemotherapy triggers programmed cell death in normal and malignant lymphocytes. *Immunity*, 1:343.
- Herrman V., Degiampieyro P., Deheim E., et al., (1987):** activties in leiomyomas and myometrium of enzymes involved in energy metabolism. *Gynecol.obst. invest.*, 23: 194.
- Hickman J.K.(1992):** Apoptosis induced by anticancer drugs. *Cancer Metastasis Rev.*, 11:121.
- Hockenbery D.M., Nunez G., Millionan C. et al.,(1990):** Bcl-2 is an inner mitochondrial membrane protein that blocks programmed Cell death. *Nature*, 348: 334.
- Hockenbery D.M., oltvai Z.N., Yin X-M., et al.,(1993):** Bcl-2 function in an antioxidant pathway to prevent apoptosis. *Cell*, 75:241.
- Hockenbery D.M., (1995):** Bcl-2, a novel regulator of cellddeath. *BioEssays*, 17:631.
- Hofman G.E., Rao, Barrows G.H., et al., (1984):** Prostaglandin E and F2 alpha receptors in human uterine leiomyomas *J.Clin.Endocr. Metab.*, 58:545.
- Hsueh A.J.W., Billing H., Tsafrini(1994):** ovarian follicle atresia a hormonally controlled apoptotic process. *Endoc.Rev.*, 15:707.
- Huet-Hudson Y.M., chakraborty C., Desk,et al., (1990):** Estrogen regulates synthesis of epidermal growth factor in mouse uterine epithelial cells. *Mol. Endoc.*, 4:510.
- Hunter (1993):** The genetic regulation of apoptosis. *Curr. Opin. Gen.Dev.*, 3:1.
-

- Insogna K.L., Stewart A.F., Morris C.A., et al.,(1988):** Native and synthetic analogue of the malignancy-associated parathyroid hormone-like protein have in vitro transforming growth factor-like properties. *J.clin.invest.*,83: 1057.
- Ishizaki Y., Burke J.F., Raff M.C.(1994):** Mechanisms of muscle wasting. The role of the ubiquitin- proteasome pathway. *J. cell biology*, 126:1069.
- Jacobson M.D., Wells M. and Raff M.C.(1997):** Programmed Cell death in animal development. *Cell*, 88: 347.
- Kagi D., Vignaux F., Ledermann B., et al.,(1994):** Fas and perforin pathways as major mechanisms of T-cell mediated cytotoxicity. *Science*, 265: 528.
- Kaplan C., Benirschke K., and Johnson K.C.(1980):**Leiomyomatosis Peritonealis disseminata with endometrium. *Obst. Gynecol.*, 55:119.
- Kawaguchi K., Fujii S., Konishi I., et al.,(1985):** ultrastructural study of cultured smooth muscle cells from uterine leiomyoma and myometrium under the influence of sex steroids. *Gynecol. Oncol.*,21:32.
- Kelekar A., and Thompson C.,(1998):** Bcl- family proteins: the role of the BH3 domains in apoptosis. *Trends cell Biol.*, 8: 324.
- Kempson R.L., and Bari W.(1970):** uterine sarcomas: classification, diagnosis and prognosis. *Hum. Pathol.*, 1:331.
- Kenigsberg D., Chapitis J., Zuna R. et al.,(1987):** Hyperprolactinemia arising from a uterine tumor (abstract 255) in; *proceedings of the thirty-fourth annual meeting of the Society for Gynecologic investigation Atlanta, Georgia-March*,18:21.
- Kerr J.F.R., Winterford C. M.,and Harmon B.(1994):** Apoptosis, its significance in cancer and cancer therapy. *Cancer*, 73:2013.
- Khurana K.K., Singh S.B, Tatum A.H.,et al.,(1999):** Maintenance of increased bcl-2 expression in uterine leiomyomas after Gn RH agonist therapy., 44:487.

REFERENCES

- Kiechle M., Sreekantaiah C., Berger C.S., et al., (1991):** Non-random cytogenetic changes in leiomyomas of the female genitourinary tract. *Cancer Genet. Cytogenet.*, 53: 125.
- Korneyi J., Csermely T., Szekely J.A., et al., (1986):** Two types nuclear E2 binding sites in human myometrium and leiomyomata during the menstrual cycle. *Exp. clin. Endocrinol.*, 87:256.
- Korsmeyer S.J. (1992a):** Bcl-2 initiate a new Category of oncogen; Regulators of cell death. *Blood*, 80: 879.
- Korsmeyer S.J. (1992b):** Bcl-2: an antidote to programmed cell death cancer surveys., 15:105.
- Kozopas K.M., Yang T., Buchan H.L., et al., (1993):** MCL-1 a gene expressed in programmed myeloid cell differentiation, has sequence similarity to bcl-2. *Proc. Natl. Acad. Sci. USA.*, 90:3516.
- Krajewski S., Blomquist C., Franssila k., et al., (1995):** Reduced expression of pro-apoptotic gene BAX is associated with poor response rates to combination chemotherapy and shorter survival in women with metastatic breast adenocarcinoma. *Cancer Res.*, 55I: 4471.
- Lee E-Y-H.P. (1991):** Tumor Suppressor genes; Anew era for molecular genetic studies of cancer. *Breast Cancer Res. Treat.*, 19:3.
- Levine B. (1993):** Conversion of lytic persistent adenovirus infection by the bcl-2 cellular oncogene. *Nature*, 361:739.
- Lin E.Y., Orlofsky A., Berger M.S., et al., (1993):** characterization of Al, a novel hemopoietic-specific early-response gene with sequence similarity to bcl-2. *J. immunol.*, 151: 1979.
- Lin E.Y., Orlofsky A., wang H.G., et al., (1996):** Al a bcl-2 family member, prologns cell survival and permits myeloid differentiation. *Blood*, 87:982.
- Lipschutz A., and vargas L. (1941):** Struture and origin of uterine and extragenital fibroids induced experimentally in guinea pig by prologed administraton of estrogens. *Cancer Res.*, 1:236.
-

- Lipschutz A., and Schwarz J.(1944):** Microscopic structure o estrogen induced uterine and other abdominal fibroid treated with progesterone. *Cancer Res.*, 4: 241.
- Lowe S.W., Rulley H.E., Jacks T. et al., (1993):** P53-dependent apoptosis modulates the cytotoxicity of anticancer agents. *Cell*, 74: 957.
- Lowe S.W, Jacks T., Housman D.E., et al.,(1994):**Abrogation of oncogene-associated apoptosis allows transformation of P53-deficient cells. *Proc.Natl. Acad. Sci. USA.*, 1:2026.
- Lu Q.L., Hanby A.M., Hajibagtieri M.A.N., et al.,(1994):** Bcl-2 protein localizes to the chromosomes of mitotic nuclei and is correlated with the cell cycle in epithelial cell lines.*J.Cell sci.*, 107:36.
- Lumbiganon P.,Rugpaos , phandu-fung S., et al.,(1995):** protective effect of depo-medroryprogesterone acctate on surgically treated uterine leiomyomas: a multicentre case-control study.*Br. J. obst. Gynecol.*, 103:909
- Malet C., Grompel A., Yaneva H., et al.,(1991):** Estradiol and progesterone receptor in cultured human breast epithelial cells and fibroblasts immunohistochemical studies. *J. clin. Endoc.Metab.*, 73:8.
- Marshal R.D., Schaenberg-ferzo M.L., Friedman A.J., et al.,(1994):** Analysis of androgen receptor DNA reveals the independent clonal origins of uterine leiomyomata and the secondary nature of cytogenetic aberrations in leiomyomata development. *Genes chromosome Cancer*,11:1.
- Marshal L.M., spiegelman D., Barbier R.L., et al., (1997):** variation in the incidence of uterine leiomyomata among premenopausal women by age and race. *Obstet. Gynecol.*, 90:967.
- Marshal M., spiegelman, marlene B., et al.,(1998) :** A prospective study of preproductive factors and oral contraceptive use in relation to the risk of uterine leiomyoma. *Fert. Steril.*, 70:432.
- Marx J.(1993):** Cell death studies yield Cancer clues. *Science*, 259:760.

REFERENCES

- Maslar I, and Riddick D.(1979):** Prolactin production by human endometrium during the normal menstrual cycle. *Am. J.obst. Gynecol.*, 135:75
- Maslar I., Kaplan B., Luciano A., et al.,(1980):** Prolactin production by the endometrium of early human pregnancy. *J. clin.Endocr. Metab.*, 51:78.
- Matsuo, maruo, and samoto (1997):** Increased expression of bcl-2 protein in human uterine leiomyoma and its up-regulation progesterone. *J.clin.Endoc. Metab.*, 82:293.
- Maung Z.T., Maclean F.R., Reid M.M., et al.,(1994):** The relationship between bcl-2 expression and response to chemotherapy in acute leukaemia. *Br.J.hematol.*, 88: 105.
- McCarthy N.J., Evan G.I.(1998):** Methods for detecting and quantifying apoptosis. *Curr.Top.Dev.Biol.*, 36:259.
- MCDonnell T.J., Troncasop , Brisbay S.M., et al.,(1992):** Expression of the proto-oncogene bcl-2 in the prostate and its association with emergence of androgen-independent prostate cancers. *Cancer Res.*, 52: 6940.
- Meyer R.(1930):** Die pathologische antamic der Gebar Matter, in Teil E., editor: *Henke-lubarsch Hand buchder spezielle pathologische Anatomic und Histologie, Bd-71 Erlin, Springer-velag, chap.5, PP. 228.*
- Miura S., and Koide S.S. (1970):** Effect of insulin and growth hormone on rat uterine RNA synthesis. *Proc.Soc.exp.Biol.Med.*, 133:882.
- Moegi M., Takayanaki N., sando Y., et al., (1993):** A Case of so-called benign metastasizing leiomyoma responsive to progesterone. *Nipponkyobu shikkan Gakkal zasshi*, 31:890.
- Mok L.L.S., Nickals G.A., Thompson, et al., (1989):** Parathyroid hormone as a smooth muscle relaxant. *Endoc.Rev.*, 10:420.
-

REFERENCES

- Morales A.R., Fine G., Morris Pardo V., et al., (1975):** The ultrastructure of smooth muscle tumors with a consideration of the possible relationship of glomangiomas, hemangiopericytomas, and Cardiamyomas *path.Annu.*, 10:65.
- More K.L.(1982):** Regulation and phystological relevance of non-pituitary prolactin *In the developing human, Philadelphia, W.B. Saunders Co., chap.5, PP.279.*
- Morris R.G., Hargreaves A.D., Duvall E., et al., (1984):** Hormone induced cell death :II surface changes in thymocytes undergoing apoptosis. *Am.J. path.*, 115:426.
- Motoyama M., Wang F.P., Roth K.A., et al., (1995):** Massive cell death of immature hematopoietic cells and neurons in bcl- X deficient mice. *science (washington D.C.)*, 267:1506.
- Mountz J.D., Wu J., cheng J., et al., (1994):** Cell death mechansim and the immune system. *Arth. Reum.*, 37: 1415.
- Muram D., Gillieson M., and walters J.H.(1980):** Myomas of the uterus in pregnancy: ultrasonographic follow-up. *Am .J. obst.Gyn.*, 138:16.
- Muram M.D., pitlik M.D., sagie M.D., et al., (1998):**Intravenous leiomyomatosis with cardiac involvement in a pregenant woman. *Am J. obst. Gynecol.*, 178: 6200.
- Murphy A.A., Kettel L.M., Morales A.J., et al., (1993):** Regression of uterine leiomyomata in response to the antiprogesterone Ru 486. *J.Clin. End. Metab.*, 76:513.
- Nagata S.(1997):** Apoptosis by death factor. *Cell*, 88:355.
- Nelson K.G., Takahashi T., Lee D.C., et al., (1992):**
Transforming growth factor α is a potential mediator of estrogen action in the mouse uterus. *Endocr.*, 131: 1657.
- Nemes Z. (1996):** Expression and activation of tissue transglutaminase in apoptotic cells involuting rodent mamary tissue. *Eur.J.cell biol.*, 70:125.

REFERENCES

- Nicholson D.W., and Thornberry N.A.(1997):** Caspases: killer proteases. *Trends Biochem. Sci.*,22: 299.
- Novak E.R.and woodruff J.D.(1974):** Pathologic features of u terine fibroids. *Gynecologic and obstetric pathology,ed.7, pheladelphia,W.B.Saunders Company, chap.4, P.P.249.*
- Novak R.A., Goad D.L., Broadus A.E., et al.,(1992):** Human myometrium and myometrial tumors express the parathyroid hormone-related peptide gene. (*Abstract 94*) *Soc.stud. Reprod*, 46 (suppl.1),74.
- Nunez G. London L., Hockenbery D., et al., (1990):** Deregulated bcl-2 gene expression seletively prolongs survival of growth factor-deprived hematopoietic cell lines.*J. Immunol.*,144:3602.
- Nunez G,Hockenbery D, McDonnell T.J. et al., (1991):** Bcl-2 maintains β cell memory.*Nature*, 353 :71.
- Oltvai Z., Milliman G., and Korsmeyer S.J. (1993):** Bcl-2 heterodimerizes in vitro with Conserved homolog, Bax, that accelerates programmed cell death. *Cell*, 74:609.
- Oppenheim R.W.(1991):** Cell death during development of the nervous system. *Annu.Rev. neurosci.*, 14:453.
- Orrenius S., Mc Conkey D.J., Bellomo G., et al., (1989):** Role of CA^{2+} in toxic cell killing. *Trends pharmacol. Sci.*, 10:281.
- Otsuka H., Shinohara M., Kashimura M., et al.,(1989):** A comparative study of the estrogen receptor ratio in myometrium and uterine leiomyomas. *Int. J., Gynecol.obst.*,29:189.
- Otubo J.A., Buttram V.C., Besch N.F., et al., (1982) :** Unconjugated steroids in leiomyomas and tumor bearing myometrium. *Am.J. obst. Gyn.*, 143:130.
- Pantaleo G., Graziosi C., and Fauci A.S.(1993):** The immunopathogenesis of human immunodeficiency virus infection. *N. Engl. J.Med.* 328:327.AM. Parfitt, *ibid*,P.S. 82.
-

REFERENCES

- Parazzini F., la vecchia c, Negri E., et al.,(1988):** Epidemiologic characteristics of women with uterine fibroids: a case- control study. *Obst. Gynecol.*, 72:853.
- Parazzini F., Evancegri M.D., Carlo V., et al.,(1992):** Oral contraceptive use and risk of uterine fibroids. *Obstet. Gynecol.*, 79:430.
- Parmely T.H.,(1975):** Histogenesis of leiomyomatosis peritonealis disseminata(disseminated fibrosing deciduosus). *Obst. Gynecol.*, 46:511.
- Peitsch M.C., Polzar ,B., Stephan H., et al., (1993):**Characterization of the endogenous deoxyribonuclease involved in nuclear DNA degradation during apoptosis (Programmed cell death). *EMBO J.*, 2:371.
- Pollow K., Sinnecker G., Boquol E., et al., (1978):** In vitro conversion of estradiol 17- β into estrogen in normal human myometrium and leiomyoma. *J.clin . chem.clin.Biochem.*, 16:493.
- Pritchard J.A., and Mac Donald P.C(Eds)(1980) :** Maternal adaptation to pregnancy .in *williams: obstetrics, 16 th editions. New yourk, Appleton. Centary. crofts, P.P.221.*
- Raff M.C.(1992):** Social Control on cell survival and cell death. *Nature, (Lond)* 356:397.
- Ramzi S., Cotran M.D., vinarkumar M.D., et al., (1996):** Benign tumors of the uterus *In pathologic basis of disease, 5th edition, chapter 9, chapter10, P.P. 1059.*
- Ramzi S., Cotran M.D., vinarkumar M.D., et al., (1999):** Cellular pathology:Cell injury and cell death. *In pathologic basis of disease 6th edition chapter 1, P.P. 19.*
- Reed J.C, Cuddy M., Haldars., et al.,(1990):** Bcl-2 mediated tumorigenicity of a human T- lymphoid cell line: synergy with myc and inhibition by bcl-2 antisense. *Proc. Natl.Acad.Sci. USA.*, 87:3660.
- Reed J.C.(1994):** Bcl-2 and the regulation of programmed cell death. *J. cell Biol.*, 124:1.

- Reed J.C., Kitada S., Takayama S., et al.,(1994):** Regulation of chemoresistance by the bcl-2 oncoprotein in non- Hodgkin's lymphoma and lymphocytic leukemia cell lines. *Ann. Oncol.*, 5:561.
- Reed J.C (1995):** Regulation of apoptosis by of bcl-2 family proteins and its role in cancer chemoresistance. *curr. opin. oncol.*, 7:541.
- Reed J.C.(1996):** Mechanisms of bcl-2 family protein function and dysfunction in health and disease. *Behring inst. Mitt .*,97:72.
- Reed J.C.(1997):** Cytochrome C,can't live with it-can't live without it. *Cell*, 91:559.
- Rein M.S., friedmanA.J., stuart J.M., et al., (1990) :** Fibroid and myometrial steroid receptors in women treated with the gonadotropin releasing hormone agonist leuprolid acetate. *fert. Steril.*, 53:1018.
- Rein M.S., Friedman A.J., Barbieri R-L., et al., (1991):** cytogenetic abnormalities in uterine leiomyomata. *Obst Gynecol.*, 77:723.
- Rein M.S., Robert L., Barbieri L., et al., (1995):** Progesterone a critical role in the pathogenesis of uterine myomas. *Am.J.obst. Gynecol.*,172:14.
- Riddick D., luciano A., Kusmik W.,et al.,(1978):** Denovo synthesis of prolactin by human decidua. *life sci.*, 23:1913.
- Robenstein A.H., Seftel H.C., Miller K., et al., (1969):**Metabolic response to oral glucose in healthy south African white, Indian, and African subjects. *Br.Med.J.*, 1:748.
- Roger S.I., and Thomas I.(1959):** Paraplegia Caused by extraspinal metastasis from a uterine fibroid. *J. Neural. Neurosurg. Psychiatry.*, 22.141.
- Roger C., and Sander M.A.(1985):** Sonographic features of benign tumors of the uterus. The principles and practice of U/S in *Obst. and Gyn. Third edition, chapter7, P.P.351.*

REFERENCES

- Rose .I., under Wood R.H., Dunning M.T., et al.,(1975):** Testosterone metabolism in benign and malignant breast lesions. *Cancer*, 36:399.
- Rose L.I., Reddy V.V., And Biondi R.(1978):** Reduction of testosterone to 5 α -dihydrotestosterone by human and rat uterine tissues. *J.clin. Endoc. Metab.*, 46:766.
- Rosenthal S.M., hsiao D., and Silverman L.A.(1994):**An isulin-like growth factor-II(IGF-II) receptors stimulates differentiation, but not IGF-1 receptor down-regulation in muscle cells. *Endocr.*, 135:38.
- Ross R.K., pike M.C., vessey M.P., et al., (1986):** Risk factors for uterine fibroids: reduced risk associated with oral contraceptives.*Br.Med.J.*,293:359.
- Ross A.J., waymire K.G., Moss J.E., et al., (1998):** Testicular degeneration in bcl-w deficient mice.*Nat. genet.*,18:251.
- Roth L.M., Reed R.J., and sternberg W.H.,(1996):** Cotylenoid dissecting leiomyoma of the uterus-the sternberg tumors. *Am.J. Surg. Pathol.*, 20:1455.
- Runnebaum B., Klingak., Von Halst T., et al., (1978):** steroids in human myometrium and peripheral blood during the menstrual cycle.*Am.J. obst. Gynecol.*, 131:628.
- Sachs and Lotom J.(1994):** Control of programmed cell death in normal and leukemic cells, New implications for therapy. *Blood.*, 82:15.
- Sadan D., Vaniddekinge B., Van Geldeven C.J., et al.,(1987):** Estrogen and progesteron receptor concentrations in leiomyoma and normal myometrium. *Ann.clin. Biochem.*, 24: 263.
- Saegusa M., Takano Y., and okagasu. I (1995):** Bcl-2 expression and its association with cell kinetics in human gastric carcinomas and intestinal metaplasia . *J.Cancer Res. clin. Oncol.*, 121:357.
-

REFERENCES

- Saito Y., Seki H., Tokahaski S., et al.,(1986):** The difference in sex hormone dependence for gap Junction formation in the muscle of uterine leiomyoma and normal myometrium. *Acta obst.Gynecol. Jpn.*, 38(4) :595.
- Salvesen G.S., and Dixit V.M.(1997):** Caspases intracellular signalling by proteolysis. *Cell*, 91:443.
- Sarraf CE., and Bowen I.D.(1986):** Kinetic studies on a murine sarcoma and an analysis of apoptosis .*Br. J. Cancer*, 54:989.
- Savil.J (1997):**Recognition and phagocytosis of cell undergoing apoptosis. *Br. Med. Bull.*, 53:491.
- Schmidt -Gallwitzer M., Eilet Z., Genz. T., et al.,(1979):** Determination of estradiol, estrone and progesterone in serum and myometrium. Correlation with the content of sex steroid receptors and 17 B-hydroxy steroid dehydrogenase activity throughout the menstrual cycle. *J.Clin.Endoc.Metab.*,49:370.
- SearleT.A., Lawson P.J., Abott B., et al., (1974):** An electron-microscope study of the mode of cell death induced by cancer-chemotherapeutic agents in poulations of proliferating normal and neoplastic cells.*J.of path.*,116: 129.
- Sedlak J.W., oltvai Z.N., Yang E., et al., (1995):** Multiple bcl-2 family members demonstrate selective dimerization with bax. *Proc. Natl. Acad. Sci. USA.*, 92:7834.
- Sentman C.L., shutter J.R., Hockenbery D., et al., (1991):**Bcl -2 inhibits multiple forms of apoptosis but not negative selection in thymocytes. *Cell*, 67:879.
- Sierra A., Liovera B., Castellsoguex, et al., (1995):** Bcl-2 expression is associated with lymph node metastasis in human ductal breast carcinoma. *Int. J. Cancer*, 60:54.
- SilvenbergS.G.(1971):** Leiomyosarcoma of the uterus: A clinico-pathologic study .*obst. Gynecol.*,38:613.
- Silvestrini R., veneroni S., Dardon M.G., et al.,(1994):** A prognostic indicator strongly related to P53 protein in lymph node negative breast cancer patients. *J. Natl. Cancer inst.*,86:499.

- Sinha Y.N., Selby P.W., Lewis B.J., et al.,(1973):** A homologous radio-immunoassay for human prolactin. *J. clin. Endoc. Metab.*, 36:509.
- Soules M.R., and McCarthy K.S.Jr (1982):** Leiomyomas :steroid receptor Content-variation within normal menstrual cycles. *Am. J. obst. Gynecol.*, 143:6.
- Spellacy W.N., and Buhi W.C.(1969):** Pituitary growth hormone and placental lactogen measured in normal term pregnancy and at the early and late post partum periods. *Am.J.obst. Gynecol.*, 105: 888.
- Stanley J., and Korsmeyer (1999):** Bcl-2 gene family and the regulation of programmed cell death. *Cancer Res. (Supp)*. 59: 1693s.
- Steiner P.(1939):** Metastasizing fibroleiomyoma of the uterus . *Am. J. path.*, 15:89.
- Stemmermann G.N.(1961):**Distribution of blood flow in myomatous uteri as measured by locally injected 133 xenon. *Am.J. obst. Gynecol.*, 82:1612.
- Stern C., Deichert U., Thode B., et al.,(1992):** Cytogenetic subtyping of 139 uterine leiomyoma. *Geburt shilfe fraven heiltd*, 52:767.
- Stewart A.F., and Broadus A.E(1990):** Clinical Review: parathyroid hormone-related proteins coming of age in the 1990's. *J. clin. Endocr. Metab.*,71:1410.
- Strasser A., Harris A.W., Bath M.L., et al.,(1990):** Novel primitive lymphord tumors induced in transgenic mice by cooperation between myc and bcl-2. *Nature*, 348:331.
- Strasser A., Huang D.C., and Vaux D.L.(1997):** The role of the bcl-2/ ced 9 gene family in cancer and general implications of defects in cell death control for tumorigenesis and resistance to chemotherapy. *Bioch. Biophy. Acta*, 1333: f 151.
- Tamaya T., fujimoto J., and Okada H.(1985) :** Comparison of cellular level of steroid receptors in uterine leiomyoma and myometrium. *Acta. obst. Gynecol .Scand*, 64:307.

- Tanaka, Ito H., Adachis, et al., (1994):** Hypoxia induces apoptosis with enhanced expression of fas antigen messenger RNA in cultured neonatal rat cardiomyocytes. *Cric. Res.*, 75: 426.
- Theide M.A., Daifotic A.G., Weir E.C., et al., (1990):** Intrauterine occupancy controls expression of the parathyroid hormone-related gene in preterm rat myometrium. *proc. Natl Acad. Sci. USA.*, 87:6969.
- Theide M.A., Harm S.C., Hasson D-M., et al., (1991):** In vivo regulation of parathyroid hormone-related peptide messenger ribonucleic acid in rat uterus by 17 β -estradiol. *Endoc.*, 128:2317.
- Tierney W.M., Ehrlich C.E., Bailey J.C., et al., (1980):** Intravenous leiomyomatosis with extension into the heart. *Am. J. med.*, 69:471.
- Tiltman A.J. (1985):** The effect of progestins on the mitotic activity of uterine fibromyomas. *Int. J. Gynecol. path.*, 4:89.
- Tomita K., Coshenj., fernandez G., et al., (1982):** Immunologic and biologic characteristics of human decidual prolactin. *Am. J. obst. Gynecol.*, 142:420.
- Townsend D.E., Sparkes R.S., Boluda M.C., et al., (1970):** Unicellular histogenesis of uterine leiomyomas as determined by electrophoresis of glucose-6-phosphate dehydrogenase. *Am. obst. Gynecol.*, 107:1168.
- Tsujimoto Y. and Croce C.M. (1986):** Analysis of the structure transcripts, and protein products of bcl-2, the gene involved in human follicular lymphoma. *proc. natl. Acad. Sci. USA.*, 83:5214.
- Vasserman J., Baracut E.C., Bonduk C., et al., (1988):** Vascularization of uterine myomata. In *Abstracts of III world congress of gynecol. and obst.*, chapter 3, P.P. 108.
- Vaux D.L., Cory S., and adams J.M. (1988):** Bcl-2 gene promotes hematopoietic cell survival and cooperates with C-myc to immortalize pre-B Cells. *Nature*, 335:440.

- Vaux D.L., Haecker G., and Strasser A.(1994):** An evolutionary perspective on apoptosis. *Cell*, 76:777.
- Vaux D.L., and strasser A.(1996):** The molecular biology of apoptosis. *proc. Natl .Acad. sci. USA.*, 93:2239.
- Verrcellini p., Bocciolone L., Comlombo A., et al., (1993):**Gonadotropin releasing hormone agonist treatment before hysterectomy for menorhagia and uterine leiomyomas.*Am.J.Obs.Gynecol.*, 72:367.
- Vogelstein B., and kinzler K.W.(1992):** P53 function and dysfunction. *Cell*, 70: 523.
- Vollenhoven B.J., Lawrence A.S., and Healy D.L. (1990):** Uterine fibroids ; a clinical review.*Br.J. obst.Gynecl.*, 97:258.
- Von Micskyl(1977):** Sonographic study of uterine fibromyomas. In *sander R., James AE., Jr (eds)U/S in obst-and Gynecl. New York Appleton, centary Crofts, chapter9, P.P525.*
- Waggnar J.D(1966):** Ultrastructure of benign peripheral nerve sheeth tumors.*Cancer*, 19: 699.
- Walters G.A., Daly D.C., Kusmik W.F., et al.,(1983):** Hormonal Control of myometrial prolactin production, *presented at the sixty-fourth Annual Meeting of the Endocrine Society, Sanfrancisco, California, Abst.,872.*
- Walton M.I., whysong D., O'connor P.M., et al., (1993):** Constitutive expression of human bcl-2 modulates nitrogen mustard and comptothecin induced apoptosis. *Cancer Res.*, 53:1853.
- Weingold A.B., and Boltuch S-M (1961):** Metastatsizing breast Carcinoma to a large solitary uterine leiomyoma. *Am.J.obst.Gynecol.*, 82:1267.
- Weir E.C., Goad D.T., Darfatis A.G., et al.,(1994):** Relative overexpression of the parathyroid hormone related protein gene in human leiomyomas. *J.Clin.Endoc. Metab.*, 78:784.

REFERENCES

- Welsh R.A., and Meyer A.T.(1969):** Ultrastructure of gastric leiomyomas. *Arch. Pathol.*, 87:71.
- West C.P., Lumsden M.A., Lawsons, et al.,(1987):** Shrinkage of uterine fibroids during therapy with goserelin : a luteinizing hormone releasing hormone agonist administered as a monthly subcutaneous depot. *fert. Steril.*, 48: 45.
- Whitfield C.R.(ed) (1986):** Benign tumors of the uterus. Dewhurst's textbook of obstetrics and Gynecology for post. graduates, 4th edn., Black well scientific publications, London , chapter 5, P.P. 726.
- Williams G.T.(1991):** Programmed cell death apoptosis and oncogenesis. *cell*, 65:1097.
- Wilson J.F. and peale R.A (1952):** Multiple peritoneal leiomyoma associated with a granulosa cell tumors of the ovary. *Am.J. obst. Gynecol.*, 64:204.
- Wilson E.A., Yang F., and Rees E.D. (1980):** Estradiol and progesterone binding in uterine leiomyomata and in normal uterine tissues. *Obst. Gynecol.*, 55:20.
- Winn K.I., woodruff J.D., and Parmley T.H.(1976):**Electron microscopic studies of leiomyomatosis peritonealis disseminata. *obst. Gynecol.*, 48:225.
- Wyllie A.H.(1980):** Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature*, 284:555.
- Wyllie A.H., Kerr J.F.R., and Currie A.R.(1980):** Cell death: the significance of apoptosis. *Int. Rev.Cytol.*, 68: 251.
- Yamamoto I.(1961):** An electron microscope study on development of uterine smooth muscle. *J. Electronmicro. (Tokyo)*, 10: 145.
- Yamamoto T., Takamoni, and okada H.(1984):** Estrogen biosynthesis in leiomyomas and myometrium of the uterus. *Horm. metab. Res.*, 16:678.