

REFERENCES

REFERENCES

- Ahokas, R.A.; Mercer, B.M. and Sibai, B.M. (1991) :**
Enhanced endothelium derived relaxing factor activity in pregnant spontaneously hypertensive rats.
Am. J. Obstet. Gynecol.; 165 : 801.
- Allen, J.; Uidbjerg, N. and Petersen, L.K. (1989) :**
Intracervical 17- β oestradiol before induction of second trimester abortion with a prostaglandin E1 analogue.
Euro. J. Obstet. Gynecol. And Repro. Biol.; 32 : 123.
- Anderson, K.E.; Forman, A. and Ulmsten, U. (1983) :**
Pharmacology of labor. Clin. Obstet. Gynaecol.; 26 : 56.
- Anggard, E.E. (1994) :**
Nitric oxide : mediator murderer and medicine.
Lancet; 343 : 1199.
- Barclay, C.G.; Brennand, J.E.; Kelly, R.W. and Calder, A.A. (1993) :**
Interleukin-8 production by the human cervix.
Am. J. Obstet. Gynaecol.; 169 : 625.
- Barik, S. and Stewart, P. (1997) :**
[Letter] oral versus vaginal misoprostol in cervical priming.
Br. J. Obstet. Gynaecol.; 104 : 644.
- Barnes, P.J. (1993) :**
Nitric oxide and airways.
Eur. Resp. J.; 6 : 163.
- Bell, R.J.; Sutton, B. and Eddiel, W. (1989) :**
Relaxin level in antenatal patient in vitro fertilization.
Fertil. Steril.; 52 : 85.
- Bibby, J.G.; Brunt, J.; Mitchell, M.D.; Anderson, A.B.M. and Turnbull, A.C. (1979) :**
The effect of cervical encircage of plasma PG concentration during early human pregnancy.
Br. J. Obstet. Gynaecol.; 86 : 19.

Blumenthal, P.D. (1988) :

Prospective comparison of Dilapan and Laminaria for pretreatment of the cervix in the second trimester of abortion. Obst. Gynecol.; 72 : 243.

Bokström, H.; Bryman, I.; Norstorm, A. and Platz-Christesen, J.J. (1995) :

Dilapan-gemeprost regimen veresus combinations of extra-amniotic Rivanol-Laminaria/ Lamicel and Oxytocin for second trimester abortion. Int. J. Obstet. Gynaecol.; 48 : 69.

Brennand, J.E.; Calder, A.A.; Leitch, C.R.; Greer, I.A.; Chou, M.M. and Mackenzie, I.Z. (1997) :

Recombinant human relaxin as a cervical ripening agent. Br. J. Obstet. Gynaecol., 104 : 775.

Buckingham, J.C.; Selden, R. and Danforth, D.N. (1962) :

Connective tissue changes in the cervix during pregnancy and labor. Ann. NY. Acad. Sci.; 97 : 733.

Buhimschi, C.; Buhimschi, I.; Yallampalli, C.; Chwalisz, K. and Garfield, R.E. (1997) :

Contrasting effects of diethylene triamine-nitric oxide a spontaneously releasing nitric oxide donor on pregnant rat uterine contractility in vitro versus in vivo. Am. J. Obstet. Gynaecol.; 177 : 690.

Buhimschi, I.; Ali, M.; Jain, V.; Chwalisz, K. and Garfield, R.E. (1996):

Differential regulation of nitric oxide in the rat uterus and cervix during pregnancy and labor. Hum. Reprod.; 11 : 17550.

Burnett, A.L.; Lowenstein, C.J.; Bredt, D.S.; Chang, T.S.K. and Snyder, S.H. (1992) :

Nitric oxide : a physiological mediator of penile erection. Science, 257 : 401.

Calder, A.A. (1980) :

Pharmacological management of the unripe cervix in the human. In Naftolin. F. and Stubblefield, P.G. : Dilatation of uterine cervix. Raver Press New York, p. 317.

Calder, A.A. and Greer, I.A. (1992) :

Cervical physiology and induction of labor : In Bonnar J. (eds).
Recent advances in obstetrics and gynecology. Edinburgh
Churchill. Livingstone, p. 33.

Calver, H.; Collier, J. and Vallance, P. (1993) :

Nitric oxide and cardiovascular control.
Exp. Physiol.; 78 : 303.

Cameron, T.; Van Papendorp, C.L.; and Palmer, R.M.J. (1993) :

Relationship between nitric oxide synthesis and increase in
systolic blood pressure in women with hypertension in
pregnancy.
Hypertens. Pregnancy; 12 : 85.

Chanrockokul, B.; Herabutya, Y. and Punyavachira, P. (2000) :

Randomized comparison of glyceryl trinitrate and
prostaglandin E₂ for cervical ripening at term.
Obstet. Gynecol.; 96 : 549.

Chard, T. and Giobbens, G.L.D. (1983) :

Spurt release of oxytocin during surgical induction of labor in
woman.
Am. J. Obstet. Gynecol.; 147 : 678.

Chudleigh, P. and Pearce, J.M. (1992) :

Obstet. ultrasound in how, why and when Ed, Chudleigh, P.
and Pearce, J.M. 2nd ed. Churchill-Livingstone, London, p. 44.

Chvapil, M.; Droegemuller, W. and Meyer, T. (1982) :

New synthetic laminaria.
Obstet. Gynecol.; 60 : 729.

Chwalisz, K. and Garfield, R.E. (1998) :

Role of nitric oxide in the uterus and cervix : implication for
the management of labor.
J. Perinat. Med.; 26 : 448.

Chwalisz, K.; Shao-Oing, S.; Garfield, R.E. and Reier, H.M. (1997) :

Cervical ripening in guinea-pigs after local application of nitric
oxide.
Hum. Reprod; 12 : 2093.

- Chwalisz, K.; Benson, M. and Schlz, P. (1994) :**
Cervical ripening with interleukin-8 (IL-8), interleukin-1B (IL-1B) and tumor necrosis factor alpha (IL-1 α) in guinea pigs.
Hum. Reprod, 9 : 2173.
- Conard, K.P.; Joffe, G.M. and Kruszyna, H. et al. (1993) :**
Identification of increased nitric oxide biosynthesis during pregnancy in rats.
FASAB, J.; 7 : 566.
- Corbett, J.A.; Khael, A.; Shimizu, J.; Fredrick, K.; Misko, T.P.; McDaniel, M.L.; Kanagawa, O. and Unanue, E.R. (1993) :**
Nitric oxide production in islets from non-obese diabetic mice : aminoguanidine-sensitive and resistant stages in the immunological diabetic process.
Proc. Natl. Acad. Sci. USA; 90 : 8992.
- Corriveau, C.C.; Madara, P.J.; Van Dervot, A.L.; Tropea, M.M.; Wesley, R.A. and Danner, R.L. (1998) :**
Effects of nitric oxide on chemotaxis and endotoxin-induced interleukin-8 production in human neutrophils.
J. Infect. Dis.; 177 : 116 (English Abstract).
- Cross, W. and Pitkin, R. (1978) :**
Laminaria as an adjunct in induction of labor.
Obstet. Gynecol.; 51 : 606.
- Cunningham, F.G.; MacDonald, P.C.; Gant, N.F. and Levenek, J. (1997) :**
Williams Obst. The Egyptian 20th edition by Mass publishing Co., Egypt. P. 579-605, 677-607-631.
- Cuthbertson, Galley, B.H. and Webster, H.F. (1997) :**
Effect of exogenous nitric oxide and superoxide on interleukin-8 from human polymorphnuclear leucocytes.
Br. J. Anaesth.; 78 : 714-18 (English abstract).
- Danforth, D.N. (1947) :**
The fibrous nature of the human cervix and its relation to the isthmic segment in gravid and non-gravid uteri. Am. J. Obstet. Gynecol; 53 : 541-60. In Olah K.S. (1996). The cervix in pregnancy and labor. In : Studd. J. (ed) : Progress in Obstetrics and Gynecology, Vol. 12 New York, Churchill, Livingstone, p. 99.

Danforth, D.N. (1983) :

The morphology of the human cervix.
Clin. Obstet. Gynecol.; 26 : 7.

Davidge, S.T.; Billiar, T.R. and Roberts, J.M. (1994) :

Plasma and nitrates stable end products of nitric oxide are elevated in women with pre-eclampsia. Proceedings of the 9th international conference ISSHP; A60-164.

Dasai, K.M.; Sessa, W.C. and Vane, J.R. (1991) :

Involvement of nitric oxide in the reflex relaxation of the stomach to accommodate food or fluid. Nature; 351 : 477.

Dimitrios, S.; Mastrogiannis, D.S. and Knuppel, R.A. (1995) :

Labor induced using methods that do not involve oxytocin.
Clin. Obstet. and Gynecol.; 38 : 259.

Dong, Y.L.; Dai, B.S. and Yallampalli, C. (1997) :

Involvement of nitric oxide pathway in prostaglandin F2 α -induced preterm labor on rats.
Am. J. Obstet. Gynecol.; 177 : 907.

Dong, Y.L.; Gangula, P.R.R. and Yallampalli, C. (1996) :

Nitric oxide synthase isoforms in the rat uterus : differential regulation during pregnancy and labor.
J. Reprod. Fertil; 107 : 249.

Dong, Y.L. Yallampalli, C. (1996) :

Interaction between nitric oxide and prostaglandin E2 pathways in pregnant rat uteri.
Am. J. Physiol.; 270 : 471.

Duncan, M.R. and Berman, B. (1991) :

Stimulation of collagen and glycosaminoglycan production in cultured adult dermal fibroblasts by recombinant human interleukin-6.
J. Invest. Dermatol.; 97 : 686.

Ekerhovd, E.; Bronnstrom, M.; Delbro, D. and Norstorm, A. (1998) :

Nitric oxide mediated inhibition of contractile activity in the human uterine cervix.
Hum. Reprod.; 4 : 915.

Ellwood, D.A.; Mitchell and Anderson, A.B.M. (1980) :

The in vitro production of prostaglandins by the human cervix during pregnancy : preliminary observations.
Br. J. Obstet. Gynecol.; 87 : 210.

El-Maradny, E.; Kanayama, N.; Halim, A.; Maehara, K.; Kobayashi, T. and Terao, T. (1996) :

Biochemical changes in the cervical mucus after application of laminaria tent.
Acta. Obstet. Gynecol. Scand.; 75 : 203.

Embrey, M.P. and Mollison, B.G. (1967) :

The unfavorable cervix and induction of labor using cervical balloon.
J. Obstet. Gynecol. Br. Common, W.; 74 : 44.

Emmi, A.M.; Sturnick, J. Goldsmith, L.T. (1991) :

Ovarian control of pituitary hormone secretion in early human pregnancy.
J. Clin. Endocrinal. Metabol.; 72 : 1351.

Ezimokhai, M. and Nwabineli, J.H. (1980) :

The use of foley's catheter in ripening the unfavorable cervix prior to induction of labor.
Br. J. Obstet. Gynecol.; 87 : 281.

Facchinetti, F.; Piccinini, F. and Volpe, A. (2000) :

Chemical ripening of the cervix with intracervical application of sodium nitroprusside : a randomized controlled trial.
Hum. Reprod.; 15 : 2224.

Fitzgerald, D.J.; Rocki, W.; Murray, R. and Fitzgerald, G.A. (1990) :

Thromboxane A2 synthesis in pregnancy induced hypertension.
Lancet.; 335 : 751.

Fitzpatrick, R.J. and Dobson, H. (1981) :

The cervix in pregnancy and labor : clinical and biochemical investigations.
Churchill livingstone, p. 40-56.

Fitzpatrick, R.J. and Liggins, G.C. (1980) :

Effects of prostaglandins on the cervix of pregnant women and sheep. In Naftolin N and stubble field PG dilatation of the uterine cervix. New York, Ra. Press, p. 287.

Fluhman, C.F. (1980) :

The nature and development of the so-called glands of the cervix uteri.

Am. J. Obstet. Gynecol.; 68 : 753.

Frostell, C.G.; Fratacci, M.D.; Wain, J.C.; Zapol, W.M. (1991) :

Inhaled nitric oxide : a selective pulmonary vasodilator reversing hypoxic pulmonary vasoconstriction.

Circulation; 83 : 2038.

Frydman, R.; Lelaidier, C.; Baton-Saint-Mleux, C. et al. (1992) :

Labor induction of women at term with mifepristone (RU 486) : a double-blind randomized placebo-controlled study.

Obstet. and Gynecol.; 80 : 972.

Furchgott, R.F. and Vanhauhte, P.M. (1989) :

Endothelium derived relaxing and contractor factors.

FASAB, J; 3 : 2007.

Garvey, E.P.; Tuttle, J.V.; Convington, K. et al. (1994) :

Purification and characterization of the constitutive nitric oxide synthase from human placenta.

Arch. Biochem. Biophys.; 311 : 235.

Gerson Weiss, M.D. (1995) :

Relaxin uses to produce the cervical ripening.

Clin. Obstet. and Gynecol.; 38 : 293.

Gilson, G.J.; Russell, D.J.; Izquierdo, L.A.; Qualls, C.R. and Curet, L.B. (1996) :

A prospective randomized evaluation of a hygroscopic cervical dilator. Dilapan in the pre-induction ripening of patients undergoing induction of labor.

Am. J. Obstet. Gynecol.; 175 : 145.

Goldstien, I.; Zimmer, E.A. and Tamir, A. (1991) :

Evaluation of normal gestational sec growth : appearance of embryonic heart beat and embryo body movements using the transvaginal technique.

Obstet. Gynecol.; 77 : 885.

Gordon, A.S. and Calder, A.A. (1977) :

Oestradiol applied locally to ripen the unfavourable cervix.

Lancet. II : 1319.

Granstrom, L.; Ekman, G. and Malmstrom, A. (1991) :

Insufficient remodelling of the uterine connective tissue in women with protracted labor.

Br. J. Obstet. Gynecol.; 98 : 1212.

Greer, I.A. and Calder, A.A. (1989) :

Pre-induction cervical ripening with extra-amniotic and vaginal PGE 2.

J. Obstet. and Gynecol.; 16 : 18.

Grimes, D.A.; Schultz, K.F.; Cates, W.J. (1984) :

Prevention of uterine perforation during curettage abortion.

JAMA; 251 : 2108.

Grimes, D.A.; Schulz, K.F. and Cates, W.J. (1984) :

Prevention of uterine perforation during curettage abortion.

J. Am. Med. Assoc.; 251 : 2108.

Grozdanovic, Z.; Mayer, B.; Baumgarten, H.C.; Burnung, G. (1994) :

Nitric oxide synthase-containing nerve fibres and neurons in the genital tract of the female mouse.

Cell. Tissue Res.; 276 : 355.

Haluska, G.J.; Cook, M.J. and Novy, M.J. (1997) :

Inhibition and augmentation of progesterone production during pregnancy : effects on parturition in rhesus monkeys.

Am. J. Obstet. Gynecol.; 176 : 682.

Henry, I.; Lepoivre, M.; Drapier, J.C.; Ducrocq, C.; Boucher, J.L. and Guissani, A. (1993) :

EPR characteristics of molecular targets for NO in mammalian cells and organelles.

FASEB J.; 1 : 1124.

HSU, C.D.; Aversa, K.; Meaddough, E.; Lee, I.S. and Copel, J.A. (1997) :

Elevated amniotic fluid nitric oxide metabolites and cyclic guanosine 3' 5'-monophosphate in regnant women with intra-amniotic infection.

Am. J. Obstet. Gynecol.; 177 : 793.

Hughesdon, P.E. (1952) :

The fibromuscular structure of the cervix and its changes during pregnancy and labor.
J. Obstet. Gynecol. Br. Emp.; 59 : 763.

Husslein, P.; Fuschs, A.R. and Fuschs, F. (1981) :

Oxytocin and the initiation of human parturition and prostaglandin release during induction of labor by oxytocin.
Am. J. Obstet. Gynecol.; 141 : 688.

Huszar, G. (1980) :

Cellular aspects of labor in premature labor. Mead Johnson symposium on perinatal and developmental medicine on 15,p.16.

Ianaro, A.; O'Donnell, C.A.; Dirosa, M. and Liew, F.Y. (1994) :

A nitric oxide synthetase inhibitor reduces inflammation down regulates inflammatory cytokines and enhances IL-10 production in carrageenin-induced mice. Immunology.;82: 370.

Isemura, M.; Sato, N.; Yamnguchi, Y.; Nunakata, H.; Hayashe, M. and Yoshizawa, M. (1987) :

Isolation and characterization of fibronectin-binding proteoglycan carrying both heparin sulphate and dermatan sulphate chains from human placenta.
J. Biol. Chem.; 262 : 8926.

Ishimi, Y.; Miyaura, C. and Jin, C.H. (1990) :

IL-6 is produced by osteoblasts and induces bone resorption.
J. Immunol.; 145 : 3297.

Ito, A.; Hiro, D.; Sakyo, K. and Mori, Y. (1987) :

The role of leukocyte factors on uterine cervical ripening and dilatation.
Hum. Reprod.; 39 : 511.

Ito, A.; Itoh, Y.; Sasaguri, Y.; Morimatsu, M. and Mori, Y. (1992) :

Effect of interleukin-6 on the metabolism of connective tissue components in rheumatoid synovial fibroblast.
Arthritis Rheum.; 35 : 1197.

- Izumi, H.; Yallampalli, C. and Garfield, R.E. (1990) :**
Gestational changes in L-arginine induced relaxation of pregnant rat human myometrial smooth muscle.
Am. J. Obstet. Gynecol.; 169 : 1327.
- Johnston, T.A.; Greer, I.A.; Kelly, R.W.; Gallacher, A. and Calder, A.A. (1990) :**
Plasma prostaglandin metabolites in human labor. 7th international conference on prostaglandins and related molecules florance, p. 165.
- Jorens, P.G.; Vermeira, P.A. and Herman, A.G. (1993) :**
L-arginine dependant nitric oxide synthetase : A new metabolic pathway in the lung and airways.
Eur. Resp. J.; 6 : 258.
- Junquiera, L.C.U.; Zugai, B.M.; Montes, G.S.; Toledo, O.M.S.; Krisztan, R.M. and Shigihara, K.M. (1980) :**
Morphologic and histochemical evidence for the occurrence of collagenolysis and for the role of neutrophilic polymorphonuclear leukocytes during cervical dilatation.
Am. J. Obstet. Gynecol.; 138 : 273.
- Karsdorp, V.H.M.; Van Vugt, J.M.G.; Jacobs, C.; Dekker, G.A. and Van Geijn, H.P. (1994) :**
Amino acids glucose and lactate concentration in umbilical blood in relation to umbilical artery flow patterns.
Eur. J. Obstet. Gynecol.; 57 : 117.
- Keirse, M.J.N.C. (1979) :**
PGs in parturition in Keirse, M.; Anderson, A.; Gravenhorst, J. (eds) : Human parturition. The Hague Netherlands Martinus Nijhoff, p. 101.
- Keirse, M.J.N.C. (1992) :**
Estrogen for cervical ripening. In Chalmers I., Enkin M., Keirse M.J.N.C. (eds) : Effective care in pregnancy and childbirth. Oxford University Press, p. 1057.
- Keirse, M.J.N.C. and Van Oppen, A.C.C. (1989) :**
Preparing the cervix for induction of labor. In : Chalmers, I.; Enkin, M. and Keirse, M.J.N.C. (eds) : Effective care in pregnancy and childbirth. Oxford University Press, p. 988.

Keirse, M.J.N.C.; Flint, A.P.F. and Turnbull, A.C. (1974) :

F prostaglandins in amniotic fluid during pregnancy and labor.
J. Obstet. Gynecol. Br. Commonw; 81 : 35.

Keirse, M.J.N.C. and Turnbull, A.C. (1973) :

E prostaglandin in amniotic fluid during pregnancy and labor.
J. Obstet. Gynecol. Br. Commonw; 80 : 970.

Keller, R.; Bastti, S.; Keist, R.L.; Mulsch, A. and Klauser, S. (1992) :

Induction of nitric oxide synthetase is a necessary precondition
for expression of tumor necrosis factor-independent lumoricidal
activity by activated macrophages : biochemi.
Biophys. Res. Commun.; 184 : 1364.

Kitagawa, H.; Takeda, F. and Kohei, H. (1989) :

Gastric mucosal protective action of endothelium-derived
relaxing factor.
Scand. J. Gastroenterol.; 162 : 131.

Kitamura, K.; Ito, A.; Mori, Y. and Hirakawa, S. (1979) :

Changes in the human uterine cervical collagenase with special
reference to cervical ripening.
Biochem. Med.; 22 : 332.

Kitamura, K.; Ito, A.; Mori, Y. and Hirakawa, S. (1980) :

Glycosaminoglycans for the human uterine cervix : Heparin
sulphate increases with reference to cervical ripening.
Biochemi. Med.; 23 : 159.

Kolb, H. (1992) :

Nitric oxide a pathogenic factor in autoimmunity.
Immunol. Tod.; 13 : 157.

Kovacs, A.G.; Makary, A.; Peto, J. and Steinmetz, G. (1994) :

Deficiency of c-GMP level in placental circulation in
pregnancy-induced hypertensive disorders : possibility of
decreased endothelium-derived relaxing factor.
Hypertens. Pregnancy; 13 : 163.

Lawrie, A.; Penney, G. and Templeton, A. (1996) :

A randomized of oral and vaginal misoprostol for cervical
priming before suction termination of pregnancy.
Br. J. Obstet. Gynecol.; 103 : 1117.

Ledingham, M.A.; Denison, F.C.; Kelly, R.W.; Young, A. and Norman, J.E. (1999) :

Nitric oxide donors stimulate prostaglandin F (2 alpha) and inhibit thromoxane B (2) production in the human cervix during first trimester of pregnancy.

Hum. Reprod., 5 : 973.

Ledingham, M.A.; Thomson, A.J.; Young, A.; Macara, L.A. and Norman, J.E. (2000) :

Changes in the expression of nitric oxide synthase in the human uterine cervix during pregnancy labor.

Mol. Hum. Reprod. Nov; 6 (11) : 1041.

Lees, C.; Campbell, S.; Jauniaux, E.; Brown, R.; Ramsay, B.; Gibb, D.; Moncada, S. and Martin, J.F. (1994) :

Arrest of preterm labor and prolongation of gestation with glyceryl trinitrate a nitric oxide donor. Lancet; 343, 1325.

Leppert, P.C. (1992) :

Cervical softening effacement and dilatation : A complex biochemical cascade.

Journal of Maternal and Fetal Medicine; 1 : 213.

Leppert, P.C. and Yu, S.Y. (1994) :

Apoptosis of cervical cells in pregnant rats in association with cervical softening.

Gynecol. Obstet. Invest.; 37 : 150.

Leppert, P.C.; Keller, S.; Cretta, J. and Mandl, I. (1982) :

Conclusive evidence of the presence of elastine in human and monkey cervix.

Am. J. Obstet. Gynecol.; 142 : 179.

Liew, F.Y. and Cox, F.E. (1981) :

Cervical ripening as an inflammatory reaction in Ellwood, D.A.; Anderson, A.B.M. editors. The cervix in pregnancy and labor : Clinical and Biochemical Investigations.

Edinburgh; p. 1-9.

Lowenstein, C.J.; Dinerman, J.L. and Synder, S.H. (1994) :

Nitric oxide a physiological messenger.

Ann. Intern. Med.; 120 : 227.

Mackenzie, I.Z. and Fry, A. (1981) :

Prostaglandin E2 pessaries to facilitate first trimester aspiration termination.

Br. J. Obstet. Gynecol.; 8 : 1033.

MacLennan, A.H. (1981) :

Cervical ripening and the induction of labor by vaginal prostaglandin F2 α and relaxin. In : Elwood, D.A.; Anderson, A.B.M. (eds.) The cervix in pregnancy and labor : clinical and biochemical investigations.

Churchil Livingstone Edinburgh. P. 187.

MacLennan, A.H.; Green, R.C. and Gant, P. (1985) :

Ripening of the human cervix and induction of labor with intra-cervical purified porcine relaxin.

Obstet. and Gynecol.; 92 : 693.

MacLennan, A.H.; Green, C.R.; Gant, P.; Grant, P. and Nicolson, R. (1986) :

Ripening of the human cervix and induction of labor with intracervical purified porcine relaxin.

Obstet. Gynecol.; 68 : 598.

Magan, E.F.; Perry, K.G.; Dockery, J.R.; Bass, J.D.; Chauhan, S.P. and Morrison, J.C. (1995) :

Cervical ripening before medical induction of labor : A comparison of prostaglandin E2 estradiol and oxytocin.

Am. J. Obstet. gynecol.; 172 : 1702.

Marill, K.A.; Ingmire, T.E. and Nelson, B.K. (1994) :

Ability of a single beta HCG measurement to evaluate for absence of ectopic pregnancy.

J. Emerg. Med.; 17 : 419.

Maruo, N.; Morita, I.; Shirao, M. and Murota, S. (1992) :

IL-6 increases endothelial permeability in vitro.

Endocrinology; 131 : 710.

Mastrogiannis, D.S. and Knuppel, R.A. (1995) :

Misoprostol to induce labor.

Clinical Obstet. and Gynecol; 38 : 259.

McInnes, D.R.; Naftolin, F. and Van der Rest, M. (1980) :

Cervical changes in pregnant woman. In Naftolin F. and Stubble field PG (eds). Dilatation of the uterine cervix. New York : Raven Press, p. 181.

Minamoto, T.; Arai, K.; Hirakawa, S. and Nagai, Y. (1987) :

Immunohistochemical studies on collagen types in the uterine cervix in pregnant and non-pregnant states.
Am. J. Obstet. Gynecol.; 156 : 138.

Mitchell, M.D.; Dudeley, D.J.; Edwin, S.S. and Lundin-Schiller, S. (1991) :

Interleukin 6 stimulates prostaglandin production by human amnion and decidual cells. Eur. J. Pharmacol.; 192 : 189.

Molnar, M.; Suto, T.; Toth, T. and Hertenlendy, F. (1994) :

Prolonged blockade of nitric oxide synthesis in gravid rats produces sustained hypertension, proteinuria, thrombocytopenia and intrauterine growth retardation.
Am. J. Obstet. Gynecol.; 170 : 1485.

Moncada, S.; Higgs, A. (1993) :

The L-arginine-nitric oxide pathway.
N. Engl. J. Med.; 329 : 2002.

Morris, N.H.; Sooranna, S.R.; Eaton, B.M. and Steer, P.J. (1993) :

NO synthase activity in placental bed and tissues from normotensive pregnant

Munsick, R.A. and Fineberg, N.S. (1996) :

Cervical dilatation from multiple laminaria tents used for abortion.
Obstet. Gynecol.; 87 : 726.

Nakatsuka, M.; Habara, T.; Kamada, Y.; Tada, K. and Kuda, T. (2000):

Elevation of total nitrite and nitrate concentration in vaginal secretions as a predictor of premature delivery.
Am. J. Obstet. Gynecol.; 182 : 644.

Nathan, C. (1992) :

Nitric oxide as a secretory product of mammalian cells.
FASAB, J.; 6 : 3051.

Nussler, A.K. and Billiar, T.R. (1993) :

Inflammation immunoregulation and inducible nitric oxide synthetase.

J. Biol. Chem.; 54 : 171.

Olah, K.S. (1994) :

Magnetic resonance imaging in the assessment of cervical hydration state.

Br. J. Obstet. Gynecol.; 101 : 255.

Olah, K.S. (1996) :

The cervix in pregnancy and labor. In : Studd J. (ed) : Progress in Obstetrics and Gynecology first edition Vol. (12). New York Churchill Livingstone; p. 99.

Osmer, R.; Rath, W.; Pflanz, M.; Kuhn, W.; Stuhlsta, H.W. and Szeverenyi, M. (1993) :

Glucosaminoglycans in cervical connective tissue during pregnancy and parturition.

Obstet. Gynecol.; 81 : 88.

Palmer, R.M.J.; Ferrige, A.G. and Moncada, S. (1987) :

Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor.

Nature; 327 : 524.

Palmer, R.M.J.; Rees, D.D.; Ashton, D.S. and Moncada, S. (1988) :

L-arginine is the physiological precursor for the formation of nitric oxide in endothelium dependant relaxation.

Biochem. Biophys. Res. Comm.; 153 : 1251.

Phyllis, C. and Leppert, M.D. (1995) :

Anatomy and physiology of cervical ripening.

Clinical Obstet. and Gynecol.; 38 : 267.

Piner, M.S.; Bolognese, R.J. and Feldman, J.D. (1967) :

Long-acting progesterone as a cause of missed abortion.

Am. J. Obstet. Gynecol.; 97 : 579.

Qing, S.; Bier, H.; Garfield, R. and Chwalisz (1996) :

Local application of nitric oxide donor induces cervical ripening.

J. Soc. Gynecol. Invest.; 3 : 462.

Raifer, J.; Aronson, W.J.; Bush, P.A.; Doren, F.S. and Ignarro, L.J. (1992) :

Nitric oxide as a mediator of relaxation of the corpus cavernosum in response to nonadrenergic non cholinergic neurotransmission.

N. Engl. Med.; 326 : 90.

Rajabi, M.; Solomon, S. and Poole, R. (1992) :

Activation of protein kinase C stimulates collagenase production by cultured cells of the cervix of pregnant guinea pig.

Am. J. Obstet. Gynecol.; 167 : 194.

Rall, T.W. and Schleifer, L.S (1985) :

Oxytocin PG ergotalkaloids and other drugs tocolytic agents. In : Gilman AG Goodman LS Rall TW Murad F (eds) : the pharmacological basis of therapeutics 7th edition. New York : macmillan Publishing.

Rath, W.; Adelman-Girill, B.C. and Pieper, U. et al. (1987) :

The role of collagenase and protease in prostaglandin induced cervical ripening.

Prostaglandins; 34 : 119.

Rayburn, W.F. (1989) :

PGE₂ gel for cervical ripening and induction of labour. A critical analysis.

Am. J. Obstet. Gynecol.; 160 : 529.

Read, M.D. and Mattock, E.J. (1982) :

Cervical ripening with PGE₂ pessaries-a question of dose.

J. Obstet. Gynaecol.; 93 : 933.

Robinson, G.; Filshie, G.M. Cooper, J.C.; Cust, M. and Macleod, F. (1991) :

A comparative study of Dilapan ceragem and Laminaria for preparation of the cervix prior to termination of first trimester pregnancy. XIII World Congress of Gynecology and Obstetrics (FIGO) Singapore; Poster no. 1586.

Rockett Awburn, M.; Aggarwal, B. and Cowden Clark (1992) :

Invivo induction of nitrite and nitrate by tumor necrosis factors lymphotoxin and interleukin-1 : possible roles in malaria.

Infect. Immun.; 60 : 3725.

Rossiant, R.; Falke, K.S.; Lopez, F.; Slamak, Pison, U. and Zapol, W.M. (1993) :

Inhaled nitric oxide for the adult respiratory distress syndrome.

N. Engl. Med.; 328 : 399.

Roubin, D. and Arias, F. (1993) :

A randomized trial of extra-amniotic saline infusion with intracervical Foley's catheter balloon versus PGE2 vaginal gel for ripening of the cervix and inducing labour in patients with unfavourable cervix.

Obstet. Gynecol.; 82 : 290.

Roztocil, A.; Miklica, J.; Koudelka, M.; Jeline, K.J.; Kutalkova, S.; Burnog, T. and Pilka, L. (1996) :

Induction of cervical ripening with hydrophilic Dilapan rods in pregnancy at term.

Zentralbl. Gynakol; 118 : 29.

Ruoslanti, E. (1988) :

Structure and biology of proteoglycans.

Ann. Rev. Cell. Biol.; 4 : 229.

Salvemini, D.; Misko, T.P.; Masferrer, J.L.; Seibert, K.; Currie, M.G. and Needleman, P. (1993) :

Nitric oxide activates cyclooxygenase enzyme.

Proc. Natl. Acad. Sci. USA.; 90 : 7240.

Sanchez-Ramos, L.; Kaunitz, A.M.; Delvalle, G.O.; Delke, I.; Schroeder, P.A. and Briones, D.K. (1993) :

Labor induction with PGE1 methyl analogue misoprostol versus oxytocin : Randomized trial.

Obstet. Gynecol.; 81 : 322.

Schluz, K.F.; Grimes, D.A. and Cates, W.Jr. (1983) :

Measures to prevent cervical injury during suction curettage abortion.

Lancet.; 1 : 118.

Seligman, S.P.; Abramson, S.B.; Young, B.K. and Buyon, J.P. (1994) :

The role of nitric oxide in the pathogenesis of pre-eclampsia.

Am. J. Obstet. Gynecol.; 172 : 944.

- Sherman, K.S.; Jayanthi, V.; Probet, C.S.; Stemart, C.R. and Mayberry, J.F. (1994) :**
Infertility Obstet. and Gynecological problems in coeliac pressure.
Digest. Dis., 12 : 186.
- Sherwood, O.P. (1988) :**
Relaxin in : Knobil. E. Nel. J. (eds) : the physiology of reproduction. New York, Raven; 585.
- Shi, L.; Shi, S.O.; Saade, G.R.; Chmalisz, K. and Garfield, R.E. (2000) :**
Studies of cervical ripening in pregnant rats : effects of various treatments.
Mol. Hum. Repord.; 6 : 382.
- Shirley, K. and William, F. (1995) :**
Outpatient cervical ripening.
Clin. Obstet. and Gynecol.; 38 : 301.
- Sitteri, P.K.; Febres, F. and Clemens, L.E. (1977) :**
Progesterone and maintenance of pregnancy : is progesterone nature's immunosuppressant?
Ann. NY. Acad. Sci.; 286 : 384.
- Snyder, S.H. (1992) :**
Nitric oxide : first in a new class of neurotransmitters.
Science; 54 : 171.
- Soloff, M.S. (1988) :**
The role of oxytocin in the initiation of labor and oxytocin-prostaglandin interactions. In : Mc. Nellis, D.; Challis, J.; MacDonald, P.; Nathaniels, Z.P.; Roberts, J. editors. The onset of labor : cellular 2nd integrative mechanisms in the onset of labor. Ithaco (NY) : Perinatology Press; p. 87.
- St. Onge, R.D. and Conners, G.T. (1995) :**
Pre-induction cervical ripening : A comparison of intracervical prostaglandin E2 gel versus the Foley catheter.
Am. J. Obstet. Gynecol.; 172 : 687.
- Stark, M.E.; Szurszewski, J.H. (1992) :**
Role of nitric oxide in gastrointestinal and hepatic function disease. Gastroenterology; 130 : 1928.

Steiner, A.L. and Cresay, R.K. (1983) :

Methods of cervical priming.

Clinical Obstet. and Gynecol.; 26 : 37.

Steinelt, B.G.; O'Bryne, E.M. and Kroe, R.L. (1980) :

The role of relaxin in cervical softening during pregnancy in mammals. In Nafolin F and Stubblefield (eds) : Dilatation of the uterine cervix. New York Raven Press; p. 155.

Stiers, J.A.; Bergsja, P. and M. Yking, O. (1984) :

Human chorionic gonadotrophin in maternal plasma after induced abortion spontaneous abortion remained ectopic pregnancy.

Obstet. Gynecol.; 64 : 391.

Stjernholm, Y.; Sahlin, L.; Akerberg, S.; Elinder, A.; Erilesson, H.A.;

Malmstrom, A. and Ekman, G. (1996) :

Cervical ripening in humans : potential roles of estrogen progesterone and insulin-like growth factor-I.

Am. J. Obstet. Gynecol.; 174 : 1065.

Sugihara, T.; Hattari, Y.; Yamamoto, Y.; Qi, F.; Ichikawa, R.; Sato, A.;

Liu, M.Y.; Abe, K. and Kanna, M. (1996) :

Treatment of preterm cervical dilatation with glyceryl trinitrate, a nitric oxide donors.

Aust. NZ. J. Obst. Gynecol.; 100 : 377.

Szalay, S.; Husslein, P. and Grunberge, W. (1981) :

Local application of prostaglandin E2 and its influence on collagenolytic activity of cervical tissue.

Sing. J. Obstet. Gynecol.; 12 : 15.

Thomson, J.; Burnett, C.; Cameron, A.D.; Cameron, II.; Greer, I.A.

and Jane, E. (1997) :

Nitric oxide donors induce ripening of the human uterine cervix.

Br. J. Obstet. Gynecol.; 104 : 1054.

Thomson, A.J.; Lman, C.B.; Cameron, A.D.; Cameron, I.T.; Greer,

I.A. and Norman, J.E. (1997) :

Nitric oxide donors induce ripening of the human uterine cervix : a randomized controlled trial.

Br. J. Obst. Gynecol.; 104 : 1054.

Thomson, A.J.; Lman, C.B.; Ledingham, M.; Hamat, R.C.; Cameron, I.T.; Greer, I.A. and Norman, J.E. (1998) :

Randomized trial of nitric oxide donor versus prostaglandin for cervical ripening before first trimester termination of pregnancy.

Lancet; 352 : 1093.

Thurnau, G.R. and Rayburn, W.F. (1992) :

Cervical ripening agents and uterine stimulants. In : Drug therapy in obstetrics and gynecology. Ed Rayburn W.F. and Zuspan, F.P. 3rd ed. P. 299. Mosby Year Book. St. Louis USA.

Trachtman, H.; Futterweit, S.; Garg, P.; Reddy, K. and Singhal, P.C. (1996) :

Nitric oxide stimulates the activity of a 72-kd a neutral matrix metalloproteinase in cultured rat mesangial cells.

Biochem. Biophys. Res. Commun.; 218 : 704.

Uldbjerg, N. (1989) :

Cervical connective tissue in relation to pregnancy labour and treatment with prostaglandin E2.

Acta. Obstet. Gynaecol. Scand. Supplement; 148.

Uldbjerg, N.; Ekman, G.; Malmstrom, A.; Olsson, K. and Ulmsten, U. (1983c) :

Ripening of the human uterine cervix related to changes in collagen glycosaminoglycans and collagenolytic activity.

Am. J. Obstet. Gynecol.; 147 : 662.

Uldbjerg, N.; Ekman, G.; Malmstrom, A.; Sporrang, B.; Ulmsten, U. and Wingerup (1981) :

Biochemical and morphological changes of human cervix after local application of PGE2 in pregnancy.

Lancet; 267.

Uldbjerg, N.; Ekman, G. and Ulmsten, U. (1983b) :

The ripening of human uterine cervix in terms of connective tissue biochemistry.

Clin. Obstet. and Gynecol.; 26 : 14.

Uldbjerg, N.; Malmstrom, A.; Ekman, G.; Sheehan, J.; Ulmsten, U. and Wingerup, L. (1983d) :

Isolation and characterization of dermatan sulphate proteoglycan from human uterine cervix.
Biochem. J.; 209 : 497.

Van Buren, G.A.; Yang, D.; Clark, K.E. and Estrogen (1992) :

Induced uterine vasodilation is antagonised by L-nitroarginine methyl ester an inhibitor of nitric oxide synthesis.
Am. J. Obstet. Gynecol.; 167 : 828.

Verma, A.; Hirsch, D.J.; Glatt, C.E.; Ronnett, G.V. and Synder, S.H. (1993) :

Carbon monoxide : a putative neural messenger.
Science; 259 : 381.

Von Malliot, K.V.; Stuhlastzh, W.; Mohanaradhakrishnan, V. and Breiling, H. (1977) :

Changes in glycosaminoglycans distribution pattern in the human uterine cervix during pregnancy and labour.
Am. J. Obstet. Gynecol.; 135 : 503.

Warren, W.B.; Timar-Trich, I. And Peisner, D.B. (1989) :

Dating the early pregnancy by sequential appearance of embryonic structures.
Am. J. Obstet. Gynecol.; (6) : 747.

Weiner, C.; Martinez, E.; Zhu, L.K.; Chodsi, A.; Chestnut, D. (1989) :

Invitro release of endothelium-derived relaxing factor by acetylcholine is increased during the guinea pig pregnancy.
Am. J. Obstet. Gynecol.; 161 : 1599.

Weletz, P.S. and Goldstein, S.R. (1991) :

Abnormal first trimester gestation in diagnostic medical sonography in Obstet. and Gynecol. Ed, Berman, M.C. L. Pipincott, Co. Publication, Philadelphia, p. 347.

Wingerup, L.; Anderson, K.E. and Ulmsten, U. (1978) :

Ripening of the uterine cervix and induction of labour at term with PGE₂ in viscous gel.
Acta. Obstet. Gynecol. Scand.; 57 : 403.

Wingerup, L.; Ulmsten, U. and Andesson, K.E. (1979) :

Ripening of the cervix by intracervical application of PGE₂ gel before termination of pregnancy with dilatation and evacuation. Acta. Obstet. Gynecol. Scand. (Suppl.); 84 : 15.

Yadava, R.P. (1990) :

Sublingual glyceryl trinitrate spray facilitates IUD insertion. Br. F. Sexual. Med.; 7 : 217.

Yallampalli, C. and Garfield, R.E. (1993) :

Inhibition of nitric oxide synthesis in rats during pregnancy produces symptoms identical to preedampsia. Am. J. Obstet. Gynecol.; 169 : 1316.

Yallampalli, C.; Garfield, R.E. and Byam-Smith, M. (1993) :

Nitric oxide inhibits uterine contractility during pregnancy but not during delivery. Endocrinology; 133 : 1899.

Yallampalli, C.; Izumi, H.; Byam-Smith, M. and Garfield, R.E. (1994) :

An L-arginine-nitric oxide-cyclic guanosine monophosphate system exists in the uterus and inhibits contractility during pregnancy. Am. J. Obstet. Gynecol.; 170 : 175.

Yamada, K.M. (1983) :

Cell surface interactions with extracellular materials. Annu. Rev. Biochem.; 52 : 761.

Zorumski, G.F. and Igumi, Y. (1993) :

Nitric oxide and hippocampal synaptic plasticity. Biochem. Pharmacol.; 46 : 777.