

REFERENCES

REFERENCES

- Abdel Salam E., Hafez A.A., Abdel Aziz S., and Elsaphori Z., (1982):
 α_1 antitrypsin in schistosomal hepatic disease.
 Egypt J. of Biochem. 1:3.
- Ala- Mouhala. I., (1990):
 Effects of exercise on glomerular passage of macromolecules in patients with diabetic nephropathy and in healthy subjects.
 Scand. J. clin. lab. invest., 50(1):27.
- Ala- Mouhala I., and Pasternack A., (1987):
 Fractional dextran and protein clearance in glomerulonephritis and in diabetic nephropathy.
 Clin. Sci., Mar., 72(3):289-296.
- Anderson J., (1981):
 Disorders of metabolism:
 Text book of adverse drug reaction, 2nd edition. Ed. by Davis D.M. University press, oxford. P.314.
- Andraoli T.E., Hoffman J.E., and Fanestil D., (1978):
 "Physiology of membrane disorder".
 2nd Ed. Plenum publishing Co., Newyork., P.63.
- Axelsen N.H.V., Bock E., and Larsen P., (1983):
 Protein in body, urine acids and tumor markers: Diagnostic and clinical aspects.
 Scand.J. immunol 17(10):87.
- Bagdasarian A., Wheeler J., Stewart G.T., Ahmed S.S, and Colman R.W., (1981):
 Isolation of α_1 protease inhibitor from human normal and malignant ovarian tissue.
 J. Clin. invest., 67:281.
- Ballard D.J., Hmpfrey L.L., Melton L.J., et al., (1988):
 Epidemiology of persistent proteinuria in type II diabetes mellitus.
 Diabetes, Apr., 37(4):405-412.
- Bending J.J., Loboyeo A., Vergani D., and Viberti G.G., (1988):
 Proteinuria and activated T.lymphocyte in diabetic nephropathy.
 Diabetes. May., 37:507-511.
- Bennett C.M., Classoc R.J., and Brenner., (1976):
 Permselectivity of the glomerular capillary wall.
 J. clin invest., 57:1287-1294.

- Berggard I., and Beam A.G., (1968):
Isolation and properties of a low molecular weight B₂ microglobulin occurring in human biological fluid.
J. Biol. Chem., 243:4095.
- Bieter R.N., (1931):
Albuminuria in glomerular and aglomerular fish.
J. pharmacol. Exp. Ther. 43:407- 419.
- Bohrer M.P., Baylis C., Robertson C.R., and Brenner B.M., (1977):
Mechanisms of the puromycin- induced defects in the transglomerular passage of water and macromolecules.
J. Clin. Invest., 69:152-161.
- Bourrillon R., Razafimahaleo E., and Parraudeau M., (1968):
Study of structure of the alpha-2- macroglobulin in human.
Biochim. Biophys. Acta., 154:405.
- Breit N., and Penny R., (1980):
The role of Alpha-1 protease inhibitor (Alpha-1 Antitrypsin) in the regulation of immunologic, inflammatory reactions.
Aust. N.Z. J. Med., 10:449-453.
- Brod J., and Sirota J.H., (1948):
Determination of serum and urine creatinine by Jaffe reaction.
J. Clin. Invest. 27:645.
- Buffone G.J., Lewis S.A., and Iose Fsohn M., (1978):
Chemical and immunological measurment of total iron binding capacity.
Clin. Chem. 24:1788-1791.
- Camerson J.S., and Blandford G., (1966):
The simple assessment of selectivity in heavy proteinuria.
Lancet 2:242.
- Carrie B.J., and Myers B.D., (1980):
Proteinuria and functional characteristics of the glomerular barrier in diabetic nephropathy.
Kidney int. May., 17(5):669-676.
- Caulfield J.P., and Farquhar M.G., (1975):
The permeability of glomerular capillaries of aminonucleoside nephrotic rats to graded dextrans.
J. Exp. Med., 142:61.
- Carlston A., and Johansson B.G., (1983):
As quoted by Gradwohl's clinical laboratory methods and diagnosis 8th edition., (1986).
The C.V. Mosby company St. Louis, Toronto, London.

- Cawley L.P.P., (1969):
Electrophoresis and Immuno-electrophoresis. Vol. I.
Little Bown and Co., Boston.
- Chang R.L.S., Ueki I.F., and Troy J.L., (1975):
Permeability of the glomerular capillary wall to
macromolecules. Experimental studies in rat using dextran.
Biophys. J., 15:887.
- Chan S.K., and Rees D.C., (1975):
Molecular basis for the alpha protease inhibitor deficiency.
Nature, 255:240.
- Chiu K.M., Mortensen R.F., Osmand A.P., and Gewarz H., (1977):
Interaction of alpha 1 acid glycoprotein with immune
system.
Immunology, 32:997.
- Christy M., Green A., Christau B., (1979):
Studies of the HLA system and IDDM.
Diabetes care., 2:209.
- Collens J.A., (1982):
Clinical judgment versus the laboratory.
N. Eng. J. Med., 306:987.
- Cooper M.E., Brien R.C., and Murray, (1989):
Intermittent diabetic microalbuminuria association with
blood pressure, glycaemic control and protein intake.
J. Diabetic Complications, 3(2):92.
- Cooper M.E., Frauman A.O., Brien R.C., et al., (1988):
Progression of proteinuria in type I and type II diabetes.
Diabetic Med. May., 5:361-368.
- Cooper M.D., and Lawton A.R., (1974):
The development of immune system.
Sci. Amr., 231:58.
- Cortney M.A., Swan L.L., and Weiss D.D., (1970):
Renal tubular protein absorption in the rat.
J. Clin. Invest., 49:1-4.
- Courtesy P.J., Lombart C., Feldmann G., Moguilevsky N., and
Rogier E., (1981):
Synchronous increase of four acute phase protein synthesized
by the same hepatocytes during the inflammatory reaction.
International Academy of pathology, 44(2):105.
- Couser W.G., Hoyer J.R., and Stilmant M.M., (1978):
Effect of aminonucleoside nephrosis on immune complex
localization in autologous immune complex nephropathy in
rats.
J. Clin. Invest., 61:561-572.

- Cudworth A.G., Gamble D.R., White G.B., et al., (1977):
Etiology of diabetes.
Lancet 1:385.
- Cudworth A.G., and Woodrow T.C., (1976):
Genetic susceptibility in type I diabetes.
Diabetologia, 12:385.
- Daniles J.e., (1975):
Carrier protein abnormalities in serum protein abnormalities. Diagnostic and chemical aspects.
Ed. by S.E. Ritzmann and J.c. Daniles, Boston, Little Brown and company P. 213-242.
- Deckert T., Poulsen J.E., and Larsen M., (1978):
Prognosis of diabetes with diabetes onset before the age of thirty-one.
Diabetologia., 14:363.
- Dermer G.B., Silverman. L.M., Gandller S.J., and Tvokes Z.A., (1980):
Incidence of a split alpha-2- glycoprotein band in electrophoresis pattern for serum of adenocarcinoma patients.
Clin. Chem., 26:392.
- Dewardner H.E., (1985):
The kidney, an outline of the normal and abnormal function 5th edition.
Language Booksociety Churchill Livingstone.
- Dickson I.R., and Manning C.W., (1972)
Changes in serum alpha-2-macroglobulin levels in children after major bone surgery.
Lancet, 2:484.
- Dillard M.G., Perce A.S., Pollak V., and Eavel Boreisha I., (1971):
Proteinuria and renal protein clearances in patients with renal tubular disorders.
J. Lab. Clin. Med., 78:203-215.
- Divall G.B., (1984):
Methods in molecular biology.
Vol. I, Ed. by Walker J. M., 12:105.
- Drysdale J.W., (1975):
Methods of protein separation, Vol. I.
Ed. by Catsim polas, New York plenum.
- Dunand P., Cropp C.J.A., and Middle tom E., (1976):
Severe obstructive lung disease in a 14 years old girl with alpha-1 antitrypsin deficiency.
J. Allergy Clin. immunol., 57:615-622.

- Edelman G.M., (1973):
Antibody structure and molecular immunology.
Science, 180:830.
- Erikson S., (1965):
Studies in α_1 antitrypsin deficiency.
Acta Med. Scand., 177:1-15.
- Fagerhal M.K., and Brand M., (1965):
Serum Prealbumin: Polymorphism in man.
Science., 149:986.
- Fajans S.S., (1981):
Etiological aspects of types of diabetes.
Diabetes Care., 4:1.
- Fajans S.S., Cloutier M.C., and Growther R.L., (1978):
Clinical and etiological heterogeneity of idiopathic
diabetes mellitus.
diabetes., 27:1112.
- Fang K.K., and Kalant N., (1972):
Phospholipid of the rat glomerular basement membrane in
experimental nephrosis.
Biochem. J., 129:733-774.
- Fawcett J.K., and Scott J.E., (1960):
Determination of serum urea by the urease method using the
Berthelot reaction.
J. Clin. Path., 13:156.
- First M.R., Patel V.B., Pesce A.J., et al., (1978):
Albumin excretion by the kidney: The effect of osmotic
diuresis.
Nephron., 20:171-175.
- Fleischman J.B., Porter R.R., and Press R.M., (1963):
The arrangement of the peptide chains in gamma-globulin.
Biochem. J., 88:220.
- Flier J.S., Khan C.R., and Roth J., (1979):
Receptors, antireceptor antibodies and mechanism of insulin
resistance.
N. Engl. J. Med., 300:413.
- Gambardella S., Morano S., and Cancelli, (1989):
Urinary IgG₄: An additional parameter in characterizing
patients with diabetic nephropathy.
Diabetes Res., 10(4):153.
- Ganrot P.O., and Schersten B., (1967):
Serum alpha 2 macroglobulin concentration and its variation
with age and sex.
Clin. Chim. Act., 15:113.

- Gatling W., Knight C. and Hibl R.D., (1985):
Renal function in diabetes.
Lancet 1:875.
- Gatherings W.E., Lorwton A.R., and Cooper M.D., (1977):
Immunofluorescent studies of the development of pre-B-cell,
B-lymphocytes, and immunoglobulin isotype diversity in
human.
Eur. J. Immunol., 7:804.
- Ghalioungi P., and Ghareeb A., (1978):
Diabetes mellitus: Endocrine disease and some metabolic
disorders. Vol. I.
Dar El-Alam El-Arabi, Cairo, Egypt.
- Ghanem M.H., Guirgis F.K., and El-sawy M., (1973):
Serum glycoprotein in schistosomal hepatic fibrosis.
J. Trop. Med. Hyg., 76:13.
- Gitlin, and Biasucci, (1969):
Development of gamma globulin, ceruloplasmin transferrin,
baptoglobin, fibrinogen, α_2 antitrypsin, orosomucoid, B-lipo
protein, α_2 macroglobulin and prealbumin in the human
conceptus.
J. Clin. Invest., 48:1433.
- Glad C., Greeb A.O., and Grimsberg., (1983): As quoted by Grad
Wohl's., (1986):
Clinical laboratory methods and diagnosis, 8th edition.
The C.V. Mosby company St. Louis- Toronto. London.
- Goetz F.C., Hartmann J.F., and Lazarow A., (1960):
Electron microscopy of the human glomerulus in early
diabetes.
J. Clin. invest., 39:991.
- Goldman A.S., and Goldblum R.M., (1977):
Primary deficiency in humoral immunity.
Pediat. clin., North Amer., 24:277.
- Goldstein I.M., Ros D., Wissman G., and Kaplan H., (1975):
Complement and immunoglobulins stimulate superoxide produc-
tion by human leukocytes independently of phagocytosis.
J. of clin. Invest., 65:1155-1163.
- Gordon S., (1978):
Regulation of enzyme secretion by mononuclear phagocytes:
Studies with macrophage plasminogen activator and lysozyme.
Fed. proc., 37:2754.
- Graham R.C., and Karnovsky M.J., (1966):
The early signs of absorption of injected horseradish
peroxidase in the proximal tubules of mouse kidney:
Ultrastructure cytochemistry by a new technique.
J. Histochem. cytochem. 14:291.

- Grundbacher F.J., and Massie A.E., (1985):
Level of immunoglobulin A. in black and white individuals.
J. Aller. Clin. Immunol., 75:656.
- Guthrow C.E., Morris M.A., Day J.F., et al., (1979):
Enhanced non enzymatic glucosylation of human serum albumin
in diabetes mellitus.
Proc. Nat. Acad. Sci. U.S.A., 76:4258.
- Hardwicke J., and Squire J.R., (1955):
The relationship between plasma albumin concentration and
protein excretion in patients with proteinuria.
J. Clin. Sci., 14:509-530.
- Harris E.D.Jr., (1987):
Disorders of the kidney and urinary tract in Harrison's
principles of internal medicine: Vol.II Ed. by Kelley W.N.,
Harris E.D. and Raddy S.
C.B. Saunders Company P.1139.
- Harrison J.F., and Barnes A.D., (1970):
The urinary excretion of lysozyme.
Clin. Sci., 38:533-547.
- Harrison J.F., and Blainey J.D., (1968):
Low molecular weight proteinuria in chronic renal disease.
Clin. Sci., 33:381.
- Harpel P.C., (1973):
Studies on human plasma alpha-2-macroglobulin enzyme
interactions.
J. Exp. Med., 138:508.
- Hause G.R., (1974):
Water transport in cell and tissues. Vol.II.
The Williams and Wilkins Co., Baltimore.
- Herve M., Garnier A., Tosi L., and Steinbuch M., (1981):
Spectroscopic and photoreduction studies on copper in
Ceruloplasmin.
European J. Bioch., 116:177-183.
- Horne C.H.W., Briggs J.D., Howie P.W., and Kennedy A.C., (1972):
Serum alpha-2-macroglobulins in renal disease and preecl-
ampsia.
J. Clin. Path., 25:590.
- Hosoi H., Asakura Y., and Irie M., (1989):
The 24 hours urinary excretion of albumin, beta-2-microglo-
bulin and N-acetyl B-D glucosaminidase activity in children
with IDDM.
Nippon-IKa-Daigaku zasshi., 56(3):241.
- Housley J., (1968):
Alpha-2-macroglobulin levels in disease in man.
J. Clin. pathol., 21:27.

- Hoyer J.R., Elema J.D., and Verpier R.L., (1976):
 Unilateral renal disease in rat: Glomerular mesangial uptake
 of colloidal carbon in unilateral aminonucleoside nephrosis
 and nephrotoxic serum nephritis.
 Lab. invest., 34:250-255.
- Ikenaka T., Ishiguro M., Isemura S., et al., (1971):
 The amino acid sequence of the carboxy-terminal CNBR-
 Fragment of alpha-1-acid glycoprotein.
 Biochemical and biophysical research communica., 42(6):1142.
- Irvine W.J., Gray R.S., McCallum C.J., and Duncan L.J., (1977):
 Clinical and pathogenic significance of pancreatic islet
 cell antibodies in diabetic treated with oral hypoglycaemic
 agents.
 Lancet, 1:1025.
- Ishii K., Koyama A., and Kobayashi, (1989):
 A study of microalbuminuria in patients with diabetes
 mellitus.
 Nippon-Jinzo-Gakkaishi., 31(8):815.
- Ishizaka T., Depernado R., and Tomioke H., (1972):
 Identification of basophil granulocytes as a site of
 allergic histamine release.
 J. immunol., 108:1000.
- Jacob A., (1973):
 Digestive factors in iron absorption.
 Progress in Gastroenterology., 2:221.
- Jarrett R.J., and AL-sayeach H., (1978):
 Detection of patients at risk of diabetic complications.
 Diabetologia, 15:243.
- Jensen K., (1970):
 Metabolism of bence Jones proteins in non myeloma patients
 with normal renal function and in renal disease.
 Scand. J. clin. Lab. invest., 25:281-289.
- Jeppson J.O., Laurell C.B., and Prenzen B., (1979): As quoted by
 Gradwohl's., (1986):
 Clinical laboratory methods and diagnosis 8th edition.
 The C.V. Mosby company- St. Louis-Toronto.
- Jerry L.M., and Kunkel H.G., (1974):
 Special characteristics of the IgA₂ subclass.
 Adv. Exp. Med. Biol., 45:151.
- Jerums G., Allen I.J., and Cooper M.E., (1989):
 Triphasic changes in selectivity with increasing proteinuria
 in type 1 and and type 2 diabetes.
 Diabetic- Med. 6(9):772.

- Johansson B.G., (1972):
 Agarose gel electrophoresis.
 Scand. J.clin. Lab. Invest., 29 (124):7.
- Jones B.M., and Hua A.S.P., (1980):
 A simple radial diffusion technique for measuring
 selectivity of proteinuria.
 J. Clin. Pathol. Jan., 33(6):598-599.
- Jones D.B., (1969):
 Mucosubstances of the glomerulus.
 Lab invest., 21:119-125.
- Jones J.M., (1972):
 Thiol reduction of human alpha-2-macroglobulin.
 Biochem. J., 127:187.
- Kettmann R., (1982): As quoted by Varley H., (1987):
 Practical clinical biochemistry 5th edition.
 Willian Heinman Medical books Ltd. London.
- Katz J., Rosenfold S., and Sellers A.L., (1960):
 Role of the kidney in plasma albumin catabolism.
 Amer. J. Physiol., 198:814-818.
- Keen H., and Chlouverakisc., (1963):
 An immunoassay method for urinary albumin at low
 concentrations.
 Lancet 2:913.
- Klein R., Klein B.E., Moss S., and Demetes D.L., (1988):
 Proteinuria in diabetes..
 Arch. intern. Med., Jan., 148(1):181-186.
- Knight K.L., and Dray S., (1968):
 Contribution of alleic genes to the formation of individual
 α_2 macroglobulin molecules.
 Biochemistry, 7:3830.
- Kohn J., J.G., (1958):
 Electrophoresis on cellulose acetate.
 Clin.Chem.Acta., 3:450.
- Koj A., Regoez E., and Toews C.J., (1978):
 Synthesis of antithrombin III and alpha-1- antitrypsin by
 the perfused rat liver.
 Biochem. Biophys. Acta, 539:496-504.
- Kreisberg J.I., and Karnovsky M.J., (1977):
 Rapid and focal loss of negative charge following
 adminstration of nephrotic serum in rats.
 Kidney int., 12:515.

- Latta H., Johnston W.H., and Stanely T.M., (1975):
Sialoglycoproteins and filtration barriers in the glomerular capillary wall.
J. ultrastruct. Res., 51:354.
- Laurell C.B., (1966):
Quantitative estimation of proteins by electrophoresis in agarose gel containing antibodies.
J. Anal. Biochem., 15:45-52.
- Laurell C.B., (1972):
Electrophoretic analysis of plasma proteins.
Scand. J. Clin. Lab. Invest., 30:233.
- Laurell C.B., (1985):
Acute phase proteins, a group of prospective proteins in Recent advances in clinical biochemistry.
Ed. by C.P. Price, K.G.M.M., Alberta, Churchill Livingstone, Edinburgh, London, Melbourne and New York.P. 108-124.
- Laurell C.B., and Jeppson J.O., (1975):
The plasma protein structure, function, and genetic control, 2nd edition. Ed. by F.W. Putnam Vol.I., New York. Academic Press. P.229.
- Lee-foon Chio, and Chory-Jin On, (1979):
Changes in serum alpha-1-antitrypsin, alpha-1-acid glycoprotein in patients with malignant hepatocellular carcinoma.
Cancer, 43:596-604.
- Lesile R.D.G., Lazarus N.R., and Verganil D., (1989):
Etiology of insulin dependent diabetes.
Br. Med. Bull., 45(1):58.
- Mamaeva G.G., Servo V.V., Spesivtseva V.G., and Muklrun N.A., (1985):
Clinico-morphophological characteristics of diabetic nephro-angiopathy in insulin dependent diabetes mellitus.
Am. J. Kidney-Dis., 57(12):12.
- Marre M., and Krempf M., (1987):
Microalbuminuria in diabetics.
Diabet. Metab. Jun., 13:232-239.
- Martin T.J., Melick R.A., and Luise M.D.E., (1969):
The effect of nephrectomy on the metabolism of labelled parathyroid hormone.
Clin. Sci., 37:137-142.
- Martin P., Walton C., and Chapman C., (1990):
Increased urinary excretion of transferrin in children with type I diabetes mellitus.
Diabetic Med., 7(1):35.

- Maunsbach A.B., (1966):
Absorption of ferritin by rat kidney proximal tubule cells.
J. Ultrastructure. Res., 16:1-12.
- McCormick J.R., Sonksen P.H., and Egdahl R.H., (1969):
Renal handling of insulin and growth hormone in hemorrhagic shock.
Surgery., 66:175-180.
- McLean P.R., and Robson J.S., (1967):
A simple method for determining selectivity of proteinuria.
Lancet. 1:539.
- Michael A.F., Blau E., and Vernier R.L., (1970):
Glomerular polyanion. Alteration in aminonucleosides nephrosis.
Lab. Invest., 23:649-657.
- Misra R.P., (1973):
Glomerular basement membrane, antigen of masugi nephritis.
Immunology, 25:267-280.
- Misra R.P., and Berman L.B., (1979):
The molecular basis of proteinuria.
Jama., 242(8):757.
- Mogensen C.E., (1986):
Microalbuminuria in "Recent advances in diabetes". Ed. by Nattrass M., Churchill living stone, Edinburgh, London Melbourne and New York.
P.139-150.
- Mogensen C.E., (1987):
Microalbuminuria as a predictor of clinical diabetic nephropathy.
Kidney Int., 31:673.
- Morgan E.M., (1974):
Transferrin and transferrin iron in biochemistry and medicine.
2nd edition. Ed. by A.Jacobs and M.Wrwod (London, Academic press).P.29.
- Morris P.J., Irvine W.J., Gray R.S., et al., (1976):
HLA and pancreatic islet cell antibodies in diabetes.
Lancet, 2:652.
- Morse J.o., Lebowitz M.D., Kundson R.J., and Burrow S.B., (1977):
Relation of protease inhibitor phenotypes to obstructive lung disease in a community.
New. Eng. J. Med., 296:1190-1194.
- Mosher D.F., and Wing, D.A., (1976):
Synthesis and secretion of α_2 macroglobulin by cultured human fibroblasts.
J. Exp. Med., 143:462.

- Mothony J.F., Wallace E.G., and Allen D.M., (1978):
A non barbital buffer for Immuno electrophoresis and zone electrophoresis in agarose gels.
J.clin. chem., 24:1825.
- Nakamura Y., and Myers. B.D., (1988):
Charge selectivity of proteinuria in diabetic glomerulopathy.
Diabetes., Sept., 37(9):1202-1211.
- National Diabetes Data Group., (1979):
Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance.
Diabetes., 28:1039.
- Natvig J.B., and Kunkel H.G.G., (1973):
Human immunoglobulins classes, subclasses genetic variants, and Idiotypes.
Adv. Immunol., 16:1.
- Nelson R.G., Petti H.D.J., Carraher M.J., Baried H.R., and Knowler W.C., (1988):
Effect of proteinuria on mortality in non-insulin dependent diabetes mellitus.
Diabetes, Nov., 37(11):1449-1504.
- Nerupt, Platz-P., and Ryder L.P., (1978):
HLA, islet cell antibodies and types of diabetes.
Diabetes., 27 (sup.1):247.
- Oberling C.H., Gautier A., and Bernard W., (1951):
The structure of the renal glomerulus.
Presse. Med., 59:938.
- Parkes D., (1984):
Wilson's disease.
Br. Med. J., 288:1180.
- Parkman R., (1984):
Immunology in shaffer's disease of new born.
3rd edition. Ed. by Avery M.E., William H., and Tauesch J. W.B. Saunders Co. Philadelphia, London, and Toronto. P.120-221.
- Parving H.H., Oxenboll B., and Svendsen P.A., (1982):
Detection of patients at risk of developing diabetic nephropathy. A longitudinal study of urinary albumin excretion.
- Parving H.H., Stroper W., and Brein R., (1976):
The effect of metabolic regulation on microvascular permeability to small and large molecules in short term Juvenile diabetes.
Diabetologia., 12:161.

- Patwardhan U.N., Maghrabi R.H., Mouser W., et al., (1971):
 Serum glycoprotein in protein-calorie deficiency disease.
 The Am. J. of Clin. Nutrition., 24:906.
 Acta Endocrinology., 100:550.
- Petersson P.A., Evrin P.E., and Berggard I., (1969):
 Differentiation of glomerular, tubular, and normal protein-
 uria: Determination of urinary excretion of B₂ microglo-
 bulin, albumin and total proteins.
 J. Clin. Invest., 84:1189-1198.
- Porte D., and Halter J.B., (1974):
 The endocrine pancreas and diabetes mellitus in text book
 of endocrinology 5th edition Ed. by Williams R.H.,
 W.B.Saunders company, Philadelphia, London, Toronto.P. 555.
- Putnam F.W., (1975):
 The plasma proteins structure, function and genetic control.
 2nd edition Ed. by F.W. Putnam, Vol.1., Academic Press.,
 P.81-83.
- Reinhold J.G., (1953):
 Standard methods in clinical chemistry, Vol.I. Ed. by M.
 Reiner, Academic Press., New york P.88.
- Rennke H.G., and Venkatachalam M.A., (1977):
 Glomerular permeability: in vivo tracer studies with poly-
 anionic and polycationic ferritins.
 Kid. Int., 11:44-53.
- Retting B., Teuch S.M., (1984):
 The incidence of end-stage renal disease in type I and type
 II diabetes mellitus.
 Diabetic nephropathy 3:26.
- Rifkin D.B., Leob J.M., and Move G., (1974):
 Properties of plasminogen activators formed by neoplastic
 human cell cultures.
 J. Exp. Med., 139:1317.
- Ringden O., Persson U., Johansson S.G., and Gahrton G., (1983):
 Markedly elevated serum IgE levels following allogenic and
 syngenic bone marrow transplantation.
 Blood., 61:1190.
- Robert F. pihs., (1974):
 The text book of physiology of the kidney and body fluids.
 3rd edition. Ed. by Vippincott company, philadelphia,
 Toronto., P.151-160.
- Rothschild M.A., Oratz M., and Schreiber S.S., (1972):
 Albumin synthesis.
 N. Engl. J. Med., 286:748-816.

- Roy L.P., Vernier R.L., and Michael A.F., (1972):
Effect of protein load proteinuria on glomerular polyanion.
Proc. Soc. Exp. Biol. Med., 141:870-874.
- Rubenstein A.H., and Spitz J., (1968):
Role of the kidney in insulin metabolism and excretion.
Diabetes., 17:161-169.
- Ruiz M., Rujutanavin R., Young R.A., et al., (1982):
Familial dysalbuminemic hyperthyroxinemia. A syndrome that
can be confused with thyrotoxicosis.
N. Eng. J. Med., 306:635.
- Salmon S., Utiger R., Parker M., and Reichlin S., (1962):
The fate of I^{125} labeled human growth hormone in the rabbit.
Endocrinology., 70:459-464.
- Scaronia F., Martino P., and Goi M., (1981):
Findings on proteinuria in selected diabetic patients.
Relation to microangiopathy and its prognostic significance.
Minerva Med., 72(9):561.
- Schultz H.E., (1960):
Molecular biology of human proteins.
Int. Med. J.1:255.
- Schultz H.E., (1962):
Influence of bound sialic acid on electrophoresis mobility
of human serum proteins.
Arch. Biochem. Biophys., 1:290.
- Schultz H.E., and Heremans J.F., (1966):
The urinary proteins in molecular biology of human proteins
with special reference to plasma proteins.
J.exp. Med. 62:699.
- Sharp H.L., (1971):
Alpha-1- antitrypsin deficiency.
Hosp. Pract., 6:83.
- Sharp H.L., Bridges R.A., Krivit W., and Frier E.F., (1969):
Cirrhrosis associated with α_1 antitrypsin deficiency: A
previously unrecognized inherited disorder.
J. Lab. Clin. Med. 73:934.
- Shibata K., Okubo H., and Ishibashi H., (1977):
Rat α_1 acid glycoprotein-purification and immunological
estimation of its serum concentration.
Biochem. Bioph. Acta., 495(1):37.
- Simanwitz M.D., Mac Gregor W.G., and Hobbs J.R., (1973):
Proteinuria in preeclampsia.
J. of Obst and Gynec. of British common wealth., 80:103-108.

- Slater L., Carter P.M., and Hobbz J.R., (1975):
 Measurment of albumin in the sera of patients.
 Ann. Clin. Biochem., 12:33.
- Smith B.J., (1984):
 Methods in molecular biology Vol.I. Ed. by Walker J.M.
 Humana press clifton, New Jersey.
- Spiegelberg H.L., (1974):
 Biological activities of immunoglobulins of different
 classes and subclasses.
 Adv. immunol., 19:259.
- Spiro R.G., (1976):
 Search for a biochemical basis of diabetic microangiopathy.
 Diabetologia, 12:1-14.
- Staprsans I., Anderson C.D., Lurz P.W., and Felts J.M., (1980):
 Separation of a lipoprotein lipase cofactor from the alpha-
 1- acid glycoprotein fraction from the urin of nephrotic
 patients.
 Biochem. Bioph. Acta., 617(3):514.
- Strober W., and Waldmann T.A., (1974):
 The role of the kidney in the metabolism of plasma proteins.
 Nephron., 13:35-66.
- Sun T., (1982):
 Physiology of immunoglobulins: Diagnostic and clinical
 aspects.
 3rd edition. Ed. by Allan R.Liss, New York., P .29.
- Sun T., and Lien Y.Y., (1980):
 Reversed polarity- immunoelectrophoresis.
 J. Clin. Chem., 26:1763.
- Talamo R.C., (1975):
 Basic and clinical aspects of the alpha 1- antitrypsin.
 Pediatric., 56:91.
- Taylor R., (1989):
 Etiology of non insulin dependent diabetes.
 Br. Med. Bull., 45(1):73.
- Thaler M.S., Klausner R.D., and Cohen H.J., (1977):
 Immunoglobulin molecule in medical immunology.Vol.I.
 Lippincott J.B. company, Philadelphia, Toronto., P. 35.
- Tietz N.W., (1987):
 Fundamental of clinical chemistry. 3rd edition.
 Sunders Co., philadelphia, London, Toronto. P 312-324.
- Trinder P., (1969): As gouted by Varley H., (1987):
 Practical clinical biochemistry 5th edition.
 William Heinmann Medical books Ltd. London.

- Tunstall A., and James K., (1974):
Preliminary studies of the synthesis of alpha-2-macroglobulin by human lymphocytes in Vitro.
Clin. Exp. immunol., 17:697.
- Valdrol-Hansen F., Jensen T., Borch, Johnsen K., and Deckert T., (1987):
Cardiovascular risk factors in type I diabetic patient with and without proteinuria.
Acta. Med. Scand., 222(5):439 444.
- Vergier M.F., and Larpent L., (1988):
Hypertensive type I diabetics have a higher level of micro-albuminuria than normotensive irrespective of blood glucose equilibrium.
Am. Med. Interne., 139:2513.
- Vernier R.L., Rapermaster B.W., and Good R.A., (1959):
Aminonucleoside nephrosis.
J. Exp. Med., 109:115-126.
- Vibertti G., and Keen H., (1984):
The pattern of proteinuria in diabetes mellitus.
Diabetes., 33:686-692.
- Vibertti G.C., Keen H., and Wiseman M.J., (1987):
Raised arterial pressure in parents of proteinuric insulin dependent diabetics.
Br. Med. J., 295:515-517.
- Vibertti G.C., Hill R.D., and Jarrett., (1982):
Microalbuminuria as predictor of clinical nephropathy in insulin dependent diabetes mellitus.
Lancet., 1:1430-1432.
- Vibertti G.C., Pickup J.C., and Jarrett R.J., (1979):
Effect of control of blood glucose on urinary excretion of albumin and B₂ microglobulin in insulin dependent diabetes.
N. Engl. Med., 300:638-641.
- Vittinghus E., and Mogensen C.E., (1982):
Grade exercise and protein excretion in diabetic man and the effect of insulin treatment.
Kidney Int., 21:725-729.
- Walker J.M., (1984):
Methods in molecular biology. Vol.I.(Proteins).
Humann Press. Clifton, New Jersey.
- Washburn E.J., and Allen N.C., (1960):
Idiopathic tropical splenomegaly syndrome.
Brit. Med. J., 4:793.

- Welborn T.A., (1984):
"The definition of diabetes" from recent advances in diabetes.
2nd edition. Ed. by Molcolm M., Julio U.S., Churchill livingstone Edinburgh London. P.15.
- W.H.O., (1980):
Expert Committee on diabetes mellitus. Second report.
W.H.O. Teck. Resp. Ser. No. 646.
- Wibell L., and Karlsson, (1976):
Urinary excretion of B₂ microglobulin after the induction of a diuresis.
Nephron., 17:343-352.
- Wochner R.D., Strober W., and Waldmann T.A., (1967):
The role of the kidney in the catabolism of bence Jones proteins and immunoglobulin fragments.
J. Exp. Med., 126:207-221.
- Woldmann T.A., Gordon R.S., and Rose W., (1964):
Studies on the metabolism of serum proteins and lipids in patients with analbuminemia.
Am. J. Med, 37:960.
- Woldmann T.A., Strober W., and Mogielnicki R.P., (1972):
The renal handling of low molecular weight proteins.
J. Clin. Invest., 51:2162-2174.
- Woo J., Floyd M., and Cannon D.C., (1981):
Albumin and B₂ microglobulin radioimmunoassay in differentiating glomerular and tubular disease.
Clin. Chem., 27:709-713.
- Yoshida.A., Ewing.C., Wessels, Lieberman J., and Giadulis L., (1977):
Molecular abnormality of variant human alpha-1-antitrypsin.
Am.J. Hum. Genetic., 29:233.