## References

Abouzahr C: Antepartum and postpartum haemorrhage. In: Murray CJ, Lopez AD, eds. Health Dimensions of Sex and Reproduction. Boston, Mass: *Harvard University Press*; 1998: 172-4.

Albers LL, Schiff M, Gorwoda JG. The Length of Active Labor in Normal Pregnancies. Obstet Gynecol 1996; 87:355-9.

Alexander JM, Grant AM, Campbell MJ. Randomised controlled trial of breast shells and Hoffman's exercises for inverted and non-protractile nipples. *British medical journal*, 1992; 304:1030-1032.

Alisjahbana A, Williams C, Dharmayanti R, Hermawan D, Kwast BE, Koblinsky M. An integrated village maternity service to improve referral patterns in a rural area in West-Java. Int J Gynecol Obstet 2001; 48 (Suppl): S83-S94.

Allahbadia GN, Vaidya PR. Why deliver in the supine position? Aust NZ J Obstet Gynaecol 1992; 32/2:104-106.

Alpay F, Sarici S, Tosuncuk HD, Serdar MA, Inanc, N, Go"kc, ay E. The value of first-day bilirubin measurement in predicting the development of significant hyperbilirubinemia in healthy term newborns. *Pediatrics*. 2000;106(2).

American Academy of Pediatrics, Breastfeeding Promotion in Physicians' Office Practices Program. Elk Grove Village, IL: American Academy of Pediatrics; 2001, 2004.

American Academy of Pediatrics, Committee on Drugs. Transfer of drugs and other chemicals into human milk. *Pediatrics*. 2001;108: 776–789.

American Academy of Pediatrics, Committee on Practice and Ambulatory Medicine. Recommendations for preventive pediatric health care. *Pediatrics*. 2000;105:645–646.

American Academy of Pediatrics, Committee on Fetus and Newborn. Controversies concerning vitamin K and the newborn. *Pediatrics*. 2003; 112:191–192.

American Academy of Pediatrics, Medical Home Initiatives for Children With Special Needs Project Advisory Committee. The medical home. *Pediatrics*. 2002;110:184–186.

American Academy of Pediatrics, Provisional Committee for Quality Improvement and Subcommittee on Hyperbilirubinemia. Practice parameter: management of hyperbilirubinemia in the healthy term newborn. *Pediatrics*. 1994;94:558–562

American Academy of Pediatrics, Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics*. 2004;114:297–316.

American Academy of Pediatrics. Work Group on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 1997;100: 1035–1039.

American Association of Blood Banks Technical Manual Committee. Perinatal issues in transfusion practice. In: Brecher M, ed. *Technical Manual*. Bethesda, MD: American Association of Blood Banks; 2002: 497-515

American Dietetic Association. Position of the American Dietetic Association: breaking the barriers to breastfeeding. *J Am Diet Assoc.* 2001;101:1213–1220.

Anderson EW, Johnstone BM, Remax DT. Breastfeeding and cognitive development: a meta-analysis. A MJ Clin Nutr 1999;70: 525-535.

Anderson PO. Alcohol and breastfeeding. J Hum Lact. 1995;11:321-323.

Arnold LDW. Recommendations for collection, storage and handling of a mother's milk for her owninfant in the hospital setting. Denver, CO: Human Milk Banking Association of North America, 1999.

Ashraf RN, Jalil F, Aperia A, Lindblad BS. Additional water is not needed for healthy breast-fed babies in a hot climate. *Acta Paediatr*. 1993;82:1007-1011.

Auerbach KG, Gartner LM. Breast feeding and human milk: their association with jaundice in the neonate. Clin Perinatol 1987; 14: 89–107.

Baby-Friendly USA. U.S. UNICEF Baby-Friendly Hospital Initiative materials. Baby-Friendly USA, Sandwich, MA, 2000.

Bachrach VR, Schwarz E, Bachrach LR. Breastfeeding and the risk of hospitalization for respiratory disease in infancy: a meta-analysis. Arch Pediatr Adolesc Med. 2003;157:237–243

Balistreri WF. Neonatal cholestasis J Pediatr, 1985; 106:171-184

Barrett JFR, Savage J, Phillips K, Lilford RJ. Randomized trial of amniotomy in labour vs the intention to leave membranes intact until the second stage. Br J Obstet Gynaecol 1992; 99:5-10.

Bastian H, Lancaster PAL. Home births in Australia 1985-1987, 1988-1990. Sidney, AIHW National Perinatal Statistics Unit 1992.

**Benaron DA, Bowen FW**. Variation in serum bilirubin with illness during first few days of life: possible evidence for a free radical scavenging reaction. *Lancet* 1991; 338(8759): 78-81

Berk MA, Mimouni F, Miodovnik M, Hertzberg V, Valuck J. Macrosomia in infants of insulin-dependent diabetic mothers. *Pediatrics*. 1989; 83:1029–1034.

Bernshow N. A Mother's Guide to Milk Expression and Breast Pumps. Schaumburg, Illinois: La Leche League International, 1996. Publication No. 30.

Bernstein HH, Spino C, Baker A, Slora EJ, Touloukian CL, McCormick MC. Postpartum discharge: do varying perceptions of readiness impact health outcomes? *Ambul Pediatr*. 2002;2:388-395.

Bhardwaj N, Kukade JA, Patil S, Bhardwaj S. Randomised controlled trial on modified squatting position of delivery. *Indian J Maternal and Child Health* 1995; 6(2):33-39.

**Bhutani VK, Johnson L, Sivieri EM.** Predictive ability of a predischarge hour-specific serum bilirubin for subsequent significant hyperbilirubinemia in healthy term and near-term newborns. *Pediatrics*. 1999;103: 6–14

**Biancuzzo M.** Selecting pumps for breastfeeding mothers. *JOGNN* 1999; 28:417-26,.

Biringer A, Davies B, Nimrod C, Sternberg C, Youens W. Attaining and maintaining best practices in the use of caesarean sections. An analysis of four Ontario hospitals. Report of the Caesarean Section Working Group of the Ontario Women's Health Council. Ottawa; 2000

Blair PS, Fleming PJ, Smith IJ, et al. Babies sleeping with parents: case-control study of factors influencing the risk of the sudden infant death syndrome. *BMJ*. 1999;319:1457–1462.

Bloomer JR, Barret PV, Rodkey L, et al. Studies on the mechanism of fasting hyperbilirubinemia. Gastroenterology. 1981;61:479-487.

Bornmann, Pricilla G. "Legal Considerations and the Lactation Consultant-USA." New York: Avery, 1996. Unit 3 of Lactation Consultant Series. La Leche League International, Inc. 18 units.

Boulvain M, Fraser W D, Brisson-Carroll G, Faron G, Wollast E. Trial of labour after caesarean section in sub-Saharan Africa: a meta-analysis. British Journal of Obstetrics and Gynaecology 1997;104:1385-1390.

Braccire RC, Buonocores G, Garesi S, Berni S. Epidemiology of neonatal hyperbilirubinemia: A study of contributing factors. Acta Paediatr Scand. 1989;360

**Bradley W.** Newborns' and Mothers' *Health Protection Act of* 1996. *Pub* L No. 104-204.

**Bratlid D.** How bilirubin gets into the brain. Clin Perinatol. 1990;17: 449-465

Bull P, Barger J. Finger feeding with the SNS. Rental Roundup 1987; 4:2-3,.

Bullough CHW, Msuku RS, Karonde L. Early suckling and postpartum haemorrhage: controlled trial in deliveries by traditional birth attendants. Lancet 1989; ii:522-525.

Centers for Disease Control and Prevention. Recommendations for using fluoride to prevent and control dental caries in the United States. MMWR Recomm Rep. 2001;50(RR-14):1-42

- Carroli G, Bergel E: Umbilical vein injection for management of retained placenta (Cochrane Review). In: *The Cochrane Library*, Issue 2. Oxford, UK: Update Software. 2002.
- Chapman DJ, Perez-Escamilla R. Identification of risk factors for delayed onset of lactation. J Am Diet Assoc. 1999;99:450-454
- Chapman MG, Jones M, Spring JE, De Swiet M, Chamberlain GVP. The use of a birthroom: a randomized trial comparing delivery with that in a labour ward. *Br J Obstet Gynaecol* 1999; 93:182-187.
- Chen DC, Nommsen-Rivers L, Dewey KG, Lonnerdal B. Stress during labor and delivery and early lactation performance. Am J Clin Nutr. 1998;68:335-344.
- Chen SZ, Aisaka K, Mori H, Kigawa T. Effects of sitting position on uterine activity during labour. Obstet Gynecol 1997; 79:67-73.
- Cohen RJ, Brown KH, Canahuati J, Rivera LL, Dewey KG. Determinants of growth from birth to 12 months among breast-fed Honduran infants in relation to age of introduction of complementary foods. *Pediatrics*. 1995;96:504–510
- Cohen R, Mrtek MB, Mrtek RG. Comparison of maternal absenteeism and infant illness rates among breast-feeding and formula-feeding women in two corporations. Am J Health Promot 1995;10(2):148-153.
- Cooper WO, Atherton HD, Kahana M, Kotagal UR. Increased incidence of severe breastfeeding malnutrition and hypernatremia in a metropolitan area. *Pediatrics*. 1995;96:957-960.
- Coutsoudis A, Rollins N. Breast-feeding and HIV transmission: the jury is still out. J Pediatr Gastroenterol Nutr. 2003;36:434-442
- COx J P. Delivery alternatives in the term breech pregnancy. ICEA Review, 1988; vol. 12, no. 4.
- Crowley P, Elbourne DR, Ashhurst H, Garcia J, Murphy D, Duignan N. Delivery in an obstetric birth chair: a randomized controlled trial. Br J Obstet Gynaecol 1991; 98:667-674.
- Crowther C, Enkin M, Keirse MJNC, Brown I. Monitoring the progress of labour. In: Chalmers I et al (eds). Effective care in pregnancy and childbirth. Oxford, Oxford University Press, 1989.

Cunningham FG, Gant NF, Leveno KJ, et al: Conduct of normal labor and delivery. In: Williams Obstetrics. 21st ed. New York, NY: McGraw-Hill; 2001: 320-5.

**Da Silva OP et al.** Effect of domperidone on milk production in mothers of premature infants: a randomized double blind, placebo-controlled trial. *Can Med Assoc J* 164:117-21,2001.

Dansereau J, Joshi AK, Helewa ME, et al: Double-blind comparison of carbetocin versus oxytocin in prevention of uterine atony after cesarean section. Am J Obstet Gynecol 1999 Mar; 180(3 Pt 1): 670-6

De Carvallo M, Robertson S, Klaus M. Fecal bilirubin excretion and serun bilirubin concentration in breast – fed infants. *J pediatr* 1985;107:786-9.

De Carvelho, Monoel, Michael Hall and David Harvy. "Effects of Water Supplementation on Physiologic Jaundice in Breast-Fed Babies." Archives of Diseases of Childhood .1991; 568-569.

**Dettwyler KA.** A time to wean: the hominid blueprint for the natural age of weaning in modern human populations. In: Stuart-Macadam P, Dettwyler KA, eds. *Breastfeeding: Biocultural Perspectives*. Hawthorne, NY: Aldine de Gruyter; 1995:39-73

Dewey KG. Growth characteristics of breast-fed compared to formula-fed infants. Biol Neonate 1998;74:94-105.

**Dewey KG, Heinig MJ, Nommsen-Rivers LA.** Differences in morbidity between breast-fed and formula-fed infants. *J Pediatr* 1995;126(5):696-702.

Dewey KG, Peerson JM, Brown KH, Krebs NF, Michaelsen KF, Persson LA, Salmenpera L, Whitehead RG, Yeung DL, WHO Working Group on Infant Growth. Growth of breast fed infants deviates from current reference data: a pooled analysis of US, Canadian, and European Data Sets. *Pediatrics*. 1995;96:495-503.

**Dickersin K.** Pharmacological control of pain during labour. In: Chalmers et al (eds). Effective care in pregnancy and childbirth. Oxford, Oxford University Press 1989.

**Domellof M, Lonnerdal B, Abrams SA, Hernell O.** Iron absorption in breast-fed infants: effects of age, iron status, iron supplements, and complementary foods. Am J Clin Nutr. 2002;76:198–204

- **Doumas BT, Ashwood E.** Performance of bilirubin determinations in US laboratories—revisited. *Clin Chem.* 2004;50:190–194
- **Drayton S, Rees C.** They know what they're doing. Do nurses know why they give pregnant women enemas? *Nurs Mirror* 1984; 5:4-8.
- Duffy LC, Byers TE, Riepenhoff-Talty M, La Scolea LJ, Zielezny M, Orga PL. The effects of infant feeding on rotavirus-induced gastroenteritis: a prospective study. Am J Public Health 1986;76(3):259-263.
- Duncan B, Ey J, Holberg CJ, Wright AL, Martinez FD, Taussig LM. Exclusive breast-feeding for at least 4 months protects against otitis media. *Pediatrics* 1993;91(5):867-872.
- Eaton AP. Early postpartum discharge: recommendations from a preliminary report to Congress. *Pediatrics*. 2001;107:400-403
- Ebbesen F. Recurrence of kernicterus in term and near-term infants in Denmark. Acta Paediatr. 2000;89:1213-1217
- Eggert P, Stick C, Schroder H. On the distribution of irradiation intensity in phototherapy. Measurements of effective irradiance in an incubator. Eur J Pediatr. 1984;142:58-61
- Elbourne DR, Prendiville WJ, Carroli G, et al: Prophylactic use of oxytocin in the third stage of labour (Cochrane Review). In: The Cochrane Library, Issue 2. Oxford, UK: Update Software. 2002.
- Enkin M, Keirse M, Renfrew M, Neilson J: A Guide to Effective Care in Pregnancy and Childbirth. 3rd ed. Oxford, UK: Oxford University Press; 2000.
- Enkin M, Keirse, M, Renfrew, M and Neilson, J: Effective Care in Pregnancy and Childbirth, 1995; 2nd ed. pp142-143.
- Fathalla MF. FIGO Workshop Conclusion. Int J Gynecol Obstet 1992; 38(Suppl):S75-S77.
- Felsher BF, Carpio NM. Caloric intake and unconjugated hyperbilirubinemia. Gastroenterology. 1985;69:42-47

Fenton TR. Neonatal jaundice in AGM Campball, and N. McIntosh (Eds.), Forfar and Arniel's Textbook of Padiatrics 14<sup>th</sup> ed. 1992; Churchill Livinstone, Edinburgh – London – Madrid, pp. 240-5

Fewtrell MS, Lucas P, Collier S. Randomized trial comparing the efficacy of a novel manual breast pump with a standard electric breast pump in mothers of term infants *Pediatrics*, 2001; 107:1291 97.

Flamm, B L: Birth After Caesarean, the Medical Facts. 1992; P51.

Francome, C: Caesarean Birth in Britain (Supplement). 1994; p1.

Fraser WD, Turcot L, Krauss I, Brisson-Carrol G. Amniotomy for shortening spontaneous labour. Cochrane Database Syst Rev 2003;(2).

Fraser W, Hatem-Asmar M, Krauss I, Maillard F, Breart G, Blais R. Comparison of midwifery care to medical care in hospitals in the Quebec pilot projects study:clinical indicators. Can J Public Health 2000;91:1-11.

Freed GL, Clark SJ, Lohr JA, Sorenson JR. Pediatrician involvement in breast-feeding promotion: a national study of residents and practitioners. *Pediatrics*. 1995;96:490–494

Freed GL, Clark SJ, Sorenson J, Lohr JA, Cefalo R, Curtis P. National assessment of physicians' breastfeeding knowledge, attitudes, training, and experience. *JAMA*. 1995;273:472-476.

Gagnon AJ. Individual or group antenatal education for childbirth/parenthood. Cochrane Database Syst Rev 2001;(3).

Gale R, Seidman DS, Dollberg S, Stevenson DK. Epidemiology of neonatal jaundice in the Jerusalem population. J Pediatr Gastroenterol Nutr. 1990;10:82-86

Garcia FJ, Nager AL. Jaundice as an early diagnostic sign of urinary tract infection in infancy. *Pediatrics*. 2002;109:846-851

Garland JS, Alex C, Deacon JS, Raab K. Treatment of infants with indirect hyperbilirubinemia. Readmission to birth hospital vs nonbirth hospital. Arch Pediatr Adolesc Med. 1994;148:1317–1321

Gartner, Lawrence and Lawrence. "BFHI and BfHI: The Baby Friendly Hospital in the U.S." ABM News and Views: The Newsletter of The Academy of Breastfeeding Medicine 1,(1) (1995) 2,6,8

Gartner LM. Breastfeeding and jaundice. J perinatal 2001; 21:s25-s29.

Gartner LM, Greer FR; American Academy of Pediatrics, Section on Breastfeeding and Committee on Nutrition. Prevention of rickets and vitamin D deficiency: new guidelines for vitamin D intake. *Pediatrics*. 2003;111:908–910

Gartner LM, Herschel M. Jaundice and breastfeeding. Pediat Clin North Am 2001; 48:389-99.

Gartner U, Goeser T, Wolkoff AW. Effect of fasting on the uptake of bilirubin and sulfobromophthalein by the isolated perfused rat liver. Gastroenterology. 2002;113:1707–1713

Giovanna Bertini, Carlo Dani, Michele Tronchin and Firmino F. Rubaltelli Is Breastfeeding Really Favoring Early Neonatal Jaundice? *Pediat.* 2001; 107-41.

Glover, Jacalynne, and Mark Sandilands. "Supplementation of Breastfeeding Infants and Weight Loss in the Hospital." *Journal of Human Lactation* 6.1990; 163-166.

**Graffy JP.** Mothers' attitudes to and experience of breast feeding: a primary care study. *British journal of general practice*, 1992; 42:61-64.

Grant A. Monitoring the fetus during labour. In: Chalmers I et al (eds). Effective care in pregnancy and childbirth. Oxford, Oxford University Press 1990.

**Goldman AS.** The immune system of human milk: antimicrobial, antiinflammatory and immunomodulating properties. *Pediatr Infect Dis J* 1993;12(8):664-672.

Gotsch G. Breastfeeding: Pure & Simple, Schaumburg, Illinois: La Leche League International, 1993.

Gottstein R, Cooke R. Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn. Arch Dis Child Fetal Neonatal Ed. 2003;88:F6-F10

Guaran RL, Drew JH, Watkins AM. Jaundice: Clinical practice in 88,000 liveborn infants. Aust NZJ Obstet Gynaecol, 1992; 32:186-192

Gupta R, Ramji S: Effect of delayed cord clamping on iron stores in infants born to anemic mothers: a randomized controlled trial. *Indian Pediatr* 2002 Feb; 39(2): 130-5

Gupta JK, Nikodem VC. Postion for women during second stage of labour. Cochrane Database Syst Rev 2002;(1).

Gunther M. Instinct and the nursing couple. Lancet. 1955;1:575-578

Habeeb SM. Three years retrospective survery study of neonatal hyperbilirubinemia in NICU, Kasr El – Aini Hosiptal. M. Sc. Pediatrics, Faculty of Medicine, Cairo University. 1994.

Hall RT, Mercer AM, Teasley SL, et al. A breast-feeding assessment score to evaluate the risk for cessation of breast-feeding by 7 to 10 days of age. *J Pediatr.* 2002;141:659–664

Hambraeus L, Forsum E, Lo"nnerdal B. Nutritional aspects of breast milk and cow's milk formulas. In: Hambraeus L, Hanson L, MacFarlane H, eds. Symposium on Food and Immunology. Stockholm, Sweden: Almqvist and Wiksell; 1975

Hamprecht K, Maschmann J, Vochem M, Dietz K, Speer CP, Jahn G. Epidemiology of transmission of cytomegalovirus from mother to preterm infant by breastfeeding. *Lancet*. 2001;357:513-518

Hammermann C, Goldstein R, Kaplan M, Eran M, Goldschmidt D, Eidelman AI. Bilirubin in the premature: toxic waste or natural defense? Clin Chem 1998; 44: 2551-3

Hammerman C, Kaplan M, Vreman HJ, Stevenson DK. Intravenous immune globulin in neonatal ABO isoimmunization: factors associated with clinical efficacy. *Biol Neonate*. 1996;70:69–74

Hansen KN, Ebbesen F. Neonatal vitamin K prophylaxis in Denmark: three years' experience with oral administration during the first three months of life compared with one oral administration at birth. *Acta Paediatr.* 1996;85:1137–1139

Hansen TWR. Acute management of extreme neonatal jaundice the potential benefits of intensified phototherapy and interruption of enterohepatic bilirubin circulation. Acta Paediatr 1997; 86: 843-6

Hansen TWR. Kernicterus in term and near-term infants—the specter walks again. Acta Paediatr 2000; 89: 1155-7

Harrison RF, Brennan M, North PM, Reed JV, Wickham EA. Is routine episiotomy necessary? Br med J 1994; 288:1971-1975

Haverkamp AD, Thompson HE, McFee JG, Cetrulo C. The evaluation of continuous fetal heart rate monitoring in high-risk pregnancy. Am J Obstet Gynecol 1976; 125:310-320.

Healthy People 2010 (Conference Edition, in Two Volumes). Washington DC: US Department of Health and Human Services, 2000

Heinig MJ, Dewey KG. Health effects of breast feeding for mothers: a critical review. Nutrition Research Reviews 1997;10:35-56.

Helm A, Windham CT, Wyse B: Dietitians in breastfeeding management: an untapped resource in the hospital. J Hum Lact. 1997;13:221-225.

**Hemminki E.** Impact of caesarean section on future pregnancy: a review of cohort studies. *Paediatric and Perinatal Epidemiology* 1996;10:366-379.

Hemminki E, Saarikoski S. Ambulation and delayed amniotomy in the first stage of labor. Eur J Obstet Gynecol Reprod Biol 1993; 15:129-139.

Hemminki E, Virta AL, Koponen P, Malin M, Kojo-Austin H, Tuimala R. A trial on continuous human support during labor: Feasibility, interventions and mothers' satisfaction. *J Psychosom Obstet Gynaecol* 1990; 11:239-250.

Henderson A, Stamp G, Pincombe J. Postpartum positioning and attachment education for increasing breastfeeding: a randomized trial. *Birth* 2001; 28(4):236-42.

Hintz SR, Gaylord TD, Stevenson DK, Lemons JA, Oh W, Fanaroff AA, et al. Breastfeeding-related jaundice and bilirubin production in term infants. Acta Paediatr 2001; 90: 776-81.

Hodnett ED. Caregiver support for women during childbirth. Cochrane Database Syst Rev 2001;(3).

Hodnett ED, Osborn RW. A randomized trial of the effect of monitrice support during labor: mothers' views two to four weeks postpartum. Birth 1989; 16:177-183.

Hodnett ED, Lowe NK, Hannah ME, Willan AR, Stevens B, Weston JA, et al. Effectiveness of nurses as providers of birth labor support in North American hospitals. JAMA 2002;288:1373-81.

Hoey C, Ware JL. Economic advantages of breast-feeding in an HMO setting: a pilot study. Am J Man Care 1997;3(6):861-865.

Hofmeyr GJ, Nikodem VC, Wolman WL, Chalmers BE, Kramer T. Companionship to modify the clinical birth environment: effects on progress and perceptions of labour, and breastfeeding. Br J Obstet Gynaecol 2001; 98:756-764.

Howard CM, Howard F, Lawrence R, Andresen E, DeBlieck E, Weitzman M. Office prenatal formula advertising and its effect on breast-feeding patterns. Obstet Gynecol. 2000;95:296-303.

Howard CR, Howard FM, Lanphear B, et al. Randomized clinical trial of pacifier use and bottle-feeding or cupfeeding and their effect on breastfeeding. *Pediatrics*. 2003;111:511-518

Horwood LJ, Darlow BA, Mogridge N. Breast milk feeding and cognitive ability at 7-8 years. Arch Dis Child Fetal Neonatal Ed. 2001;84: F23-F27

Howie PW, Forsyth JS, Ogston SA, Clark A, du V Florey C. Protective effect of breast feeding against infection. *BMJ* 1990;300:11-16.

Huffman SL, Ford K, Allen H, Streble P. Nutrition and fertility in Bangladesh: breastfeeding and post partum amenorrhoea. *Popul Stud Camb*). 1987;41:447–462.

Hundley VA, Cruickshank FM, Lang GD, Glazener CMA, Milne JM, Turner M, Blyth D, Mollison J, Donaldson C. Midwife managed delivery unit: a randomised controlled comparison with consultant care. Br Med J 2000; 309:1400-1404.

- Ibrahim SA, Omer MIA, Amin IK, Babiker AG, Rushwan H. The role of the village midwife in detection of high risk pregnancies and newborns. Int J Gynecol Obstet 1992; 39:117-122.
- ICN. Reducing the impact of HIV/AIDS on Nursing/Midwifery personnel: guidelines for National Nurses' Associations and others. International Council of Nurses. Geneva. 1996.
- Ip S, Chung M, Kulig J. et al. An evidence-based review of important issues concerning neonatal hyperbilirubinemia. *Pediatrics*. 2004;113(6).
- Inch S, Garforth S. Establishing and maintaining breastfeeding. In: Effective care in pregnancy and childbirth. Oxford, Oxford University Press, 1989:1359-1374.
- Institute of Medicine, Committee on Nutritional Status During Pregnancy and Lactation. Nutrition During Lactation. Washington, DC: National Academy Press. 1991; 24-25, 161-171, 197-200
- Institute of Medicine. Crossing the Quality Chasm: A New Health System for the 21st Century. Washington, DC: National Academy Press; 2001.
- Jackson GL, Kennedy KA, Sendelbach DM, et al. Problem identification in apparently well neonates: implications for early discharge. Clin Pediatr (Phila). 2000;39:581-590.
- Jackson KW Jr, Allbert JR, Schemmer GK, et al: A randomized controlled trial comparing oxytocin administration before and after placental delivery in the prevention of postpartum hemorrhage. Am J Obstet Gynecol 2001 Oct; 185(4): 873-7.
- Jacobson SW, Chiodo LM, Jacobson JL. Breast- feeding effects on intelligence quotient in 4- and 11-year-old children. Pediatrics 1999;103(5):71.
- Jensen RG. Lipids in human milk. Lipids 1999;34(12):1243-1271.
- Johnson L, Boggs TR. Bilirubin-dependent brain damage: incidence and indications for treatment. In: Odell GB, Schaffer R, Simopoulos AP, eds. *Phototherapy in the Newborn: An Overview*. Washington, DC: *National Academy of Sciences*. 1974; 122–149.

Johnson LH, Bhutani VK, Brown AK. System-based approach to management of neonatal jaundice and prevention of kernicterus. J Pediatr. 2002;140:396-403.

**Johnson L.** Hyperbilirubinemia in the term infant: when to worry, when to treat. NY State *J Med* 1991; 91: 483–9.

Johnson R, Slade P. Does fear of childbirth during pregnancy predict emergency caesarean section? *BJOG* 2002;109:1213-21.

Jones E, Dimmock PW, Spencer SA. A randomized controlled trial to compare methods of milk expression after preterm delivery. Arch Dis Child Fetal Neonatal Ed, 2001; 85:F91-5.

Janke JR. Breastfeeding duration following cesarean and vaginal births. J Nurs Midwifery, 1988; 33(4): 159-64.

**Kaplan M, Hammerman C.** Severe neonatal hyperbilirubinemia: a potential complication of glucose-6-phosphate dehydrogenase deficiency. *Clin Perinatol.* 1998;25:575–590.

Kantor HI, Rember R, Tabio P, Buchanon R. Value of shaving the pudendal-perineal area in delivery preparation. *Obstet Gynecol* 1965; 25:509-512.

Kappas A, Drummond GS, Munson DP, Marshall JR. Sn-mesoporphyrin interdiction of severe hyperbilirubinemia in Jehovah's Witness newborns as an alternative to exchange transfusion. *Pediatrics*. 2001;108: 1374–1377.

**Kearney MH, Cronenwett LR, Reinhardt R.** Cesarean delivery and breastfeeding outcomes. Birth 17(2): 97 - 103, 1990.

Kersting M, Dulon M. Assessment of breast-feeding promotion in hospitals and follow-up survey of mother-infant pairs in Germany: the SuSe Study. *Public Health* Nutr 2002;5:547-52.

Khan GQ, John IS, Wani S, et al: Controlled cord traction versus minimal intervention techniques in delivery of the placenta: a randomized controlled trial. Am J Obstet Gynecol 1997 Oct; 177(4): 770-4.

Klein M, Papageorgiu AN, Westreich R, Spector-Dunsky L, Elkins V, Kramer MS, Gelfand MM. Care in a birthroom vs a conventional setting: a controlled trial. Can Med Assoc J 1984; 131: 1461-1466.

Kramer LI. Advancement of dermal icterus in the jaundiced newborn. Am J Dis Child. 1969;118:454-458.

Kramer MS, Barr RG, Dagenais S, et al. Pacifier use, early weaning, and cry/fuss behavior: a randomized controlled trial. *JAMA*. 2001;286: 322–326

Kramer MS, Chalmers B, Hodnett ED, et al. Promotion of Breastfeeding Intervention Trial (PROBIT): a randomized trial in the Republic of Belarus. *JAMA*. 2001;285:413-420.

Kuan LW, Britto M, Decolongon J, Schoettker PJ, Atherton HD, Kotagal UR. Health system factors contributing to breastfeeding success. *Pediatrics*. 1999 104(3):e28.

Kuczmarksi RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, Mei Z, Curtin LR, Roche AF, Johnson CL. CDC growth charts: United States. Advance data from vital and health statistics: no. 314. Hyattsville, Maryland: National Center for Health Statistics, 2000.

Kuzela AL, Stifter CA, Worobey J. Breastfeeding and mother-infant interactions. J Reprod Infant Psychol 1990;8:185-194.

Kwast BE. Midwives: key rural health workers in maternity care. Int J Gynaecol Obstet 1992; 38(Suppl):S9-S15.

Kwast BE. Maternity care in developing countries. In: Health matters. Public health in North-South perspective. Van der Velden K et al (eds). Houten, Bohn Stafleu Van Loghum 1995.

La Leche League International. Positioning Your Baby at the Breast. Schaumburg, Illinois, 1988; Publication No. 107.

Lang S, Lawrence CJ, Orme RL'E. Cup feeding: an alternative method of infant feeding. Archives disease in childhood, 1994; 71:365-369.

Lawrence, Ruth A. and Lawrence, Robert M. Breastfeeding: A Guide for the Medical Profession. 5th ed. New York; C.V. Mosby, 1999.

Leung GM, Lam T-H, Ho L-M. Breast-feeding and its relation to smoking and mode of delivery. Obstet Gynecol 2002;99:785-94.

Linn S, Schoenbaum SC, Monson RR, Rosner B, Stubblefield PG, Ryan KJ. Epidemiology of neonatal hyperbilirubinemia. *Pediatrics*. 1995;75: 770-774.

Liu LL, Clemens CJ, Shay DK, Davis RL, Novack AH. The safety of newborn early discharge: the Washington State experience. *JAMA*. 1997; 278:293–298.

Lopez-Zeno JA, Peaceman AM, Adashek JA, Socol ML. A controlled trial of a program for the active management of labor. N Engl J Med 1992; 326:450-454.

Lowe NK. Amazed or appalled, apathy or action? [editorial]. JOGN Nurs 2003; 32: 281-82.

Lydon-Rochelle M, Holt VL, Easterling TR, Martin DP. Risk of uterine rupture during labour among women with a prior cesarean delivery. New England Journal of Medicine 2001;345:3-8.

Macmahon JR, David R, Stevenson, Oski FI.: Neonatal Jaundice Avery s diseases of the neonate neonatal jaundice 1999; 995:1035

MacVicar J, Dobbie G, Owen-Johnstone L, Jagger C, Hopkins M, Kennedy J. Simulated home delivery in hospital: a randomised controlled trial. Br J Obstet Gynaecol 1993; 100:316-323.

**Madlon-Kay DJ.** Identifying ABO incompatibility in newborns: selective vs automatic testing. *J Fam Pract*. 1992;35:278–280

MAIN Collaborative Group Preparing for Breast Feeding. Treatment of inverted and non protractile nipples in pregnancy. *Midwif* 1994; 10.200-14.

Maisels MJ. Epidemiology of neonatal jaundice. In: Maisels MJ, Watchko JF, editors. Neonatal jaundice. Amsterdam: Harwood Academic Publishers, 2000; 37–49.

Maisels MJ. Why use homeopathic doses of phototherapy? *Pediatrics*. 1996;98:283–287.

Maisels J, Gifford K. Normal serum bilirubin levels in the newborn and the effect of breast-feeding. *Pediatrics*. 1996;78:837–843.

Maisels J, Gifford K, Antle CE, Leib GR. Jaundice in the healthy newborn. Newborn infant: a new approach to an old problem. *Pediatrics*. 1998;81: 505-511.

Maisels MJ, Kring EA. Length of stay, jaundice, and hospital readmission. *Pediatrics*. 1998;101:995–998.

Maisels MJ, Newman TB. Kernicterus in otherwise healthy, breast-fed term newborns. *Pediatrics*. 1995;96:730–733.

Maisels MJ, Newman TB: Predicting hyperbilirubinemia in newborns: the importance of timing. *Pediatrics* 1999 Feb; 103(2): 493-5.

Malkin JD. Not as much as you think: toward a truer estimate of the difference in direct medical costs between vaginal and cesarean deliveries. *Birth* 2001;28:208–9.

Malkin JD, Garber S, Broder MS, Keeler E. Infant mortality and early postpartum discharge. Obstet Gynecol. 2000;96:183–188.

Manganaro R, Mami C, Marrone T, Marseglia L, Gemelli M. Incidence of dehydration and hypernatremia in exclusively breast-fed infants. *J Pediatr*. 2001;139:673–675.

Mann Mertz P.Intervention: deressing effects on wound healing. New Directions in Wound Healing. Cinvatec, 1990; 83.

Marasco, Lisa and Jan Barger. "Examining the Evidence for Cue Feeding of Breastfed Infants." Breastfeeding Abstracts, June 1999.

Martinez JC, Maisels MJ, Otheguy L, et al. Hyperbilirubinemia in the breast-fed newborn: a controlled trial of four interventions. *Pediatrics*. 1993;91:470–473.

Maternity Center Association. How do I get the labor support I need? Labor support options. Retrieved 2001; March 20, 2003.

Mbizvo MT, Fawcus S, Lindmark G, Nyström L, and the Maternal Mortality Study Group. Operational factors of maternal mortality in Zimbabwe. *Health Pol and Plann* 1993; 8(4):369-378.

McCandlish R. National Perinatal Epidiology Unit, Oxford, 1996, Personal Communication.

McDonald S, Prendiville WJ, Elbourne D: Prophylactic syntometrine versus oxytocin for delivery of the placenta (Cochrane Review). In: *The Cochrane Library, Issue 2. Oxford, UK:* Update Software. 2002.

Mckay S, Mahan C. Modifying the stomach contents of labouring women: why, how, with what success, and at what risks? How can aspiration of vomitus in obstetrics best be prevented? *Birth* 1988; 15(4):213-221.

Meier PP, Brown LP, Hurst NM. Breastfeeding the preterm infant. Chapter 14 In Auerback K, Riordan J. Breastfeeding. Gaithersburg, MD: Aspen, 1998, pp. 449-480.

Melzack R, Belanger E, Lacroix R. Labor pain, effect of maternal position on front and back pain. *J Pain symptom Manegem* 1991; 6:476-480.

Menticoglou SM, Manning F, Harman C, Morrison I. Perinatal outcome in relation to second-stage duration. Am J Obstet Gynecol 1995; 173:906-912.

Michaelsen KF, Milman N, Samuelson G. A longitudinal study of iron status in healthy Danish infants: effects of early iron status, growth velocityand dietary factors. *Acta Paediatr* 1995; 84:1035-1044.

Mikiel-Kostyra K, Mazur J, Boltruszko I. Effect of early skin-to-skin contact after delivery on duration of breastfeeding: a prospective cohort study. *Acta Paediatr*. 2002;91:1301–1306.

Milligan RA, Flenniken PM, Pugh LC. Positioning intervention to minimize fatigue in breasting women. Appl Nurs Res 1996; 9(2):67-70.

Mohrbacher N and Stock J. The Breastfeeding Answer Book, Third Revised ed. Schaumburg, Illinois: La Leche League International, 2003, p. 148-149.

Montgomery DL, Splett PL. Economic benefit of breast-feeding infants enrolled in WIC. J Am Diet Assoc 1997;97:379-385.

Morse JM, Jehle C, Gamble D. Initiating breastfeeding: a world survey of the timing of postpartum breastfeeding. *Breastfeeding review*, 1992; May:210-216.

Moyer VA, Ahn C, Sneed S. Accuracy of clinical judgment in neonatal jaundice. Arch Pediatr Adolesc Med. 2000;154:391-394.

Neifert MA. Early assessment of the breastfeeding infant. Contemp Pediatr. 1996;13:142-166.

Neifert, Marianne, and Joy Seacat. "Milk Yield and Prolactin Rise With Simultaneous Breastpump" Abstract from the Ambulatory Pediatric Association Annual Meeting; Washington DC, 1985; May 7-10.

Neifert, M and Seacat, J. "Increased Serum Prolactin Levels With Double Breast Pumping" Abstract from the Ambulatory Pediatric Association Annual Meeting: Washington DC, 1985; May 7-10.

Nelson VR. The effect of newborn early discharge follow-up program on pediatric urgent care utilization. J Pediatr Health Care. 1999;13:58-61.

Nelson NM, Enkin MW, Saigal S, Bennett KJ, Milner R, Sackett DL. A randomized clinical trial of the Leboyer approach to childbirth. New Engl J Med 1990; 302:655-660.

Newman J. Finger feeding. J Hum Lact, 2000; 12(2): 129-35.

Newman TB, Liljestrand P, Escobar GJ. Jaundice noted in the first 24 hours after birth in a managed care organization. Arch Pediatr Adolesc Med. 2002;156:1244–1250.

Newman TB, Xiong B, Gonzales VM, Escobar GJ: Prediction and prevention of extreme neonatal hyperbilirubinemia in a mature health maintenance organization. *Arch Pediatr Adolesc Med* 2000 Nov; 154(11): 1140-7.

Newcomb PA, Egan KM, Titus-Ernstoff L, et al. Lactation in relation to postmenopausal breast cancer. Am J Epidemiol 1999; 150:174-182.

Nichols, F. H., & Humenick, S. S. (Eds.). Childbirth education: Practice, research and theory (2nd ed.). *Philadel phia: W. B.* Saunders Company.2000.

Nicoll A, Ginsburg R, Tripp JH. Supplementary feeding and jaundice in newborns. Acta Paediatr Scand. 1982;71:759–761.

Nielsen HE, Haase P, Blaabjerg J, Stryhn H, Hilden J. Risk factors and sib correlation in physiological neonatal jaundice. *Acta Paediatr Scand.* 1997;76:504-511.

Nikodem VC. Immersion in water in pregnancy labour and birth. Cochrane Database Syst Rev 2003;(1).

Notorianni L. Plasma protein binding of drugs in pregnancy and in neonates. Clin pharmacokinet, 1990; 18 (1): 20-36.

Novotny R, Hla MM, Kieffer EC, Park CB, Mor J, Thiele M. Breastfeeding duration in a multiethnic population in Hawaii. *Birth*. 2000;27(2):91-96.

O'Cathain A, Walters SJ, Nicholl JP, Thomas KJ, Kirkham M. Use of evidence based leaflets to promote informed choice in maternity care: randomised controlled trial in everyday practice. *BMJ* 2002;324:643.

O'Driscoll K, Meagher D. Active management of labour: the Dublin experience. 2nd ed. Baillière Tindall, London 1986.

O'Heir J, From Kobe to Oslo; the WHO Midwifery Education Modules. Proceedings of the 24<sup>th</sup> triennial International Confederation of Midwives Congress, *ICM*, Oslo 1996, pp. 520-522.

Ohlsson G, Buchhave P, Leandersson U, Nordstrom L, Rydhstrom H, Sjolin I. Warm tub bathing during labour: maternal and neonatal effects. Acta Obstet Gynecol Scand 2001;80:311-14.

Osborn LM, Bolus R. Breast feeding and jaundice in the first week of life. J Fam Pract. 1985;20:475-480.

Osborn LM, Reiff MI, Bolus R. Jaundice in the full-term neonate. Pediatrics. 1994;73:520-525

Oski FI. Bilirubin metabolism . Pediatrics. 1999; 108:31-39.

Ottervanger HP, Keirse MJNC, Smit W, Holm JP. Controlled comparison of induction versus expectant care for prelabor rupture of the membranes at term. *J Perinat Med* 1996 (in press).

Ozmert E, Erdem G, Topcu M. Long-term follow-up of indirect hyperbilirubinemia in full-term Turkish infants. *Acta Paediatr*. 1996;85: 1440–1444.

Perez-Escamilla, R et al. The Association Between Cesarean Delivery and Breast-Feeding Outcomes among Mexican Women. Am J Public Health. June 1996; 86(6):832-836.

**Peters MH.** Midwives and the achievement of safer motherhood. Int J Gynecol Obstet 2001; 50(Suppl.2):S89-S92.

Philipp BL, Merewood A, Miller LW, Chawla N, Murphy-Smith MM, Gomes JS, Cimo S, Cook JT. Baby-friendly hospital initiative improves breastfeeding initiation rates in a US hospital setting. *Pediatrics* 2001;108:677-81.

Philipsen T, Jensen NH. Maternal opinion about analgesia in labour and delivery. A comparison of epidural blockade and intramuscular pethidine. Eur J Obstet Gynecol Reprod Biol 1990; 34;205-210.

Picciano MF. Human milk: nutritional aspects of a dynamic food. Biol Neonate 1998;74:84-93.

**Pisacane A.** Neonatal prevention of iron deficiency. Placental transfusion is a cheap and physiological solution. *Br Med J* 1996; 312:136-137.

Potter BJ, Berck BD. Uptake of bilirubin at the hepatocyte. Hepatology 1994; 20: 1078-1085

Radford N and Thompson A. A Pathway to Practice: Direct Entry Midwifery. Department of Health and Social Services. HMSO. 1987.

Ramsey M, Gisel E. Neonatal sucking and maternal feeding practices. Dev Med Child Neurol, 1996; 26(3):34-47.

Ransjo-Arvidson AB, Matthiesen AS, Lilja G, Nissen E, Widstrom AM, Uvnas-Moberg K. Maternal analgesia during labor disturbs newborn behavior: effects on breastfeeding, temperature, and crying. *Birth.* 2001;28:5–12

Ryan AS, Wenjun Z, Acosta A. Breastfeeding continues to increase into the new millennium. *Pediatrics*. 2002;110:1103–1109

Renfrew M, Fisher C, Arms S. Best feeding: Getting Breastfeeding Right for You. Berkeley, California: Celestial Arts, 1990.

Renfrew MJ, Lang S, Martin L, Woolridge MW. Feeding schedules in hospitals for newborn infants. Cochrane Database Syst Rev 2000.

Righard L, Alade M. Sucking technique and its effect on success of breast feeding. Birth 1994; 185-89, 1992.

Righard L, Alade MO. Effect of delivery room routine on success of first breast-feed. Lancet. 1990;336:1105-1107.

Riordan Jan. Breast feeding and human lactation, third edition, vol.2, chapter 11. 2004;311-312.

Riordan Jan, and Auerbach Kathleen G., eds. Breastfeeding and Human Lactation. 2nd ed. Boston MA; Jones and Bartlett, 1993.

Riordan J, Bibb D, Miller M, Rawlins T. Predicting breastfeeding duration using the LATCH breastfeeding assessment tool. J Hum Lact. 2001;17:20-23.

Roberts, LJ. Elective section after two sections - where's the evidence? British journal of Obstetrics and Gynaecology, 1991; vol. 98, pp 1199-1202.

Romney ML, Gordon H. Is your enema really necessary? Br Med J 1981; 282:1269-1271.

Rooks JP, Weatherby NL, Ernst EKM, Stapleton S, Rosen D, Rosenfield A. Outcomes of care in birth centers: the National Birth Center Study. N Engl J Med 1989; 321:1804-1811.

Rosenblatt KA, Thomas DB, and the WHO Collaborative Study of Neoplasia and Steroid Contraceptives. Lactation and the risk of epithelial ovarian cancer. *Int J Epidemiol* 1993;22(2):192-197.

Royal Collage of Midwives. Successful breastfeeding. London: Churchill Livingstone, 1991; pp.18-19,.

Rubaltelli FF et al. Unconjugated and conjugated bilirubin pigments during perinatal development IV. The influence of of breast feeding on neonatal hyperbilirubinemia. *Biol Neonate* 1993; 64:104-9.

Rubaltelli FF, Gourley GR, Loskamp N, et al. Transcutaneous bilirubin measurement: a multicenter evaluation of a new device. *Pediatrics*. 2001;107:1264–1271.

Rutter N, Spencer A, Mann N, Smith M. Glucose during labour. Lancet 1990; 2:155.

Ryan AS, Wenjun Z, Acosta A. Breastfeeding continues to increase into the new millennium. *Pediatrics*. 2002;110:1103-1109.

Scariati PD, Grummer-Strawn LM, Fein SB. A longitudinal analysis of infant morbidity and the extent of breastfeeding in the United States. *Pediatrics* 1997;99(6):5.

Schanler RJ. The use of human milk for premature infants. *Pediatr Clin North Am.* 2001;48:207–219.

Schanler, Richard and Nancy Hurst. "Human Milk for the Hospitalized Preterm Infant." Seminars in Perinatology 1994; 6:476-484.

Schanler RJ, O'Connor KG, Lawrence RA. Pediatricians' practices and attitudes regarding breastfeeding promotion. *Pediatrics*. 1999;103(3):e35.

Schmid R. Bilirubin metabolism: state of art. Gastroentrology 1978; 74: 1307

Schneider AP. Breast milk jaundice in the newborn. JAMA. 1996;255: 3270–3274.

Schutte MF, Treffers PE, Kloosterman GJ, Soepmati S. Management of premature rupture of the membranes: the risk of vaginal examination to the infant. Am J Obstet Gynecol 1983; 146:395-400.

Schwarcz RL, Diaz AG, Fescina RH, Belitzky R, Rossello JLD. Lowrisk pregnancy anddelivery. Guidelines for the management of antenatal care, labor and delivery. Latin American Center for Perinatology and Human Development. CLAP Scientific Publication No 1321.02. Montevideo, Pan American Health Organization - World Health Organization 1995.

Senden IPM, Wetering MD, Eskes TKAB, Bierkens PB, Laube DW, Pitkin RM. Labor pain: A comparison of parturients in a Dutch and an American teaching hospital. Obstet Gynecol 1998; 71:541-544.

Simkin P. Non-pharmacological methods of pain relief during labour. In: Chalmers I et al (eds). mEffective care in pregnancy and childbirth. Oxford, Oxford University Press 1989.

Sleep J, Roberts J, Chalmers I. Care during the second stage of labour. In: Chalmers I et al (eds). Effective care in pregnancy and childbirth. Oxford, Oxford University Press 1999.

Sluka, K. A. and D. Walsh. "Transcutaneous electrical nerve stimulation: basic science mechanisms and clinical effectiveness." *J Pain* 2003; 4(3): 109-21.

Slusser, Wendelin, and Nancy G. Powers. "Breastfeeding Update 1: Clinical Lactation Management." Pediatrics in Review Vol 18, No 4 (April 1997): 111-119.

Soskolne EI, Schumacher R, Fyock C, Young ML, Schork A. The effect of early discharge and other factors on readmission rates of newborns. Arch Pediatr Adolesc Med. 1996;150:373-379.

Stapleton H. Qualitative study of evidence based leaflets in maternity care. BMJ 2002;324:639.

Stevenson DK, Bortoletti AL, Ostrander CR, et al. Pulmonary excretion of carbon monoxide in the human infant as an index of bilirubin production: effects of breast-feeding and caloric intake in the first postnatal week. *Pediatrics*. 2000; 65:1170–1172.

Stevenson DK, Fanaroff AA, Maisels MJ, et al. Prediction of hyperbilirubinemia in near-term and term infants. *Pediatrics*. 2001;108:31-39

**Stokamer CL**. Breastfeeding promotion efforts: why some do not work. *International journal of gynecology and obstetrics*, 1990; 31(Suppl 1):61-65.

Stoker R, Yamamoto Y. Bilirubin is an antioxidant of possible physiologic importance. Science 1997; 235:1043-1046.

Sugarman M, Kendall-Tackett KA. Weaning ages in a sample of American women who practice extended breastfeeding. Clin Pediatr (Phila). 1995;34:642-647

Swanstrom S, Bratteby LE. Metabolic effects of obstetric regional analgesia and of asphyxia in the newborn infant during the first two hours after birth. *Acta Paed Scand* 1991; 70:791-800.

Tarnow-Mordi WO, Shaw JCL, Liu D, Gardner DA, Flynn FV. Iatrogenic hyponatraemia of the newborn due to maternal fluid overload; a prospective study. *Br Med J* 1991; 283:639-642.

Taylor PM, Maloni JA, Brown DR. Early suckling and prolonged breastfeeding. American journal of diseases of children, 1986; 140:151-154.

Thacker SB, Stroup DF. Continuous electronic heart rate monitoring for fetal assessment during labor. Cochrane Database Syst Rev 2000;(1).

The Global Criteria for the WHO/UNICEF Baby Friendly Hospital Initiative. In: Baby Friendly Hospital Initiative. Part II. Hospital level implementation. WHO/UNICEF, 1992.

The Innocenti Declaration was adopted by the Forty-fifth World Health Assembly in May 1992 in Resolution WHA 45.34.

Thomas IL, Tyle V, Webster J, Neilson A. An evaluation of transcutaneous electrical nerve stimulation for pain relief in labour. Aust NZ J Obstet Gynaecol 1988; 28:182-189.

Thomson AM. Pushing techniques in the second stage of labour. J Adv Nurs 1993; 18:171-177.

Thorp JA, Hu DH, Albin RM, McNitt J, Meyer BA, Cohen GR, Yeast JD. The effect of intrapartum epidural analgesia on nulliparous labor: a randomized, controlled, prospective trial. Am J Obstet Gynecol 1993; 169:851-858.

Treffers PE. Obstetric care in developed countries: the case of the Netherlands. In: Health matters. Public health in North-South perspective. Van der Velden K et al (eds). Houten, Bohn Stafleu Van Loghum 1995.

US Department of Health and Human Services. Healthy People 2010: Conference Edition—Volumes I and II. Washington, DC: US Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health; 2000:47–48.

Valaes T. Severe neonatal jaundice associated with glucose-6-phosphate dehydrogenase deficiency: pathogenesis and global epidemiology. *Acta Paediatr Suppl.* 1994;394:58–76.

Vestermark V, Hogdall CK, Birch M, Plenov G, Toftager-Larsen K. Influence of the mode of delivery on initiation of breast-feeding. Eur J Obstet Gynecol Reprod Biol 1990;38:33-8.

Victora CG et al. Pacifier-use and short breastfeeding duration: cause, consequence or coincidence? *Pediatrics*, 1997; 99(3):445-453.

Visness CM, Kennedy KI. Maternal employment and breast-feeding: findings from the 1988 National Maternal and Infant Health Survey. *Amer J Public Health*. 1997;87(6):945-950.

Volpe JJ. Neurology of the Newborn. 4th ed. Philadelphia, PA: W. B. Saunders; 2001.

Waldenström U, Gottvall K. A randomized trial of birthing stool or conventional semirecumbent position for second-stage labour. Birth 1991; 18:1:5-10.

Waldenström U, Nilsson CA. Experience of childbirth in birth center care. A randomized controlled study. *Acta Obstet Gynecol Scand* 1994; 73:547-554.

Waldenström U, Nilsson CA, Winbladh B. The Stockholm birth center trial. Maternal and infant outcome. *Brit J Obstet Gynae* 1996 (in press).

Wennberg RP. Cellular basis of bilirubin toxicity. N Y State J Med. 1991;91:493–496.

Westphal MF et al. Breast-feeding training for health professionals and resultant institutional changes. *Bulletin of the World Health Organization*, 1995; 73(4):461-468.

Whitmer DI, Gollan JL. Mechanism and significance of fasting and dietary hyperbilirubinemia. Semin liver Dis 1983; 3:42-51.

Widstrom A-M et al. Short-term effects of early suckling and touch of the nipple on maternal behaviour. Early human development, 1990; 21:153-163.

Widstrom A-M, Thingstrom-Paulsson J. The position of the tongue during rooting reflexes elicited in newborn infants before the first suckle. *Acta Paediatr.* 1993;82:281–283.

Wiessinger D. A breastfeeding teaching tool using a sandwich analogy for latch on. J Hum Lact 14(1):51-56,1998.

Wilkinson C, Enkin MW: Manual removal of placenta at caesarean section (Cochrane Review). In: The Cochrane Library, Issue 2. Oxford, UK: Update Software. 2002.

Winikoff B et al. Dynamics of infant feeding: Mothers, professionals, and the institutional context in a large urban hospital. *Pediatrics*, 1986; 77(3):357-365.

Woolridge MW. Problems of establishing lactation. Food and nutrition bulletin, 1996; 17(4):316-323.

Woolridge, Michael.W., J.D. Baum and R.F. Drewett. Effect of Traditional and a New Nipple Shield on Sucking Patterns and Milk Flow. EarlyHumanDevelopment1980;357-364.

Woolley RJ. Benefits and risks of episiotomy: a review of the English-language literature since 1980. Obstet Gynecol Surv 1995; 50:806-820, 821-835.

World Health Organization. Mother-Baby Package: Implementing safe motherhood in countries. Unpublished document FHE/MSM/99.11. Geneva, World Health Organization 2000.

World Health Organization. HIV and Infant Feeding: A Guide for Health Care Managers and Supervisors. Publication Nos. WHO/FRH/NUT/ 98.2, UNAIDS/98.4, UNICEF/PD/NUT/(J)98.2. Geneva, Switzerland: World Health Organization; 1998.

Yamauchi Y, Yamanouchi I. Breast-feeding frequency during the first 24 hours after birth in full-term neonates. *Pediatrics*. 1990;86:171–175.

Yamauchi Y, Yamanouchi I. Difference in TcB readings between full term newborn infants born vaginally and by cesarean section. *Acta Paediatr Scand.* 1999;78:824–828

Yancey MK, Herpolsheimer A, Jordan GD, Benson WL, Brady K. Maternal and neonatal effects of outlet forceps delivery compared with spontaneous vaginal delivery in term pregnancies. *Obstet Gynecol* 1991; 78:646-650.

Zetterstro"m R. Breastfeeding and infant-mother interaction. Acta Paediatr 1999; 88 (Suppl): 1-6.

**Zheng Qiao-ling, Zhang Xiao-min.** Analysis of 39 cases of maternal deaths caused by incorrect use of oxytocin. *Chin J Obstet Gynecol.* 1994; 29:317.