

References

- Abdel sattar , N.A; Sadek , S.H, , Sallam M.S.and Abdel Dayem , M.K. (1986) :**
 Apo-B and certain other lipid parameters in coronary atherosclerosis.
Thesis submitted for partial fulfillment of M.D. of clinical pathology . 1986 .
- Abe , M.; On,J.;Sago Y.; Okeda , T. and Takaki,R.(1990) :**
 Effects of glucose and insulin on cultured human microvascular endothelial cells .
Diabetes , Res . Clin . Pract .; 9 (3) : 287-950
- Abrams J.J. , Ginsberg H. and Grundy S.M. (1982) :**
 Metabolism of cholesterol and plasma triglycerides in non ketotic D.M. 31:903 .
- Agardh C-D, Santor G,Nilsson -Ehle P.(1983) :**
 plasma high density lipoproteins and lipolytic enzyme activities in diabetic patients.
Acta Med Scand 213:123-128.
- Agardh .E;Tallroth G.;Bauer B, et al (1987) :**
 Retinopathy and nephropathy in insulin dependent diabetics : An inconsistent relationship ? *Diabetic Med .4:248.*
- Albrechtsen , O.K. (1957):**
 The fibrinolytic activity of human tissue .
Brit . J.Haematol .; 3:284 .
- Albrink , M.J. and Man , E.B, (1959) :**
 Serum triglycerides in coronary artery disease .
Arch . Intern . Med .; 103 .4.
- Allen , R.A.; and Pepper , D.S.(1981):**
 Isolation and properties of human vascular plasminogen activator .
Thromb . Haemost .; 45:43.
- Almer , L.O.; and Nilsson I.M.(1975):**
 On fibrinolysis in diabetes mellitus .
Acta .Med .Scand . 192,101 .

Anderson , L.O.; Barrowcliffe , T.W.; Holmer , E.; Johnson , E.A. and Smis ,G.E.C. (1976):

Anticoagulant properties of heparin fractioned by affinity chromatography on matrix bound antithrombin III and by gelfiltration.

Aoki , N.; and kawano , T. (1972):

Inhibition of plasminogen activators by naturally occurring inhibitors in man.

Am .J. physiology , 223 (6): 1334.

Aoki , N. (1979):

Natural inhibitors of fibrinolysis .

Prog . Car . Vas . Dis . ; 21(4) : 267 .

Astrup , T. and Thorsen , s. (1972):

The physiology of fibrinolysis .

Med . Clin .J. Am . ; 65:153.

Attman P.O. , Nyberg G., Olsson T.W.; Gibson C.K. and Alaupovic P. (1992).

Dyslipoproteinaemia in diabetic renal failure .

Kidney International 42:1381-1389 .

Averna , M.R; Rini G.B.; Difede , G., Montalto , G .; Novo S.; and Natarbartolo , A . (1982) :

Apolipoprotein pattern in type II diabetics .

Boll . Soc . Ital . . Biol . Sper . ; 14 : 914 .

Avogaro , P. Bon G.B. ; Gazzolato G . and Quinci G.B. (1979) .

Are apolipoproteins better discriminators than lipids for atherosclerosis ?

Lancet , 1:901 .

Avogaro P .; Bon G.B; Gazzolato , G.and Rorai E, (1980) .

Relationship between apolipoproteins and chemical components of lipoproteins in survivors of myocardial infarction .

Atherosclerosis , 37:69 .

Ayrdult , J.M.; Alix J.F. and Polonovski , J.(1978) :

Some Considerations of methodology and Standardization of apo-A-I immunoassay . *Biochemic*, 60:65 .

Quoted from Clin . Chem . ; 29,3:415, 1983.

Babier , B.A.and Stossel , T.P.(1984):

Haematology : A pathophysiological approach . 1st edition .

Churchill Livingstone . Inc . N.Y.;P.171.

- Bagdasarian , A- and Colman, R..W. (1978) :**
Subcellular localization and purification of platelet antitrypsin .
Blood , 51:139 .
- Bagdade JD, Subbaiah P.V. (1986) :**
Atherosclerosis in type I diabetic female : does her HDL protect or predispose ? *Diabetes* 35 (Suppl . 1) : 67A .
- Baily .K.; and Bellelheim , F.R. (1955) :**
The clotting of fibrinogen .The liberation of peptide material .
Biochem .Biophys . Acta , 18:495 .
- Baker , N.N.; Delahunty , T. and Jakson , R-L. (1974):**
The primary structure of human plasma HDL Glutamine I (apo A-I).
Proc . Natl . Acad . Sci . USA . 71:3631 .
- Bannererjee, R.N.; Sahri , A-L.; and Kumar ,V.; (1979) :**
Antithrombin III deficiency in maturity onset diabetes and atherosclerosis .
Thromb . Diath . Haemorrh . 31,399 .
- Barlow , G.H. (1976) :**
Urinary and kidney cell plasminogen activator (urokinase) .
Methods Enzymol . ; 45:239 .
- Barter , P.J.; Hopkins , G.J. and Gorjatschkol , A. (1985) :**
Lipoprotein substrates for plasma cholesterol esterification .
Influence of particle size and composition of high density lipoprotein subfraction 3 .
Atherosclerosis 58:97 .
- Barton , P. G.; Jackson , C.M.; Hanahan , D.J. (1967) :**
Relationship between factor V and activated factor X in the generation of prothrombokinase .
Nature (London) , 214 :923
- Bassiouny , A-R.; Rosenberg , H.; and Mc Donald , T.L. (1983) :**
Glycosylated Collagen is antigenic
Diabetes , 32 : 1182-4.
- Behringwerke , A.G.(1991) :**
"Determination of clotting factors VIII and IX"
Edition october 1991 , Hoechst company . Marburg , W.Germany .
- Behringwerke , A-G. (1991) :**
" Determination of coagulation factor VII"
Edition July 1991 , Hoechst company . Marburg , W.Germany .

Behringwerke , A-G. (1992) :

"Turbiquant for the quantitative determination of human serum proteins with the turbidimeter "

Edition March , 1992 .Hoechst company .Marburg , W.Germany .

Beisiegel U.; Weber W.; Hainga J-R-(1988) :

Apolipoprotein E-binding protein isolated from dog and human liver.

Arteriosclerosis , 8 : 288.

Bending J-J.; Viberti G.C.; Watkins P.J. et al , (1986):

Intermittent clinical proteinuria and renal function in diabetes.

Evolution and effect of glycaemic Control .

*Br.Med .J.*292-83 .

Bending J.J, Dodds R,Keen H, et al (1986) :

Lowering protein intake and the progression of diabetic renal failure.

Diabetologica 29:516A .

Berglund , J.; Lins P-E, Adamson U.; et al , (1987) :

Microalbuminuria in long term insulin - dependent diabetes mellitus.

Prevalance and clinical characteristics in a normotensive population.

Acta Med .Scand .222 :333.

Berman ,M.; Hall , M.; Levy , R-I.; Eisenberg , S.; Bilheimer , D-W.; Phain , R.D- and Goebel , R.H. (1978):

Metabolism of apo-B and apo C-lipoprotein in man - Kinetic studies in normal and hyperlipoproteinaemic subjects .

J.lipid Res .; 19:38.

Bern MM., Cassani MP, Horton J. Rand L, Davis G. (1980) :

Changes of fibrinolysis and factor VIII coagulant , antigen and ristocetin cofactor in diabetes mellitus and atherosclerosis .

Thromb , Res , 998 : 999 .

Berrettini , M,(1986) :

Detection of in vitro and in vivo cleavage of high molecular weight Kininogen in human plasma by immunoblotting with monoclonal antibodies .

Blood , 68 455 .

Betteridge J. (1990) :

Lipid disorders in diabetes.In Tattersall R.B.and Gale E.A-M .(ed .)

Diabetes clinical management .churchill living stone .

Edinburgh london Melbourne and New York .PP:333-345.

- Brewer , H.B.; Lux,S.E.; Ranon , R.; and John , K.M. (1972):**
 Amino acid sequence of human apo- LP.Gln -II (apo A-II) , an apolipoprotein isolated from high density lipoprotein complex .
Proc .Natl . Acad .Sci USA 69 : 1304 .
- Briones ER, Mao SJT, Palumbo PJ, O'fallon WM, Chenoweth W,Kottke BA.(1984):**
 Analysis of plasma lipids and apolipoproteins in insulin - dependent and non-insulin dependent diabetics .
Metabolism 33:42-49.
- Brodoff , B.N.; Bleicher , J. (1982) :**
 Diabetes Mellitas . In : Brodoff , B .N. (ed): Diabetes Mellitus and obesity .
Baltimore , Williams and Kilkins PP 735 .
- Brodows , R. G.; Nichlos , D .; Shaker G .; et al (1986) :**
 Evaluation of a new radio-immunoassay for urinary albumin .
Diabetes Care 9:189 .
- Brommer , E.J.P.; Brakman , P. Haver Kate , F.; kluft , C.; Traas , D. and wijngaards , G.(1981):**
 Progress in fibrinolysis .In : Poller, L.(Ed.) ,Recent advances in blood coagulation .
Churchill Livingstone , Edinburgh, 3:125-149.
- Brownlee M .; Valssara , H.; and Labic D. and Basselte , (1984) :**
 The biochemistry of the vascular diabetic complications .
Clin . Chem . Acta , 7 . 502-504 .
- Brownlee . M.; Cerami A.; Vlassara H . (1988) :**
 Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications .
N. Engl . J . Med . 318 : 1315-1321 .
- Brunzell , J. D .; Chait A.; and Bierman E.L . (1978) :**
 Pathophysiology of lipoprotein transport .
Metabolism , 27 : 1109 .
- Brunzell JD, Chait A.; Bierman EL (1985):**
 Plasma lipoprotein in human diabetes mellitus .In : Alberti KGMM, krall LP (eds) The diabetes annual .Elsevier Science Publishers , Amsterdam , PP 463-479.
- Bucolo, G.and David , H.(1973):**
Clin . chem . 19,475.
 Quoted from pamphlet of triglycerides Kit . Sclavo diagnostica.

Burstein , M. and Legmann , P.(1982) :

Lipoprotein precipitation .In clarkson , T.B.; Kritchevsky , D. and Pollak , O. J.(eds):

Monographs on Atherosclerosis , Vol .II .Basel,S.Karger A.G.

Burton , B.T.; Hirschman , G.H.(1980):

Diabetes in the USA : a demographic overview , In : Friedman , E.A.;L'Esperance , F.A.Jr . (eds); Diabetic renal . retinal syndrome .
New York, Grune and stratton Inc -PP 5-18.

Busch , C. and Owen , W.G.(1982) :

Identification in vitro of an endothelial cell surface cofactor for Antithrombin III .parallel studies with isolated perfused rat heart and micro carrier cultures of bovine endothelium .

J .Clin Invest .; 69 :726 .

Calbreath D.F.(1992):

Clinical chemistery .A fundamental textbook - W.B.

Saunders Company PP 280-298.

Campbell P.N.; and Smith A.D. (1988) :

Biochemistry illustrated .2nd ed - churchill living stone .

Edinburgh London Melborne and New York .P.218.

Carew T.E.; Koschensky T.; Hayes SB.(1976) :

A mechanism by which high density lipoprotein may slow the atherogenic process -*Lancet .1:1315-1317 .*

Carlson L.A. and Bottiger L.E(1972) :

Ischemic heart disease in relation to fasting values of plasma triglycerides and cholesterol .

Lancet , 22 : 865 .

Castel, M.; Horellou ,M.H.; Condard , J . and Samoma , M.M. (1983):

Immunochemical determination of histidine rich glycoprotein in healtly subjects and in a clinical population ,In : Davidson , J.F.; Bachmann , F.; Bouvier , C.A . and kruithof , E.K.O. (Eds) :
Progress in fibrinolysis ,.vol .VI .

Churchill Livingstone , Edenburgh m P. 370- 373.

Castelli , W.P .; Doyle J.T.; Gordon T.; Hames C.G.; Hyortland , M.C.; Hully S.B.; et al .; (1977) :

HDL-cholesterol and other lipids in coronary heart disease . The cooperative lipoprotein phenotyping study .

Circulation 55 : 767 .

Ceriello A, (1987) :

Induced hyperglycaemia alters AT-III activity but not its plasma conc. in healthy normal subjects .

Diabetes 36 , 320-323 .

Ceriello A .; Giugliano D .; Quatraro A.; Marchi E .; Barbanti M.; and Le Febre P. (1990) .

Evidence for a hyperglycemia - dependent decrease of antithrombin III-thrombin complex formation in humans .

Diabetologica 33:163-167 .

Chan , T.K. and chan ,V. (1981) :

Antithrombin III, The major modulator of intravascular coagulation is synthesized by human endothelial cells .

Thromb .Haemost. ; 46:504 .

Chait A.; Bierman EL.; Brunzel KD. (1985) :

Diabetic macroangiopathy . In Alberti KGMM , Krall LP (eds) .
Diabetes annual I . Amsterdam :

Elsevier, PP 323-48 .

Charles , R. (1988) :

Diabetes Mellitus ; Defination , classification and diagnosis . In :
Galloway J.A.(ed) Diabetes Mellitus .

Eli Lilly and company ,Indianapolis , Indiana , PP 2-13 .

Chen ,R.; Doolittle .; R.F. and Rosen , F.L. (1969) :

Identification of the polypeptide chains involved in the cross -
linkage of fibrin .

Proc .Nat .Acad .Sci (USA) , 63:420 .

Chihara J, Takebayashi S, Taguchi T. et al (1986):

Glomerulonephritis in diabetic patients and its effect on the
prognosis.

Nephron 43:45.

Christe M .; Fritschi J .; Lammle B.; Tran TH., Marbet GA., Duckert F. (1984) :

F.Fifteen Coagulation and fibrinolysis parameters in diabetes
mellitus and in patients with vasculopathy .

Thromb Harnosasis .; 52:138-43 .

Christensen , C.K. (1984) :

Abnormal albuminuria and blood pressure rise in incipient diabetic
nephropathy induced by excercise .

Kidney Int .25:819.

Christensen , C.K.; Mogensen C.E. (1985)

The course of incipient diabetic nephropathy : studies of albumin excretion and blood pressure . *Diabetic Med* .2:97.

Christensen Ck., Mogensen CE.(1986) :

Acute and long term effect of antihypertensive treatment on exercise-induced albuminuria in incipient diabetic nephropathy .
Scand .J.Clin - Lab . Invest . 46:553 .

Christensen CK, Mogensen (1987):

Antihypertensive treatment : Long term reversal of progression of albuminuria in incipient diabetic nephropathy -A longitudinal study of renal Function .

J.Diabetic Complications . 1 : 45.

Christiansen , J.S. (1988) :

Early renal hyperfunction and hypertrophy in insulin - dependent patients:Changes found at diagnosis and early in the course of diabetes . In Mogensen C.E. (ed) : The kidney and hypertension in Diabetes Mellitus.

Boston , Martinus Nijhoff Publishing .PP 157 .

Clemmenson , I .; Thorsen , S.; and Mullertz ,S. (1976):

Purification of a plasminogen activator inhibitor indistinguishable from α_2 antitrypsin and urokinase inhibitor in pregnancy plasma .
Hemostasis , 5:218 .

Cohen D.,Dodds R, Viberti GC.,(1987) :

Effect of protein restriction in insulin - dependent diabetics at risk of nephropathy .

Br .Med .J. 294 : 794 .

Coller , B.S.; Frank , D.N. and Milton , R.C. (1978) :

Plasma cofactors of platelet function ,correlation with diabetic retinopathy.

Annals .Intern .Med .88,311 .

Collen ,D.(1980) :

The regulation and control of fibrinolysis .

Thromb.Haemost .; 43:77.

Colman , R.W.; Hirsh , J.;Marder , V.J.; et al (1987) :

Haemostasis and thrombosis : Basic principles and clinical practice.

2nd edition , J.B. Lippincott company .philadelphia .

Colwell , J.A.; Halushka , P.V.; Sarji ,K, Levine ,J and Nair , R.M.G. (1976):

Altered platelet function in diabetes mellitus .

Diabetes 25,826.

Colwell ,J.A.; Winocour , P. D. and Halushka , P.V.(1983):

Do platelets have anything to do with diabetic microvascular disease?

Diabetes , 32 (2) : 14-19.

Colwell ,J.A.; Bingham , S.E.; Abaira , C.; Anderson , J.W.; Comstock ,T.;Kwaan , H.C.; Nattal , F. ; and the cooperative Study group (1986) .

Veterans administration cooperative study on antiplatelet agents in diabetic patients after amputation for gangrene : II Effects of aspirin and dipyridamole on athero-sclerotic vascular disease rates .

Diabetes care , 9 : 140 -8 .

Comp .; P.C.; and Esmon , C.T. (1979) :

Activated protein C.inhibits platelet prothrombin converting activity.

Blood , 54 : 1272 -1281 .

Comp , P.C. and Esmon , C.T. (1981) :

Generation of fibrinolytic activity by infusion of activated protein C into dogs .

J.Clin . Invest . ; 68 :1221 .

Comp , P.C.; Nixon , R. R. and Esmon , C.T.(1984) :

Determination of functional levels of protein C, an antithrombotic protein , using thrombin - thrombo-modulin complex .

Blood , 63:15-21.

Comp .; P.C.; Thurnau , G.R. and welsh , J. (1986):

Functional and Immunologic protein S levels are decreased during pregnancy .

Blood , 68:881 .

Coppo R.; Amore A.; Roccatello D. et al , (1987) :

A solid phase enzyme immunoassay for the measurement of urinary albumin and the detection of microalbuminuria .

J.Diabetic Complications 1:58 .

Corbella , E.; Miraglitten ,G.; and Masperi R.(1979):

Platelet aggregation and antithrombin III levels in diabetic children.

Hemostasis , 8 , 30 .

Craighead , J.E. (1981) :

Viral diabetes mellitus in man and experimental animals . In :Skyler, J-S. and cahill , G.F. Jr. (eds) : *Diabetes Mellitus* .

New York , Medical Books ,PP 23-30 .

- Creagh , M.D. , Roberts , I.P.; clark , D.J. and Preston , P.E. (1991) :**
 Familial AT III deficiency and mycoplasma pneumonia -
J.Clin .Pathol .; 44:870-871 .
- Curry , M.D. and Gustafson A.; (1978) :**
 Electroimmunoassay and radial immuno- diffusion assay evaluated
 for quantification of Human apolipoprotein B.clin .
Chem . 24:280 -286.
- Cuttiss L.K. and Witztum J.K.(1983) :**
 A noval method for generating region-specific monoclonal
 antibodies to modified proteins .
J-Clinic . Invest , 72:1427 .
- Curtiss LK., Witztum JL(1985) :**
 Plasma Apolipoprotein AI , AII , B ,CI & E are glycosylated in
 hyperglycemic diabetic Subjects .
Diabetes 34 : 454-461 .
- Davie , E.W. and Ratnoff , O.D. (1964):**
 Waterfall sequence for intrinsic blood clotting .
Science , 145:1310-12.
- Davie , E.W. (1979) :**
 The role serine protease in blood coagulation cascade .
Adv . Enzymol .; 48:277.
- Dahlback , B.(1983):**
 Purification of humane vitamin K dependent protein S .and its
 limited proteolysis by thrombin .
*Biochem .J.;*209:837 .
- Dahi -Jorgensen K.(1987) :**
 Near normoglycemia and late diabetic Complications .The Oslo
 study .
Acta Endocrinol (Copenh) 144 (Suppl) 284:1.
- Dahi -Jorgensen K., Hanssen KF, Kierulf P. et al ,(1988):**
 Reduction of urinary albumin excretion after 4 years of continous
 Subcutaneous insulin infusion in insulin - dependent diabetes
 mellitus.
Acta Endocrinol (Copenh) 117:19.
- Deckert ,T.; (1988) :**
 Glycemic control and Complications . In Alberti KGMM .Krall L P
 (eds):The Diabetes Annual .4 . Elsevier . Science publishers , BV .

Denson , K.W. E. (1976) :

Natural and pathological inhibitors of blood coagulation .In : Hemostasis and thrombosis . Biggs,T. (Ed.) , Blackwell Scientific publications oxford .P.143 .

De Fronzo , R.A.; Ferrannini , E. and Koivisto , V. (1983) :

New Concepts in the pathogenesis and of non insulin - dependent diabetes mellitus .

Am -J- Med .; 74 (1A) : 52-81 .

De Fronzo , R.A . (1985) :

Diabetes and the kidney : An update . In Olefsky , J.M. and Sherwin, R.S. (eds) : Diabetes Mellitus ; Management and complications.

chap . 5 , PP 159-221 .

Devonne , T.L.; and Mouray , H. (1978) :

Human macroglobulin and its antitrypsin and antithrombin activities in serum and plasma .

Clin . chem . Acta , 90 : 83 .

Discipio , R.G. and Davis , E.W. (1979):

Characterization of protein S, a γ -carboxy- glutamic acid containing protein from bovine and human plasma .

Biochemistry , 18:899.

Ditzel , J. and Junker ,K.(1972):

Abnormal glomerular filtration rate , renal plasma flow and renal protein excretion in recent and short-term diabetics .

Brit .Med .J. , 2 :13-19 .

Donaldson , V.H. and Ratnoff , O.D.(1965):

Hageman factor : Alterations in physical properties during activation.

Science ,150:754 .

Dullaart R.P.F; Dikkeschei L.D, and Doorenbas H.(1989) :

Alterations in serum lipids and apolipoproteins in male type I (In sulin-dependent) diabetic patients with microalbuminuria .

Diabetologica 32:685-689 .

Dunn FL., Pietri A., Raskin P. (1981) :

Plasma lipid and lipoprotein levels with continuous Subcutaneous insulin infusion in type I diabetes mellitus .

Ann - Intern - Med .95:426-431 .

Durrington P.N.(1982):

High density lipoprotein cholesterol - Methods and clinical significance .

CRC Crit Rev .Clin Lab .Sci 18:31-78 .

Eckel RH, Albers JJ, Cheung MC, Wahl P.W , Lindgren F.T, Bierman EL.(1981):

High density lipoprotein composition in insulin- dependent diabetes mellitus .

Diabetes 30 : 132-138 .

Egeberg , O. (1963):

The blood coagulability in diabetic retinopathy . *Scand .J.Clin .Lab Invest .* 15,522 .

Eisenberg , S.; Bilheimer , D.W. and Levy , R.I. (1972):

The metabolism of VLDL lipoprotein protein studies on the transference of apolipoproteins between plasma lipoprotein. *Biochem . Biophys . Acta* ,280:94.

Eisenberg , S. ; Bilheimer , D.W. and Levy R.I.(1973 a) :

Metabolic conversion of human plasma VLDL Low LDL .
Biochim . Biophys . Acta , 326 : 361 .

Eisenberg , S. and Levy , R.I. (1975):

lipoprotein metabolism .

Adv . Lipid . Res .; 13;1 .

Eisenberg , S.; Patsch , J.R.; Sparrow ,J.T.; Gotto , A.M. and Olivercrona ,T.; (1978) :

VLDL : Removal of apo c-II and C-III during lipolysis in vitro .
J- Biol - Chem .; 254: 12603 .

Eisenberg , S.(1983):

lipoprotein and lipoprotein metabolism .

Klin . wachenschr 61 : 119 .

Elder , G.E; Mayne E.E.; Daly , J.G.; and Kennedy A.L.(1980) :

Antithrombin III activity and other Coagulation changes in proliferative diabetic retinopathy .

Haemostasis 9,288.

EL -Hawary , M.B.; Khayyal , M.T and Isaak , Z.(1981) :

Handbook of pharmacology chapter 31,P.346.

EL .Nahas A.M.& Coles GA.(1986) :

Dietary treatment of chronic renal failure :10 Unanswered questions.

Lancet 1;597.

- Feldt-Rasmussen B., Dinesen , B., Deckert M.(1985) :**
 Enzyme immuno-assay an improved determination of urinary albumin in diabetics with incipient nephropathy .
Scand . J. Clin . Lab . Invest .
 45 : 539 - 44 .
- Feldt-Ramussen B.(1986) :**
 Increased transcapillary escape rate of albumin in type I (insulin-dependent) diabetic patients with microalbuminuria .
Diabetologica 29 : 283 - 286 .
- Feldt - Rasmussen B.; Mathiesen E.; Deckert T.; (1986):**
 Effect of 2 years of strict metabolic control on the progression of incipient nephropathy in insulin - dependent diabetes .
Lancet 1:1300 .
- Feldt-Rasmussen B.; Mathiesen ER.; Deckert T.; et al .;(1987) :**
 Central role for sodium in the pathogenesis of blood pressure changes independent of angiotensin , aldosterone , and catecholamines in type I .(insulin-dependent) diabetes mellitus
Diabetologica 30 : 610 .
- Fergusson , J.E.; Mackay , N.; philip J.A.D. (1975) :**
 Determination of platelet and fibrinogen half-life with selenomethionine : studies on normal and in diabetic subjects .
Clin .Sci .Mol .Med . 49,115 .
- Fey , G.H. (1987):**
 Regulation of acute phase gene expression by inflammatory mediators .
Mol . Biol .Med .; 4:323 .
- Finley P.R.et al , (1978) :**
Clin-Chem-24 , 3 : 931 .
 Quoted from pamphlet of CHOL-HDL .Kit , Sclavo diagnostica .
- Forbes , C.D.(1970) :**
 Inhibition of activated hageman factor and activated plasma thromboplastin antecedent by purified serum C₁ inactivator .
J. Lab .Clin . Med .; 76:809 .
- Fraber , S.D.; Fraber , M.O.; Brewer , G.; Magnes , C.T. and peterson ,R.G.(1991) :**
 Oxygen affinity of hemoglobin and peripheral nerve degeneration in experimental diabetes .
J-Neurol -Scil.; 102(2):204-7.

Francis ,R.B. and Patch , M.J. (1983):

A Functional assay for protein C . in human plasma .
Thromb . Res . ; 32:605-613 .

Freinkel , N.; Josimovich , J.and Session , V. (1980):

Report of workshop chairmen . Workshop .conference on
gestational diabetes . Summary and recommendations .
Diabetes Care , 3 : 499-501 .

Friedewald , W. T.; Levy R.I. and Fredrickson , D.S.(1972) :

Estimation of the concentration of low - density lipoprotein
cholesterol in plasma without use of the preparative ultracentrifuge .
Clin . Chem . ; 18:499 .

Fujikawa , K.; Legaz , M.E. and Davie , E.W. (1972):

Bovine factor X₁ and X₂ (Stuart factor) isolation and
characterization .
Biochemistry , 11:4882.

**Fujikawa , K.; Thompson , A.R.; Legaz . M.E.; Meyer , R.G. and
Davie , E.W. (1973) :**

Isolation and characterization of bovine factor IX .
(christmas factor). *Biochemistry , 12 .*

Fuller , H.; Richter O.; Stein . R.S. (1979) :

Haemostatic Variables associated with diabetes and its
complications
Brit . Med -J-, 2,964.

Fuller P.; Deckmyn H.; and Bertele V. (1979) :

Clotting factors and coagulation inhibitors in diabetic retinopathy .
Thromb . Res . 30 : 407-413 .

Ganda O.P. (1985) :

Pathogenesis of macrovascular disease including the influence of
lipids .

In Joslin's diabetes mellitus , 12 th edn.

philadelphia : Lea and Febiger PP:217-50

Gandolfo , G.; Colwell , J.A. , Levin J. (1974) :

Platelet factor 3 availability in patients affected by pancreatic
diabetes .

Haematologica 59 , 316 .

Gandolfo G.M.; Deangelis A.; Torresi M.; (1980):

Determination of antithrombin III activity by different methods in
diabetic patients .

Haemostasis 9 , 15.

- Ganrot , P.O.; Gydell , K.; and Ekelund , H.(1967):**
 Serum concentration of α_2 - macroglobulin , haptoglobulin , and α_1 -antitrypsin in diabetes mellitus .
Acta Endocrinol . 55, 537 .
- Ganrot , P.O.; and Bjerre , B. (1967):**
 α_1 -antitrypsin and α_1 macroglobulin concentration in serum during pregnancy .
Acta obst . Gynaecol . Scand .; 46:126 .
- Garg , S.K.; Lackner , H.; and Karparkin S.(1972):**
 The increased percentage of megathrombocytes in various clinical disorders .
Annales intern . Med .77, 316.
- Gatling W.; Rowe D.J.F.; Hill R.D. (1988) :**
 Microalbuminuria:An appraisal of assay techniques and urine collection procedures for measuring urinary albumin at low concentrations ,In Mogensen CE(ed) : The kidney and Hypertension in Diabetes Mellitus. Boslon , Martinus Nijhoff publishing PP 41 .
- Gensini , G. F.; Abbate , R. ; Farilla , S. ; and Neri , G.G. (1979) ;**
 Changes in platelet function and blood clotting in diabetes mellitus .
Thrombos. haemostas . 42,983 .
- Ghishan , F.K. and Greene , H. L . (1988) :**
 Liver disease in children with α_1 antitrypsin deficiency .
Hepatology , 8 (2) : 307 - 310 .
- Gill , D. S.; Thompson , C.S. and Dandona P. (1990) :**
 Histamine synthesis and catabolism in various tissues in diabetic rats.
Metabolism , 39 (8) : 815-8 .
- Glonsset JA .; Norum , K. R.; Gjone E. (1983) :**
 Familial lecithin : Cholesterol acyltransferase deficiency . In : Stanburg J.B. wyngaarder JB , Fredricjson DS (eds) . The metabolic basis of inherited disease 5 th ed , Nrew York . McGraw-Hill . 643 .
- Gomen B .; Baensiger , J. and Schonfeld G . (1981) :**
 Nonenzymatic glycosylation of low density lipoproteins in vitro . Effects on cell interactive properties .
Diabetes , 30 : 875 .

Gonen B., White N, Sch onfeld G ., Stor D, Miller P , Santiago J (1985):

Plasma levels of apoprotein B in patients with diabetes mellitus :
The effect of glycemic control .

Metabolism 34:675-679 .

Gonzalez , K .; Colwell , J . A.; Sarji , K . E. and Nair , R . M . (1980):

Effect of metabolic control with insulin on plasma Von Willebrand
Factor activity in diabetes mellitus .

Thrombosis .Res . 17,261 .

Gonzales , G.; Bamtt , P .; Aguayo , J . Chenbeg . H. M . and chlack, L.T.Jr.(1984) :

Direct measurement of polyol pathway - activity in the ocular lens
Diabetes , 22 : 196-9 .

Gordon D.J. and Rifkind BM. (1989) :

High density lipoprotein . the clinical implications of recent studies .
N.Engl-J.Med . 321:1311-1316 .

**Gorsuch , A . N .; Spencer , K . M .; Lister , j .; Mc Nolly , j .M .
Deam , B.M.; Bottago , G . F. and Cudworth , A .G. (1981) :**

Natural history of type I (Insulin - dependent) diabetes mellitus ;
Evidence for a long pre-diabetic peroid .

Lancet , 2 : 1363-5 .

Gotto , A . M.; Gorrry G.A. and Thompson J .A . (1977) :

Relationship between plasma lipid concentration and coronary
artery disease in 496 patients .

Circulation , 56:875 .

Gray , C . H.; Howorth , P . J.N. and Rinsler , M.G. (1985) :

Clinical chemical pathology loth . ed Edward Aenold . Ltd .
Ch . 9 . PP . 152 .

Greaves M.; Preston FE. (1984) :

Haemostatic abnormalities in diabetics .

In jarrett R J (ed.) Diabetes and heart disease . Amsterdam :
Elsevier, PP:45-80 .

Greaves , M.; Pickering C , Knight G . et al .; (1987) :

Changes in the factor VIII complex in diabetic ketoacidosis :
evidence of endothelial cell damage ?

Diabetologica , 30 : 160 - 5 .

Greenfield M. ; Kolterman , O. ; Olefsky , S. and Reaven , G.(1980):
Mechanism of hypertriglyceridaemia in diabetic patients with fasting hyperglycaemia.

Diabetologica 18 : 441 .

Groop , I. C. ; Bottazzo , G.F. and Doniach , D. (1986) :

Islet cell antibodies identify latent type I diabetes in patients aged 35-75 years at diagnosis .

Diabetes , 35 : 237-41 .

Grundy . S.M.(1984) :

Low density lipoprotein cholesterol and apolipoprotein B. Mechanism of atherogenicity .

Am . J . Cardiol , 54:200 .

Groop, L.C.; Bottazzo, G.F. and Doniach, D, (1986):

Islet cell antibodies identify latent type I diabetes in patients aged 35-75 years at diagnosis.

Diabetes, 35:237-41 .

Gruppo , A. R. ; Leimer , P. ; Francis R. ; and Marlar , R.A. (1988) :

Protein C deficiency resulting from possible double heterozygosity and its response to danazol .

Blood , 71 (2) 370-374 .

Harker , L.A.; (1974) :

Haemostasis .Ed . by Davis , F.A.; Co . 2nd edition . Philadelphia P . 19-40 .

Harmoinen , A. ; Vuorinen P. ; Jokela H. (1987) :

Prevalence and causes of microscopic hematuria in type I (Insulin-dependent) diabetic patients with persistent proteinuria .

Diabetologica 30 : 627 .

Harris , M.I. ; Hadden W.C. ; Knowler W.C. ; Bennett P.H. (1985):

International criteria for the diagnosis of diabetes and impaired glucose tolerance .

Diabetic care , 8:562-7 .

Harper , P.L. ; Daly . M. ; Price . J . ; Edgors , P.P and Carrel , R.W.(1991):

Screening for heparin binding variants of antithrombin .

J . clin . Pathol . ; 44 : 477-479 .

Havel , R.J. ; Eder , H.A. and Bragdon J.H. (1955) :

The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum .

- J.clin - Invest* . 34 : 1345 .
- Havel , R.J.; Goldstein , j. L. and Brown , M.S.(1980) :**
 Lipoprotein and lipid transport . In Bondy . P.K. , and Rosenberg ,
 L.I(eds) : Metabolic control and disease . 8th ed .; philadelphia ,
 W.E. Saunders Co .; 393-494 .
- Havel . K . J. (1982) :**
 Familial dysbeta-lipoproteinaemia .
The Medical clinics of North Asmerica , 66(2) :319 .
- Hedner , U.(1973) :**
 Studies on an inhibitor of plasminogen activation in human serum .
Thromb . Diath . Haemorrh . ; 4:12 .
- Hemiark , R. (1980) :**
 Surface activation of blood Coagulation , fibrinolysis and Kinin
 formation .
Nature , 286:31 .
- Hender , u . and Martinson , G . (1978) :**
 Inhibition of activated hageman factor (factor XIIa) by an
 inhibitor of the plasminogen activation (P.A.inhibitor) .
Thromb . Res. ; 12:1015 .
- Hirsh m J . and Brain , E , (1983) :**
 Haemostasis and thrombosis . 2nd edition .
Churchill Livingstone , New York , P.10 .
- Hoffbrand , A . V, and , Lewis, SM. (1983) :**
 Postgraduate Heamatology . 2nd edition . William Heinmann .
Medical Book .ltd .London . P. 603 .
- Hoffbrand, A.V. and Pettit, J.E.(1985) :**
 Essentil Haematology . 2nd edition .
Blackwell Scientific Publications .P.87 .
- Holmquist L . (1980) :**
 Quantitation of human very low density apolipoproteins C1 , C2 ,
 C3 and E by enzyme immunoassay .
J.Immunol . Methods , 34;2234-251 .
- Holmquist L . (1982) ;**
 Quantitation of human serum apolipoprotein B by enzyme
 immunoassay .
Clin . chem . Acta . ; 121 : 327-336 .
- Hommel E., Parving H.H, Mathiesen E, et al (1986) :**
 Effect of captopril on kidney function in insulin-dependent diabetic
 patients with nephropathy .

- Br. Med. J.* 293:467.
- Hormila, p.(1985) :**
Ischemic heart disease and its risk factors in insulin-dependent diabetes.
Academic dissertation . University of Helsinki .
- Hostetter, T. H.; Rennke , H.G. and Brenner, B.M. (1982) :**
The case for intrarenal hypertension in the initiation and progression of diabetic and glomerulopathies .
Am - J. Med .; 72:375-80 .
- Howard B.V. (1987) :**
Lipoprotein metabolism in diabetes mellitus .
J-lipid Res . (in press) .
- Huff , M. W.; Fidge ,N.H. Nestel , P. J.; Billington , T. and Watson, B. (1981) :**
Metabolism of C-apolipoproteins :
Kinetics of C-II, CIII and C-III2 and VLDL - apolipoprotein B. in normal and hyperlipoproteinaemic Subjects .
J. Lipid . Res . 22 : 1235 .
- Hui , B.Y.; Innerariyt , T. L and Mahely , R. W. (1981) :**
Lipoprotein binding to canine hepatic membranes metabolically distinct apo-E and apo-B , E receptors .
J. Biol - chem .; 256:5646 .
- Hui DY.; Brecht WJ.; Hall EA . (1986) :**
Isolation and characterization of the apolipoprotein E receptor from kanine and Human liver .
J . Biol . chem .; 261:4256 .
- Husdan , H. et al .; (1968) :**
Clin . Chem . 14,222-238 .
Quoted from pamphlet of creatinine kit . bio Merieux , laboratory reagents and instruments .
- Hutton , R. A. (1989) :**
Normal Haemostasis . In : postgraduate Haematology . Ed . by Hoffbrand , A.V. and Lewis . SM .; heinamann professional .
Publishing , London ., P.583 .
- Inomata . S' Oosawa Y .; itoh M.; et al (1987) :**
Analysis of urinary proteins in diabetes mellitus , with reference to the relationship between microalbuminuria and diabetic renal lesions.
J.Japan Diab . Soc . 30 : 429 .

- Irvine ; W.J. and Teviot , O . (1979) :**
Immunology of Diabetes Mellitus , Edinburgh . Scientific
puplication.
- Jackson , R. L.; Morrisett , J . D.; Sparrow , J . T.; Segrest , J
.P.;Pownall , H.J.; Smih , I.C.; Hoff , HTF . and Gotto , A . M.
(1974):**
The interaction of apolipoprotein - serine phosphotidycholine .
J.Biol . Chem . 249:5314 .
- Jackson . R.L.; Morrisett , J.D. Gotto , J. (1976) :**
Lipoprotein structure and metabolism .
Physiol . Rev . ; 56:259 .
- Jackson , C. (1980) :**
Chemistry of blood coagulation .
Ann . Rev . Biochem . ; 49:765-811 .
- Jackson . R.L. (1982) :**
Lipid transport and atherosclerosis . Role of apolipoproteins in
vascular smooth nuscles . In : krass III , MF , and Barnes CD (eds).
Topics in physiology , metabolic , ionic and contractile mechanisms.
2nd edition , New York , Academic Press , P.163 .
- Jaspan , J.B.; Herold , K. and Bartleus C.; (1985) :**
Effects of sorbinil therapy in diabetic patients with painful
neuropathy and autonomic neuropathy .
Am .J. Med . ; 79 (5A) : 24-37 .
- Jay R . H.; Jones S.L.; Hill C.E; Richmond W.; Viberti G.C.;
Rampling M.W. and Betteridge D.J.(1991) :**
Blood Rhealogy and cardiovasclar risk facotrs in type -I-diabetes:
Relatlionship with microalbuminuria .
Daibetic . Med . ; 8 (7) : 662-7 .
- Jensen , T. ; Stender S.; and Deckert T. (1988)**
Abnormalities in plasma concentrations of lipoproteins and
fibrinogen in type I (insulin - dependent) diabetic patients with
incereased urinary albumin excretion .
Diabetologica , 31(3) : P:142-5 .
- Jensen , T. ; Richter E.A.; Feldt . Rasmussen B; et al (1988):**
Impaired aerobic work capacity in type I (Insulin-dependent)
diabetic patients with increased urinary albumin excretion .
Br. med . J.

- Johnosn , A. M.; Alper , CA .; Rosen , F.S. and Criag , J.M. (1977) :**
 C1 inhibitor evidence for decreased hepatic synthesis in hereditary angioneurotic oedema .
Science , 173:553-554 .
- Jones , R. L .; and Peterson , C.M. (1979) :**
 Reduced fibrinogen survival in diabetes mellitus : A reversible phenomenon .
J. Clin. Invest. 63,485 .
- Jones , R.L.; and Peterson , C.M.; (1981) :**
 The fluid phase of coagulation and accelerated atherosclerosis of diabetes .
Diabetes 30 (suppl.)233 .
- Jones , R.L. ; pardise , C. and Petersson , C.M. (1981) ;**
 Platelet survival in patients with diabetes mellitus :
Diabetes 30 , 486 .
- Jonse E.W., Mitchell JRA.(1983):**
 Venous thrombosis in diabetes mellitus .
Diabetologica 25:502-5 .
- Jones , R. R.; and Peterson , C.M.(1984) :**
 The platelet and fluid phase of coagulation :Relationship to diabetes. In Recent advances in diabetes 1st Edition . Eds: M. Nattrass. And V.J. Santiago . Churchill livingstone , Edinburgh, London , Melbourne and . New York , P.181 .
- Jones D.Y.; Judd J.T.; Taylor .; Campbell W.S.; Nair P.P. (1988):**
 Menstrual Cycle effct on plasma lipids .
Metabolism 37:1-2 .
- Jones , S.L.; Colse C.F.; Mattock M.B. ; Jarrett R .J.; Keen H.; and Viberti G.C.(1989):**
 Plasma lipid and coagulation facotr concentrations in insulin-dependent diabetics with microalbuminuria .
BMJ. Feb . 25;298(6672) P: 487-90 .
- Kahn.; C.R.; Flier J.S.; Bar, R.S.; Archer, J.A.; Gorden, P.; martin, M.M. and Roth, J. (1976):**
 The syndromes of insulin resistance and acanthosis negricans:Insulin disorders in man.
N.Engl. J. med.; 294:739-45.

- Kane, J.P.; Sato, T.; Hamilton, R.L.; and Havel, R.J. 91985):**
 Apoprotein composition of very low density lipoproteins of human serum.
J.Clin. Invest.; 56:1622.
- Kane, J.P.; Hardman, D.A. and Paulus, H.E. (1980):**
 Heterogeneity of apolipoprotein B isolation of a new species from human chylomicrons.
Proc.Natl. Acad. Sci : USA 77 : 2465
- Kane J.P and Malloy M.J. (1991):**
 Disorders of lipoprotein metabolism. In: Greenspan F.S. (ed.) Basic and clinical endocrinology. 3rd ed. Librairie du liban PP3.
- Karlin J.B.; Juhn, D.J.; Goldberg R. et al (1978):**
 Recent progress in the development of radioimmunoassays for human serum lipoproteins.
Am. Clin. Lab. Sci.; 8 : 142-154.
- Kapelrud, H.; Bdgstad H.J.; Dahi-Jorgensen K.; Berg K. and Hanssen K.F. (1991):**
 Serum LP (a) lipoprotein concentrations in insulin-dependent diabetic patients with microalbuminuria.
B.M.J. 21; 303 (6804):75-8.
- Kaplan, A.P. and Silverberg, M. (1987):**
 The coagulation kinin pathway of human plasma.
Blood, 70 (1) : 1-15.
- Kashyap M.L.; Srivastava L.S.; Chen C.Y. (1977):**
 Radioimmunoassay of human apolipoprotein CII: A study of normal and hypertriglyceridemic subjects.
J. Clin. Invest. 60:171-180.
- Kawano, T.; Morimoto, K. and Uemura, Y. (1968):**
 Urokinase inhibitor in human placenta.
J.Biol. Chem.; 236:8140.
- Kaye, J.P. and Galton D.S. (1975):**
 Triglyceride-production rates in patients with type IV hypertriglyceridaemia.
Lancet, 1:1005.
- Kazmier, F.; Bowie, W.O.; Fallen, W.M.; and Zimmerman, B.R. (1981):**
 A prospective study of peripheral occlusive arterial disease in diabetes IV platelet and plasma functions.
Myoclinical proceedings 16,243.

Keen, H.; and Jarrett, P.G. (1979):

The WHO multinational study of vascular disease in diabetes;
Macrovascular disease prevalence.
Diabetic Care, 2:187-95.

Keen, H. (1988):

Chronic Complications of diabetes mellitus. In : Galloway, T.A;
Potrin, J.H and shuman, C.R. (eds): Diabetes mellitus, 9th ed. by Eli
Lilly and company, Indianapolis, Inadian PP: 278-305.

Keller H, (1986):

Klinisch. Chemische Labordiagnostik fur die praxis, 91-148, thieme
verlag, stuttgart.

Kelly. D.A.; and Summerfield J.A. (1987):

Haemstasis in liver disease.
Semin. Liv. dis.; 7 : 182.

Kernoff, P.B. and Mc Nicol, G.P. (1977):

Normal and abnormal fibrinolysis.
Brit. Med. Bull.; 33 (3) : 239-244.

Kessaniemi Y.A. and Grundy S.M. (1983):

Overproduction of low density lipoprotein associated with coronary
heart disease.
Arteriosclerosis 3 (1):40.

Kimura, S.; Tamaki, T. and Aoki. N. (1985):

Acceleration of fibrinolysis by the N-terminal peptide of α_2 plasmin
inhibitor.
Blood, (1) : 157:160.

Kissebah A.H; Alfarsi, S.; Evans D.J. and Adams P.W. (1982):

Intergrated regulation of very low density lipoprotein triglyceride
and apolipoprotein-B kinetics in non-insulin-dependent diabetes
mellitus.
Diabetes, 31:217.

Kissebah, A.H; Alfarsi, S.; Evans D.J. and Adams P.W. (1983):

Plasma LDL transport. Kinetics in NIDDM.
J.Clin. Einvest.; 71:655.

Kluft, C. (1978):

Blood fibrinolysis. Proactivators and activators in human plasma.
Ph; D.; Thesis, Dutch Efficiency Burreau, Pijnacker.
Quated from Thomson. (1980).

Kluft, C. (1980):

Plasminogen proactivators and activators in human plasma. factor XII dependent fibrinolysis. Peeters, H. (Ed.), Properties of the Biological fluids, Vol. XX VIII pergamon press, Oxford : 375-378.

Kluft, C. (1981):

Quantitation and behaviour of extrinsic plasminogen activator in blood. progress/in Fibrinolysis churchill livingstone, Edenbergh, 3 : 24-30.

Koivisto, V.A. Huttanen, N.P.; Vierikko, P. (1981):

Continuous subcutaneous insulin infusion corrects exercise - induced albuminuria in Juvenile diabetes.

Brit. Med. J.; 282:778-9.

Koj, A. (1974):

Acute phase reactions. In : Structure and function of plasma proteins. Ed. by Allison, A.C. Plenum, New york, P. 73-131.

Korninger, C.; Stassen, J.M. and collen, D. (1982):

turnover of human extrinsic (tissue type) Plasminogen activator in rabbits.

Thromb. Haemost.; 46:658-661.

Kostner, G. (1974):

Studies on the composition and structure of human serum lipoproteins. Isolation and partial characterization of apolipoprotein A-III.

Biochem. Biophys. Acta, 336:383.

Kostner G.M. (1976):

Clin. Chem. 27:25.

Quoted from the pamphlet of chol-HDL Kit. Sclavo diagnostica.

Kramer, B.K.; Jesse U.; Ress K.M.; et al, (1987):

Enzyme-linked immunosorbent assay for urinary albumin at low concentrations.

Clin. Chem. 33:609.

Kreisberger, A. (1983):

Lipids, Lipoproteins, apoproteins and atherosclerosis.

Ann. Int. med. 99 (5) : 713.

Kroc Collaborative study Group (1984):

Blood glucose control and the evolution of diabetic retinopathy and albuminuria. A preliminary multicenter trial.

N. Engl. J. Med. 311:365.

Krzymicr F.J.; Bowie E.J.W, O'Fallon W.M, Zimmerman B.R, O'Smcendson P.T, palumbo P.J. (1981):

A prospective study of peripheral occlusive arterial disease in Diabetes. IV. Platelet and plasma Functions.

Mayo clin. Proc. 56:243-53.

Krzywanek, H.J.; Breddin, K.; Schoffting K.; and Uberla, K. (1984):

Platelet aggregation as a risk factor in the Diabetic. Preliminary results of a prospective study on enhanced platelet aggregation of vascular disease in VIII International congress on thrombosis. The mediterranean league against thromboembolic diseases. P. 261.

Kupin WL, Cortes P, Dumler F, et al (1987):

Effect on renal function of a change from high to moderate protein intake Effect on renal function of a change from high to moderate protein intake in type I diabetic patients. *Diabetes* 3 : 73.

Kwaan H.C.; Colwell F.A.; Suwanwela N. (1972):

Disseminated intravascular coagulation in diabetes mellitus with reference to the role of increased platelet aggregation.

Diabetes 21:108-13.

Laak, K.; and Vennerod, A.M. (1974):

Factor XII induced fibrinolysis. Studies on separation of prekallikrein plasminogen proactivators and factor XI in human plasma.

Thromb. Res-4; 255-302.

Laakso M.; Pyorala K.; sarlund H.; Voutiainen E. (1986):

Lipid and lipoprotein abnormalities associated with coronary heart disease in patients with insulin dependent diabetes mellitus.

Arteriosclerosis 6 : 679-684.

Latner, A.L. (1975):

"Clinical Biochemistry" 7th ed. WB. Saunders company. philadelphia, London, Toronto ch.2 PP. 94.

Lee J.K. and Carrico R.T. (1983):

Substrate labeled fluorescence immunoassay for the determination of serum protein. In lippel. K.(ed) : proceedings of the workshop on the Apolipoprotein Quantification. U.S. Department of Health and Human Services, National Institutes of Health publication No. 83-12 P. 332 - 342.

- Leebeek, F.W.; Kluft, C.; knot E.A.; Los, P.; Cohen, A.F. and Six, A.J. (1990):**
 Plasmin inhibitors in the prevention of systemic effects during thrombolytic therapy : specific role of the plasminogen binding form of α_2 antiplasmin.
J.Am. Coll. cardiol.; 15 (6) : 1212-20 (Abstract).
- Leswk, A.; Terminella, A. and Groff. C.D. (1967):**
 Study of human urokinase.
Thromb. Dith. Haemorrh.; 18 : 293.
- Leung, L.; Harpel, P.; Nachman, R. and Kbellina, E. (1983):**
 Histidine rich glycoprotein is present in human platelet and is released following thrombin stimulation.
Blood, 62 : 101-17.
- Levin, E.G. (1983):**
 Latent tissue plasminogen activator produced by human endothelial cells in culture: Evidence for an enzyme inhibitor complex.
Proc. Nat. Acad. Sci.; USA.; 80 : 6804.
- Lijnen, H.R; Hoylaerts, M. and Collen, D. (1980):**
 Isolation and characterization of a human plasma protein with affinity for the lysine binding sites in plasminogen.
J.Biol. Chem.; 255 : 10214-22
- Lijnen, H.R. and Collen, D. (1984):**
 Molecule mechanism of fibrinolysis.
Adv. Exp. Med. Biol.; 17 : 217.
- Lindahl, U.; Backstorm, G.; Hook, M.; Thunberg, L. Fransson L. A. Linker, A. (1979):**
 Structure of the antithrombin binding site in heparin.
Proceedings of the National Academy of science, USA, 79 : 3198-3202.
- Lindgren F.T.; Jensen L.C. and Hatch F.R. (1972):**
 The isolation and quantitative analysis of serum lipoproteins. In Nelson, G.J. (ed) : Blood lipids and lipoproteins-Quantitation, composition and metabolism. New York, Interscience.
- Lipid Research clinics program (1984):**
 The lipid Research clinics cornory primary prevention trial Results.
JAMA 251 : 351-374.
- Lippel K.; Tyroler, H.; Eder H.; Gatto, A.Jr. and Vahouny, G. (1981):**
 Relationship of hypertriglyceridaemia to atherosclerosis.

- Arteriosclerosis*, 1 (6) : 406.
- Lloyd J. and Reckless J.P. (1993):**
Lipid abnormalities in diabetes.
Postgraduate doctor, 16 (2) : 66-72
- Loller, P. and Owen, W.G. (1980):**
Clearance of thrombin from circulation in rabbits by high affinity binding sites on endothelium. Possible role in the inactivation of thrombin by antithrombin III.
J. Clin-Invest.; 66 : 1222.
- Lopes-Virella M.F.; Wohltmann H.J.; Mayfield R.K.; Loadholt C.B.; Colewell J.A. (1983):**
Effect of metabolic control on lipid, lipoprotein and apolipoprotein levels in 55 insulin dependent diabetic patients. A longitudinal study.
Diabetes 32 : 20-25.
- Lorand, L.; Rule, N.G.; Forlanetto, R.; Jacobson, A.; Downey, J. Oner, N. and Brunner, J. (1968):**
Amino specificity in transpeptidation. Inhibition of fibrin cross-linking.
Biochemistry, 7 : 1214.
- Lyons T.J.; Klein R.L.; Baynes J.; Lopes-Virella M.F. (1986):**
VLDL and LDL from diabetic patients stimulate increased cholesteryl ester (CE) synthesis in human monocyte derived macrophages.
Diabetes 35 (Suppl. 1) : 67A.
- Lyons J.J.; Klein R.L.; Baynes J.W.; Stevenson H.C, Lopes-virella M.F. (1987):**
Stimulation of cholesteryl ester synthesis in human monocyte-derived macrophages by low-density lipoproteins from type I (insulin-dependent) diabetic patients: the influence of non-enzymatic glycosylation of low density lipoproteins.
Diabetologica. 30 : 916-923.
- Macfarlane, R.G. (1964):**
An enzyme cascade in the blood clotting mechanism and its function as a biochemical amplifier.
Nature, 202 : 498-499
- Mackey, N.; and Hume, R. (1964):**
Fibrinolytic activity in diabetes mellitus.
Scotl. med. J. 9, 359

- Mackie, M.; Bennett, B.; Ogston D.; and Douglas, A.S. (1978):**
 Familial thrombosis : Inherited deficiency of antithrombin III.
Brit. Med. J.; 1 : 13-138.
- Mahley, R.W. (1981):**
 Cellular and molecular biology of lipoprotein metabolism in atherosclerosis.
Diabetes, 30 (Suppl. 2) : 60.
- Mahley, R.W.; and Innerarity T.L. (1983):**
 Lipoprotein receptors and cholesterol homestasis.
Biochim. Biophys. Acta 737 : 197.
- Mahley, R.W.; Innerarity T.L.; Rall S.C. Jr.; Weisgraber K.H; (1984):**
 Plasma lipoproteins; apolipoprotein structure and function
J. Lipid Res. 25 : 1277.
- Mahly R.W. (1990):**
 Biochemistry and phsyology of lipid and lipoprotein metabolism. In:
 Becker K.L. (ed.) principles and practice of endocrinology and metabolism J.B.Lippincott company. Philadelphia PP.1219.
- Malmendier C.L.; Lontie J.F.; Grutmen G.A.; Delcroix C. (1988):**
 Metabolism of apolipoprotein C.III in normolipemic human subjects.
Atherosclerosis 69 : 51.
- Mann, K.G.; Heldebrant, C.M. and Fass, D.N. (1973):**
 Multiple active froms of thrombin. polypeptide chain location of active site serine and carbohydrate.
J. Biol. Chem. 248 : 1868.
- Marlar, R.A.; Klies, A.J. and Griffin, J.H. (1982):**
 Mechanism of action of human activated protein C.; a thrombin-dependent anticoagulant enzyme.
Blood, 59 : 1067-1072.
- Marre M.; Leblanc H.; Suarez L. et al (1987):**
 Converting enzyme. inhibition and kidney function in normotensive diabetic patients with persistent microalbminuria.
Br. Med. J. 294 : 1184.
- Masperi. R.; Mussoni, L; Bini, A.; Donartic, M.B.; and Chiumello, G. (1980):**
 Haemostatic variables in uncomplicated diabetic children.
Thromb. haemost. 43 , 227.

- Mathiesen, E.R.; Saurbrey N.; Hommel E.; et al, (1986):**
Prevalence of microalbuminuria in children with type I (Insulin-dependent) diabetes mellitus.
Diabetologica 29 : 640.
- Mathews, J.H.; O'conner, J.F.; Hearnshaw, J.R.; and Wood, J.K. (1979):**
Beta-thromboglobulin and glycosylated haemoglobin in diabetes mellitus.
Scand. J. H. aematol. 23, 421.
- Macser, S.; Steffes, M.; Brown, D. (1981):**
The kidney in diabetes.
Am. J. med.; 70 : 603-12.
- Mauer, S.; Steffes, M.; Ellis, E.N.; et al, (1984):**
Structural, Functional relationships in diabetic nephropathy.
J. Clin. Invest.; 74:1143-55.
- Mayne, J.M.; Bridges, J.M.; and Weewor J.A. (1970):**
Platelet plasma fibrinogen and factor VIII levels in diabetes.
Diabetologica , 436.
- Mc Conathy, W.J. and Alaupovic, P. (1976):**
Studies on the isolation and partial characterization of apolipoprotein D.
In human plasma Biochemistry, 15 : 515.
- Mckee, P.A.; Mattock, P. and Hill, R.L. (1970):**
Subunit structure of human fibrinogen, Soluble fibrin and cross-linked insoluble fibrin.
Proc. Nat. Acad. Sci. USA.; : 738.
- Mc Millan, D.E. (1976):**
Plasma protein changes, Blood viscosity and diabetic microangiopathy.
Diabetes 25 (Suppl. 2), 58.
- Meiattini F, Et al.; (1978):**
Clin. Chem. 24 : 2161.
Quoted from pamphlet of cholestest kit. Sclavo diagnostica.
- Menzel, H.J.; Kladetzky, H.G. and Assman, G. (1882):**
Apolipoprotein A.IV isoform, isoelectric focusing identification.
J. Lipid Res.; 23 : 915.
- Messmore,, H.L. Jr. (1982):**
Natural inhibitors of the coagulation system.
Semin. Thromb. Haemost.; 8 : 267.

Metzmann, E, (1985):

Protein Quantitation on both branches of the Heidelberger curve by monitoring the kinetic of immunoprecipitation.

Behring institute Mitteilungen 78, 167-175.

Microalbuminuria Collaborative Study Group UK (1987):

Microalbuminuria and glycemic control. In Diabetic complications '87. Multicenter and/or prospective study session on Diabetic complications. Consiglio Nazionale delle Ricerche, Rome. Italy.

Miller, N.E.; Forde O.H.; Thelle, P.S. and Mjas, O.D. (1977):

The Tromso heart study. High density lipoprotein and coronary heart disease : a prospective case-control study.

Lancet, 1 : 965.

Mogensen C.E, (1971):

Kidney function and glomerular permeability to macromolecules in early juvenile diabetes.

Scand. J. Clin. Lab. Invest. 28 : 79.

Mogensen, C.E; Steffes, N.W.; Deckert, T.; et al.; (1981):

Functional and morphological renal manifestations in diabetes mellitus.

Diabetologica, 21 : 89-93.

Mogensen C.E. (1982):

Long term antihypertensive treatment inhibiting progression of diabetic nephropathy.

Br. Med. J. 1 : 2.

Mogensen, C.E. (1983):

A complete screening of urinary albumin concentration in an unselected diabetic outpatient clinic population.

Diabetic Nephropathy 2 (4) : II.

Mogensen C.E.; Christensen C.K.; Vittinghus E. (1983):

The stages in diabetic renal disease, with emphasis on the stage of incipient diabetic nephropathy.

Diabetes. 32 : 4.

Mogensen C.E. (1984):

Microalbuminuria predicts clinical proteinuria and early mortality in maturity onset diabetes.

N. Engl. J. Med. 310 : 356.

Mogensen, C.E; Christensen C.K. (1984):

Predicting diabetic nephropathy in insulin-dependent patients.

N. Engl. J. Med. 311 : 89.

- Mogensen, C.E.; (1986):**
Early glomerular hyperfiltration in insulin-dependent diabetics and late nephropathy.
Scand. J. Clin. Lab. Invest. 46 : 201.
- Mogensen, C.E.; Chachati A.; Christensen C.K.; et al, (1986):**
Microalbuminuria : An early marker of renal involvement in diabetes.
Uremia Invest. 9 : 85.
- Mogensen, C.E. (1987):**
Microalbuminuria as a predictor of clinical diabetic nephropathy.
Kidney Int. 31 : 673.
- Mogensen C.E. (1988):**
Management of diabetic renal involvement and disease.
Lancet 1 : 867.
- Mogensen C.E.; Christensen C.K.; Christiansen J.S. et al (1988):**
Early hyperfiltration and late renal damage in insulin dependent diabetes.
Pediatr. Adolesc Endocrinol 17 : 197.
- Mogensen C.E. , and Schmitz S.F. , (1988) :**
The Kidney and hypertension in Diabetes Mellitus. Boston, Martinus Nijhoff Publishing.
- Moroi, M. and Aoki, N. (1976 a):**
Isolation and characterization of α_2 plasmin inhibitor from human plasma.
J. Biol. Chem.; 251 : 395.
- Moroi, M. and Aoki; N. (1976 b):**
Isolation and characterization of α_2 plasmin inhibitor from human plasma. A novel proteinase inhibitor which inhibits activated induced clot lysis.
J. Biol. Chem.; 251 : 5956.
- Morrisett, J.D.; David, J.S.K; Pownall, H.J. and Gatto, A.M. (1973):**
Interaction of an apolipoprotein (apo LP-alanine) with phosphatidyl choline.
Biochemistry, 12 : 1290.
- Morrisett, J.D.; Jackson, R.L and Gatto, A.M.; (1975):**
Lipoprotein structure and function.
Annu. Rev. biochem.; 44 : 183.
- Moorhead J.F.; Chan M.K.; El-Nahas M.; Varghese Z. (1982):**
Lipid nephrotoxicity in chronic progressive glomerular and tubulointerstitial disease.

- Lancet* 11 : 1309-1311.
- Muhlhauser, I.; Schernthaner, G.; Silberboauer, k.; and Kaliman, J. (1980):**
Platelet proteins (beta - TG and PE4) in atherosclerosis and related disease.
Artery 8,73.
- Muller-Esterl, W. and fritz, H. (1984):**
Human Kininogens and their function in the kallikrein-Kinin systems.
Adv. Exp. med. Biol.; 167 : 41.
- Muramatsu, S.; Mizuno, Y.; Murayama, H.; and Ikemoto, S. (1990):**
Hereditary AT III deficiency with superior sagittal sinus thrombosis : evidence for a possible mutation starting in the mother of propositus.
Thromb. Res.; 57 (4) : 593-660 (Abstract).
- Mulec H.; Johnson S.A. and Bjork S. (1990):**
Relation between serum cholesterol and diabetic nephropathy.
Lancet, 335 : 1537-38.
- National Diabetes Data Group, (1979):**
Classification and diagnosis of diabetes mellitus other categories of glucose intolerance.
Diabetes, 28 : 1039-57.
- Nelsestuen, G.L.; and Suttie, T.W, (1972):**
Purification and properties of abnormal prothrombin; protein produced by dicumarol treated Cow. A comparison to normal prothrombin.
J. Biol. Chem.; 247 : 9176.
- Niazy. S.; Feldt-Rasmussen, B.; Deckert T.; (1987):**
Microalbuminuria in insulin-dependent diabetes: Prevalance and practical consequences.
J. Diabetic Complications 1 : 76.
- Nikkila E.A; Huttunen, J.K. and Enholm, C. (1977):**
Postheparin plasma lipoprotein lipase and hepdtic lipase in D.M. Relationship to plasma triglyceride metabolism.
Diabetes, 2 : 11.
- Nikkila. E.A.; (1978):**
Triglyceride metabolism in D.M.
Prog. Biochem. Pharmacol.; 8 : 271.

Nikkila E.A.; and Hormila P. (1978):

Serum lipids and lipoproteins in insulin treated diabetes.
Demonstration of increased high density lipoprotein concentrations.
Diabetes 27 : 1078-1086.

Nikkila E.A. (1981):

High density lipoproteins in diabetes.
Diabetes 30 : 82-87.

Nikkila E.A. (1983):

Familial lipoprotein deficiency and related disorders of metabolism.
In: Stanburg JB, Wyngaarden JB, Fredrickson DS. (eds). The metabolic basis of inherited disease, 5th ed. New York : MCGraw-Hill, 622.

Nikkila E.A. (1984):

Plasma lipid and lipoprotein abnormalities in diabetes. In : Jarrett (ed). Diabetes and heart disease. Elsevier Science Publishers, Amsterdam. PP:133-167.

Nikkila, E.A; Hormila P. and Gibbons G.F. (1986):

Hyperlipidemia of diabetes.
Clin. Sci.; 71 : 477-86.

Nikkila E.A.; and Helve E.; (1986):

Antiatherogenic changes in lipoprotein induced by continuous insulin infusion therapy.
Diabetologica 29 : 576 A.

Nilson, E.; Carfinkel, A.S. and Schotz, M. (1980):

Lipolytic enzymes and plasma lipoproteins metabolism.
Ann. Rev. Biochem. 49 : 668.

Noma A.; Hate, Y.; and Goto Y.; (1991):

Quantitation of serum apolipoprotein A.I, A-II, B, CII, C-III and E in healthy Japanese by turbidimetric immunoassay : Reference values, and age - and sex-related differences.
Clinica chemical Acta, 199:147-158.

Nordoy, A.; and Rudset, J.M. (1970):

platelet phospholipids and their function in patients with juvenile diabetes and maturity onset diabetes.
Diabetes 19, 698.

Nossel, H.L. (1976):

The contact system, human blood coagulation, haemostasis and thrombosis. Biggs, R. (Ed) Blackwell Scientific Publications, Oxford, P. 81.

Ockelford, P.A.; Carter, C.J.; Cerkus, A.; Smith, C.A.; and Hirsh, J. (1982):

Comparison of the in vivo haemorrhagic and antithrombotic activity of a low antithrombin III affinity heparin Fraction.

Thromb. Res.; 27 : 679-690.

O'Donnell, M.J.; Le-Guen-CA.; Lawson. N.; Gyde. O.H.; and Barnett A.H.; (1991):

Platelet behaviour and haemostatic variables in type I (Insulin-dependent) Diabetic patients with and without albuminuria.

Diabetic. Med. 8 (7) : 624-8.

Ogston, D.; Bennett. B.; Herbert, P. and Douglas, A.S.; (1973):

The inhibition of urokinase by α_2 macroglobulin.

Clin. Sci.; 44 : 73.

Ogston, D. and Bennett, B. (1977):

Naturally occurring inhibitors of coagulation. In : Haemostasis Biochemistry, Physiology, and pathology. (Eds.). Ogston, D. and Bennett, B.; London. P. 202.

Olofson, S.; McConathy, W.J. and Alaupovic, P. (1978):

Isolation and partial characterization of a new acidic apolipoprotein (apolipoprotein F). From high density lipoprotein of human plasma
Biochemistry, 17 (6) : 1032.

Parker , C.J.; Frame , R.N.; and EL-Stod , M.R.; (1988) :

Vitronectin (Protein S) augments the functional activity of monocyte receptors for IgG and Complement C3b .

Blood , 71 (1) : 86-93 .

Parving H.H. (1986) :

Letter . *Br. Med. J.* 292:558 .

Parving H.H.; Andersen A.R.; Smidt U.M. , et al (1987) :

Effect of antihypertensive treatment on kidney function in diabetic nephropathy .

Br. Med. J. 294:1443 .

Parving H.H.; Hommel E.; Mathiesen E.; et al (1988) :

Prevalence of microalbuminuria , arterial hypertension , retinopathy and neuropathy in patients with insulin-dependent diabetes .

Br. Med. J. 296:156 .

Patrassi , G.M.; Vettor R. and Padovan D. (1982) :

Contact phase of blood coagulation in D.M.

Europ. J. Clin. Invest. ; 12:307-11

Patrassi G.M.; Vettor R. , and Padovean (1982) :

Contact phase of blood coagulation in D.M. Europ .

J.Clin . Invest . ; 12:307-11 .

Patton C.J.and Crouch S.R.(1977) :

Anal chem . ; 49,464-469 .

Quoted from pamphlet of urea kit.bio Merieux laboratory reagents and instruments .

Paul , S.B.; Robert , I. L.; and Basil , M.R. (1991) :

Lipids and Dyslipoproteinemia. In John B.H(eds. Clinicial , Diagnosis and Management by laboratory methods . 18th ed W. B . Saunders Company . Philadelphia , London Tokyo , PP.193 .

Pietri A.O.; Dunn F.L. , Grundy S.M.; Raskin P. (1983) :

The effect of continous subcutaneous insulin infusion on very -low-density lipoprotein triglyceride metabolism in type I diabetes mellitus.

Diabetes 32:75-81 .

Pirart , J. (1978):

Diabetes Mellitus and its degenerative complications : a prospective study of 4,400 patients observed between 1947 and 1973 .

Diabetes Care, 1 : 168:88 (Part 1) , 252-63 (Part. 2) .

Prager R.; Schernthaner , G.; Kostner G.M.; Muhlhauser , I .; Zechner, R. and Dorda W. (1982) :

Effect of bezafibrate on plasma lipids, Apolipoprotein AI , AII and B and LCAT ativity in hyperlipidemic non insulin dependent diabetics .

Atherosclerosis , 43:321 .

Prowse, C.V. and Cash, J.D. (1984):

Physiologic and pharmacologic enhancement of fibrinolysis.

Seminar in thrombosis and Haemostasis, 4 : 64-68.

Pyorala K.; Laakso M.; Uusitupa M. (1987):

Diabetes and atherosclerosis : an epdidemiologic view.

Diabetes.Metab. Rev. 3 : 463-524.

Radcliffe , R. , and Heinze , J. (1978) :

Isolation of plasminogen activator from human plasma by chromatography on lysine -sepharose .

Arch. Biochem . Biophys.; 189:185 .

- Pak K., Beck P.; Udvardy M.; Pfliegler G.; Misz M.; Boda Z.(1983):**
 Plasma levels of of beta-thromboglobulin and factor VIII-related antigen in diabetic childrens and adults ,
Thromb . Res .; 29:155-62 .
- Ramirez L.; Rosenstock J.; Arauz . C.; Hellenbrand D.; and Raskin-P . (1991) :**
 Low prevalence of microalbuminuria in normotensive patients with insulin-dependent diabetes mellitus .
Diabetic - Res - Clin-Pract - 12(2) : 58-90 .
- Raskin ,P.; and Rosenstock , J. (1986) :**
 Blood glucose control and diabetic Complications .
Am. Intern. Med.; 105 : 254-63 .
- Ratnoff , O.D.; Pensky , J.; Ogeston , D. and Naff , G.B. (1969) :**
 Inhibition of plasmin , plasma kallikrein plasma permeability factor and CI subcomplement of the first component of complement by CI esterase imhibitor .
J.Exp . Med .; 129 :315 .
- Raum , D.; Marcus . D.; Alper C.A.; Levey , R.; Taylor , R.D.; and Starzl . T.E. (1980) :**
 Synthesis of human plasminogen by the liver .
Science , 208 : 1036-37 .
- Reaven , G.M.; and Steiner , G. (1981) :**
 Proceedings of a conference on diabetes and atherosclerosis .
Diabetes, 30 (Supp. 2) P.1-110 .
- Revak, S.D.; Coahrane , C.G.; Johnston, A. R. and Hugh, J.E. (1974):**
 Structural changes accompanying enzymatic activations of human Hageman factor .
J. Chin. Invest. 54:619 .
- Ridley , O.D.; Ledingham , S.J.; Lennox , S.C.; Burman , J.F.; Chang , H.I.; Sheffield , E.A.; Talbot , S. and Bevan, D. (1990):**
 Protein C deficiency associated with massive cerebral thrombosis following open heart sugrery .
J.Cardiovasc . Surg . (Torino), 31 (2) : 249-51(Abstract) .
- Robbins , K.C.; ummaria . L .; Hisiech ,B.; and Shalsa . R.H. (1967):**
 Mechanism of activation of human plasminogen to plasmin .
J.Biol Chem.; 242:2333.
- Robbins, K.C. (1982) :**
 Fibrinolysis In : K waan , H. C.and Bowie , E.J. (Eds .),London ,

- W.B. Saunders company . PP . 23 .
- Ross R., (1986) :**
The pathogenesis of atherosclerosis : an update .
N. Engl . J. Med 314-488-500 .
- Rosseneu M.; Versament R.; Steinberg K.K. et al .; (1983) :**
Some Consideration of methodology and standarization of apo-lipoprotein B immunoassay .
Clin. Chem. 29 : 427 - 433 .
- Rpssemeu M.; (1983) :**
Immunonephlometric assay of apolipoproteins .
In : proceedings of workshop on apolipoprotein quantification . NIH Publication No.80-1266. Bethesda, 305 - 313
- Sakata , J. and Aoki , N. (1982) :**
Significance of cross-linking of α_2 plasmin inhibitor to fibrin inhibition of fibrinolysis and in haemostasis .
J. Clin. Invest . ; 60 : 536 .
- Salans , L. B.; Knitte , J.L.; Hirsch , J.(1983) :**
Obesity glucose intolerance and diabetes Mellitus . In :-In Ellenderg , M.; Rifkin H. (eds) ; *Diabetes Mellitus : Theory and practice* , 3rd ed . New Hyde Park , Ny , Medical Examination publishing Co.; Inc .; Chap. 22 , PP 469-79 .
- Saudek , C.D. and Young N.L. (1981) :**
Cholesterol metabolism in diabetes mellitus. The role of diet .
Diabetes , 30 (Suppl . 2) : 76.
- Schaffer , E.J.; Eisenberg , S. and Levy , R.I. (1978 a) :**
Lipoprotein apoprotein metabolism .
J. lipid. Res . ; 19:667 .
- Scherntz G.; Kostner G.M.; Kieplinger H., Prager R., Muhlhauser (1983):**
Apolipoproteins (A-I .; A-II,B), Lp (a) lipoprotein and lecithin cholesterol acyltransferase activity in diabetes mellitus .
Atherosclerosis 49:277-293 .
- Schiff, L. and Schiff , E.R. (1982) :**
Disordered haemostasis in hepatic disease . In : *Disease of the liver* , 5th edition , JB. Lippincott , Co . , philadelphia , Toronto , P.237-258 .
- Schmit , J.M.; (1964) :**
Thesis. Lyon. Quoted from pamphlet of total lipid Kit . Borhringer Mannheim . GmbH Diagnostica . 1990 .

Schmidt , S.B.; Wasserman A.G.; Meusing ,R.A.; schlesselman , S.E. Larosa , J.C. and Ross , A.M. (1985) :

Lipoprotein and apolipoprotein levels in angiography defined coronary atherosclerosis .

Am .J.Cardiology , 55 : 1459 .

Schmitz . A.; and Ingerslev J. (1990):

Haemostatic measures in type 2 diabetic patients with microalbuminuria .

Diabetic . Med . J . ; (6); P 521-5 .

Schrier , S.L. (1981) :

Disorders of haemostasis and Coagulation .

Scientific American Medicine , IV:1-36 .

Scott , C.F.; Silver , L.D .; Schapira , M. and Cloman R.W. (1984) :

Cleavage of human high molecular weight kininogen markedly enhances its coagulant activity. Evidence that this molecule exists as a pro-cofactor.

J.Clin .Invest .; 73:954 .

Seidel , D. (1982) :

Serum triacylglycerol in coronary artery disease .

Ann . Clin . Biochem .; 14:278 .

Shahian , D.M.and Lveine , J.D. (1990):

Open heart surgery in a patient with homozygous α_2 antiplasmin deficiency . Perioperative / strategies in the first reported case .

Chest , 97(6) : 1488-90 .

Shepherd J. and Packard C.J. (1987) :

Lipid transport through the plasma , the metabolic basis of hyperlipidemia .

Bailliere's clinical endocrinology and metabolism.I (3):495 .

Shernthaner , G; Muhlhauser , I.; and Silberbauer , K.; (1979):

Beta thromboglobulin lowered by dipyridamole in diabetes .

Letter , Lancet .; 2 , 748 .

Singh , R.G.; Gambhir , I.S.; Dube , B.; and Gupta , S.(1982):

The fibrinolytic system pattern in diabetic nephropathy .

Asian Med .J.25(4), 301 .

Sinderman A.D.; Wolfson , C. and Teng B.(1982) :

Association of hyperapobetalipoproteinemia with endogenous hypertriglyceridemia and atherosclerosis .

Ann . Intern . Med.; 97:833 .

Sosenko , J.M.; Breslow , J. L .; Miettinen O.S.and Gabbay K.H.(1980):

Hyperglycaemia and plasma lipid levels .

A prospective study of young insulin-dependent diabetic patients .

N. Engl. J. Med. 302 : 650 .

Spiro , M.J.(1984):

Effect of diabetes on the Sugar nucleotides in several tissues of the Rat.

Diabetologica , 26:70 -5 .

Stehower , C.D.A. ; Stroes , E.S.G.; Hackeng W.A.L.; Mulder ; P.G.H. and Holander , G.J.H(1991):

Von-Villbrand factor and development of diabetic nephropathy in IDDM.

Diabetes , 40:971-76 .

Stenibrecher , U.P and Witztum J.L.(1984) .

Glycosylation of LDL to an extent comparable to that seen in diabetes slows their catabolism .

Diabetes , 33 : 130 .

Stein E.A .and Pesce A.(1981):

Enzyme linked immunoassays for apolipoproteins : Advantages , problems and prototype assay . In Lippel .K.(ed.) : Proceedings of the workshop on Apolipoprotein Quantification .U.S.Department of Health and human services .

National Institutes of health publication No . 83-1266,PP 319-331 .

Stein E.A.(1987):

Lipids , Lipoproteins and Aplipoproteins. In :Tietz N..(ed .) : Fundamentals of clinical chemistry 3rd ed .WB.Saunders Co.philadelphia london .P.448 .

Steinberg K.K.; Cooper G.R.; Graiser S.R. et al (1983):

Some considerations of methodology and standardization of apolipoprotein A-I immunoassays .

Clin . Chem .; 29:415-426 .

Steiner G . and Reardon M.F. (1983) :

A new model of human VLDL metabolism based on simultaneous studies on its apo-B and triglycerides .

Metabolism , 32:342 .

Stenflo , J.(1976):

A new vitamin K-dependent protein purification from bovine plasma and preliminary characterization .

- J.Biol.Chem.* ; 251:355-363 .
- Stiller , C.R.; Dupre , J.; Gent , M.; Jenner , M.R.; Keown , P.A.; Laupacis , A.; Martell , R. ; et al (1984) :**
Effects of cyclosporine immunosuppression in insulin - dependent diabetes mellitus of recent onset .
Science . 223 : 1362-7 .
- Straight , D.L. and Mckee , P. A.(1984) :**
Characterization of thrombin binding to α_2 macroglobulin .
J.Biol.Chem. ; 259:1272 .
- Sun T.(1991):**
Lipoprotein analysis . In: Interpretation of protein and isoenzyme patterns in body fluids .IGaku -Shoin .
New York ,Tokyo , PP : 172-196 .
- Tan,M.H.; Wilmhurst , E.G.; Gleason ,R.E, and Soeldner , J.S. (1973):**
Effect of posture on serum lipids .
N.Engl J-Med. ; 289:416 .
- Tall , A.R.; and Small D.M.(1978):**
Plasma high density lipoproteins .
N.Engl. J.Med. 299:1232-1236 .
- Taskinen M.R.& Nikkila E.A. (1979):**
Lipoprotein lipase activity of adipose tissue and skeletal muscle in insulin-deficient human diabetes .Relation to high-density and very low-density lipoproteins and response to treatment .
Diabetologica 17:351-356.
- Taskinen M.R. , Nikkila E.A. & Ollus A. (1983):**
Diabetes Care 6:224 .
- Taskinen M.R., Kuusi T.; Nikkila E.A. (1985):**
Regulation of HDL- and its subfractions in chronically insulin treated patients with type I diabetes .In : Crepaldi G et al (eds)Diabetes,Obesity and hyperlipidemias .Elsevier Science publishers , Amsterdam , PP 251-259 .
- Taskinen M.R. (1987):**
Lipoprotein lipase in diabetes .
Diabetes . Metab-Rev (in press).
- Taskinen M.R. (1992) :**
Quantitative and Qualitative lipoprotein abnormalities in diabetes mellitus .

- Diabetes* 41 (suppl 2) : 12 - 17 .
- Taskinen M.R.; Kuusi T. and Nikkila E.A.(1987):**
 Serum lipoproteins and atherosclerosis in type I and type II diabetes.
- Thaler , E. ; and Lechner , K.(1981):**
 Antithrombin III deficiency and thromboembolism .
Clin . Haematol . ; 10:369-390 .
- Thompson , R.B.(1977):**
 Disorders of blood Clotting .A textbook of clinical haematology .
 Churchill Livingstone .P.734-768 .
- Thuesen , L.; Christiansen J.S.; Mogensen C.E .; et al ,(1988):**
 Cardiac hyperfunction in insulin -dependent diabetic patients developing microvascular complications .
Diabetes 10 : 369 .
- Tietz , N.W.; Aldrich , J.; Bhagavan , N.V.; Coiner , D.; Conn , R.B.; Kowalczyk , R.S.; Lehmann , H.R .; Pruden , E.L.; and Whitleg , R.J.(1987):**
 "Fundamentals of Clinical Chemistry "3rd ed . W.B. Saunders Company, Philadelphia , London , Tornto , Ch .14.PP 448 .
- Titani , K.; Hermodson , M.A.; Fujikawa , K.; Ericsson , L.H; Waish ,K.A.; Neurath , H. and Davie , E.W. (1972):**
 Bovine factor X2 (activated stuart factor) : evidence of homology with mammalian serine proteases.
Biochemistry , 11 :4899 .
- Trinder P.(1969):**
Ann . Clin .Biochem . 6,24-27 .
 Quoted from pamphlet of test combination glucose . Boehringer Mannheim G mbH Diagnostica . 1990 .
- Tripodi , A.; Franchi , F.; Krachmalincoff , A.; and Mannucci , P.M.(1990):**
 A symptomatic homozygous protein C deficiency .
Acta haematol . (Basel) , 83(3): 152-5 .
- Uterman , G . and Beispiegel , U. (1979) ;**
 Apoliporotein A.IV,a protein occuring in human mesenteric lymph chylomicrons and free in plasma , isolation and quantification.
Eur . J . Biochem . , 99:333 .

Van Hinsbergh , V.W.; Bertina , R.M.; Emeis , J.J. and Haverkate , F. (1985):

Activated protein C decreases plasminogen activator inhibitor activity in endothelial cell Conditioned medium .

Blood ,65(2):444-451 .

Vannini P., Ciavarella A., Flammini M.; et al , (1984):

Lipid abnormalities in insulin-dependent diabetic patients with albuminuria .

Diabetic Care , 7(2)151 .

Van-Wersch J.W; Donders S.H.; Wasterhuis L.W.; and Venekamp W.J. (1991) :

Microalbuminuria in diabetic patients : Relationship to lipid , glyco-metabolic , coagulation and fibrinolysis parameters .

Eur -J-Clin-Chem-Clin - Biochem . 29(3) : 493 - 8 .

Viberti , G.C.; Jarrett , R.J. and Keen H.;(1979):

Effect of control of blood glucose on urinary excretion of albumin and B2- microglobulin in insulin-dependent diabetes .

N.Engl .J. Med .; 300-461 .

Viberti , G.C.; Hill , R.D.; Jarrett , R.J. et al ,(1982) :

Microalbuminuria as a predictor of clinical nephropathy in insulin-dependent diabetes mellitus .

Lancet , 1 :1430-2 .

Viberti G.C. , Dodds R.A., Bending J.J. et al ,(1983):

Long term correction of hyperglycemia and progression renal failure in insulin - dependent diabetes .

*Br .Med .J.*286:598 .

Viberti G.C.; and Keen ,H. (1984) :

The patterns of proteinuria in diabetes mellitus : Relevance to pathogenesis and prevention of diabetic nephropathy .

Diabetes , 33:686 -92 .

Viberti G.C., Dodds R.A., Bending J.J. et al (1988):

Non glycemic intervention in diabetic nephropathy : The role of dietary protein intake .In Mogensen CE(ed): The Kidney and hypertension in Diabetes Mellitus .Boston ,Martinus Nijhoff Publishing (P 205).

Vigano - D'Angelo , S.; D'Angelo , A. and Kaufman , C.E.(1987):

Protein S deficiency occurs in nephrotic syndrom .

Ann . Intern .Med .; 107:42 .

Vigstrup J.; Mogensen C.E.; (1985):

Proliferative diabetic retinopathy : At risk patients identified by early detection of microalbuminuria .

Acta Ophthalmol .63:530 .

Vlachokosta E.V., Asmal A.C, Ganda O.P., Aoki T.T. (1983) :

The effect of strict control with the artificial B-cell on plasma lipid levels in insulin -dependent diabetes .

Diabetes Care 6:351-355 .

Vlassara , H.; Brownlee , M. and Ceramli , A. (1986) :

Nonenzymatic glycosylation . Role in the pathogenesis of diabetic complications.

Clin . Chem . 32 110 : B37 .

Vukovich T.N., Schernthaner G .; and Seebacher G. (1984) :

Factor VIII complex , protein C. and fibrinolysis in well-controlled type -I diabetics .

Eur.J. Clin - Invest .; 12:336-7 .

Waife , S.D. (1980):

Diabetes Mellitus .Idianapolis by Eli Lilly and Company .

PP 1000 .

Walden C.E., Knopp R.H., Wahl P.W., Beach K.W., Strandness E. (1984):

Sex differences in the effect of diabetes mellitus on lipoprotein triglyceride and cholesterol Concentration .

N.Engl .J. Med .311:953 - 959.

Walker , F.J.(1981):

Regulation of activated protein C by protein S . The role of phospholipid in factor Va inactivation .

J.Biol . Chem .; 256:11128 .

Walker , F.J. (1984):

Protein S and the regulation of APC .

Semin - Thromb .Haemat .;10:131-136 .

Walker , F.J.; Chavin , S.I. and Fay , P.J. (1987):

Inactivation of factor VIII by activated protein C. and protein S.

Arch . Biochem . Biophys .; 252:322 .

Walker , F.J.; Scandella , D. and Fay , P.J. (1990):

Identification of the binding site for activated protein C.on the light chain of factors V and VIII.

J.Biol .Chem .; 25, 265(3):1484-9.

Watt G.F.; Naumova R.; Slavin B.M. Morris R.W.; Houlston; Kubal C.; Shaw K.M. (1989):

Serum lipids and lipoproteins in insulin-dependent diabetic patients with persistent microalbuminuria.

Diabetic Med.; 6 (1), P : 25-30.

Weiland, D.; Monden C.E. and Reoven G.M. (1980):

Evidence for multiple causality in development of diabetic hypertriglyceridaemia.

Diabetologica, 18 : 335.

Weiman, W.; Stibbe, J.; Vonseyan, A.J. and collen, D. (1981):

Specific lysis of an iliofemoral thrombus by administration of extrinsic (Tissue type) plasminogen activator.

Lancet, ii : 1018-1020.

Weinstock N.; Bartholome M.; and Seidel D. (1981):

Determination of apolipoprotein A.I by Kinetic nephelometry.

Biochim. Biophys. Acta, 663:279-288.

Weisgraber, K.H. and Mahley, R.W. (1978):

Apolipoprotein (E-AII) Complex of plasma lipoproteins. I. Characterization of this mixed disulfide and its identification in a high density lipoprotein subfraction.

J. Biol. Chem. 253:6281.

Weisgraber, K.H.; Roll, S.C. and Mahley, R.W. (1981):

Human E apolipoprotein heterogeneity.

J. Biol. Chem.; 256:9077.

Weiss, H.J.; Hoyer, L.W.; and Rickles, F.R. (1973):

Qualitative assay of a plasma factor deficient in Von Willebrand's disease that is necessary for platelet aggregation.

J. Clin. Invest. 52,2708.

Whitby, L.G.; Smith, A.F.; and Beckett, G.J. (1988):

"Lecture Notes on clinical chemistry" 4th ed. Blackwell scientific publications. Oxford London Edinburgh Ch.12 PP 233.

White, G.C.; and Maranf, A.A. (1981):

Platelet factor 4 levels in patients with artery disease.

J. Lab. Clin. Med.

Wiggins, R.C. (1983):

Kinin release from high molecular weight kininogen by the action of Hageman factor in the absence of kallikrein.

J. Biol. Chem. 258:8963.

Williams, W.J. and Norris, D.G. (1966):

Purifications of bovine plasma protein (Factor VII) which is required for the activity of lung microsomes in / blood Coagulation.
J. Biol. Chem. ; 24:1847.

Wilson, C.; Walker, I.D. and Davidson, J.F. (1987):

Mesentric venous thrombosis and antithrombin III deficiency .
J. Clin. Pathol. ; 40:406-408 .

Wilson, P.W.; Carrison, R.J.; Castelli W.P.; Feinleib, M.; McNamara, P.M. and Kannel, W.B. (1980):

Prevalance of coronary heart disease in framingham offspring study : Role of lipoprotein cholesterols.
Am. J. cardiol., 46 : 49

Windmueller, H.G.; Herbert, P.N. and Levy, R.J. (1973):

Biosynthesis of lymph and plasma lipoprotein apoprotein by isolated perfused rat liver and intestine .
J. lipid Res. ; 14:215 .

Winocour P.D.; Lopes-Virella M.; Laimins M.; Colwell J.A.; (1983):

Time course of changes in vitro platelet function and plasma von willebrand factor activity (VIII R :vwF) and factor VIII - related anigen (VIII R : Ag) in the diabetic rat.
J. Lab. Clin. Med., 102 : 795-804.

Winocour P.H. , Durrington P.N. ; Ishola M.; Anderson D.C. (1986):

Lipoprotein abnormalities in insulin dependent diabetes mellitus .
Lancet ii 1176-1178 .

Winocour P.H.; Durrington P.N.; Ishola, M; Anderson D.C.; and Choen H. (1987):

Influence of proteinuria on vascular disease, blood pressure and lipoproteins in insulin dependent diabetes mellitus.
BMJ., 294 (27) : 1648-1651.

Wintrobe, M.M.; Lee, G.R.; Bogg, D.R.; Bithel, T.C.; Foerster ; J.; Athens J.W. and Lukens, J.N. (1981):

Clinical Haematology . 8th edition . (Eds).
Lea and Febiger, Philadelphia , P.29 .

Wiseman M.; Viberti G.; Mackintosh D. et al, (1984):

Glycemia , arterial pressure and microalbuminuria in type I (Insulin -dependent) diabetes mellitus .
Diabetologica 26 : 401 .

- Wiseman, M.J.; Saunders, A.J.; Keen, H.; et al (1985):**
Effect of blood glucose control on increased glomerular filtration rate and kidney size in insulin-dependent diabetes.
N. Engl. J. Med.; 312 : 617-21.
- Wood F.C., and Bierman E.L. (1986):**
Is diet the Cornerstone in management of diabetes ?
N. Engl. J. Med. 315 : 1224.
- World Health Organization Expert committee (1980):**
Second report on diabetes mellitus.
Technical report series No. 646. Geneva, Switzerland.
- World Health Organization study Group (1985):**
Technical report series No. 727.
Geneva, Switzerland.
- Wuepper, K.D.; and Cochrance, V.G.; (1972):**
Plasma prekallikrein isolation, characterization and mechanism of action.
J. Exp. Med., 92 : 1359.
- Yaari S.; and Goldbourt U.; (1981):**
Associations of serum HDL and total cholesterol with total, cardiovascular, and cancer mortality in a 7-year. Prospective study of 10,000 men.
Lancet, 1 : 1011.
- Zannis, V.I.; and Berslow, J.L. (1980):**
Characterization of a unique human apolipoprotein E Variant associated with type III hyperlipoproteinaemia.
J. Biol. Chem., 255 (5) : 1759.
- Zannis, V.I.; and Breslow, J.L.; (1981):**
Human very low density lipoprotein apolipoprotein E isoprotein polymorphism is explained by genetic variation and post-translational modification.
Biochemistry, 20 : 1033.
- Zhao, J.B.; Mikata, A. and Azuma (1990):**
Immunoglobulin deposits in diabetic microangiopathy. Observations in autopsy materials.
Acta. Pathol. Jpn., 40 (10) : 729-34.
- Zilva, J.F.; and Pannall, P.R. (1984):**
" Clinical Chemistry in Diagnosis and treatment : 4th ed. Lloyd-luke (Medical Books) LTD, London.
Ch. X, PP. 232.

Zubay G.; (1988):

Cholesterol and related derivations. In : biochemistry. 2nd ed.
MaGmillan publishing Company.
New York. PP. 735.