

REFERENCES

- Aagaard –Tillery K.M, and Nuthalapaty, F.S., (2005):** Impaired collagen chaperone results in preterm PROM. Am J Perinatol. 22, 287-297.
- Akercan F, Cirpan T, Kazandi M, et al., (2005):** The value of the insulin-like growth factor binding protein-1 in the cervical-vaigal secretion detected by immunochromatographic dipstick test in the prediction of delivery in women with clinically unconfirmed preterm premature rupture of membranes. Eur. J. Obstet Gynecol Reprod Biol. 1; 121 (2): 159-63.
- Al-Awaysheh F, and Al-Qa'Qa K, (2005):** Neonatal outcome and prenatal antibiotic treatment in premature pupture of membranes. Pak J Med Sci. vol. 21 No. 4.
- Alexander JM, Mercer BM, Miodovnik M, (2000):** The impact of digital cervical examination on expectantly managed preterm rupture of membranes. Am J Obstet Gynecol. 183: 1003-7.
- Alfirevic Z, (2000):** Oral misoprostol for induction of labor (Cochrane review). The Cochrane library 4, CD001338.
- Alfirevic Z. (2004):** Oral misoprostol for induction of labour (Cochrane Review): In: The Cochrane library, issue 3, chichester, UK: John Wiley & Sons, Ltd.
- Alger LS, Lovchik JC, Hebel JR, et al., (1988):** The association of Chlamydia trachmatis, Neisseria gonorrhoeae, and group B streptococci with preterm rupture of the membranes and pregnancy outcome. Am J Obstet Gynecol. 159: 397.

- Al-Hussaini TK, Abdel-Aal SA, Youssef MA. (2003):*** Oral misoprostol vs. Intravenous oxytocin for labor induction in women with prelabor rupture of membranes at term. *Int J Gynaecol Obstet.* 82: 73-5.
- American College of Obstetricians and Gynecologists (1989):*** Dystocia. Technical Bulletin No. 137.
- American College of Obstetricians and Gynecologists (1995a):*** Dystocia and the augmentation of labor. Technical Bulletin No. 218.
- American Collage of Obstetrician and Gynecologists (1998):*** Premature rupture of membranes. Clinical management guidelines for obstetrician-gynecologists. ACOG practice bulletin no. 1 *Int. J Gynecol.* 101: 178-93.
- American College of obstetricians and gynecologists (1999a):*** Induction of labor. Practice Bulletin No. 10.
- American College of obstetricians and gynecologists (1999b):*** Induction of labor with misoprostol. Committee opintion No. 228.
- American College of Obstetricians and Gynecologists (2000):*** Induction of pregnancy. ACOG committee opinion 234. Washington, DC; ACOG .
- American College of Obstetricains and Gynecologists (2003):*** Use of progesterone to reduce preterm birth. Committee Opinion No. 291.
- Anai T, Tanaka Y, Hirota Y, et al., (1997):*** Vaginal fluid hCG levels for detection of PROM. *Obstet Gynecol.* 89 (2): 261-4.
- Ananth VS, Ogelese Y, Srinivas N, (2004):*** PPRM, intrauterine infection, and oligohydramnios. *Obstet Gynecol.* 104: 71-77.
- Ansari S and Nili F, (2003):*** Neonatal complication of pre mature repture of membranes. *Acta Medic Iranica.* 41 (3): 175-179.

- Arulkumaran S, Koh Ch, Ingemarsson I, et al., (1987):** Augmentation of labour-mode of delivery related to cervimetric progress. Aust N Z J Obstet Gynaecol. 27: 304.
- Azargoon A, and Darvishzadeh S. (2006):** Association of bacterial vaginosis, trichomonas vaginalis, and vaginal acidity with outcome of pregnancy. Arch Iran Med. 9 (3): 213-7.
- Bar J, Metsger MA, Hodmoshe et al., (2000):** Effect of antibiotic therapy PPRM on neonatal mortality and morbidity. American Journal of Peinatology/ vol. 17, No. 5.
- Becker C, August Fuhr N, van Baalen A, et al., (2006):** Antibiotic therapy for preterm premature rupture of membranes –results of a multicenter study. J Perinat Med. 34 (3): 203-6.
- Biggio JR, Ramsey PS, Cliver SP, et al., (2005):** Mid trimester amniotic fluid matrix metalloproteinase -8 (MMP-8) levels above the 90th percentile are a marker for subsequent preterm rupture of membranes. Am J. Obstet Gynecol. 192-13.
- Blackwell SC, and Berry SM. (1999):** Role of amniocentesis for the diagnosis of subclinical intra-amniotic infection in PROM. Curr Opin Obstet Gynecol. 11: 541-547.
- Bolnick JM, Velazquez MD, Gonzalez JL, et al., (2004):** Randomized trial between two active labor management protocols in the presence of an unfavorable cervix. Am J Obstet Gynecol. 190: 124.
- Brian M, and Mercer D, (2004):** Preterm premature rupture of the membranes: Diagnosis and Management. Clin Perinatol. 31: 765-782.
- Brian M, and Mercer, MD (2005):** Preterm premature rupture of the membranes: Current Approaches to evaluation and Management. Obstet Gynecol Clin N Am. 32: 411-428.

- Burchard E.G, Ziv E, Coyle N, et al., (2003):** Impaired collagen chaperone results in preterm PROM. *N Engl J Med.* 348, 1170-1175.
- Burquet A, Kaminski M, Abraham –Lerat L, et al., (2004):** The complex relationship between smoking in pregnancy and very preterm delivery. Results of the Epipage study. *BJOG.* 111(3): 258-65.
- Burkman RT, Tonascia JA, Atienza MF, et al. (1976):** Untreated endocervical gonorrhea and endometritis following elective abortion. *Am J Obstet Gynecol.* 126: 648.
- Butt KD, Bennett KA, Crane JM, et al., (1999):** Randomized comparison of oral misoprostol and oxytocin for labor induction in term prelabor membrane rupture. *Obstet Gynecol.* 94: 994-9.
- Buyukbayrak EE, Turan C, Unal O, et al., (2004):** Diagnostic power of the vaginal wash fluid prolactin assay as an alternative method for the diagnosis of premature rupture of membranes. *J Matern Fetal Neonatal Med.* 15 (2): 120-5.
- Calado E and Ayres-de-Campos D. (2007):** Premature Rupture of membranes at 20 weeks: reports of a successful outcome after Transcervical Application of fibrin Glue. *Fetal Diagn Ther.* 21; 22(1): 14-17.
- Canavan PT, and Simhan NH, (2005):** Premature rupture of membranes: diagnosis, evaluation and management strategies. *BJOG.* 112: (1) pp 32-37.
- Carlan SJ, O'Brien WF, Parsons MT et al., (1993):** PPRM: a randomized study of home versus hospital management. *Obstet Gynecol.* 81: 61-4.

- Casey ML, MacDonald PC: (1997):** Lysyl oxidase expression in human amnion: Ontogeny and cellular localization. *J Clin Endocrinol Metab.* 82: 167.
- Catalano A, and Zardini E, (1994):** PROM. Spontaneous course of the event and the therapeutic approach with a human fibrin glue. *Minerva Ginecol.* 46 (12): 675-80.
- Centers for Disease Control and Prevention (2002d):** Sexually transmitted diseases treatment guideline 51: 6.
- Chane JM, Delaney T, Hutchens D. (2003):** Oral misoprostol for premature rupture of membranes at term. *Am J Obstet Gynecol.* 189: 720-4
- Chang J, Tracy TF, Carr SR, et al., (2006):** Port insertion and removal techniques to minimize premature rupture of the membranes in endoscopic fetal surgery. *J Pediatr Surg.* 41 (5): 905-9.
- Chen M, Hsieh CY, Cameron AD et al., (2005):** Management of oligohydramnios with antepartum amnioinfusion, amniopatch and cerclage. *Taiwanese J Obstet Gynecol.* 44(4): 347-352.
- Christensen KK, Ingemarsson I, Leideman T, et al., (1980):** Effect of ritodrine on labor after premature rupture of the membranes. *Obstet Gynecol.* 55: 187-190.
- Clark R, Andrew C, Michelle McCune, RN, et al., (2004):** Aggressive tocolysis does not prolong pregnancy or reduce neonatal morbidity after preterm premature rupture of the membranes. *Am. J Obstet Gynecol.* 190: 1723-31.
- Clausen HV, Jorgensen JC, Ottesen B. (1999):** Stem villous arteries from the placentas of heavy smokers: functional and mechanical properties. *Am J Obstet Gynecol.* 180 476-82.

- Cnattinguis S, Granath F Kypklund-blomberg (2005):** Maternal smoking and causes of very preterm birth. *Acta Obstet Gynecol Scand.* 84: 572-577.
- Combs CA, McCune M, Clark R, et al., (2004):** Aggressive tocolysis does not prolong pregnancy or reduce neonatal morbidity after PPRM. *Am. J Obstet Gynecol.* 190: 1728-31.
- Conner EM, Sperling RS, Gelber R, et al., (1994):** Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. *N Eng J Med;* 331: 1173- 80.
- Contino B, Armellino F, Brokaj L, (2004):** Amniopatch, a repairing technique for premature rupture of amniotic membranes in second trimester. *Acta Biomed Ateneo Parmense.* 75 Suppl. 1: 27-30.
- Copper RL, Goldenberg RL, Mercer BM, (1996):** The preterm prediction study: fetal fibronectin testing and spontaneous preterm birth. NICHD maternal fetal medicine units network. *Obstet Gynecol.* 87: 643-8.
- Cotch MF, Pastorek JG II, Nugent RP, et al., (1997):** *Trichomonas vaginalis* associated with low birth weight and preterm delivery. *Sex Transm Dis.* 24: 353.
- Cousins ML, Smok PD, Lovett MS, (2005):** Amnisure placental alpha microglobulin-1 rapid immunoassay versus standard diagnostic methods for detection of rupture of membranes. *Am J Perinatology.* 22: 0735-1631.
- Cox SM, and Alexander JM, (1996):** Clinical course of premature rupture of the membranes. *Semin Perinatol.* 20: 369-374.

- Crane JMG and Young DC (2003):** Induction of labor with a favourable cervix and/or prelabour PROM. *Best prac Res Clin Obstet Gynecol.* 17: 795-809.
- Craven CM, Zhao L., Ward K. (2000):** Lateral placental growth occurs by trophoblast cell invasion of decidual veins. *Placenta* 21: 160.
- Dare MR, Middleton P, Crowther CA, et al., (2006):** Planned early birth versus expectant management (waiting) for prelabour rupture of membranes at term (37 weeks or more). *Cochrane Database Syst Rev.* 25; (1): CD005302.
- Darj E, and Lyrenas S, (1999):** Insulin-like growth factor binding protein-1, a quick way to detect amniotic fluid. *Acta Obstet Gynecol Scand.* 78 (6): 557-8.
- De Hann HH, Offermans PM, Smits F, et al., (1994):** Value of the fern test to confirm or reject the diagnosis of ruptured membranes in modest in nonlaboring women presenting with nonspecific vaginal fluid loss. *Am J Perinatol.* 11: 46-50.
- De Meeus JB, Sima Ole B, Bascou V et al., (1997):** Biological diagnosis of premature rupture of membranes: respective values of diamine oxidase activity compared to vaginal fluid pH (Amnicator) *J Gynecol obstet boil Reprod (paris).* 26 (7): 730-3.
- Dibble LA and Elliott JP (1992):** Possible amniotic fluid embolism associated with amnioinfusion. *J Matern Fetal Med.* 1: 263-6.
- Dinsmoor MJ, Bachman R, Haney EI et al., (2003):** Outcome after expectant management of extremely preterm premature rupture of the membranes. *Am J obstet Gynecol.* 190: 183-7.

- Dragich DA, Ross AF, Chestnut DH, et al., (1991):*** Respiratory failure associated with amnioinfusion during labor. *Anesth Analg.* 72: 549-51.
- Du Vigneaud V, Ressler C Swan JM et al., (1955):*** The synthesis of oxytocin. *J Am Chem Soc.* 75: 4879.
- Dunlop PDM, Crowley PA, Lamont RF, et al., (1986):*** Preterm ruptured membranes, no contractions. *J Obstet Gynecol.* 7: 92-6.
- Dye T, Aubry R, Gross et al., (1994):*** Amnioinfusion and the intrauterine prevention of meconium aspiration. *Am J Obstet Gynecol.* 171: 1601-5.
- Edmonds K (2007):*** Dewhurst's textbook of obstetric gynaecology-7th ed. Oxford: Black well science. P. 210.
- Egarter C, Leitich H, Karas H, et al., (1996):*** Antibiotic treatment in preterm premature rupture of the membranes. *Lancet.* 174: 589-597.
- Ekwo EE, Gosselink CA, Woolson R, (1993):*** Risks for premature rupture of amniotic membranes. *Int J Epidemiol.* 22: 495-503.
- Esim E, Turan C, Unal O, et al., (2003):*** Diagnosis of PROM by identification of beta-HCG in vaginal washing fluid. *Eur J Obstet Gynecol Repord Biol.* 107 (1): 37-40.
- Feng LH, Yu LZ, Ni Cy, Jia WX, (2003):*** Practicability of using vaginal fluid markers in detecting PROM. *Ann Clin Biochem.* 40 (5): 542-5.
- Ferguson SE, Smith GN, Salenieks M, et al., (2002):*** Preterm premature rupture of membranes: nutritional and socioeconomic factors. *Obstet Gynecol.* 100: 1250-6.
- Flannelly GM Turner MJ, Rassmussen MJ, et al., (1993):*** Rupture of the uterus in Dublin: An update. *J Obstet Gynecol.* 13: 440.

- Fontenot T, and Lewis DF, (2001):** Tocolytic therapy with preterm premature rupture of membranes. Clin perinatol. 28: 787-96.
- Fortunato SJ and Menon R. Distinct (2001):** Distinct molecular events suggest different pathways for preterm labor and premature rupture of membranes. Am J obstet Gynecol 184: 1399-405.
- French JI and McGregor JA. (1996):** The pathobiology of PROM. Semin perinatol. 20: 344-68.
- French JI. And Mc Gregor JA, (1997):** Evidence based prevention of preterm birth and rupture of membranes: infection and inflammation. J Obstet Gynecol Can. 19: 835-52.
- Frieden E. (1980):** Caeruloplasmin: a multi-functional metalloprotein of vertebrate plasma. In: Biological roles of copper. Chiba foundation symposium. New York: Elsevier/ North-Holland. 93-124.
- Frigoletto FD, Lieberman E, Lang JM, et al., (1995):** A clinical trial of active management of labor. N Engl J Med. 333: 745.
- Furman B, Shoham –Vardi I, Basiri A, et al., (2000):** Clinical significance and outcome of preterm prelabor rupture of membranes: population–based study. Eur J Obstet Gynecol Reprod Biol. 92: 209-216.
- Furman B, Shoham-Vardi I, Bashiri A, et al., (2001):** Preterm premature rupture of membranes is not an independent risk factor for neonatal morbidity. J. Matern Fetal med. 10 (2): 107-111.
- Gabbe SG, Ettinger BB, Freeman KR, et al., (1976):** Umbilical cord compression associated with amniotomy: laboratory observations. Am J. Obstet Gynecol. 126: 353-5.

- Gardman MA, Low DE, Saginur R, (1998):** Group B streptococcal necrotizing fascitis and streptococcal. Arch Intern Med. 158: 1704-8.
- Garzetti GG, Ciavattini A, De Cristofaro F, et al., (1997):** Prophylatic transabdominal amnioinfusion in oligohydramnios for PPRM: increase of amniotic fluid index during latency period. Gynecol Obstet Invest. 44 (4): 249-54.
- Gaucheranad P, Guibaud S, Awada A, et al., (1995):** Comparative study of three amniotic fluid markers in premature rupture of membranes: fetal fibronectin, alpha-fetaoprotein, diamino-oxydase.
- Gibble JW and Ness PM. (1990):** Fibrin glue: the prefect operative sealant? Transfusioin. Am J Obstet Gynecol. 30: 741-7.
- Gibbs R, Romero R, Hillier S, et al., (1992):** A review of premature birth and subclinical infection. Am J Obstet Gynecol. 166: 1515.
- Glantz JC and Letteney DL. (1996):** Pumps and warmers during amnioinfusion: is either necessary Obstet Gynecol. 87: 150-4.
- Goldberg AB, Greenberg MB & Darney PD (2001):** Misoprostol and pregnancy. N Engl J Med. 344: 38-47.
- Goldman-Wohl DS, Ariel I, Greenfield C, et al., (2000):** Antigen-G expression in extravillous trophoblasts is associated with peeclampsia. Mol Hum Reprod. 6: 88.
- Gomez R, Romero R, Nien JK et al., (2005):** Idiopathic vaginal bleeding during pregnancy as the only clinical manifestation of intrauterine infection. J Matern fetal Neonatal Med. 18(1): 31-7.

- Gopalani S, Krohn M, Meyn L, (2005):** Contemporary management of preterm premature rupture of membranes: determinants of latency and neonatal outcome. *Obstet Gynecol.* 60: 16-7.
- Gramellini D, Fieni S, Kaihura C et al., (2003):** Antepartum amnioinfusion: a review. *The Journal of Maternal-Fetal and Neonatal Medicine.* 14: 291-296.
- Gramellini D, Fieni S, Kaihura C et al., (2003):** Transabdominal antepartum amnioinfusion. *International J Gynecol Obstet.* 83 (2): 171-178.
- Gyr TN, Malek A, Mathez L, et al., (1994):** Premeation of human chorioamniotic membranes by escherichia coli in vitro. *Am J Obstet Gynecol.* 170: 223-7.
- Hill A, and Medina MT (2006):** Preterm premature rupture of membranes: diagnosis and management. vol. 73 No. 4.
- Hadley CB, Main DM, Gabbe SG. (1990):** Risk factors for preterm premature rupture of the fetal membranes. *Am J Perinatol.* 7: 374-9.
- Han-Kim Y, Wonpark Y, Sungkwon H, (2005):** Vaginal fluid β -human chorionic gonadotropin level in the diagnosis of premature rupture of membranes. Vol. 84 Issue 8. p 802.
- Hannah ME, Ohlsson A, Farine D et al., (1996):** Induction of labor compared with expectant management for PROM at term. *N Engl J Med.* 334: 1005-10.
- Hant MD, Mercer BM, Thurnau G, et al., (2005):** Perinatal outcomes in women with PROM between 24 and 32 weeks of gestation and a history of vaginal bleeding. *Am J Obstet Gynecol.* 193 (1): 164-8.
- Harger JH. (2002):** Cerclage and cervical insufficiency: an evidence-based analysis. *Obstet Gynecol.* 100: 1313-27.

- Harmanlio-Wapner R and Lontz J. (1998):** Human egaculate: Effect on the biochemical properties of the human chorioamniotic membranes. J Reprod Med. 43: 779.
- Hellberg D, Nilsson S, Mardh PA. (2000):** Bacterial vaginosis and smoking. Int J STD AIDS. 11: 603-6.
- Hiyamuta S, Shimizu K, Aoki T. (1993):** Early Diagnosis of Wilson's disease. Lancet. 342: 56-57.
- Hoffman Ra, Anthony J, Fawcus JA, (2001):** Oral misoprostol vs. placebo in the management of prelabor rupture of membranes at term. Int J Obstet Gynaecol. 72: 215-21.
- Hofmeyr GJ, Gulmezoglu AM, Nikodem VC et al., (1996):** Amnioinfusion. Eur J Obstet Gynecol. Reprod Biol. 64: 159-65.
- Hofmeyr GJ and Gulmezoglu AM (2002):** Vaginal misoprostol for cervical ripening and induction of labour (Cochrane review). The Cochrane library 1, CD000941.
- Hofmeyr GJ and Gulmezoglu AM (2003):** Vaginal misoprostol for cervical ripening and induction of labor. Cochrane Database Syst Rev. (3): CD 000941.
- Hofmeyr GJ, and Gulmezoglu AM, (2004):** Vaginal misoprostol for cervical ripening and induction of labor (Cochrane Review). In: The Cochrane library, Issue 3. CD000134.
- Hofmeyr GJ (2007):** Amnioinfusion for PROM cochrane data base of systemic reviews Issue 2. CD000150.
- How HY, Cook CR, Cook VD, Miles DE, Spinnato JA. (1998):** Preterm premature rupture of membranes: aggressive tocolysis versus expectant management. J Matern Fetal Med. 7: 8-12.

- Iams JD, Goldenberg RL, Meis PG, et al., (1996):** Transvaginal cervical sonography and the risk of spontaneous preterm. *N Engl J Med.* 334: 567-72.
- Ioannidis J.P.A., Ntzani E.E. Trikalino T, (2004):** Impaired collagen chaperone results in preterm PROM. *Nat Genet.* 36: 1312-1318.
- Jackson GM, Edwin SS, Varner MW, et al., (1996):** Regulation of fetal fibronectin production in human amnion cells. *J Soc Gynecol Investig.* 3: 85-88.
- Jacobson GF, Autry AM, Kirby RS, et al., (2001):** A randomized controlled trial comparing amoxicillin and azithromycin for the treatment of *Chlamydia trachomatis* in pregnancy. *Am J Obstet Gynecol.* 184: 1352.
- Janek L, Krizko M, Ferianec V, et al., (2002):** Premature fetal membrane rupture after amniocentesis in the second trimester and successful use of fibrin glue case report. *Ceska Gynekol.* 67 (2): 71-4.
- Jazayeri A, Jazayeri MK, Sutkin G. (2003):** Tocolysis does not improve neonatal outcome in patients with PROM. *Am J Perinatol.* 20: 189-93.
- Jazayeri A, Jazayeri MK, Sutkin G, (2006):** Tocolysis does not improve neonatal outcome in patients with preterm rupture of membranes. *Am J Perinatol.* 20 (4): 189-93.
- Johnson JWC, Daikoku NH, Niebyl RJ, et al., (1981):** PROM and prolonged latency. *Obstet Gynecol.* 57: 547-56.
- Kacmar J, Cheh E, Montagno A, et al., (2001):** A randomized trial of azithromycin versus amoxicillin for the treatment of *Chlamydia trachomatis* in pregnancy. *Infect Dis Obstet Gynecol.* 9: 197.

- Kafali H, and Oksuzler C. (2006):** Vaginal fluid urea and creatinine in diagnosis of premature rupture of membrane. Arch Gynecol Obstet. 12.
- Karger AG, and Basel S, (2007):** Comparative Study of three amniotic fluid markers in PPRM: prolactin, Beta Subunit of human Chorionic Gonadotropin, and Alpha-fetoprotein. Gynecol Obstet Invest. 63 (4): 195-199.
- Katz VL and Bowes WA: (1992):** Meconium aspiration syndrome; reflections on a murky subject. Am J Obstet Gynecol. 166: 171-83.
- Kerise MJ. (1993):** Prostaglandins in preinduction cervical ripening: meta-analysis of worldwide clinical experience. J Reprod Med. 38 suppl: 89-100.
- Khadija A.H. AL Saad A.H., Afifi NM et al., (2004):** Obstet and Gynecol Department, human medical corporation. Biomedical science program Qatar University. 13: 1.
- Killbride HW, and Thibeault DW, (2001):** Neonatal complications of preterm premature rupture of membranes, pathophysiology and management. Clinics in Perinatology. 28: 761-785.
- Killbride HW (2001):** Neonatal complications of preterm premature rupture of membranes, pathophysiology and Management. Clin perinatol. 29:800-805.
- Kilpatrick SJ, Patil R, Connell J, et al., (2006):** Risk factors for previable premature rupture of membranes or advanced cervical dilation: a case control study. Am J Obstet Gynecol. 194 (4): 1168-74.
- Krauss C, (2004):** Ehlers-Danlos Syndrome. American collage of medical Genetics. 7: 1-3.

- Lamont RF (2003):** Recent evidence associated with the condition of preterm prelabour rupture of the membranes. *Curr Opin Obstet Gynecol.* 15: 91-9.
- Laudy JA, Tribboel D, Robben SG, et al., (2002):** Prenatal prediction of pulmonary hypoplasia: clinical, biometric and Doppler velocity correlates. *Pediatrics.* 109: 250-8.
- Lee T, and Silver H. (2001):** Etiology and epidemiology of preterm premature rupture of membranes. *Clinics in perinatology.* 28: 721-734.
- Lee MJ, Davies J, Guinn D, et al., (2004):** Single versus weekly courses of antenatal corticosteroids in preterm premature rupture of membranes. *Obstet Gynecol.* 103 (2): 274-81.
- Leeson S.C., Maresh, M.J.A. Martindale E.A. (1996):** Detection of fetal fibronectin as a predictor of preterm delivery in high risk asymptomatic pregnancies. *British Journal of Obstetrics and Gynaecology.* Vol. 103, pp. 48-53.
- Levy D, and Warsof SL. (1985):** oral ritodrine and preterm premature rupture of membranes. *Obstet Gynecol.* 66: 621-623.
- Lewis, FD. (2003):** Antibiotic therapy in preterm rupture of membranes. Are seven days necessary? A preliminary randomized clinical trial. *Am J Obstet Gynecol.* 188 (6): 1413-6.
- Li HY, and Change TS, (2000):** Vaginal fluid creatinine, human chorionic gonadotropin and alpha-fetoprotein levels for detecting premature rupture of membranes. *Zhonghua Yi Xue Zhi (Taipei)* 63 (90): 686-90.
- Lieman JM, Brumfield CG, Carlo W, et al., (2005):** Preterm premature rupture of membranes: is there an optimal gestational age for delivery? *Obstet Gynecol.* 105: 12-7.

- Lo JY, Alexander JM, McIntire DD, et al., (2003):** Misoprostol and oxytocin for induction of cervical ripening and labor in patients with term pregnancy and premature membrane rupture (in Spanish). *Ginecol Obstet Mex.* 70: 469-76.
- Locatelli A, Vergani P, Di Pirro G, et al., (2000):** Role of amnioinfusion in the management of PPRM at < 26 weeks' gestation. *Am J Obstet Gynecol.* 183 (4): 878-82.
- Loderio JG, Hsieh KA, Byers et al., (1989):** The fingerprint, a false-positive fern test. *Obstet Gynecol.* 73: 873-874.
- Lopez-Zeno JA, Peaceman AM, Adashek JA, et al., (1992):** A controlled trial of a program for the active management of labor. *N Engl J Med.* 326: 450.
- Marcher S, Sipurzynski-Budrass S, Haeusler M. et al., (2006):** Successful treatment of PROM after genetic amniocentesis by intra-amniotic injection of platelets and cryoprecipitate (amniopatch): a case report. *Vox Sang.* 91 (1): 88-90.
- MacKenzie IZ and Bruns E (1997):** Randomized trial of one versus two doses of prostaglandin E2 for induction of labor. 1 clinical outcome. *BJOG* 104, 1062-7.
- Macones GA, Parry S, Elkousy M, et al., (2004):** A polymorphism in the promoter region of TNF and bacterial vaginosis: preliminary evidence of gene-environment interaction in the etiology of spontaneous preterm birth. *Am J Obstet Gynecol.* 190: 1504-8.
- Martas H. (1985):** Fibrin sea of the art. *J Oral Maxillofac. Surg.* 13: 605-11.

- Matonhodze BB, Katsoulis LC and Hofmeyr GJ (2002):** Labor induction and meconium. in vitro effects of oxytocin, dinoprostone and misoprostol on rat ileum relative to myometrium. J perinatal Med. 30: 405-10.
- Maxwell GL, and Watson WJ. (1992):** Preterm premature rupture of membranes: Results of expectant management in patients with cervical cultures positive for group B streptococcus or Neisseria gonorrhoeae. Am J. Obstet Gynecol. 166: 9.
- Maxwell G.L. (1993):** PPRM. Obstet Gynecol Surv. 48:576-83.
- McDonald SD, Perkins SL, Jodouin CA, et al., (2002):** Folate levels in pregnant women who smoke: an important gene/environment interaction. Am. J. Obstet Gynecol. 187: 620-5.
- McParland PC, Taylor DJ, Bell SC. (2000):** Myofibroblast differentiation in the connective tissue of the amnion and chorion of term human fetal membranes: Placenta. 21: 44.
- Menon R, McIntyre JO, Matrisian LM, (2006):** Salivary proteinase activity: a potential biomarker for preterm premature rupture of the membranes. Am J Obstet Gynecol. 194 (6): 1609-15.
- Mercer BM, Goldenberg RL, Meis PJ et al., (2000):** The preterm prediction study: prediction of preterm premature rupture of membranes through clinical findings and ancillary testing. Am J Obstet Gynecol. 183: 738-45.
- Mercer BM. (2003):** Preterm premature rupture of the membranes. Obstet Gynecol. 101: 178-93.
- Mercer BM, (2005):** Preterm premature rupture of the membranes: current approaches to evaluation and management. Obstet Gynecol Clin North Am. 32 (3): 411-28.

- Merrill DC, and Zlatnik FJ, (1999):** Randomized, double-masked comparison of oxytocin dosage in induction and augmentation of labor. *Obstet Gynecol.* 94: 455.
- Miller JM, Maupin RT, Mestad RE, et al., (2003):** Initial and repeated screening for gonorrhea during pregnancy. *Sex Transm Dis.* 30: 728.
- Milluzzi C, Mercer B, Collin M, (2005):** Periviable birth at 20 to 26 weeks of gestation: proximate causes, previous obstetric history and recurrence risk. *Am J Obstet Gynecol.* 193 (3 Pt 2): 1175-80.
- Milnerowicz H, Zalewski J, Geneja R, et al., (2000):** Effects of exposure to tobacco smoke in pregnancies complicated by oligohydramnios and PROM. I. Concentration of Cd and Pb in blood and Zn, Cu, Cd and Pb in amniotic fluid. *Int J Occup Med Environ Health.* 13 (3): 185-93.
- Moawad AH. Mercer BM, Meis PJ et al., (2000):** The preterm prediction study: prediction of PPROM through clinical findings and ancillary testing. *Am J. Obstet Gynecol.* 183: 738-45.
- Modena AB, Kaihura C, Fieni S (2004):** Prelabour rupture of the membranes: recent evidence. *Acta biomed ateneo parmense.* 75 (suppl 1): 5-10.
- Morales WJ, and Talley T. (1993):** PROM at < 25 weeks: a management dilemma. *Am J Obstet Gynecol.* 168: 503-7.
- Morantz AC, (2004):** Practice Guidelines ACOG release guidelines for prophylactic Antibiotic Use in Labor and Delivery. *Practice Guidelines- February 1. American Family Physician.* Vol. 69. No. 3.
- Morgan Ortiz F, Baez Barraza J, Quevedo Castro E, et al., (2002):** Misoprostol and oxytocin for induction of cervical ripening

and labor in patients with term pregnancy and PROM in Spanish. *Ginecol Obstet Mex.* 70: 469-76.

Morris JM, Roberts, CL, Crowther CA et al., (2006): Protocol for the immediate delivery versus expectant care of women with preterm prelabor rupture of the membranes close to term *BMC pregnancy child birth.* 23: 6-9.

Mozurkewich E, Horrocks J, Daley S, et al., (2002): The Miso PROM study: a multicenter randomized of oral misoprostol and oxytocin for premature rupture of membranes at term. *Am J Obstet Gynecol.* 189: 1026-30.

Naef RW III, Allbert JR, Ross EL, et al., (1998): Premature rupture of membranes at 34 to 37 weeks' gestation: aggressive versus conservative management. *Am J Obstet Gynecol.* 178(1 pt1): 126-30.

Naidu KA, (2003): Vitamin C in human health and disease is still a mystery? An overview, *Nutr J.* 2: 7-16.

National Institutes of Health NIH (1994): Effect of corticosteroids for fetal maturation on perinatal outcome. consensus statement. 12: 1-24.

National Institutes of Child Health and Human Development Trial NIH (1997): Consensus Development statement: Effect of corticosteroids for fetal maturation on perinatal outcomes, February 28- March 2. *Am J Obstet Gynecol.* 173: 246-52.

Naylor CS, Gregory K, Hobel C, (2001): Premature rupture of the membranes: an evidence –based approach to clinical care. *Am. J. Perinatol.* 18: 397-413.

Negeotte MP, Bertucci L, Towers CV, et al., (1991): Prophylactic amnioinfusion in pregnancies complicated by

- oligohydramnios: a prospective study. *Obstet Gynecol.* 77: 677-80.
- Ngai SW, To WK, Lao T, (1996):** Cervical priming with oral misoprostol in pre-labor rupture of membranes at term. *Obstet Gynecol.* 87: 923-6.
- Ngai SW, Chan YM, Lam SW, et al., (2000):** Labor characteristics and uterine activity: misoprostol compared with oxytocin in women at term with prelabour rupture of the membranes. *BJOG.* 107: 222-7.
- Ni CY, Jia WX, Yi WM, et al., (2003):** Practicability of using vaginal fluid markers in detecting premature rupture of membranes. *Ann Clin Biochem.* 40 (pt 5): 542-5.
- Nicaise C, (2003):** Neonatal complications of preterm premature rupture of membranes. Pathophysiology and management. *Clin perinatol.* 28: 761-85.
- Nimrod C, Davies D, Iwanick S, et al., (1986):** Ultrasound prediction of pulmonary hypoplasia. *Obstet Gynecol* 68: 495.
- Nourse CB, and Steer PA. (1997):** Prenatal outcome following conservative management of mid-trimester pre-labor rupture of the membranes. *J Paediatr Child Health.* 33: 125-130.
- Ogino M, Hiyamuta S, Takatsuji-Okawa M, et al., (2005):** Establishment of a prediction method for premature rupture of membranes in term pregnancy using active ceruloplasmin in cervicovaginal section as a clinical marker. *J Gynaecol Obstet.* Issue (5): 421-6.
- Ogunyemi D, and Thompson W. (2002):** A case controlled study of serial transabdominal amniotomies in the management of second trimester oligohydramnios due to PROM. *Eur. J*

- Obstet. Gynecol and Reproductive Biology. 102 (2): 167-172.
- Ozden S, Delikara MN, Avci A, et al., (2002):** Intravaginal misoprostol vs. Expectant management in premature rupture with low bishop scores at term. Int J Gynaecol Obstet. 77: 109-15.
- Parry, S and Strauss III. FJ (1998):** "Premature rupture of the fetal membranes: mechanisms of disease" the new England journal of medicine 338, no. 10 (5 Mar): 663.
- Rabello YA, Mercer BM, Thurnau GR, et al., (2006):** The NICHD-MFMU antibiotic treatment of preterm PROM study: impact of initial amniotic fluid volume on pregnancy outcome. Am J Obstet Gynecol. 194 (2): 438-45.
- Rama Sastry BV, Hemontolor ME, Olenick M. (1999):** Prostaglandin E2 in human placenta: its vascular effects and activation of prostaglandin E2 formation by nicotine and cotinine. Pharmacology. 58: 70-86.
- Ramsey PS, Ogburn PL, Harris DY et al., (2002):** Effect of vaginal PH on efficacy of the dinoprostone gel for cervical ripening/labor induction: Am J Obstet Gynecol. 187: 843.
- Ramsey PS, Lieman JM, Brumfield CG, (2005):** Chorioamnionitis increases neonatal morbidity in pregnancies complicated by preterm rupture on membranes. Am J Obstet Gynecol. 192 (4): 1162-6.
- Ramus R, Sheffield JS, Mayfield JA, et al., (2001):** A randomized trial that compared oral cefixime and intramuscular ceftriaxone for the treatment of gonorrhea in pregnancy. Am J Obstet Gynecol. 185: 629.
- Randelovic G, Kocic B, Miljkovic –Selimovic B, (2006):** High –density cervical ureaplasma urealyticum colonization in pregnant

women as a risk factor for premature rupture of membranes.
Vojnosanit Pregl. 63 (8): 737 -41.

Read SJ, and Klebanoff MA, (1993): Sexual intercourse during pregnancy and preterm delivery: Effects of vaginal microorganisms. Am J Obstet Gynecol. 168: 514-519.

Reiner, A.P. Ziv E, Lind D.L., et al., (2005): Impaired collagen chaperone results in preterm PROM. Am J Hum Genet. 76, 463-477.

Riggs MA, and Klebanoff MA, (2004): Treatment of vaginal infections to prevent preterm birth: a meta-analysis. Clin Obstet Gynecol. 47: 796-807.

Rizzo G, Capponi A, Angelini E et al., (2000): Blood flow velocity waveforms from fetal peripheral pulmonary arteries in pregnancies with PPRM: relationship with pulmonary hypoplasia. Ultrasound Obstet Gynecol. 15: 98-103.

Robert L, Goldenberg, J, Flatow C, (2003): Infection as a cause of preterm birth Clin Perinatol. 30: 677-700.

Rosemond RL, Lombardi SJ, Boehm FH (1990): Ferning of amniotic fluid contaminated with blood. Obstet Gynecol. 75: 338-340.

Rouse DJ, Owen J, Hauth JC, (1999): Active-phase labor arrest: Oxytocin augmentation for at least 4 hours. Obstet Gynecol. 93: 323.

Rouse DJ, Owen J, Savage KG, et al., (2001): Active phase labor arrest: Revisiting the 2 –hour minimum. Obstet Gynecol. 98: 550.

Royal College of Obstetricians and Gynaecologists (2002): Tocolytic drugs for women in preterm Labour. Green-top Guideline. No. 36 London: RCOG.

- Royal College of Obstetricians and Gynaecologists (2003a):** Induction of labor. Evidence based clinical guideline (no. 9) London. RCOG. Press.
- Royal College of Obstetricians and Gynaecologists (2003b):** Prevention of Early Onset Neonatal Group B Streptococcal Disease. Green-top Guideline No. 36 London: RCOG.
- Royal College of Obstetricians and Gynaecologists (2004):** Antenatal Corticosteroids to prevent respiratory distress syndrome Green-top Guideline No. 7. London: RCOG.
- Salkeld G, Buchanan LS, Crowther AC, et al., (2006):** Protocol for the immediate delivery versus expectant care of women with preterm prelabor rupture of the membranes close to term trial. BMC pregnancy child birth. 6:9.
- Salle B, Gaucherand P, Sergeant P, et al., (1997):** Comparative study of three vaginal markers of the premature rupture of membranes. Insulin like growth factor binding protein, 1 diamine-oxidase, and PH. Acta Obstet Gynecol Scand. 76: 536-540.
- Sanchez Ramos L, Chen AH, Kaunitz AM, et al., (1997):** Labor induction with intravaginal misoprostol in term premature rupture of membranes: a randomized study. Obstet Gynecol. 89: 909-12.
- Sang QA (1995):** Specific proteolysis of ceruloplasmin by leukocyte elastase. Biochem Mol Biol Int. 37: 573-581.
- Satin AJ, Leveno KJ, Sherman ML, et al., (1994):** High-dose oxytocin: 20-versus 40-minute dosage interval. Obstet Gynecol. 83: 234.

- Sciscione AG, Manley JS, Pollock M, et al., (2001):** Intracervical fibrin sealants: a potential treatment for early PPRM. Am J Obstet Gynecol. 184 (3): 368-73.
- Segel SY, Miles AM, Clothier B, et al., (2003):** Duration of antibiotic therapy after preterm premature rupture of fetal membranes. Am J Obstet Gynecol. 189-799-802.
- Sherman Michael P, (2003):** Maternal Chorioamnionitis. January 31, (4): 1-24.
- Shetly A, Stewart K, Stewart G, (2002):** Active management of term prelabour rupture of membranes rupture of membranes with oral misoprostol. BJOG. 109: 1354-9.
- Siege-Riz AM, Promislow JH, Savitz DA, et al., (2003):** Vitamin C intake and the risk of preterm delivery. Am J Obstet Gynecol. 189: 519-25.
- Silverman NS, Sullivan M, Hochman M, et al., (1994):** A randomized prospective trial comparing amoxicillin and erythromycin for the treatment of Chlamydia trachomatis in pregnancy. Am J Obstet Gynecol. 170: 829.
- Simhan HN, Caritis SN, Krohn MA, et al., (2005):** The vaginal inflammatory milieu and the risk of early premature preterm rupture of membranes. Am J Obstet Gynecol. 192: 213-8.
- Smith M.W., Patterson N, Lautenberger J.A. et al., (2004):** Impaired collagen chaperone results in preterm PROM. Am J Hum Genet. 74, 1001-1013.
- Soziol J, and Ness RB, (1998):** Chlamydial lower genital tract infection and spontaneous abortion. Infect Dis Obstet Gynecol. 6: 8.
- Strauss III, FJ and Parry S, (2006):** Premature rupture of the fetal membranes. Mechanism of Disease. N Eng J Med. Vol. 338. No. 10.

- Stringer M, Miesnik SR, Brown L, et al., (2004):*** Nursing care of the patient with preterm premature rupture of membranes. MCN. Am. J. Matern Child Nurs. May-Jun. 29 (3): 144-50.
- Strong TH, Hetzler G, Sarno AP, et al., (1990):*** Prophylactic intrapartum amnioinfusion: a randomized clinical trial. Am J Obstet Gynecol. 162: 1370-5.
- Stuart GS, and Castano PM, (2003):*** Sexually transmitted infections and contraceptives: Selective issues. Obstet Gynecol Clin North Am. 30: 795.
- Stuart EL, Evans GS, Lin YS, et al., (2005):*** Reduced collagen and ascorbic acid concentrations and increased proteolytic susceptibility with prelabor fetal membrane rupture in women. Biol Reprod. 72: 230-5.
- Svigos JM, Robinson JS, Vigne S (1999):*** Prelabor rupture of the membranes. In: James DK, Steer PJ, Weiner CP, Gonik B (eds). High risk pregnancy management options. 2nd edition. W.B. Saunders. Lond. 1015-1025.
- Tan LK, Kumar S, Jolly M et al., (2003):*** Test amnioinfusion to determine suitability for serial therapeutic amnioinfusion in midtrimester PROM. Fetal diagnosis and Therapy. 18 (3): 183-189.
- Taylor H.K, Robert D.S Jeremy F.T, (2006):*** Impaired collagen chaperone results in preterm PROM. Proceedings of the National Academy of Sciences of the United States of America (PNAS). Vol. 103. No. 36: 13267-13268.
- Tejero E, Perichart O, Pfeffer F, et al (2003):*** Collagen synthesis during pregnancy, vitamin C availability and risk of premature rupture of fetal membranes. Int J Gynecol Obstet. 81: 29-34.

- Tina Mason C, Loeb J. Gaither, K et al., (2006):** Outcome in gestations between 20 and 25 weeks with preterm premature rupture of membranes Southern Medical Journal –Abstract. Vol. 99 (7): 709-712.
- Tranquilli AL, Giannubilo SR, Bezzeccheri V et al., (2005):** Transabdominal amnioinfusion in PPRM: a randomized controlled trial. BJOG: an international journal of Obstetrics and Gynaecology. 112: 759: 763.
- Turhan NO, and Atacan N. (2002):** Antepartum prophylactic transabdominal amnioinfusion in preterm pregnancies complicated by oligohydramnios. Int J Obstet Gynecol. 76 (1): 15-21.
- Turner G, Robichaux A, Droste S, (2003):** The MisoPROM study: a multicenter randomized comparison of oral misoprostol and oxytocin for premature rupture of membranes at term. Am J Obstet Gynecol. 189: 1026-30.
- Ugwumadu A, Manyonda I, Reid F, et al., (2003):** Effect of early oral clindamycin on late miscarriage and preterm delivery in asymptomatic women with abnormal vaginal flora and bacterial vaginosis: A randomized controlled trial. Lancet. 361: 983.
- Ugwumadu A, Reid F, Hay P, et al., (2004):** Natural history of bacterial vaginosis and intermediate flora in pregnancy and effect of oral clindamycin. Obstet gynecol. 104: 114.
- Vergani P, Locatelli A, Verderio M et al., (2004):** PROM at < 26 weeks' gestation: role of amnioinfusion in the management of oligohydramnios. Acta Bio-Medica de I Ateneo Parmense. 75 Suppl. 1: 62-66.

- Vidaeff AC, Doyle NM, Gilstrap LC III. (2003):** Antenatal corticosteroids for fetal maturation in women at risk for preterm delivery. Clin Perinatol. 30: 825-40. vii.
- Weiner CP, Penk K, Klugman M. (1988):** The therapeutic efficacy and cost-effectiveness of aggressive tocolysis for premature labor associated with PROM. Am J Obstet Gynecol. 159: 216-22.
- Weismiller GD and David G. (1998):** Transcervical amnioinfusion. American Family Physician. Vol. 57. No. 3.
- Wen T, Beceir A, Xenakis E et al., (2001):** Is there a maximum effective dose of pitocin? Am J Obstet Gynecol. 185: S212.
- Wenstrom K, Andrews WW, Maher JE, (1995):** Amnioinfusion survey: prevalence, protocols and complications. Obstet Gynecol. 86: 572-6.
- Whang JD, Roh CR, Yang SH et al., (2001):** Prenatal outcome with active expectant management of second trimester rupture of membranes. Korean J Obstet Gynecol. 44: 348-54.
- Whitelam A, and Thoresen M. (2000):** Antenatal steroids and the developing brain. Arch Dis Child Fetal Neonatal Ed. 83 (2): F154-7.
- Wing DA, and Paul RH, (1998):** Induction of labor with misoprostol for PROM beyond thirty-six weeks' gestation. Am J Obstet Gynecol. 179: 94-9.
- Wing DA (2002):** A benefit-risk assessment of misoprostol for cervical ripening and labor induction. Drug Safety. 25: 665-76.
- Woyton J, Klosek A, Zimmer M, et al., (1999):** Insulin-like growth factor binding protein-1, in vaginal secretion as a marker of PROM Ginekol Pol. 70 (11): 809-14.
- Wu YW, and Colford JM (2000):** Chorioamnionitis as a risk factor for cerebral palsy: a meta-analysis. JAMA. 284: 1417-1427.

- Xenakis EMJ, Langer O, Piper JM, et al., (1995):** Low-dose versus high-dose oxytocin augmentation of labor- a randomized trial. Am J Obstet Gynecol. 173: 1874.
- Yogev Y, Glickman H, Hod Moshe G. et al., (2004):** Mode of delivery in pregnancies with premature rupture of membranes at or before term following induction of labor with vaginal prostaglandin E2. Am Journal of Perinatology. Vo. 21, N. 5.
- Yoon BH, Kim YA Romero R, et al. (1999):** Association of oligohydramnios in women with premature rupture of membranes with an inflammatory response in fetal, amniotic, and maternal compartments. Am J Obset Gynecol. 181: 784-788.
- Yoon BH, Park CW, Chaiworapongsa T. (2003):** Intrauterine infection and the development of cerebral palsy. BJOG; 110: Suppl 20: 124-7.
- Young BK, Mackenzie AP, Roman AS, et al., (2004):** Endoscopic closure of fetal membrane defects: comparing iatrogenic versus spontaneous rupture cases. J Matern Fetal Neonatal Med. 16 (4): 235-40.
- Zimmermann R, Aline W, Ursula M, (2006):** Neonatal Mortality and Morbidity after Aggressive Long-term tocolysis for preterm premature rupture of the membranes. Fetal Diagnosis and Therapy. 21: 366-373.