
REFERENCES

Aantaa, R.; Kanto and Scheinin, M. (1990):

Assessment of the sedative effects of dexmedetomidine, on $\alpha 2$ -adrenoceptor agonist, in minor gynecologic operations. *Anesth. Analg.*, 26: 407.

Aldrete, J.A. and Kroulik, D. (1970):

A post anesthetic recovery score. *Anesth. Analg.*, 49: 924.

Aldrete, J.A. (1995):

The post anesthesia recovery score revisited. *Clinical Anesthesiology*, 7: 89-91.

Alon, E. and Himmelseher, S. (1998):

Ondasteron in the treatment of postoperative vomiting, a randomized double-blind comparison with droperidol and metoclopramide. *Anesthesia Analgesia*, 14: 509.

Anderson, H.J. (1992):

Surgery delivery systems change as more patient go home the same day. *Hospitals*, 6: 33.

Apfelbaum, J.; Roizen, M.F. and Robinson, D. (1993):

How frequently do asymptomatic patients benefit from the pursuit of abnormalities in their preoperative tests results. *Anesthesiology*, 10: 1254.

Ashburn, M.A.; Lind, G.H. and Gillie, M.H. (1993):

Oral transmucosal fentanyl citrate for the treatment of postoperative pain. *Anesth. Analg.*, 30: 377.

Bailey, P.L.; Stiesand, J.B. and Pace, N.L. (1990):

Transdermal scopolamine reduces nausea and vomiting after outpatient laparoscopy. *Anesthesiology*, 21: 277.

Baily, P.L.; Egan, T.D. and Stanley, T.H. (2000):

Intravenous opioid anesthetics. In Miller R.D., *Anesthesia*, 5th ed, London, Churchill Livingstone, 273-277.

Barbara, A.C. (1997):

Opioids. In: Barash PG, Cullen BG and Stoelting RK (eds), Clinical Anesthesiology, 3rd ed., Philadelphia, New York, Lippincott-Raven, 329-358.

Basta, S.J.; Savarese, J.J. and Ali, H.H. (1992):

Clinical pharmacology of a tracurium besylate. Anesthesia Analgesia, 24: 723.

Basta, S.J. (1992):

Clinical pharmacology of mivacurium. Clin. Anesth., 12: 153.

Berkoff, M.J. and Pangrazio, J.R. (1986):

Planning and designing ambulatory surgical facilities. In: Ambulatory Surgery, Davis JE (ed), Baltimore, Williams & Wilkins, 3: 172.

Bernard, J.; Faintreny, A.; Haut, P. and Pinaud, M. (1993):

Patient controlled premedication by I.V. midazolam for ambulatory anesthesia. Anesthesiology, 11: 119.

Berry, A.J.; Briggs, T.P.; Parker, C. and Milder, R.A. (1991):

Beds or day case unit: your choice? Ann. R. Coll. Surg. Eng., 8: 114.

Bitner, R.L. (1983): Awareness during anesthesia. In: Complications of anesthesia. Orkin FK & Cooperman LH (eds), Philadelphia, JB Lippincott, 35: 349.

Boore, J. (1987):

Nursing the physically ill adult. In: Medical Surgical Nursing. London, Churchill Livingstone, 15: 158.

Borgeat, A.; Wilder-Smith, O.H.G. and Suter, P.M. (1994):

The non-hypnotic therapeutic applications of propofol. Anesthesiology, 11: 642.

Borgeat, A.; Wilder-Smith, O.H.G. and Rifat, K. (1992):

Subhypnotic doses of propofol possess antiemetic properties and relieve pruritus induced by epidural and spinal morphine. *Anesth. Analg.*, 27: 539.

Borgeat, A.; Wilder-Smith, O.H.G. and Suter, P.M. (2000):

The non hypnotic therapeutic applications of propofol. *Anesthesiology*, 11: 688.

Brandner, B.; Blagrove, M.; McCallum, G. and Bromley, L.M. (1997):

Comparative study of the recovery of propofol vs desflurane. *Anesthesiol.*, 52: 750.

Brown, A.R.; Weiss, R. and Greenberg, C.P. (1993):

Interscalene block for shoulder arthroscopy, A comparison with general anesthesia. *J. Arthro. Related. Surgery*, 25: 295.

Cade, L. and Ashley, J. (1993):

Prophylactic paracetamol for analgesia after vaginal termination of pregnancy. *Anesth. Inten. Care*, 11: 93.

Caplan, R.A.; Posner, K.L.; Ward, R.J. and Cheney, F.W. (1990):

Adverse respiratory events in anesthesia. *Anesthesiology*, 12: 828.

Cattalini, D. (1993):

Best practices in ambulatory surgery. *Healthcare Forum. J.*, 6: 28.

Chicoine, R.E.; Lilly, R.B.; Manzi, D.J.; Gannon, R.H. (1993):

Oral ibuprofen is less expensive and as effective as parenteral Ketorolac in ambulatory surgical patients. *Anesthesiology*, 11: 254.

Chung, F.; Ong, D. and Seyone, C. (1991):

PADS-A discriminative discharge index for ambulatory surgery. *Anesthesiology*, 34: 1105.

Chung, F. (1993):

Are discharge criteria changing? J. Clin. Anesth., 25: 647.

Chung, F.F.; Chan, V.W.S. and Ong, D. (1993):

A postanesthetic discharge scoring system for home readiness after ambulatory surgery. Amb. Surg., 6: 189.

Cohen, D. and Dillon, J. (1966):

Anesthesia for outpatient surgery. JAMA, 8: 98.

Cohen, M.M.; Duncan, B.G.; DeBoer, D.P. and Tweed, W.A. (1994):

The postoperative interview, assessing risk factors for nausea and vomiting. Anesth. Analg., 22: 7.

Cohen, S.E.; Woods, W.A. and Wyner, J. (1984):

Antiemetic efficacy of droperidol and metoclopramide. Anesthesiology, 11: 67.

Cros, A.M. (1994):

Surgical procedures in outpatient unit. Chaiers d'Anesth., 1: 15.

Davis, J.E. (1995):

Ambulatory surgery. How far we can go? Med. Clinics N. Amer., 3: 77.

de Bros, F. (1994):

Pharmacokinetic evaluation mivacurium during infusion in neurosurgical patient. Anesthesiology, 24: 1185.

Delahunt, B. and Turnbull, P.R.G. (1980):

How cost effective are routine preoperative investigations? J. Clin. Anesth., 10: 430.

Denny, N.M. and Gadelrab, R. (1980):

Complications following general anesthesia for cataract surgery, a comparison of the laryngeal mask airway with tracheal intubation. Anesthesiology, 12: 512.

Detmer, D.E. (1981):

Ambulatory surgery. N. Eng. Med. J., 8: 305.

Devitt, J.H.; Wenstone, R.; Noel, A.C. and O'Donnel, M.P. (1994):
The laryngeal mask airway and positive pressure ventilation. *Anesthesiology*, 11: 550.

Diamond, M.J. and Kee-i-Szanto, M. (1980):
Reduction of postoperative vomiting by preoperative administration of oral metoclopramide. *Can. Anesth. Soc. J.*, 11: 36.

Ding, Y. and White, P.F. (1992):
Comparative effects of Ketorolac and fentanyl as adjuvants during outpatient anesthesia. *Anesth. Analg.*, 7: 566.

Ding, Y.; Freedman, B. and White, P.F. (1993):
Use of ketorolac and fentanyl during outpatient gynecologic operations. *Anesth. Analg.*, 23: 205.

Ding, Y.; Freedman, B. and White (1994):
Use of mivacurium during laparoscopic surgery, effect of reversal drugs on postoperative recovery. *Anesth. Analg.*, 24: 450.

Dobbs, A.R. (1984):
Patient learning in ambulatory unit. *Anesthesiology*, 2: 36.

Duncan, P.C.; Cohen, M.M. and Tweed, W.A. (1992):
Are anesthetic complications predictable in day surgical practice? *Can. J. Anesth.*, 27: 440.

Dyck, J.B. and Chung, F. (1991):
A comparison of propranolol and diazepam for preoperative anxiolysis. *Can. J. Anesth.*, 11: 704.

Eddy, D.M.; Hasselblad, V.; McGivney and Hendee, W. (1988):
The value of mammography screening in women under age 50 years. *Anesthesiology*, 10: 259.

Fancourt-Smith, P.F. and Merrick, P. (1992):
Anesthesia for outpatients. *Can. Med. Assoc.*, 39: 893.

Fang, Z.X.; Eger, E.L. and Laster, M.J. (1995):

Carbon monoxide production from degradation of desflurane, enflurane, isoflurane and sevoflurane by soda lime and baralyme. *Anesth. Analg.*, 22: 1186.

Feldman, M.; Pasternak, R. and Paul, S. (1994):

Morbidity and mortality after ambulatory surgery. *Anesthesiology*, 9: 217.

Forrest, P.; Galletly, K.E. and Yee, P. (1987):

Placebo controlled comparison of midazolam, triazolam and diazepam as oral premedicants for outpatient anesthesia. *Anesth. Intern. Care*, 11: 296.

Foster, W.E.; Griffith, K.E. and Beck, P.M. (1993):

Computer assisted preoperative assessment for ambulatory anesthesia. *Anesthesiology*, 2: 54.

Freeman, L.N.; Schachat, A.P. and Manolio, T.A. (1988):

Multivariate analysis of factors associated with unplanned admission in outpatient ophthalmic surgery. *Ophthalmic Surg.*, 33: 719.

Fu, C.; Mishra, Y. and Ramzan, I. (1994):

Omeprazole potentiates atracurium and succinylcholine paralysis in vivo in rats. *Anesth. Analg.*, 11: 527.

Ghoneim, M.M. and Block, R.I. (1992):

Learning and consciousness during general anesthesia. *Anesthesiology*, 35: 279.

Ghoneim, M.M.; Block, R.I. and Sum Ping, S.T. (1993):

The interactions of midazolam and flumazenil on human memory and cognition. *Anesthesiology*, 21: 1183.

Gibby, G.L.; Gravenstein, Layone, A.J. and Jackson, K.I. (1992):

How often does the preoperative interview change anesthetic management? *Anesthesiology*, 10: 325.

Glass, P.S.A.; Shafer, S.L.; Jacobs, J.B. and Reves, J.G. (1994):

Intravenous drug delivery systems. In: Anesthesia, RD Miller (Ed.), New York, Churchill Livingstone, 20: 389.

Hannallah, R.S. and Epstein, B.S. (1991):

The pediatric patient. In: Anesthesia for ambulatory surgery, BV Wetchler (Ed.), Philadelphia, JB Lippincott, 39: 131.

Hannallah, R.S. and Patel, R.I. (1995):

Pediatric considerations. In: The ambulatory anesthesia handbook, RS Twersky (Ed.), St Louis, CV Mosby, 39: 145.

Hayward, R.S.A.; Steinberg, E.P.; Ford, D.E.; Roizen, M.F. and Roach, K.W. (1991):

Preventive care guidelines. Ann. Intern. Med., 10: 114.

Heizmann, P.; Eckert, M. and Zeigler, W.H. (1983):

Pharmacokinetics and bioavailability of midazolam in man. J. Clin. Pharmacol., 16: 435.

Henderson, J.A. (1991):

Ambulatory surgery: past, present & future. In: Anesthesia for ambulatory surgery. Wetchler BV (Ed), JB Lippincott, Philadelphia, 1: 27.

Hertz, C.M.; Sibert, K.; Huttemeier, P. and Ginsberg, B. (1993):

Oral administration of ranitidine for premedication in outpatient anesthesia. Anesthesiology, 11: 115.

Hickock, C. and Naulty, J. (1994):

Recovery from desflurane after induction with propofol vs thiopental. Anesth. Analg., 23: 158.

Higgins, M.S.; Givogre, J.L. and Marco, A.P. (1994):

Recovery from outpatient laparoscopic tubal ligation is not improved by preoperative administration of ketorolac or ibuprofen. Anesth. Analg., 23: 274.

Hodsman, N.B.A.; Burns, J. and Blyth, A. (1987):

The morphine sparing effects of diclofenac following abdominal surgery. Anesthesia, 42: 1005.

Holley, F.O. and Van Steennis, C. (1988):

Postoperative analgesia with fentanyl; iv vs transdermal delivery. Br. J. Anesthesia, 30: 608.

Jaakola, M.L.; Ali-Melkkila, T. and Kanto, J. (1992):

Dexmedetomidine reduces intraocular pressure, intubation responses and anesthetic requirements in patients undergoing ophthalmic surgery. Br. J. Anesth., 26: 570.

Jakobsen, C.J.; Jensen, J.J.; Hansen, W. and Grabe, N. (1986):

Oral lorazepam in premedication. A comparison with diazepam. Anesthesia, 11: 457.

Jarrett, P.E.M. (1994):

Some economic aspects of day surgery. Amb. Surg., 2: 102.

J. Lance, Lichtor and Bernard, V. Wetchler (1997):

Anesthesia for ambulatory surgery. In: Clinical Anesthesia, Paul G. Barash, Cullen BF and Stoelting RK (eds), 3rd Ed., Philadelphia.

Jeffery, L. Apfelbaum (2001):

Current controversies in adult outpatient anesthesia. In ASA; 3: 151.

Joshi, G.P. (1994):

Postoperative pain management-new drugs and techniques. Curr. Anesth. Crit. Care, 5: 150.

Kallar, S.K. and Chung, F. (1992):

Practical application of postanesthetic discharge scoring system. Anesthesiology, 21: 345.

Kallar, S.K. and Everett, L.L. (1993):

Potential risks and preventive measures for pulmonary aspiration: new concepts in preoperative fasting guidelines. Anesth. Analg., 11: 171.

Kallio, A.; Salonen, M. and Forssell, H. (1990):

Medetomidine premedication in dental surgery. Acta. Anesthesiol., Scand., 11: 184.

Kapur, P.A. (1994):

Pharmacy acquisition costs in ambulatory anesthesia. *Anesth. Analg.*, 44: 617.

Karp, K.B. (1995):

Anesthetic recovery after ambulatory anesthesia. *Am. Assoc. Nurse Anesthet.*, 8: 109.

Kaufman, E.; Hargreaves, K.M. and Dionne, R.A. (1993):

Comparison of oral triazolam and nitrous oxide with placebo and intravenous diazepam for outpatient premedication. *Oral Surg. Oral. Med. Oral. Path.*, 11: 156.

Kehlet, H. and Dahl, J.B. (1993):

The value of Balanced Analgesia in postoperative pain treatment. *Anesth. Analg.*, 32: 1258.

Kitz, D.S.; Slusary-Ladden, C. and Lecky, J.H. (1990):

Hospital resources used for inpatients and ambulatory surgery. *Anesthesiology*, 3: 256.

Korttila, K. (1990):

Recovery and home readiness after anesthesia for ambulatory surgery. *Anesthesiology*, 9: 182.

Kovac, A.; Cakins, J.; Joslyn, A. and Cimfel, C. (1994):

Recovery from ambulatory surgery of male outpatients receiving ondansetron. *Anesth. Analg.*, 9: 220.

Leider, L.K.; Mannix, P.M. and Frost, A.B. (1991):

A successful facility. In: *Anesthesia for ambulatory surgery*. BV Wetchler (Ed), Philadelphia, JB Lippincott, 6: 452.

Lieber, C.P.; Seinige, U.L. and Sataloff, D.M. (1990):

Choosing the site of surgery an overview of ambulatory surgery in geriatric patients. *Clin. Geriat. Med.*, 8: 425.

Llewellyn, J.G. (1991):

Short stay surgery. *Can. J. Anesth.*, 6: 112.

Loeffler, P.M. (1992):

Oral benzodiazepines and conscious sedation. J. Oral Maxillofac. Surg., 11: 656.

Lumsdon, K.; Anderson, H.J. and Burke, M. (1992):

New surgical technologies reshape hospital strategies. Hospitals, 6: 30.

Lynch, J. (1993):

Free-standing surgery centers, the wave of the future. Anesthesiology, 18: 42.

Mackenzie, J.W. (1993):

A novel approach to anxiolytic premedication for day-case patients. J.P. Soc. Med., 11: 652.

Madej, T.H. and Simpson, K.H. (1986):

Comparison of the use of domperidone, droperidol and metoclopramide in the prevention of nausea and vomiting following gynecologic surgery in day-cases. Br. J. Anesth., 12: 853.

Manchikanti, L.; Grow, J.B.; Collvier, J.A.; Hadley, C.H. and Hohlbein, L.J. (1985):

Sodium citrate and metoclopramide in outpatient anesthesia for prophylaxis against aspiration pneumonitis. Anesthesiology, 26: 378.

Mangano, D.T. (1990):

Perioperative cardiac morbidity. Anesthesiology, 9: 153.

McKenzie, R.; Shrift-Azad, S. and Dershwietz, M. (1998):

A randomized study examination of the use of intravenous ondansetron in the prevention of postoperative nausea and vomiting in female outpatients. Clin. Anesth., 5: 30.

Melnick, B.M.; Sawyer, R. and Karembelkar, D. (1989):

Delayed side effect of droperidol after ambulatory general anesthesia. Anesth. Analg., 11: 784.

Meridy, H.W. (1982):

Criteria for selection of ambulatory surgical patients and guidelines for anesthetic management, a retrospective study of 1533 patients. *Anesth. Analg.*, 12: 921.

Meyer, P.; Thisted, R. and Roizen, M.F. (1991):

Improving preoperative lab test selection. *Anesth. Analg.*, 8:365.

Mike, N.; Hiraiwa, K. and Shimada, Y. (1994):

Effects of oral clonidine premedication on hemodynamic and intraocular pressure changes in cataract surgery with local anesthesia. *Anesth. Analg.*, 8: 458.

Millar, J.M. and Hall, P.J. (1987):

Nausea and vomiting after low dose droperidol in cases of termination of pregnancy. *Anesthesia*, 42: 613.

Miller, C.D. and Anderson, W.G. (1988):

Silent regurgitation in day case gynecological patients. *Anesthesia*, 43: 321.

Moote, C. (1992):

Efficacy of non-steroidal anti-inflammatory drugs in the management of postoperative pain. *Drugs*, 15: 265.

Naito, Y.; Shingu, K. and Fujimori, R. (1991):

Comparison between sevoflurane and halothane for pediatric ambulatory anesthesia. *Br. J. Anesth.*, 22: 387.

Nathanson, M.H.; Fredman, Smith, I. And White, P.F. (1995):

Sevoflurane vs desflurane for outpatient anesthesia. *Anesth. Analg.*, 22: 1186.

Natof, H.E. (1984):

Pre-existing diseases in ambulatory surgery. *Anesthesia*, 54: 899.

Nicoll, J.H. (1990):

The surgery of infancy. *Br. Med. J.*, 2: 63.

O'Hara, D.A.; Fragen, R.J. and Kinzer, M. (1987):

Ketorolac as compared with morphine for treatment of postoperative pain. Clin. Pharmacol. Ther., 23: 556.

Ogg, T.W. (1972):

An assessment of postoperative outpatient cases. Br. Med. J., 34: 573.

Olin, B.R. (1993):

Facts and comparisons about drugs for ambulatory anesthesia. Clin. Pharmacol. Ther., 54: 684.

Ong, B.Y.; Palahnuik, R.J. and Cumming, M. (1978):

Gastric volume and pH in outpatients. Can. Anesth. Soc. J., 11: 36.

Orkin, F.K. (1995):

Quality improvement. In Ambulatory anesthesiology, KE McGoldrick (Ed). Baltimore, Williams & Wilkins, 46: 682.

Orkin, F.K. (1992):

What do patients want? Preference for immediate postoperative recovery. Anesth. Analg., 27: 225.

Osborne, G.A. and Rudkin, G.E. (1993):

Outcome after day-case surgery in a major teaching hospital. Anesth. Intens. Care, 21: 822.

Ostman, P.L. and White, P.F. (1994):

Outpatient anesthesia. In: Anesthesia, RD Miller (Ed). New York, Churchill Livingstone, 32: 2213.

Ostman, P.L. and White, P.F. (2000):

Outpatient anesthesia. In: Anesthesia; RD Miller (Ed.) New York, Churchill Livingstone, 32: 2260.

Pandit, S.K.; Kothary, S.P.; Pandit, U.A. and Mirakhur, R.K. (1986):

Premedication with cimetidine and metoclopramide. Anesthesia, 41: 486.

Pandit, S.K.; Kothary, S.P.; Pandit, U.A.; Pandel, G. and Levy, L. (1989):

Dose-response study of droperidol and metoclopramide as antiemetics for outpatient anesthesia. *Anesth. Analg.*, 28: 798.

Park Ridge (1996):

Standards of basic anesthetic monitoring. In *ASA*, 13: 785.

Park Ridge (1999):

Guidelines for ambulatory anesthesia. In *ASA*, 3: 385.

Parnass, S.M.; McCarthy, R.J. and Bach, B.R. (1993):

Beneficial impact of epidural anesthesia on recovery after outpatient arthroscopy. *Arthroscopy*, 9: 91.

Parnis, S.J.; Foate, J.A. and Short, T. (1992):

Oral midazolam is an effective premedication for children having day-stay anesthesia. *Anesth. Intens. Care*, 11: 9.

Pasternak, D.P.; Smith, H.L. and Piland, N.F. (1996):

Critical issues surrounding the evolution of ambulatory surgery. *J. Amb. Care Manage.*, 9: 171.

Phillip Scuderi and Winston Salem (2001):

Management of postoperative nausea and vomiting. In *ASA*, 6: 223.

Pittoni, G.; Toffoletto, F.; Galcarella, G. and Giron, G.P. (1995):

Spinal anesthesia in outpatient knee surgery. *Anesth. Analg.*, 25: 81.

Power, I.; Noble, D.W.; Douglas, E. and Spence, E.E. (1990):

Comparison of IM Ketorolac and morphine for pain relief after cholecystectomy. *Br. J. Anesth.*, 23: 448.

Rabkin, S.W. and Horne, J.M. (1983):

Preoperative electrocardiography, effect of new abnormalities on clinical decisions. *Can. Med. Assoc. J.*, 10: 146.

Rasmussen, G.L.; Steckner, K.; Hogue, C. et al. (2001):

Intravenous parecoxib sodium after orthopedic knee surgery. American Pain Society.

Raybould, D. and Bradshaw, E.G. (1987):

Premedication for day-case surgery. *Anesthesia*, 11: 591.

Reasbeck, P.G.; Rice, M.L. and Reasbeck, J.C. (1982):

Double blind controlled trial of indomethacin as an adjunct to narcotic analgesia after major abdominal surgery. *Anesthesiology*, 23: 568.

Reed, W.A. and Ford, J.L. (1969):

The surgicenter is an innovation in the delivery and cost of medical care. *Ariz. Med. J.*, 26: 801.

Rebecca, S. Twersky and Brooklyn (2001):

Recovery and discharge of ambulatory anesthesia patient. In *ASA*, 4: 275.

Reich, D.L.; Konstadt, S.M. and Stanely, T.E. (1995):

The cardiovascular effects of cisatracurium in patients with coronary artery diseases. *Anesth. Analg.*, 24: 385.

Reves, J.G.; Fragen, R.J.; Vink, R. and Greenblatt, D.J. (1985):

Pharmacology and uses of midazolam. *Anesthesiology*, 62: 310.

Rigg, J.D.; Wilson, A.C. and Pollard, P.J. (1997):

Mivacurium or Vecuronium for muscular relaxation in day-case surgery. *Eur. J. Anesthesiol.*, 14: 630.

Roberson, C.R. and Walker, R.T. (1993):

Comparative value of ondansetron and metoclopramide in the prevention postoperative nausea and vomiting. *Anesth. Analg.*, 11: 350.

Roizen, M.F.; Kaplan, E.B. and Schreider, B.D. (1987):

The relative roles of the history and physical examination, and lab testing in preoperative evaluation for outpatient surgery. *Anesthesiol. Clin. North Am.*, 1: 15.

Rosenberg, M.K.; Bridge, P. and Brown, M. (1994):

Cost comparison, a desflurane vs a propofol-based general anesthesia. *Anesth. Analg.*, 44: 852.

Rossello, P.J.; Cruz, A.R. and Mayol (1980):

Routine lab tests for elective surgery in pediatric patients. *Bol. Asoc. Med. J.*, 10: 614.

Roth, J.H.; Richards, R.S. and MacLeod, M.D. (1994):

Endoscopic Carpal Tunnel release. *Can. J. Surg.*, 21: 189.

Rudkin, G.E.; Osborne, G.A. and Doyle, C.E. (1995):

Assessment and selection of patients for day-case surgery. *Med. J. Aust.*, 8: 308.

Saint-Maurice, C.; Kong-ky, B.H. and Mathey, S. (1995):

Pediatric surgery and anesthesia in day-surgery hospital. *Cahier's d'Anesth.*, 41: 407.

Sandin, R. and Nordstrom, O. (1993):

Awareness during total intravenous anesthesia. *Br. J. Anesth.*, 35: 762.

Sandler, G. (1979):

Costs of unnecessary tests. *Br. Med. J.*, 2: 21.

Schenckman, Z.; Shir, Y. and Brodsky, J.B. (1993):

Perioperative management for obese patient. *Br. J. Anesth.*, 12: 356.

Schreiner, M.S.; Nicholson, S.C. and Martin, T. (1992):

Should children drink before discharge from day-surgery? *Anesthesiology*, 34: 528.

Schuind, F.; Dap, F.; Farcot, J.M.; Jost, H. and Foucher, G. (1986):

Outpatient hand surgery. *Ann. Hand Surg.*, 5: 122.

Schwid, H.A. and O'Donnell, D. (1992):

Anestheisologist's management of simulated critical incidents. *Anesthesiology*, 52: 496.

Scott, R.P.F.; Savarese, J.J. and Basta, S.J. (1986):

Clinical pharmacology of atracurium given at high doses.
Br. J. Anesth., 24: 851.

Scuderi, P.E. (2000):

Multimodal management prevents early postoperative vomiting after outpatient laparoscopy. Anesth. Analg., 41: 1408-14.

Seeberger, M.D.; Lang, M.J. and Drewe, J. (1994):

Comparison of spinal and epidural anesthesia in patients younger than 50 years of age. Anesth. Analg., 12: 667.

Sevarino, F.B.; Sinatra, R.S. and Paige, D. (1992):

The efficacy of intramuscular ketorolac in combination with intravenous PCA morphine for postoperative pain relief. J. Clin. Anesth., 4: 285.

Shafer, A.; White, P.F.; Urquhart, M.L. and Doze, V.A. (1989):

Use of midazolam and opioid analgesics in premedication of outpatients. Anesthesiology, 23: 742.

Shanks, C.A.; Fragen, R.J. and Ling, D. (1993):

Continuous intravenous infusion of Rocuronium in patient receiving balanced, enflurane, or isoflurane anesthesia. Anesthesiology, 24: 649.

Short, T.C. and Galletly, D.C. (1989):

Double-blind comparison of midazolam and temazepam as oral premedicants for outpatient anesthesia. Anesth. Inten. Care, 17: 151.

Smith, G.B.; Hughes, D.G. and Kumar, V. (1985):

Temazepam in fast dispensing doses form as premedicant for children. Anesthesia, 11: 368.

Smith, I. And Joshi, G. (1993):

The laryngeal mask airway for outpatient anesthesia. J. Clin. Anesth., 5: 22.

Smith, I. And White, P.F. (1994):

Anesthetic considerations for laparoscopic surgery. Seminars in Laparoscopic Surgery, 1: 17.

Smith, I.; Van Hemelrijck, J.; White, P.F. and Shively, R. (1991):

Effects of local anesthesia on recovery after outpatient arthroscopy. Anesth. Analg., 25: 538.

Smith, I.; White, P.F.; Nathanson, M. and Gouldson, R. (1994):

Propofol, an update on its clinical uses. Anesthesiology, 45: 1005.

Smith, I.; Zelcer, J. and White, P.F. (1993):

Anesthesia for outpatient surgery. Curr. Opin. Anesth., 5: 356.

Souter, A.; Fredman, B. and White, P.F. (1994):

Controversies in the perioperative use of non-steroidal anti-inflammatory drugs. Anesth. Analg., 42: 1178.

Striebel, W.H.; Pommerening, J. and Reiger, A. (1993):

Intransal fentanyl titration for postoperative pain management of unselected population. Anesthesia, 30: 753.

Sukhani, R.; Lurie, J. and Jabamoni, R. (1994):

Propofol for ambulatory gynecologic laparoscopy. Anesth. Analg., 31: 831.

Sung, Y.F.; Reiss, N. and Tillette, T. (1991):

The differential cost of anesthesia and recovery with propofol-nitrous oxide versus thiopental-isoflurane-nitrous oxide anesthesia. J. Clin. Anesth., 3: 391.

Talke, P.O. and Solanki, D.R. (1993):

Dose-response study of oral famotidine for reduction of gastric acidity and volume in outpatients and inpatients. Anesth. Analg., 31: 1124.

Tang, J. (2001):

Fast-track office based anesthesia. Anesthesia Analgesia, 92: 95-99.

Tarikilla, P.; Tuominen, M. (1997):

Anesthesia with small doses of hyperbaric 0.5% bupivacaine. *Anesthesia*, 52: 1157.

Thornberry, E.A. and Thomas, T.A. (1988):

Posture and post-spinal headache. *Br. J. Anesth.*, 7: 155.

Thwaites, A.; Edmends, S. and Smith, I. (1997):

Inhalational induction with sevoflurane. *Br. J. Anesth.*, 24: 356.

Turner, N.; Scarpace, P.J. and Lowenthal, D.T. (1992):

Geriatric pharmacology, basic and clinical considerations. *Ann. Rev. Pharmacol. Toxicol.*, 12: 271.

Vaghadia, H. and Smythe, S. (1994):

Long-term follow up after ambulatory surgery. *Abstracts of Ambulatory Anesthesia Meeting Chicago*, 35: 1.

Valanne, J. and Korttila, K. (1985):

Effect of small dose of droperidol on nausea, vomiting and recovery after outpatient enflurane anesthesia. *Acta. Anesthesiol. Scand.*, 27: 359.

van Ee, R.; Hemrika, D.J. and van der Linden, C.T.M. (1993):

Pain relief following day-case diagnostic hysteroscopy, laparoscopy for infertility. *Anesthesia*, 12: 538.

Vaughn, R.W.; Aluise, J.J. and McLaughlin, C.P. (1991):

Ambulatory surgery and the hospital. *Health Care Manage Rev.*, 16: 15.

Vickery, R.G.; Sheridan, B.C. and Maze, M. (1988):

Anesthetic and hemodynamic effects of the stereoisomers of medetomidine in halothane anesthetized dogs. *Anesth. Analg.*, 11: 611.

Warner, M.; Shields, S.E. and Chute, C.G. (1993):

Major morbidity and mortality within 1 month of ambulatory surgery and anesthesia. *JAMA*, 48: 1437.

Watcha, M.F. and White, P.F. (1992):

Postoperative nausea and vomiting; its etiology, treatment and prevention. *Anesthesiology*, 12: 167.

Watcha, M.F.; Safavi, F.Z.; McCulloch, D.A.; Tan, T.S.H. and White, P.F. (1995):

Effect of antagonism of mivacurium-induced neuromuscular block on postoperative emesis in children. *Anesth. Analg.*, 37: 713.

Waters, R.M. (1991):

The downtown anesthesia. *Clinic. Am. J. Surg.*, 21: 271.

Wetchler, B.V. (1990):

Problem solving in postanesthesia care unit. In: *Anesthesia for ambulatory surgery*, BV Watcha (ed) Philadelphia-Lippincott, 43: 451.

Wetchler, B.V. (1988):

The geriatric outpatient. In: *Problems in anesthesia*. BV Wetchler (Ed.) Philadelphia, JB Lippincott, 8: 278.

White, P.F. and Smith, I. (1993):

Impact of newer drugs and techniques on the quality of ambulatory anesthesia. *J. Clin. Anesth.*, 3: 35.

White, P.F. (Ed.) (1994):

Anesthesia for ambulatory surgery. International Anesthesia Clinics. Boston, Little Brown, 1: 1.

White, P.F. (1986):

Pharmacologic and clinical aspects of preoperative medication. *Anesth. Analg.*, 32: 965.

Wierda, J.M.; Wit, A.P.; Kuizanga, K. and Agoston, S. (1990):

Clinical Observations on the neuromuscular blocking actions of Rocuronium. *Br. J. Anesth.*, 24: 521.

Wolfson, J.; Walker, G. and Levin, P.J. (1993):

Freestanding ambulatory surgery. Cost containment winner?
Healthcare Financial Manage, 6: 27.

Wood, C. (1994):

An evaluation of day-surgery telephone follow up. Journal
of One Day Surgery, 4: 35.

Zasa, R.J. (1999):

Ambulatory surgery; critical management indicators. J.
Amb. Care Manage., 6: 69.