

REFERENCES

Aldrte J.A. (1995):

The post anesthesia recovery score revisited.
J. Clin. Anesth. 7: 89-91.

Anand KJS. Spell WG. and Aynsley – Green A (1987):

Randomized trial of fentanyl anesthesia in preterm babies undergoing surgery: Effects on the stress response. Lancet 1: 243-247.

Arendt-Nielsen L, Nielsen JC and Bjerring P. (1991):

Double blind placebo controlled comparison of paracetamol and paracetamol plus codeine – a quantitative evaluation by laser – induced pain. Eur. J. Clin. Pharmacol. 40: 241-247.

Armitage EN (1986):

Local anesthetic techniques for prevention of postoperative pain. Br. J. Anaesth. 58: 790-800.

Audrey G, Johanet S, Achrafi, H., Lupoid M and Gruner M (1994):

The risk of wound infection after inguinal incision in pediatric outpatient surgery. Eur. J. paediatr. Sur. 4: 87-89.

Bander NH (1992):

Epidural agents for postoperative analgesia.
Anesthesiol. Clin. North Am 10: 321-337

Batra MS (1992):

Epidural and spinal analgesia and anesthesia:
Contemporary issues.
Anesthesiol. Clin. North. Am. Vol. 10.

Beebe DS, Belani KG, Change PN, et al. (1992):

Effectiveness of preoperative sedation with rectal midazolam, Ketamine or their combination in young children. Anesth. Analg. 75: 880-884.

Bell C, Kain ZN (1997):

The Pediatric Anesthesia Handbook, 2nd ed. Mosby, 1997.

Berde C, Ablin a glazer J et al., (1990):

American Academy of Pediatrics report of the subcommittee on disease related pain childhood cancer. Pediatrics 86: 818-825.

Bissonnette B (1992):

Temperature monitoring in pediatric anesthesia. Int. Anesthesiol Clin. 30: 63.

Bone ME and Fell D (1988):

A comparison of rectal Dicolfenac with intramuscular papaveretum or placebo for pain relief following tonsillectomy. Anaesthesia 43: 277-280.

Broadman LM, Hannallah, RS, Belman B et al., (1987):

Post circumcision analgesia – a prospective evaluation of subcutaneous ring block of the penis. Anesthesiology 67: 399-402.

Colan, S., Fujii, A.; Borow, K. (1983):

Non invasive determination of systolic, diastolic and end systolic blood pressure in neonates, infants and young children. Am. J. Cardiol., 52: 867-875.

Cote C.G.; Rolf, N.; Liu, L.M.P. et al., (1991):

A single blind study of combined pulse oximetry and capnography in children. Anesthesiology, 74: 980-987.

Cote C.J.; Goldstein, E.A.; Cote, M.A. et al. (1988):

A single-blinded study of pulse oximetry in children. Anesthesiology, 68: 184.

Cote CJ, Zaslavsky A, Dore CJ et al., (1995):

Postoperative apnea in former preterm infants after inguinal herniorrhaphy. *Anesthesiology* 82: 809-822.

Cros, A.M. (1994):

Surgical procedures in outpatient unit. *Chaiers d'Anesth.*, 1:15.

Dahl JB and Kehlet H (1991):

Non-steroidal anti-inflammatory drugs: Rationale for use in severe postoperative pain. *Br. J. Anesth.* 66: 703-712.

Darrow EJ, Badgwell, JM and Heavner JE (1994):

Bupivacaine and epinephrine act synergistically to produce ventricular arrhythmias in young pigs *Anesthesiology* 81: A 1357.

Davis P, Tome JA McGowan FX et al. (1995):

Preanesthetic medication with intranasal midazolam for brief pediatric surgical procedure: effect on recovery and hospital discharge times *Anesthesiology* 82: 2-5.

Davis P.J.; Lerman J. Sureh S et al., (1997):

Randomized multicenter study of propofol compared with alfentanil, isoflurane, or propofol in anesthetized pediatric patients undergoing elective strabismus surgery. *Anesth. Analg*; 84: 982-990.

Dexter F, et al., (1999):

Computer simulation to determine how rapid anesthetic recovery protocols to decrease the time of emergence or increase the phase I post anesthesia care unit bypass rate affect staffing of an ambulatory surgery center. *Anesth Analg* 88: 1053-63.

Doi M and Ikeda K (1993):

Airway irritation produced by volatile anesthetics during brief inhalation: comparison of halothane, enflurane, isoflurane and sevoflurane. *Can. J. Anaesth.* 40: 122-126

Ecoffey C and McIlvaine WB (1991):

Techniques of regional anesthesia in children. In PP Raj (ed.), Clinical practice of regional Anesthesia, pp. 365-388. New York: Churchill Livingstone.

Epstein B, Levy ML, Thien M and Coakley C (1985):

Evaluation of fentanyl as an adjunct to thiopental-nitrous oxide-oxygen anesthesia for short procedures. Anesthesiol. Rev. 2: 24-29.

Feld LH, Negus JB and White PF (1990):

Oral midazolam preanesthetic medication in pediatric outpatients. Anesthesiology 73: 831-834.

Ferrari LR and Donlon JV (1992):

Metoclopramide reduce the incidence of vomiting after tonsillectomy in children. Anesth. Analg. 75: 351-354.

Fischer QA, McComiskey Cm, Hill JL et al. (1993):

Postoperative voiding interval and duration of analgesia following peripheral or caudal nerve blocks in children. Anesth. Analg. 76: 173-177.

Fontaine P, Dittberner D & Scheltema K (1994):

The safety of dorsal penile nerve block for neonatal circumcision. J. Fam. Pract. 39: 243-248.

Friesen RH and Lockhart CH (1992):

Oral transmucosal fentanyl citrate for preanesthetic medication of pediatric day surgery patients with or without droperidol as a prophylactic antileptics. Anesthesiology 76: 46-51.

Friesen RH and Morrison JE (1994):

The role of ketamine in the current practice of pediatric anesthesia. Pediatr. Anaesth. 4: 79-82.

Friedman WF, Hirschklau MJ, Printz MP et al., (1976):

Pharmacologic closure of patient ductus arteriosus in the premature infant. N. Engl. J. Med 295: 526.

- Greenspun J, Hannallah R, Welborn L and Norden J (1995):**
Comparison of sevoflurane and halothane anesthesia in children undergoing outpatient ENT surgery.
J. Clin. Anesth. 7: 398-402.
- Gunter JB, Dunn CM, Bennie JB, et al., (1982):**
Concentration of bupivacaine for combined caudal general anesthesia in pediatric patients.
Anesth analg. 66: 995-998.
- Gustein HB, Johnson KL, Heard MB and Gregory GA (1992):**
Oral ketamine preanesthetic medication in children.
Anesthesiology 76: 28-33.
- Hannallah RS; Broadman LM, Belman AB et al. (1987):**
Comparison of caudal and ilioinguinal/iliohypogastric nerve blocks for control of post-orchidopexy pain in pediatric ambulatory surgery. Anesthesiology 66: 832-834.
- Hannallah, RS, and Epstein, B.S. (1991):**
The pediatric patient. In: Anesthesia for ambulatory surgery, BV Wetchler (Ed), Philadelphia, JB Lippincott, 39: 131.
- Hannallah RS, Baker SB, Casey W et al., (1991):**
Propofol effective dose and induction characteristics in unpremedicated children. Anesthesiology 47: 217-219.
- Hannallah RS and Epstein BS (1994):**
Outpatient anesthesia in GA Gregory (ed.) Pediatric anesthesia, pp. 773-804. New York: Churchill Livingstone.
- Hannallah RS and Patel RI (1995):**
Pediatric considerations. In RS Twersky (ed.), the ambulatory anesthesia handbook, pp. 145-170. St Louis: CV Mosby.

Hatch DJ, Hunter JM (1999):

The paediatric patient Br. J. Anaesthesia; 83: 1.

Henretig FM, Selbst SM, Forrest C et al., (1989):

Repeated acetaminophen overdosing. Causing hepatotoxicity in children. Clin. Ped 28: 525-528.

Hollingworth P. (1993):

The use of non-steroidal anti-inflammatory drugs in pediatric rheumatic disease. Br. J. Rheum. 32: 73-77.

Hopkins CS, Underhill S and Booker PD (1990):

Pharmacokinetics of paracetamol after cardiac surgery. Arch. Dis. In child. 65: 971-976.

Hvorka J, Kortilla and Erkola O (1990):

Gastric aspiration at the end of anesthesia does not decrease postoperative nausea and vomiting. Anaesth. Intens. Care. 18: 58-61.

Jeffery L. Apfelbaum. (2001):

Current controversies in adult outpatient anesthesia: In ASA; 3: 151.

Juggins H and McQuade B (1992):

Ondansetron as prophylaxis for chemotherapy and radiotherapy induced emesis in children Oncology 49: 279-285.

Kain, ZN, Mayes, LC, Caramico, LA Hofstader, MB (1998):

Parental presence during induction of anesthesia versus sedative premedication: which intervention is more efficient? Anesthesiology. 89: 1147-1156.

Kaplan R, Garcia M and Hannallah R (1995):

Mivacurium induced neuromuscular blockade during sevoflurane and halothane anesthesia in children. Can. J. Anaesth. 42: 16-20.

Karl HW, Keifer AT, Rosenberg JL, et al., (1992):

Comparison of safety and efficacy of itransal midazolam or sufentanil for preinduction of anesthesia in pediatric patients. *Anesthesiology* 76: 209-215.

Kelly RE, Hardman GS, Embree PB et al., (1993):

Inhaled induction and emergence from desflurane anesthesia in ambulatory surgical patient: the effect of premedication. *Anesth. Analg.* 77: 540-543.

Kirkpatrick SE, Pitlick PT, Nablioff J et al., (1976):

Frank-Starling relationship as an important determinant of fetal cardiac output. *Am J pPhysiol* 231: 495.

Korttila K, Nuotto EJ, Lichtor JL, et al., (1992):

Clinical recovery and psychomotor function after brief anesthesia with propofol or thiopental. *Anesthesiology*, 76: 676-681.

Krishnan A, Tolhurst CL and Kay B (1985):

Controlled comparison of nalbuphine and morphine for post tonsillectomy pain. *Anaesthesia* 40: 1178-1181.

Larsson S, Asgeirsson B and Magnusson J (1992):

Propofol-Fentanyl anesthesia compared to thiopental-halothane with special reference to recovery and vomiting after pediatric strabismus surgery. *Acta. Anesthesiol. Scan.* 36: 182-186.

Lee JJ and Rubin AP (1994):

Comparison of bupivacaine clonidine mixture with plain bupivacaine for caudal analgesia in children. *Br. J. Anaesth.* 72: 258-262.

Lerman J, Sikich N, Kleinman S and Yentis S (1994):

The pharmacology of sevoflurane in infants and children. *Anesthesiology* 80: 814-824.

Letty, M.P. Liu, Phuip, L. Liu (1992):

Monitoring in pediatric anesthesia.

Seminars in anesthesia, Vol. XI, No. 3, pp. 243-251.

Lindeman KS (1995):

Anesthesia, airways, and asthma. Semin in Anesth 1995;
14: 221-5.

Litman RS, Perkins FM and Dawson SC (1993):

Parental knowledge and attitude toward discussing the
risk of death from anesthesia. Anesth. Analg. 77: 256-
260.

Litman RS, Wu CL and Catanzaro RA (1994):

Ondansetron decreases emesis after tonsillectomy in
children. Anesth. Analg. 78: 478-481.

Lynch, J. (1993):

Free-standing surgery centers, the wave of the future.
Anesthesiology, 18: 42.

Lysell L and Anzen B (1992):

Pain control after third molar surgery, a comparative
study of ibuprofen and a paracetamol/codeine
combination. Swed. Dent. J. 16: 151-160.

Marshall SI, et al., (1999):

Discharge criteria and complication after ambulatory
surgery. Anesth Analg. 88: 508-17.

Maunuksela EL, Kokki H and Bullingham RES (1992):

Comparison of intravenous ketolac with morphine for
postoperative pain in children. Clin. Pharmacol. Ther.
52: 436-443.

Maxwell LG, Martin LD and Yaster M (1994):

Bupivacaine induced cardiac toxicity in neonates:
successful treatment with intravenous phenytoin.
Anesthesiology 80: 682-686.

Mc Cluskey A and Meakin GH (1994):

Oral administration of midazolam as a premedicant for pediatric day case anesthesia. *Anaesthesia* 49: 782-785.

Mc Graw T (1994):

Preparing children for the operating room: Psychological issues. *J. Anaesth.* 41: 1094-1103.

Miguel R (1994):

Eutectic mixture of local anesthetics: not just for kids. *Reg. Anesth* 19: 75-76.

Miller BR and Friesen RH (1988):

Oral atropine in infants attenuates cardiovascular depression during halothane anesthesia. *Anesth. Analg.* 67: 180-185.

Montgomery CJ, McCormack TP, Reichert CC and Marsland CP (1995):

Plasma concentrations after high-dose (45mg/kg) rectal acetaminophen in children. *Can. J. Anesth.* 42: 982-986.

Moore MA, Wandles JG and Fell D (1990):

Pediatric postoperative analgesia: A comparison of rectal diclofenac with caudal bupivacaine after inguinal herniotomy. *Anaesthesia* 45: 156-158.

Moote C (1992):

Efficacy of non steroidal anti-inflammatory drugs in the management of postoperative pain. *Drugs* 44 (Suppl. 5): 14-30.

Murray JP, Geiduschek JM, Caplan RA, et al., (1993):

A comparison of pediatric and adult anesthesia closed malpractice claims. *Anesthesiology*; 78: 461-467.

Murray JP, Geiduschek JM, Ramamoorthy MB, et al., (2000):

Anesthesia-related cardiac arrest in children. *Anesthesiology*; 93: 6-14.

Munro HM, Riegger LQ, Reynolds PI et al., (1994):

Comparison of the analgesic and emetic properties of ketorolac and morphine for pediatric out strabismus surgery. Br. J. Anaesth. 72: 624-628.

Nelson PS, Streisand JB, Mulder SM, Pace NL and Stanley TH (1989):

Comparison of oral transmucosal fentanyl citrate and an oral solution of meperidine, diazepam, and atropine for premedication in children. Anesthesiology 70: 616-612.

Ostman, P.L. and White, P.F. (1994):

Outpatient anesthesia In: Anesthesia; RD Miller (Ed) New York, Churchill livingstone, 32: 22/3.

Ostman, P.L. and White, P.F. (2000):

Outpatient anesthesia. In: Anesthesia; RD Miller (Ed.) New York, Churchill Livingstone, 32: 2260.

Pasternak, D.P. Smith, H.L. and Piland, N.F. (1996):

Critical issues surrounding the evolution of ambulatory surgery. J. Amb Care Manage., 9: 171.

Pavlin DJ, et al., (1998):

Factors affecting discharge time in adult outpatients. Anesth Analg 87: 816-26.

Penna A and Buchanan N (1991):

Paracetamol poisoning in children and hepatotoxicity. Br. J. Clin. Pharmacol. 32: 143-149.

Phillip Scuderi and Winston Salem. (2001):

Management of post operative Nausea and vomiting. In ASA., 6 : 223.

Pollard JB, Zboray AL, Mazze RI (1996):

Economic benefits attributed to opening a preoperative evaluation clinic for outpatients. Anesth. Analg. 83: 407-410.

Pullerits J and Hoizman RS (1993):

Pediatric neuro-axial blockade. Clin. Anesth. 5: 342-354.

Quinding H, Olsson GL, Boreus ET AL. (1992):

Infants and young children metabolise codeine to morphine a study after single and repeated rectal administration. Br. J. Clin. Pharmacol., 33: 45-49.

Raafat, S.H. (1992)

Ambulatory anesthesia in children. Seminars in anesthesia, 4: 303-308.

Rice LJ (1996):

Regional anesthesia and analgesia. In E Motoyama & P Davis (ed.), Smith's Anesthesia for Infants and Children, 6th edn, pp. 403-442. St Louis: CV Mosby (in Press).

Romangoli A and Keats AS (1980):

Ceiling effect for respiratory depression by nalbuphine. Pharmacol. Ther. 27: 478-485.

Romero T, Covell J. Friedman WF (1972):

A comparison of pressure-volume relations of the fetal, newborn, and adult heart. AM. J. Physiol 222: 495.

Rudolph AM (1970):

The changes in the circulation after birth: their importance in congenital heart disease. Circulation 41: 343.

Ryhanen R, Adamski J and Puhakka K (1994):

Postoperative pain relief in children, a comparison between caudal bupivacaine and intramuscular Diclofenac sodium Anaesthesia 49: 57-61.

Saint-Maurice, C.; Kong-ky, B.H. and Mathey, S. (1995):

Pediatric surgery and anesthesia in day-surgery hospital. Chaier's d'Anesth., 41: 407.

Sarner JB, Levine M, Davis PJ, et al. (1995):

Clinical characteristics of sevoflurane in children – a comparison with halothane, *Anesthesiology* 83: 38-46.

Schachtel BP and Thoden WR (1993):

Clinical trials and therapeutics. A placebo – controlled model for assay in systemic analgesic in children. *Clin. Pharmacol. Ther* 53: 593-601.

Scher C., Arnar D, Ginsburg I et al. (1992):

Post dural puncture headache in children with cancer. *Anesthesiology* 77: A 1183.

Schrum SF, Hannallah RS, Verghese PM et al. (1994):

Comparison of propofol and thiopental for rapid anesthesia induction in infants. *Anesth. Analg.* 78: 482-485.

Song D, et al. (1997):

Titration of volatile anesthetics using bispectral index facilitates recovery after ambulatory anesthesia. *Anesthesiology* 87: 842-8.

Sonr D, et al. (1998):

Fast-Tracking eligibility after ambulatory nesthesia: a comparison of desflurane, sevolfurane, and propofol. *Anesth Analg* 86:267-73.

Sossai R, Jar M, Kistler W et al. (1993):

Postoperative vomiting in children. A persisting unsolved problem. *Eur. J. Pediatr. Surg.* 3: 206-208.

Souter A, Fredman B and White PF (1994):

Controversies in the perioperative – use of non-steroidal anti-inflammatory drugs. *Anesths. Analg.* 79: 1178-1190.

Southhall DP, Richards, JM, Johnston PGB, Shinebourne EA (1979):

Study of cardiac rhythm in healthy newborn infants. Br. Heart J; 41: 382.

Spear RM, Vaster M, Berkowitz ID et al. (1991):

Preinduction of anesthesia in children with rectally administered midazolam. Anesthesiology 47: 670-674.

Sullivan M, Thompson WK and Hill CD (1994):

Succinylcholine-induced cardiac arrest in children with undiagnosed myopathy. Can. J. Anaesth. 41: 497-501.

Sureh S, Cohen I, Maluszczak M et al., (1998):

Dos ranging safety, and efficacy of a new oral midazolam syrup in children. Anesthesiology

Tait A, Pandit U, Voepel-Lewis T, et al., (1998):

Use of the laryngeal mask airway in children with upper respiratory tract infection-a comparison with endotracheal intubation. Anesth Analg; 86: 706-11.

Thwaites A, Edmonds S, Smith I. (1997):

Inhalation induction with sevoflurane-A double-blind Anesthesia, 78: 356-361.

Twersky RS (1994):

The post-anesthesia care unit and ambulatory care. Balliere's Clin Anesth 8 (4); 873-96.

Vaughn, R.W., Aluise, J.J. and McLaughlin, C.P. (1991):

Ambulatory surgery and the hospital. Health Care Manage Rev., 16: 15.

Warner M.A. Caplan RA., Epstein BS, et al., (1999):

Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: Application to healthy patients undergoing elective procedures. Anesthesiology; 90: 896-905.

Watcha MF, Simeon RM, White PF et al., (1991):

Effect of propofol on the incidence of postoperative vomiting after strabismus surgery in pediatric outpatients. *Anesthesiology* 75: 204-209.

Watcha MF, Jones MB, Lagueruda RG, Schweiger C and White PF (1992):

Comparison of ketolac and morphine as adjuvants during pediatric surgery. *Anesthesiology* 79: 368-372.

Watcha, M.F.; Safavi, F.Z.; McCulloch, D.A. Tan, T.S.H. and White, P.F. (1995):

Effect of antagonism of mivacurium-induced neuromuscular block on postoperative emesis in children. *Anesth. Analg.*, 37: 713.

Watcha MF, Ramirez-Ruiz M, White PF et al., (1992):

Perioperative effects of oral ketolac and acetaminophen in children undergoing bilateral myringotomy. *Can. J. Anaesth.* 39: 649-654.

Watcha MF and White PF (1992):

Postoperative nausea and vomiting: its etiology, treatment, and prevention. *Anesthesiology* 77: 162-184.

Weinstein MS, Nicolson SC and Schreiner MS (1994):

A single dose of morphine sulphate increases the incidence of vomiting after outpatient inguinal surgery in children *Anesthesiology* 81: 572-577.

White PF, et al., (1999):

New criteria for fast tracking after outpatient anesthesia: a comparison with the modified Alderete's scoring system. *Anesth. Analg.* 88: 1069-72.

Wolfson, J.; Walker, G. and Levin, P.J. (1993):

Freestanding ambulatory surgery. Cost containment winner?

Health Financial Manage, 6: 27.

Yapster M and Maxwell L (1993):

Opioid agonists and antagonists. In NL Schechter, CB Berde & M Yaster (eds), Pain in Infants, children and Adolescents, pp. 145-172.

Zuurmond, W.W.A.; Van Leeuwen, L., Helmers, J.H.J.H. (1989):

Recovery from fixed dose midazolam induced anesthesia and antagonism with flumazenil for outpatient arthroscopy.

Acta. Anesthesiol. Scand., 33: 160.