

REFERENCES

REFERENCES

A

- Abou-Farha, K.M.; Menheere, P.F.; Bruins, J.L.; Kester, A.D. and Janknegt, F. (1992):** The role of serum laminin P₁ in the diagnosis of transitional cell carcinoma of the bladder. *Eur. Urol.*, 21(3):240.
- Adelmann, T. and Grill, B. (1982):** Collagen degradation. In: *Connective tissue of the normal and fibroic human liver*. Gelach; V.; Pott, G. and Rautterberg, J. (Eds.), p. 18.
- Akiyama, S.K. and Yamada, K.M. (1983):** FN in diseases. In: *Connective tissue disease*. Edited by Wagner, B.M.; Fleishmajer, R. and Kafman, N. Published by Williams and Wilkins, P. 55.
- Alcolado, J.C. and Alcolado, R. (1991):** Importance of maternal history of non-insulin dependent diabetes mellitus patients. *B.M.J.*, 302:1178.
- Ali, T.V. and Hynes, R.O. (1978):** Effects of LETS glycoprotein on cell motility. *Cell*, 4:439.
- Amenta, P.S.; Clark, C.C. and Martinez, H.A. (1983):** Deposition of fibronectin and laminin in the basement membrane of rats parietal yolk sac: immunohistochemical and biosynthetic studies. *J. Cell Biol.*, 96:104.
- American College of Physicians, American Diabetes Associations and American Academy of Ophthalmology (1992):** Screening guide lines for diabetic retinopathy. *Ann. of Int. Med.*, 116(8):683.
- Amrani, D.L.; Falk, K.J. and Mosesson, H.W. (1985):** Studies of FN synthesized by cultured chick hepatocytes. *Exp. Cell. Res.*, 160:171.

- Andersen, A.R.; Christainsen, T.S. and Andersen, J.K. (1983):** Diabetic nephropathy in type I diabetes. An epidemiological study. *Diabetologia*, 25:496.
- Anderson, D.H.; Stern, W.H.; Fisher, S.K.; Erickson, P.A. and Borgula, G.A. (1983):** Retinal detachment in the cat: the PE photoreceptor interface. *Invest. Ophthalmol. Vis. Sci.*, 24:906.
- Andreassen, T.; Seyer-Hansen, K. and Bailey, A.J. (1981):** Thermal stability, mechanical properties and reducible cross-links of rat tail tendon in experimental diabetes. *Biochim. Biophys. Acta.*, 677:313.
- Anttinen, T. (1977):** Collagen hydroxylsilyl glycosyl transferases. *Acta. Univ. Oul. A*:51, *Biochem.*, 13:1.
- Aumailley, M.; Nurcombe, V.; Edgar, D.; Paulsson, M. and Timpl, R. (1987):** The cellular interaction of laminin fragments cell adhesion correlates with two fragment specific high affinity-binding sites. *J. Biol. Chem.*, 262:11532.
- Aumailley, M.; Wiedemann, H.; Mann, K. and Timpl, R. (1989):** Binding of nidogen and the laminin-nidogen complex to basement membrane collagen type IV. *Eur. J. Biochem.*, 184:241.
- Aristidis, S.; Choroni, T.; Effie, C. and Tsilibary, D. (1992):** Structural and functional changes of laminin and type IV collagen after non-enzymatic glycation. *Diabetes*, 41 (Suppl. 2): 49.
- Ashton, N. (1963):** Studies of the retinal capillaries in relation to diabetic and other retinopathies. *Br. J. Ophth.*, 47:521.
- Ayo, S.H.; Radnik, R.A.; Garoni, J.A.; Glass, W.F. and Kreisberge, J.I. (1990):** High glucose causes an increase in extracellular matrix protein in cultured mesangial cells. *Am. J. Pathol.*, 136:1339.

Ayo, S.H.; Radnik, R.A.; Glass, W.F.; Garoni, J.A.; Rampt, E.R.; Appling, D.R. and Kreisberg, J.I. (1991): Increased extracellular matrix synthesis and mRNA in mesangial cells grown in high glucose medium. *Am. J. Physiol.*, 260:185.

B

Baker, J.R.; Metcalf, P.A.; Holdaway, I.M. and Johnson, R.N. (1985): Serum fructosamine concentration as a measure of blood glucose control in type I diabetes mellitus. *Br. Med. J.*, 290:352.

Banatvala, J.E.; Bryart, J.; Borkenstein, M.; Brown, D.; Menser, M.A.; Desilva, L.M. and Silink, M. (1985): Coxsackie B, mumps, rubella and cytomegalovirus specific IgM responses in patients with juvenile onset IDDM in Britain, Australia and Austria. *Lancet*, June, 1409.

Bar, T. and Wolff, J.R. (1972): The formation of capillary basement membrane during internal vascularization of the rat's cerebral cortex. *Z. Zellforsch. Mikrosk. Anat.*, 133:231.

Beck, K.; Hunter, I. and Engel, J. (1990): Structure and function of lam.: anatomy of a multidomain glycoprotein. *FASEB J.*, 4:148.

Beisswenger, P.d.; Moore, L.L.; Brinck, J. and Curphey, J.T. (1993): Increased collagen linked pentosidine levels and advanced glycosylation end-products in early diabetic nephropathy. *J. Clin. Invest.*, 92(1):212.

Bender, B.L.; Jaffe, R. and Chung, A.E. (1981): Immunolocalization of entactin, a sulfated basement membrane component in rodent tissues and comparison with GP-2 (laminin). *Am. J. Patholog.*, 103:419.

- Benson, W.E.; Brown, G.C. and Tasman, W. (1988):** Diabetic retinopathy: clinical findings, from diabetes and its vascular complications. Saunders, W.B. Company, Harcourt Brace Javonovich, Inc.
- Bensusan, H.; Koh, T. and Henry, G. (1978):** Evidence that FN is the collagen receptor on the platelets membrane. *Proc. Natl. Acad. Sci., (USA)*, 75:5864.
- Bohlen, H.G. and Hankins, K.d. (1983):** Early microvascular pathology during hyperglycemia in rats. *Blood Vessels.*, 20:213.
- Bohnsack, J.; Ashea, J.; Takahashi, T. and Brown, E. (1985):** FN enhanced phagocytosis of an alternative pathway activator by human culture-derived macrophage is mediated by C₄b/C₃b complement receptor. *J. Immunol. Oct.*, 135(4):2680.
- Borch, I.; Johnsen, K.; Andersen, P.K. and Deckert, T. (1985):** The effect of proteinuria on relative mortality in type I diabetes mellitus. *Diabetologia*, 28:590.
- Bornstein, P. and Traub, W. (1979):** The proteins. Hans, N.; Robert, L.H. and Carol-Leigh, B. (Eds.), Vol. IV, Academic Press, New York, San Francisco, London, p. 412.
- Bosch, M. and Ponce, S. (1986):** Fructosamine calorimetric determination. *Analisis Clinicos*, 11:124.
- Bowersox, J.C. and Sorgento, N. (1982):** Chemotaxis of aortic endothelial cells in response to fibronectin. *Cancer Res.*, 42:5247.
- Brocks, D.G.; Stecker, H.; Neubauer, H.P. and Timpl R. (1986):** Radioimmunoassay of laminin in serum and its application to cancer patients. *Clin. Chem.*, 32:787.

- Brownlee, M. and Cerami, A. (1981):** The biochemistry of the complications of diabetes mellitus. *Ann. Rev. Biochem.*, 50:385.
- Brownlee, M.; Vlassara, H. and Cerami, A. (1984):** Non-enzymatic glycosylation and the pathogenesis of diabetic complications. *Ann. Int. Med.*, 101:527.
- Brownlee, M.; Vlassara, H.; Kooney, A.; Ulrich, P. and Cerami, A. (1986):** Aminoguanidine prevents diabetes induced arterial wall protein cross-linking. *Science*, 232:1629.
- Brownlee, M.; Cerami, A. and Vlassara, H. (1988):** Advanced glycosylation end products in tissue and the biochemical basis of diabetic complication. *N. Eng. J. Med.*, 318:1315.
- Bruch, M.; Landwehr R. and Engel, J. (1989):** Dissection of laminin by cathepsin G into its long arm and short arm structures and localization of regions involved in calcium dependent stabilization and self-association. *Eur. J. Biochem.*, 185, 271.
- Brumfeld, V. and Werber M.M. (1993):** Studies on fibronectin (FN) and its domains. Secondary structure and spatial configuration of FN and of its domains. *Arch. Biochem. Biophys.*, 302(1):134.
- Burditt, A.F.; Gaird, F.L. and Draper, G. (1968):** The natural history of diabetic retinopathy. *J. Ophth.*, 7:732.
- Burns, G.F.; Cosgrove, L.; Triglia, T.; Beall, J.A.; Lopez, A.F.; Werkmeister, J.A.; Begley, C.G.; Haddad, F.P., D'apice, A.J.F. and Vadas, M.A. (1986):** the IIb-IIIa glycoprotein complex that mediates platelet aggregation is directly implicated in leukocyte adhesion. *Cell.*, 45:269.



- Cagliero, E.; Roth, T.; Roy, S. and Lorenzi, M. (1991):** Characteristics and mechanisms of high-glucose induced over-expression of BM components in cultured human endothelial cells. *Diabetes*, 40:102.
- Cahill, G.F., and McDevitt, H.O. (1981):** Insulin dependent diabetes mellitus, the initial lesion. *N. Eng. J. Med.*, 304:1454.
- Campochoiro, P.A.; Jerdan, J.A.; Glasser, B.M.; Cardin, A. and Michel, R.G. (1985):** Vitreous aspirates from patients with proliferative vitreoretinopathy stimulate retinal pigment epithelial cell migration. *Arch. Ophthalmol.*, 103:1403.
- Campochoiaro, P.A. and Hackett, S.F. (1993):** Corneal endothelial cell matrix promotes expression of differentiated features of RPE cells: implication of laminin and basic fibroblast growth factor as active components. *Exp. Eye. Res.*, 57:539.
- Casaroli-Narano, P.R.; Preissner, K.T. and Vilaro, S. (1995):** Fibronectin, laminin, vitronectin and their receptors as newly-formed capillaries in proliferative diabetic retinopathy. *Exp. Eye. Res. Jan.*, 60(1):5.
- Caulfield, J.P. and Farquhar, M.G. (1974):** The permeability of glomerular capillaries to graded dextrans. *J. Cell Biol.*, 63:883.
- Cerami, A.; Steven, V.J. and Monnier, V.M. (1979):** Role of non-enzymatic glycosylation in the development of the sequelae of diabetes mellitus. *Metabolism*, 28:431.

- Charonis, A.S.; Tsilibary, E.G.; Yurchenco, P. and Furthmaer, H. (1985):** Binding of laminin to type IV collagen: a morphological study. *J. Cell. Biol.*, 100:1848.
- Charonis, A.S. and Tsilibary, E.C. (1992):** Structural and functional changes of laminin and type IV collagene after non-enzymatic glycation. *Diabetes*, 41 (Suppl.), 49.
- Cheah, K.S. (1985):** Collagen genes and inherited connective tissue disease. *Biochem. J.*, 229:287.
- Checovich, W.J. and Mosher, D.F. (1993):** Lysophosphatidic acid enhances fibronectin binding to adherent cells. *Arterioscler. Thromb.*, 13(11):1662.
- Chen, A.B.; Mossson, M.W.; and Solish, G.I. (1976):** Identification of the cold-insoluble globulin of plasma in amniotic fluid. *Am. J. Obstet. Gynecol.*, 125:958.
- Chen, A.B.; Maitland, N.; Gallimore, P.H. and McDougall, J.K. (1977):** Detection of large external transformation sensitive protein on some epithelial cells. *Exp. Cell. Res.*, 106:39.
- Cheng, C.Y., Leggelt, C.G. and Reese, A.A. (1988):** Role of prostaglandins in controlling plasma fibronectin levels. *Proc. Soc. Exp. Biol. Med. Sep.*; 188(4):444.
- Chung, A.E.; Jaffe, R.; Freeman, I.L.; Vergns, J.P. and Braginsky, J.E. and Carlin, B. (1979):** Properties of a BM related glycoprotein synthesized in culture by a mouse embryonal carcinoma derived cell line. *Cell.*, 16:277.
- Clark, R.A.F.; Qinn, J.H. and Winn H.J., Lanigan J.M.; Dellepella, P. and Colvin, P.B. (1982):** FN is produced by blood vessels in responses to injury. *J. Exp. Med.*, (156):646.

- Clements, R.S. (1982):** The role of abnormal polyol metabolism in diabetic complications. In: *Diabetes mellitus and obesity*. Brodof, B.N. and Bleicher, S.. (Eds.), Williams and Wilkins, Baltimore, p. 117.
- Cliff, W.F. (1965):** Kinetics of wound healing in rabbit ear chambers, a time lapse cinemicroscopic study. *J. Exp. Physiol.*, 50:78.
- Cogan, D.g.; Toussaint, D. and Kuwabara, T. (1961):** Retinal vascular patterns: diabetic retinopathy. *Arch. Opth.*, 66:366.
- Cohen, M.P.; Saini, R.; Klepser, H. and Vasanthi, L.G. (1987):** Fibronectin binding to glomerular basement membrane is altered in diabetes. *Diabetes*, 36:758.
- Collazos, J.; Diaz, F. and Genolla, J. (1993):** Serum concentrations of laminin in cirrhosis of the liver. *Gut.*, 34(7):974.
- Colvin, R.B. and Dvorak, A.F. (1975):** Role of the clotting system in cell-mediated hypersensitivity: Kinetics of fibrinogen accumulation and vascular permeability changes in tuberculin and cutaneous basophil hypersensitivity reaction. *J. Immunol.*, 114:377.
- Cooper, A.R.; Kurkinen, M.; Taylor, A. and Hogan, B.L.M. (1981):** Studies on the biosynthesis of laminin by murine parietal endoderm cells. *Eur. J. Biochem.*, 119:189.
- Critchley, D.R.; England, M.A.; Wakely, J. and Hynes, R.O. (1979):** Distribution of FN in the ectoplasm of gastrulating chick embryos. *Nature*, 280:498.
- Cudworth, A.G.; Bodansky, H.J. and West, K.M. (1982):** Genetic and metabolic factors in relationship to the prevalence and severity of diabetic complications in: *Complications of diabetes*. Keen, H. and Jarrett, D. (Eds.), 2nd ed., Edward Arnold Publishers, London, p. 1.

D

- Danne, T.; Spiro, M.J. and Spiro, R.G. (1993):** Effect of high glucose on type IV collagen in cultured rat glomerular epithelial endothelial and mesangial cells. *Diabetes*, 42:170.
- Darwin, J.; Prockop, L.; Kivirikko, L. and Norbotro, S. (1979):** The biosynthesis of collagen and its disorders. *JAMA*, 301(1):13.
- Das, A.; Frank, R.N.; Zhang, N. L. and Samadani, E. (1990):** Increases in collagen type IV and Laminin in galactose induced retinal capillary basement membrane thickening prevention by an aldose reductase inhibitor. *Exp. Eye Res.*, 50(3):269.
- Dauhs, T.g. (1975):** Decreased corneal sensitivity among diabetic patients. *Am. J. Ophth.*, 91:174.
- David, R.S. and Vincent, M.M. (1990):** End-stage renal disease and diabetes catalyze, the formation of a pentoze-derived crosslink from aging human collagen. *J. Clin. Invest.*, 85:380.
- Davidson, J.M.; McEneamy, L.S. and Bornstein, P. (1975):** Intermediates in the limited proteolytic conversion of procollagen to collagne. *Biochemistry*, 14:5188.
- Davis, B.H. and Madri, J.A. (1987):** An immunohistochemical and serum ELISA study of type I and III procollagen aminopropeptides in primary biliary cirrhosis. *Am. J. Pathol.*, 128:265.
- Davis, M.; Rayman, G. and Day, J. (1992):** Increased incidence of coronary disease in people with impaired glucose tolerance: link with increased lipoprotein (a) concentration. *B.M.J.*, 304:1610.

- Davis, R.; Critchley, S. and Margorie, A. (1986):** Role of FN in cell migration. *Nature*, 4(11):60.
- Davis, T.M.E.; Moore, J.C. and Turner, R.c. (1985):** Plasma fibronectin, factor VIII-related antigen and fibrinogen concentrations and diabetic retinopathy. *Diabete & Metabolisme (Paris)*, 11:147.
- Day, J.F.; Thorpe, S.R. and Baynes, J.W. (1979):** Non-enzymatic glycosylation of albumin: In vitro preparation and isolation from normal human serum. *J. Biochem. Chem.*, 254:595.
- Day, J.F.; Ingebretsen, G.; Ingebretsen, V.R.; Baynas, J.W. and Thorpe, S.R. (1980):** Non-enzymatic glycosylation of serum proteins and Hb: response to changes in blood glucose level in diabetic rats. *Diabetes*, 29:524.
- Dean, B.M.; Backer, F. and Mc-Nally, J.M. (1986):** Insulin autoantibodies in the prediabetic period correlation with islet cells antibodies and development of diabetes. *Diabetologia*, 29:339.
- Deane, F. and Mosher, M.d. (1984):** Physiology of fibronectin. *An. Rev. Med.*, 35:561.
- Deckert, T.; Poulsen, J. and Larsen, M. (1978):** Prognosis of diabetics with diabetes onset before the age of 30 on factors influencing the prognosis. *Diabetologia*, 14:371.
- DeFronzo, R.A. and Ferrannimi, E. (1983):** New concept in the pathogenesis and treatment of non-insulin dependent diabetes mellitus. *Diabetes*, 33:950.
- De-Giorgio, L.A.; Seghieri, G.; Gironi, A.; Mammini, P.; Bartoli, V. and Bartolome, G. (1984):** Raised plasma fibronectin concentration is related to the prevence of diabetic retinopathy. *Acta Diabetol. Lat. Jul-Sep.*: 21(3):251.

- De-Giorgio, L.A.; Bartolome, G.; Gironi, A.; Caselli, P. and Seghieri, A. (1988):** Increased plasma FN concentration in diabetic patients with micro-albuminuria. *Diabetes Care*: Jul-Aug., 11(7):527.
- Dehm, P.; Kejalides, N.A. and Lian, G. (1978):** The collagenous component of lens basement membrane. The isolation and characterization of an α -chain size C collagenous peptide and its relationship to newly synthesized lens components. *J. Bio. Chem.*, 253:6680.
- De-Jard, A.; Anderson, T. and Gluud, C. (1986):** The influence of insulin on the raised plasma FN concentrations in human obesity. *Acta. Med. Scand.*, 220(3):269.
- Deno, D.C.; Saba, T.M. and Lewis, E.P. (1983):** Kinetics of endogenously labelled plasma FN: incorporation into tissues. *Am. J. Physiol.*, 15:345.
- Deschampes, (1980):** HLA genotype studies in juvenile insulin dependent diabetes. *Diabetologia*, 19:189.
- Diaz, F.; Collazos, J. and Genolla, J. (1993):** Study of serum procollagen III peptide in patients with hepatic cirrhosis from a clinical point of view. *J. Clin. Invest.*, 71(5):416.
- Di-Girolamo, N.; Underwood, A.; McCluskey, P.J. and Wakefield, D. (1993):** Functional activity of plasma FN in patients with diabetes mellitus. *Diabetes*, 42, Nov.: 1606.
- Dixon, A.J.; Burns, J.; Dunnill, M.S. and Jo'D McGee (1980):** Distribution of FN in normal and diseased human kidneys. *J. Clin. Pathol.*, 33:1021.

Dominiczak, M.H.; Bell, J.; Cox, N.H.; McCruden, D.C.; Jones, S.K.; Finlay, A.Y.; Percy-Robb, I.W. and Frier, B.M. (1990): Increased collagen-linked fluorescence in skin of young patients with type I diabetes mellitus. *Diabetes Care*, 13:468.

Drury, M.I. (1986): Diabetes mellitus. Osney Mead, (Ed.),. Blackwell Scientific, London.

Drury, P.L.; Watkins, P.J. and Vibertig, G. (1980): Diabetic nephropathy. *Br. Med. J.*, 42:127.

Dunn, M.A.; Rojkind, M. and Warren, K.S. (1977): Liver collagen synthesis in murine schistosomiasis. *J. Clin. Inv.*, 59:660.



Edgar, D.; Timpl, R. and Thoenen, H. (1988): Structural requirements for the stimulation of neurite outgrowth by two variants of laminin and their inhibition by antibodies. *J. Cell. Biol.*, 106:1299.

Eknoyan, G. and Knochel, J.P. (1984): The systemic consequences of renal failure. Grune & Stratton Inc., Orlando, Fl., 1:587.

Eli, A. and Friedman, S. (1990): Diabetic renal disease in diabetes mellitus therapy and practice. McGraw Hill, 1st Ed., Ch. 24, p. 135.

Ellis, D.; Becker, D.J.; Damesen, D. and Cobes, L. (1983): Proteinuria in children with type I diabetes. Relationship to duration, metabolic control, retinal changes. *J. Ped.*, 102:673 .

Engel, J.; Bruckner, P. and Becker, U. (1977): Physical properties of the amino-terminal precursor specific protein type I procollagen. *Biochemistry*, 16:4026.

Engel, J.; Odermatt, E.; Engel, A.; Madri D. and Furthmayr, H. (1981): Shapes, domain organizations and flexibility of laminin and FN, two multifunctional proteins of the extracellular matrix. *J. Mol. Biol.*, 150:97.

Engel, J. and Furthmayr, H. (1987): Electron microscopy and other physical methods for the characterization of extracellular matrix components: Laminin, FN, Collagen IV and proteoglycans. *Methods Enzymol.*, 145:3.

Engerman, R.L. (1989): Perspectives in diabetes: Pathogenesis of diabetic retinopathy. *Diabetes*, 38:1203.

Engvall, E. and Ruoslahti, E. (1977): Binding of soluble form of fibroblast surface protein. *Int. J. Cancer*, 20:1.

Enrico-Cagliero, M.; Maiello, M.; Boeri, D.; Roy, S. and Lorenz, M. (1988): Increased expression of basement membrane components in human endothelial cells cultured in high glucose. *J. Clin. Invest.*, 82:735.

Erickson, H.P.; Carell, N.; McDonagh, J. (1981): Fibronectin molecule visualized in electron microscopy: a long thin flexible strand. *J. Cell. Biol.*, 91:673.

Everhart, J.E.; Pettit, D.J.; Benett, P.H. and Knowler, W.C. (1992): Duration of obesity increases the incidence of non-insulin dependent diabetes mellitus. *Diabetes*, 41(2):235.



Fajans, S.S. (1985): Comperimental medicine and biology. Uranic, M. (Ed.), Plenum Publishing Corp., 65.

- Fajans, S.S. (1990):** Classification and diagnosis of diabetes. In: Diabetes mellitus, theory and practice. Fourth edition, Rifkin, H. and Porte, D. (Eds.), P. 346-355. Elsevier, New York, Amsterdam, London.
- Forquhar, M.G. (1981):** The glomerular basement membrane: A selective macromolecular filter; in Cell biology of extracellular matrix. Hay, E.D. Editor, Plenum Publishing Company, New York, p. 335.
- Felden, F.; Renkes, P.; Fremont, S.; Chambre, J.F.; Champigneulle, B.; Gaucher, P. and Gueant, J.L. (1991):** Serum laminin P₁ in metastatic colon carcinoma. Clin. Chemistry, (37), 10:1795.
- Fessler, J.H. and Fessler, L.I. (1978):** Biosynthesis of procollagen. Ann. Rev. Biochem., 47:129.
- Fleischmajer, R.; Timpl, R.; Tuderhass, L.; Raisher, L.; Wiestner, M.; Perlish, T.S. and Graves, P.N. (1981):** Ultrastructural identification of extension aminopeptides of types I and II collagens in human skin. Proc. Natl. Acad. Sci., U.S.A., 78:7360.
- Foidart, J.M.; Bere, E.W.; Yaar, M.; Rennard, S.I.; Gullino, M.; Martin, G.R. and Katz, S.I. (1980):** Distribution and immunoelectron-microscopic localization of laminin, a non-collagenous basement membrane glycoprotein. Lab. Invest., 3:336.
- Forsyth, K.D., and Levinsky, R.J. (1990):** FN degradation: An in vitro model of neutrophil mediated endothelial cell damage. J. of Pathology, 161:313.
- Fraidakis, M.; Mantzavinos, T.; Dalamanga, A.; Gargaropoulos, A. and Lourlas, P.A. (1992):** Plasma and amniotic fluid concentration of FN during normal and diabetic pregnancy. Clin. Exp. Obstet. Gynecol., 19(20):28.

- Frenette, G.P.; Ruddon, R.W.; Krzesicki, R.F.; Naser, J.A. and Peters, B.P. (1989):** Biosynthesis and deposition of a non-covalent laminin-heparan sulfate proteoglycan complex and other basal lamina components by a human malignant cell line. *J. Biol. Chem.*, 264:3078.
- Friedman, J.M. and Fialkour, J.M. (1982):** Diabetes mellitus and obesity. Brodoff, B.B. and Bleicher, S.J. (eds.), Baltimore, Williams and Wilkins, P. 142.
- Friedman, R. and Gross, J.L. (1991):** Evaluation of glomerular filtration rate in proteinuric type II diabetic patients. *Diabetes, Care*, 14:355.
- Freinkel, N.; Josinovich, J. and Session, V. (1980):** Report of workshop chairman. Workshop Conference on gestational diabetes, Summary and Recommendation. *Diabetes Care*, 3:499.
- Friend, J.; Kiorpes, T.c. and Thoft, R.A. (1981):** Diabetes mellitus and the rabbit corneal epithelium. *Invest. Ophthalmol. Vis. Sci.*, 21:2, 317.
- Fukuda, M. and Hakomori, S. (1979):** Carbohydrate structure of galactoprotein A, a major transformation sensitive glycoprotein released from hamster embro fibroblasts. *J. Biol. Chem.*, 254:5451.
- Fuller, G.c. (1981):** Perspectives for the use of collagen synthesis inhibitors as antifibrotic agents. *J. Med. Chem.*, 24:651.
- Fyrand, O. (1979):** Studies on fibronectin in the skin. Indirect immunofluorescence studies in normal human skin. *Br. J. Dermatol.*, 101:263.

G

- Gahmberg, C.G.; Kiehn, D. and Hakomori, S. (1974):** Changes in a surface-labelled galactoprotein and in glycolipid concentrations in cells transformed by a temperature-sensitive polyoma virus mutant. *Nature*, 248:413.
- Gallop, P.M. and Paz, H.P. (1975):** Post-translational protein modifications with special attention to collagen and elastin. *Physiol. Rev.*, 55:418.
- Gay, S.; Martin, G.R. and Muller, R. (1979):** Simultaneous synthesis of type I and type III collagen by fibroblasts in culture. *Proc. Natl. Acad. Sci., U.S.A.*, 73:4037.
- Gay, S. and Miller, E.J. (1983):** What is collagen, what is not. *Ultrastruct. Pathol.*, 4:365.
- Gepts, W. (1965):** Pathologic anatomy of the pancreas in juvenile diabetes mellitus. *Diabetes*, 14:619.
- Gilchrest, B.A.; Rowe, J.W. and Mihm, M.C. (1980):** Clinical and histologic cutaneous findings in chronic renal failure: evidence for a dialysis-resistant transplant responsive microangiopathy. *Lancet*, 11:1271.
- Giugliano, D.; Torella, R.; Cacciapuoti, F.; Gentile, S.; Verza, M. and Varricchio, M. (1980):** Impairment of insulin secretion in man by nifedipine. *Eur. J. Clin. Pharmacol.*, 18:395.
- Goodman, S.L. and Newgreen, D. (1985):** Do cells show an inverse locomotory response to FN and laminin substrates. *EMBO J.*, 4:2769.

- Goodman, S.L. and Deutzmann, R. and Vonder-Mark H. (1987):** Two distinct cell-binding domains in laminin can independently promote non-neuronal cell adhesion and spreading. *J. Cell. Biol.*, 105:589.
- Grant, M.E. and Prockop, I.D. (1972):** The biosynthesis of collagen. *N. Engl. J. Med.*, 286:194-199, 242-249, 291.
- Grant, M.E.; Harwood, R. and Williams, I.F. (1975):** The biosynthesis of basement membrane collagen by isolated rat glomeruli. *Eur. J. Biochem.*, 54:531.
- Greene, D.A.; Littmer, S.A. and Sima, A.A. (1988):** Pathogenesis and prevention of diabetic neuropathy. *Diab. Metab. Rev.*, 4:201.
- Gressner, A.m. and Wallraff, P. (1981):** Laser nephelometric quantitation of FN in human C.S.F. and seminal plasma. *Clin. Chim Acta*, 113:207.
- Gressner, A.M.; Tittor, W. and Negwer, A. (1986):** Serum concentration of laminin and aminoterminal propeptide of type III procollagen in relation to the portal venous pressure of fibrotic liver diseases. *Clin. Chim. Acta*, 161:249.
- Grinnell, F. (1976):** The serum dependence of baby hamster kidney cell attachment to a substratum. *Exp. Cell. Res.*, 97:265.
- Grisanti, S.; Heimann, K. and Wiedmann, P. (1993):** Origin of FN in epiretinal membranes of proliferative vitreoretinopathy and proliferative diabetic retinopathy. *Br. J. Ophthalmol.*, 77(4):238.
- Grunwald, J.E.; Riva, C.E. and Sinclairs, H. (1986):** Laser doppler velocimetry study of the retinal circulation in diabetes mellitus. *Arch. Ophth.*, 104:991.

Gundersen, H.T. (1974): Small pupils with long term diabetes. *Diabetologia*, 10:366.

H

Hahn, E.; Wick, G.; Pencev, D. and Timpl, R. (1980): Distribution of basement membrane proteins in normal and fibrotic human liver: Collagen type IV, laminin and FN. *Gut* 21:63.

Haneda, M.; Rikkawa, R.; Horide, N.; Togava, M.; Koyo, D.; Majiwara, N.; Ooshima, A. and Shigeta, Y. (1991): Glucose enhances type IV collagen production in cultured rat glomerular mesangial cells. *Diabetologia*, 34:1198.

Harold, R. and Daniel, J.R. (1990): Classification of diabetes mellitus: in diabetes mellitus theory and practice, 4th edition, Elsevier Science Publishing Co., New York, p. 346.

Harris, S. and Chill, T. (1979): Impaired glucose tolerance and non-insulin dependent diabetes mellitus. *Diabetes*, 28:1039.

Hartl, L.; Oberbaumer, I. and Deutzman, R. (1988): The N-terminus of laminin A chain is homologous to the B-chains. *Eur. J. Biochem.*, 173:629.

Hasseli, J.R.; Pennypacker, J.P.; Yamada, K.M. and Pratt, R.M. (1978): Changes in cell surface proteins during normal and vitamin A inhibited chondrogenesis in vitro. *Ann. N.Y. Acad. Sci.*, 312:406.

Hasslacher, C.; Reichenbacher, R.; Gechter, F. and Timpl, R. (1984): Glomerular BM synthesis and serum concentrations of type IV collagen in streptozotocin diabetic rats. *Diabetologia*, 26:150.

- Hayashi, Y.; Makino, H. and Ota, B. (1992):** Serum and urinary concentrations of type IV collagen and laminin as a marker of microangiopathy in diabetes. *Diabet. Med.*, 9(4):366.
- Heggeness, M.H.; Ash, J.P. and Singer, S.. (1978):** Transmembrane linkage of FN to intracellular actin containing filaments in cultured fibroblasts. *Ann. Acad. Soc.*, 312:414.
- Henry, R.J. (1974):** Clinical chemistry, Principles and Techniques. Harper, A. and Row, P. (Eds.), 2nd. Ed., p. 525.
- Hess, L.; Obrien, W. and Holmberg, I. (1986):** Plasma and amniotic fluid concentration of FN during normal pregnancy. *Obstet. Gynecol.*, 68(1):25.
- Hoffman, P.R. and Hynes, R.O. (1979):** Haemostasis: the role of FN in the adhesion of platelets to collagen. *Clin. Res.*, 27:297.
- Hogan, B. (1981):** Laminin and epithelial cell attachment. *Nature*, 290:737.
- Humphrey, L.L.; Ballard, P.J. and Frohwert, P.P. (1989):** Chronic renal failure in NIDDM. *Ann. Int. Med.*, 111:788.
- Hyndink, R.A.; Karuzain, E. and Schulfzro, S. (1979):** Neuropathic and ulceration in diabetics with decreased corneal sensitivity. *Arch. Ophth.*, 95:2193.
- Hynes, R.O. (1985):** Molecular biology of FN. *Ann. Rev. Cell of Biol.*, 1:::67.
- Hynes, R.O. (1986):** Fibronectin. *J. Scientific Am. Soc.*, 254:32.

I

- Ingham, K.C.; Landwehr, R. and Engel, J. (1985):** Interaction of FN with C_{1q} and collagen: effects of ionic strength and denaturation of the collagenous component. *Eur. J. Biochem.*, 148:219.
- Inoguchi, T.; Vneda, F.; Watnabe, J. and Wagadat, T. (1986):** Plasma FN and platelet aggregation in diabetes. *Diabet. Res. Clin. Pract.*, 2(2):69.
- Ireland, J.T.; Vibert, G.G. and Watkins, P.J. (1982):** The kidney and the urinary tract. In: *Complications of diabetes*. Keen, H. and Jarrett, J. (Eds.), Edward Arnold, London, p. 137.
- Isobe, H.; Iwata, S.; Fujime, M. and Kitagawa, R. (1993):** Serm concentration of type IV collagen in urological cancer-comparison with serum concentration of laminin. *Nippon. Hinyokika. Gakkai. Zasshi*, 84(3):457.

J

- Jaffe, E.A. and Mosher, D.F. (1979):** Synthesis of FN by cultured human endothelial cells. *J. Exp. Med.*, 147:1779.
- Jakus, M. (1964):** *Ocular fine structure*. Little, Brown, Boston.
- Jeraj, K.P.; Michael, A.F. and Mower, S.M. (1983):** Glycosylated and normal human or rat albumin, don't bind to renal basement membranes of diabetic and control rats. *Diabetes*; 32:380.

Jiminez-Barbero, J.; Petersen, T.E. and Linus, M. (1992): The two polypeptide chains in FN are joined in antiparallel fashion. NMR structural characterization. *Biochemistry*, Oct., 20; 31(41):9927.


Johansson, S.; Rubin, K.; Hook, M.; Ahlgren, T. and Seljelid, R. (1979): In vitro: biosynthesis of cold insoluble globulin (FN) by mouse peritoneal macrophages. *FEBS Lett.*, 105:313.

Johnson, R.N.; Metcalf, P.A. and Baker, J.R. (1983): Fructosamine: a new approach to the estimation of serum glycosyl protein. An index of diabetic control. *Clin. Chim Acta.*, 127:87.

Jones, R.L. (1981): Hematologic alteration in diabets mellitus. *Am. J. Med.*, 70:339.

Joseph, F.; Frasio, B.W.; Thomas, D.; Rhode, W.M.; Rupp, H.P. and Leo T. (1985): Non-enzymatic glycation of FN and alteration in the molecualr association of cell matrix and basement membrane components in diabetes mellitus. *Diabetes*, May, 34:477.

Julkunen, I.; Hautanen, A. and Keski-Oja, J. (1983): Interaction of viral envelope glycoproteins with FN. *Infect. Immun.*, 40:876.



Kamegaya, K. (1982): Some aspects of collagen liver formation and degradation in hepatic fibrosis. *Acta. Pathol. Japan.*, 12:123.

Kanwar, Y.S. and Farquhar, M.G. (1979): Increased biosynthesis and processing of FN in fibroblasts from diabetic mice. *J. Cell. Biol.*, 81:137. Quoted from: *Proc. Natl. Acad. Sci., U.S.A.*, 84(7):1911.

- Katayama, M.; Kamihagi, K.; Hirai, S.; Kurome, T.; Murakami, K. and Hinof, I. (1992):** Urinary laminin fragments as a tumour marker potentially reflecting BM destruction. *Br. J. Cancer*, 65(4):509.
- Kefalides, N.A.; Cameron, J.D.; Tomichak, E.A. and Yanoff, M. (1976):** Biosynthesis of basement membrane collagen by rabbit corneal endothelium in vitro. *J. Biol. Chem.*, 251:730.
- Kefalides, N.A. (1978):** Current status of chemistry and structure of basement membranes; in Kefalides, *Biology and Chemistry of B. Ms.*, p. 215-228, Academic Press, New York.
- Kefalides, N.A.; Alper, R. and Clark, C.C. (1979):** Biochemistry and metabolism of basement membranes. *Int. Rev. Cytol.*, 61:167.
- Kefalides, N.A. (1980):** Chemistry of basement membranes: Structure and Biosynthesis; in *Advances in microcirculation*, Vol. 9, p. 295, (Karger, Basel).
- Kern, P.; Maczar, M. and Robert, L. (1979):** Increased biosynthesis and processing of FN in fibroblasts from diabetic mice. *Biochem. J.*, 182:337. Quoted from, *Proc. Natl. Acad. Sci., U.S.A.*, 84(7):1911.
- Kerr, S. (1961):** Pupil reflex is spared in diabetic III nerve paralysis. *J. Neurol. Neurosurg., Psych.*, 27:473.
- Keski-Oja, J.; Mosher, D.F. and Vaheri, A. (1976):** Cross-linking of a major fibroblast surface-associated glycoprotein (FN) catalyzed by blood coagulation factor XIII. *Cell.*, 9:29.
- Keski-Oja, J. and Yamada, K.M. (1981):** Isolation of acting binding fragment of FN. *Biochem. J.*, 193:615.

- Kjellstrom, T.J. and Malmquist, (1984):** Insulin effects on collagen and protein production in cultured human skin fibroblasts, from diabetic and non-diabetic subjects. *Horm. Metab. Res.*, 16:168.
- Klein, R.; Klein, B.E.; Moss, S.E.; Davis, M.D. and De-Mets, D.L. (1984a):** The Wisconsin Epidemiologic study of diabetic retinopathy, Prevalence and risk of diabetic retinopathy when age at diagnosis is less than 30 years. *Arch. Oph.*, 102:520.
- Klein, R.; Klein, B.E.; Moss, S.E.; Davis, M.D. and Demets, D.L. (1984b):** The Wisconsin epidemiologic study of diabetic retinopathy: Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch. Ophthalmol.*, 102:527.
- Kleinman, H.K.; Cannon, F.B.; Larie, G.W.; Hassell, J.r.; Aumailley, M.; Terranova, V.R.; Martin, G.r. and Duboisdale, M. (1985):** The biological activities of laminin. *J. Cell Biocheem.*, 27:317.
- Knee, H. and Jarret, R.G. (1979):** The WHO multinational study of vascular disease in diabetes macro-vascular disease prevalence. *Diabetes Care*, 2:187.
- Kohner, E.M. and Dollery, C.t. (1975):** Diabetic retinopathy. In: *Complications of diabetes*. Keen, H. and Jarret, M. (Eds.), Edward Arnold, London, 1.
- Kohno, T.; Sorgente, N., Goodnight, R. and Ryan, S.J. (1987):** Alterations in the distribution of FN and laminin in diabetic human eye. *Invest. Ophthalmol. Vis. Sci.*, Mar., 28(3):515.

- Kolbe, M.; Kaufman, J.L.; Friedman, J.; Dinerstein, C.; Mackenzie, J.W. and Boyd, C.D. (1990):** Changes in steady state levels of mRNAs coding for type IV collagen, Laminin and FN following capillary BM thickening in human adult onset diabetes. *Connective Tissue Research*, 25:77.
- Koteliansky, V.E.; Leytin, V.L.; Suiridov, D.D.; Repin, V.S., and Smirnov, V.N. (1981):** Human plasma FN promotes the adhesion and spreading of platelets on surfaces coated with fibrillar collagen. *FEBS Letters*, 123:59.
- Krause, W.J. and Cutts, J.H. (Eds.) (1983):** *Connective tissue in: Concise text of histology.* Williams and Wilkins, Baltimore, London.
- Kukkonen, M.; Raunio T.; Virkola, R.; Lahteenmaki K.; Makelä P.H.; Klemm, P. and Clegg Korhonen, K. (1993):** Basement membrane carbohydrate as a target for bacterial adhesion: binding of type I fimbriae of *Salmonella enterica* and *Escherichia coli* to laminin. *Mol. Microbiol.*, 7(12):229.
- Kurkinen, M. and Alitalo, K. (1979):** FN and procollagen produced by a clonal line of Schwann cells. *FEBS Lett.*, 102:64.
- Kuusela, P.; Ruoslahti, E.; Engvall, E. and Vaheri, A. (1976):** Immunological interspecies cross-reactions of fibroblast surface antigen (FN). *Immunochemistry*, 13:639.
- Kuusela, P.; Vaheri, A.; Polo, J. and Ruoslahti, E. (1978):** Demonstration of FN in human C.S.F. *J. Lab. Clin. Med.*, 92:595.
- Kwan, A.P.; Freemont, A.J. and Grant, M.E. (1986):** Immunoperoxidase localization of type X collagen in chick tibiae. *Bio. Scien. Reports*, 6:155.



- Labat-Robert, J.; Leutenegger, M.; Liopis, G.; Ricard, Y. and Derouette, J.C. (1984):** Plasma and tissue FN in diabetes. *Clin. Physiol. Biochem.*, 2(1):39.
- Labat Robert, J.; Kern, P. and Robert, L. (1991):** Modifications of the biosynthesis of type I and type III collagens and FN during diabetes and atherosclerosis. *Z. Gerontol.* Mars-Apr., 24(2):66.
- LaFleur, M.; Bealie, A.d.; Kreis, C. and Poubella, P. (1987):** FN gene expression in polymorphonuclear leukocytes accumulation of mRNA in inflammatory cells. *J. Biol. Chem.*, 262:2111.
- Lamberton, R.P.; Goodman, A.D., Kassoff, A.; Rubin, C.L., Treble, D.H.; Saba, T.M., Merimee, T.J. and Dodds, W.J. (1984):** Von Willebrand factor, FN and insulin-like growth Factors I and II in diabetic retinopathy and nephropathy. *Diabetess*, Feb., 33(2):125.
- LaPiere, C.M.; Lenaer, A. and Kohn, L.D. (1971):** Procollagen peptidase: An enzyme excising the coordination peptides of procollagen. *Proc. Natl. Acad. Sci., U.S.A.*, 68:3054.
- Laurie, G.W.; Bing, J.T.; Kleinman, H.K.; Hassell, J.R.; Aumailley, M.; Martin, G.R., Martin, G.R. and Feldman, R.J. (1986):** Localization of binding sites for laminin, heparan sulfate proteoglycan and fibronectin type IV collagen. *J. Mol. Biol.*, 189:205.
- Lauritzen, T.; Frost-Larsen, K.; Larsen, H.W. and Deckert, T. (1983):** Effect of 1 year of near-normal blood glucose level on retinopathy in insulin dependent diabetics. *Lancet*, 1:200.

- Lazarus, G.S.; Hatcher, V.. and LeViene, N. (1975):** Lysosomes and the skin. *Invest. Dermatol.*, 265:259.
- Lee, I.K.; Park, N.Y., Oh, H.K., Parh, R.W. and Jo J.s. (1994):** Plasma type IV collagen and FN concentrations in diabetic patients with microangiopathy. *J. Koream. Med. Sci.* Aug.; 9(4):341.
- Lerat, H.; Lissitzky, J.C.; Singer, J.A.; Kating, A.; Herve, P. and Charbord, P. (1993):** Role of stromal cells and macrophages in FN biosynthesis and matrix assembly in human long-term marrow cultures. *Blood*, 5:1480.
- L'Esperance, F.A. (1990):** The eye and diabetes mellitus in: *Diabetes mellitus, theory and practice*, fourth edition, Rifkin, H. and Porte, D., Elsevier, New York, Amsterdam, London, p. 661.
- Liesi, P.; Kirkwood, T. and Valieri, A. (1986):** FN is expressed by astrocyte cultures from embryonic and early postnatal rat brain. *Exp. Cell. Res.*, 163:175.
- Liesi, P.; Narvanen, A.; Soos, J.; Saviola, H. and Snounou, G. (1989):** Identification of a neurite outgrowth-promoting domain of laminin using synthetic peptides. *FEBS Lett.*, 244:141.
- Linder, E.; Vaheri, A.; Roushanti, E. and Wartiovara, J. (1975):** Distribution of fibroblast surface antigen in the developing chick embryo. *J. Exp. Med.*, 124:41.
- Lubow, M. and Mackley, T.A. (1971):** Papilledema induced by hyperglycemia in juvenile diabetes. *Arch. Ophth.*, 85:417.
- Lyons, T.J.; Thorpe, S.R. and Baynes, J.W. (1990):** Glycation and oxidation of proteins in aging diabetes. In: *Glucose metabolism, diabetes and the vascular wall*. Ruderman, N.; Brownle, M. and Williamson, J.R. (Eds.), Oxford University Press, New York.

Lyons, T.J.; Bailie, K.E.; Dyer, D.G.; Dunn, J.A. and Baynes, J.W. (1991): Decrease in skin collagen glycation with improved glycemic control in patients with insulin-dependent diabetes mellitus. *J. Clin. Invest.*, 87:1910.

M

Mac-Cafferty, M.; Lepow M. and Sabat, J. (1983): Normal FN levels as a function of age in the pediatric population. *Pediatric Res. Jun.*, 17(6):482.

Mardi, J.A.; Dreyer, B.; Pitlich, T. and Furthmahyer, H. (1980): The collagenous component of the subendothelium. Correlation of structure and function. *Lab. Invest.*, 43:303.

Mark, K. and Kuhl, V. (1985): Laminin and its receptor. *Biochim. Biophys. Acta.*, 828:147.

Martin, D.W. (1983): Contractile and structural proteins in review of biochemistry, 19th ed., Chichral Com., New York, p. 477.

Matsuda, M.; Yamanaka, T. and Matsuda, A. (1982): Distribution of fibronectin in plasma and liver disease. *Clin. Chem. Acta*, 118:191.

McKee, P.F.; Maharaj, D.P.; Garden, O.J. and Carter, D.C. (1992): FN as a prognostic indicator in portal hypertension. *H.B.P. Surg.*, 50(4):229.

McKeown, L.P.J. and Mosher, D.F. (1983): Binding plasma FN to cell laers of human skin fibroblast. *J. Cell. Biol.*, 97:466.

Mc-Millan, D.E.; Utterback, N.G. and Lapuma, J. (1978): Reduced erythrocyte deformability in diabetes. *Diabetes*, 27:895.

- Mecham, R.P.; Hineck, A.; Griffin, G.L.; Senior, R.M. and Liotta, L.A. (1989):** The elastin receptor shows structural and functional similarities to the 67 KDa tumor cell laminin receptor. *J. Biol. Chem.*, 264:16652.
- Medina, J.L.; Gastro-Correia, J.; Figueiredo, S.; Guerra, F.; Mgalhaes, S.I.; Torres, I.; Avarez, D.; Cruz, C. and Maia, J. (1991):** Glycemic control and diabetic retinopathy. *Medicographia* 13:49.
- Mendlovic, D.B.; Whitehouse, F.W. and Foreback, C.C. (1992):** The why and where for fructosamine. *Henry Ford. Hosp. Med. J.*, 40(1-2):149.
- Michael, B.; Helen, M.D. and Anthony, D. (1984):** Non-enzymatic glycosylation and the pathogenesis of diabetic complications. *Annals of Int. Med.*, 101:527.
- Migdalís, I.N.; Kalogeropoulou, K.; Zachariadis, D.; Koutoulidis, K. and Samartzis, M. (1994):** Serum levels of type III procollagen peptide and peripheral vascular disease in diabetic patients. *Diabetes, Res., Clin. Pract.*, 23(3):179.
- Miki, A.L.; Sandor, J.J. and Frountkey, D.M. (1972):** Complications of diabetes mellitus. *N. Eng. J. Med.*, 289:136.
- Miyake, H.; Nagashimo, K.; Yagi, H. and Onigatak, M. (1993):** Urinary laminin P₁ as an index of glycemic control in children with IDDM. *Diabetes. Res.*, 23(3):131.
- Mogenson, C.E. (1976):** Renal functional changes in diabetes. *Diabetes*, 25:782.
- Mogensen, C.E. (1982):** Long term and hypertensive treatment in the progression of diabetic nephropathy. *Br. Med. J.*, 282:685.

- Monnier, V.M.; Kohn, R.R. and Cerami, A. (1984):** Accelerated age-related browning of human collagen in diabetes mellitus. *Proc. Natl. Acad. Sci., U.S.A.* 81:583.
- Monnier, V.M.; Sell, D.R.; Abdl-Karim, F.W. and Emancipator, S.N. (1988):** Collagen browning and cross-linking are increased in chronic experimental hyperglycemia: relevance to diabetes and aging. *Diabetes*, 37:868.
- Morrison, P.R.; Edsall, J.T. and Miller, S.G. (1948):** Preparation and properties of serum and plasma proteins XVIII. The separation of purified fibrinogen from fraction I of human plasma. *J. Am. Chem. Soc.*, 70:3103.
- Mosesson, M.W. and Umfleet, R.A. (1970):** The cold insoluble globulin of human plasma purification primary characterization, and relationship to fibrinogen and other cold insoluble fraction components. *J. Biol. Chem.*, 245:5728.
- Mosesson, M.W.; Chen, A.B. and Huseby, R.M. (1975):** The cold insoluble globulin of human plasma: studies of its essential structural features. *Biochim Biophys. Acta*, 386:509.
- Mosesson, M.W. and Amrani, D.L. (1980):** The structure and biologic activities of plasma FN. *Blood*, 56(2):145.
- Mosher, D.F (1975):** Cross-linking of cold insoluble globulin by fibrin-stabilizing factor. *J. Biol. Chem.*, 250:6614.
- Mosher, D.F.; Sehad, P.F. and Van, J.M. (1980):** Cross-linking of collagen and FN by factor XIIIa. *J. Biol. Chem.*, 255:1181.
- Mosher, D.F. (1984):** Physiology of FN. *Ann. Rev. Med.*, 35:561.

Mostafa, M.S.; Barhooma, G.A. and Tadros, M. (1976): Pathological changes in conjunctival biopsy from complicated diabetic patients. Bull Ophthalmol. Soc. Egypt, 69:713.

Movat, H.Z. and Fernando, N.V. (1964): The fine structure of the terminal vascular bed. The venules and their perivascular cells (pericytes and adventitial cells). Exp. Molec. Path., 3:98.

Mushar, V.; Graninger, W.; Schober, E. and Schuster, E. (1984): FN in children with diabetes mellitus. Arch. Dis. Child, Oct.; 59(1):992.

Munjal, I.D.; McLean, N.V., Grant, M.B. and Blake, D.A. (1994): Differences in the synthesis of secreted proteins in human retinal endothelial cells of diabetic and non-diabetic origin. Curr. Eye. Res. Apr., 13(4):303.

Myllyla, A. (1976): Collagen glucosyl transferase: purification and characterization of the enzyme from chick embryos. Acta, Univer. Out., A44, Biochem., 8:13.

Mynderse, L.A.; Hassel, J.; Kleinman, H.K.; Martin, G.R. and Martinez-Hernandez, A. (1983): Loss of heparan sulphate proteoglycan from glomerular basement membrane of nephrotic rats. Lab. Invest., 48:292.

N

Nardelli, G.M.; Guastamacchia, E.; Paolo, S.; Lacasella, R.; Balico, A.; Montedoro, P.; Cospite, M.R. and Giorgino, R. (1987): Plasmatic levels of FN in diabetics with and without retinopathy. Correlation with some hormonal and metabolic parameters. Acta Diabetol. Lat.: Jul-Sep., 24(3):255.

Narita, T.; Funahashi, H.; Satoh, Y. and Kakagi, H. (1993): PIIINP and type IV collagen 75 domain in the sera of breast cancer patients. *Surg. Today*, 32(8):6882.

National Diabetes Data Group of The National Institute of Health (1979): Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes*, 28:1039.

Niemela, D. (1985): Radioimmunoassays for aminoterminal peptides in human type III procollagen. *Clin. Chem.*, 31:1301.

Niemela, O.; Risteli, E.A.; Sotaniemi, J. and Risteli, J. (1985): Type IV collagen and laminin-related antigens in human serum. *Eur. J. Clin. Invest.*, 15:132.



Oh, E.; Piershbacher, M. and Ruoslahti, E. (1981): Deposition of plasma FN in tissues. *Proc. Natl. Acad. Sci., U.S.A.*, 78:3218.

Ohno, M.; Ohno, N. and Kefalides, N.A. (1991): Studies on human laminin and laminin collagen complexes. *Con. Tis. Res.*, 25:251.

Okazaki, R.; Matsuaka, K.; Horiuchi, A.; Maruyama, K. and Okazaki, I. (1988): Assays of serum laminin and type III procollagen peptide for monitoring, The clinical course of diabetic microangiopathy. *Diabetes Res. Clin. Pract.*, 5(3):163.

Olefsky, J.M. (1992): Diabetes mellitus. In: *Cecil Textbook of Medicine*, 19th edition. Wyngaarden, J.B.; Smith, L.H. and Benett, J.C. (Eds.), Vol. 3, p. 1291-1310. W.B. Saunders Company.

- Ordinas, A.; Diaz-Ricart, M.; Bastida, E.; Escolar, G.; Gastillo, R.; Tandon, N.N. and Jamieson, G.A. (1992):** The role of subendothelial laminin and platelet laminin receptor in haemostasis. *Nouv. Rev. Fr. Hematol.*, 3491:61.
- Oslen, B.R. (1981):** Collagen biosynthesis. In: *Cell biology of extracellular matrix*. Elizabeth, H. (Ed.), Elsevier Science, (Pub.), p. 139.
- Osterby, R. and Gundersen, S. (1975):** Glomerular size and structure in diabetes mellitus early abnormality. *Diabetologia*, 11:295.
- Osterby, R. (1983):** Basement membrane morphology in diabetes mellitus: In *Diabetes Mellitus, theory and practice*, 3rd Edition, Ellenberg, M. and Rifkin H., editors. Medical examination Publishing Co., New Hyde-Park, 323.
- Osterby, R. (1990):** Basement membrane morphology in diabetes mellitus; in *Diabetes Mellitus, theory and practice*; Fourth edition. Rifkin, H. and Porte, D. (Eds.), p. 220-233. Elsevier, New York, Amsterdam, London.
- Ott, V.; Odermatt, E.; Engel, J.; Furthmay, H. and Timpl, R. (1982):** Protease resistance and conformation of laminin. *Eur. J. Biochem.*, 123:63.
- P**
- Pagani, A.; Zigliani, P.; Pranzo, A. and Galante, T. (1992):** Determination of serum procollagen III peptide in chronic liver diseases. *Minerva Med.*, 83(11):715.

- Palmberg, P.; Smith, M. and Waltman, S. (1981):** The natural history of retinopathy in insulin dependent juvenile onset diabetes. *Ophthalmology*, 88:613.
- Panayotou, G.; End, P.; Aumailley, M.; Timpl, R. and Engel, J. (1989):** Domains of laminin with growth factor activity. *Cell*, 56:93.
- Patz, A.; Schatz, H. and Berkow, J.W. (1973):** Macular oedema on overlooked complication of diabetic retinopathy. *Trans. Am. Acad. Ophth. Otolaryngol.*, 77:34.
- Paul, B. and Wolfie, T. (1979):** The proteins. Hans, N.; Robert, L.H. and Carol-Leigh, B. (Eds.), Vol. IV, Academic Press, New York, San Francisco, London, p. 442.
- Paulsson, M.; Duetzmann, R.; Timpl, R.; Dalzopoo, D.; Odermatt, E. and Engel, J. (1985):** Evidence for coiled α helical regions in the long arm of laminin. *EMBO J.*, (4):307.
- Paulsson, M.; Aumailley, M.; Deutzmann, R.; Timpl, R.; Beck, K. and Enge J. (1987):** Laminin-nidogen complex extraction with chelating agents and structural characterization. *Eur J. Biochem.*, 166:11.
- Paulsson, M. (1988):** The role of Ca^{++} binding in the self-aggregation of laminin-nidogen complexes. *J. Biol. Chem.*, 263:5425.
- Paulus, W.; Roggendorf, W. and Schuppan, D. (1988):** Immunohistochemical investigation of collagen subtypes in human glycoblastomas. *Virchows Arch. Pathol. Anat. Histol.*, 413:325.
- Pearlstein, E. (1976):** Plasma membrane glycoprotein which mediates adhesion of fibroblast to collagen. *Nature*, 162:497.

- Perez-Arellano, J.L.; Pedraz, M.J.; Fuertes, A.; DeLa Cruz, J.L.; Gonzalez, De Buitrago, J.M. and Jimenez, A. (1993):** Laminin fragment P¹ is increased in the lower respiratory tract of patients with diffuse interstitial lung disease. *Chest*, 104(4):1163.
- Peterson, C.M. and Jone, R.L. (1980):** Minor haemoglobin, diabetic control and diseases of post-synthetic protein modification. *Ann. Intern. Med.*, 87:489.
- Peyrol, S.; Hugues, B.; Grimaud, J.A. and Mornex, R. (1984):** Collagen heterogeneity of thickened BM of human diabetic dermis: tissular immunolabelling of collagen types I, III and IV by light and electron microscopy. *Cell. Mol. Biol.*, 30:43.
- Phan-Thanh, L.; Robert, L.; Derouette, J. and Labat-Robert, J. (1987):** Increased biosynthesis and processing of fibronectin in fibroblasts from diabetic mice. *Proc. Natl. Acad. Sci., U.S.A.*, 84:1911.
- Pierschbacher, M.D. and Ruoslahti, E. (1984):** Cell attachment activity of FN can be duplicated by small synthetic fragments of the molecule. *Nature*, 309:30.
- Pietschmann, P.; Schernthaner, G.; Schnack, C.H. and Gaube, S. (1988):** Serum concentrations of laminin P¹ in diabetics with advanced nephropathy. *J. Clin. Pathol.*, 41(9):929.
- Pikkarainen, T.; Kallunki, T. and Tryggvason, K. (1987):** Human Laminin B₂ chain. *J. Biol. Chem.*, 263:6751.
- Poduslo, F.J. and Curran, L.G. (1992):** Increased permeability across the blood-herve barrier of albumin glycated in vitro and in vivo from patients with diabetic polyneuropathy. *Proc. Natl. Acad. Sci., U.S.A.* 89:2218.

- Popova, V.A.; Ostashevskaja, M.I. and Podgorodnichentso, V.K. (1990):** Clinico-diagnostic significance of FN determination in diabetes mellitus in children. *Pediatricia*, 6:9.
- Porte, D.J. and Halter, J.B. (1981):** The endocrine pancreas and diabetes mellitus. In: William, S.R.H. (Ed.): *Textbook of endocrinology*, 6th ed., p. 716-843, Saunders, Philadelphia.
- Poulson, R.; Kurkinen, M.; Prockop, D.J. and Boothand, F.R. (1988):** Increased steady state levels of laminin B₁ mRNA in kidneys of long-term streptozotocin-diabetic rats. *J. Biol. Chem.*, 263:10072.
- Price, T.M.; Rudee, M.L.; Pierschbaker, M. and Ruoslahti, E. (1982):** Structure of FN and its fragments in electron microscopy. *Eur. J. Bio. Chem.*, 129:359.
- Prince, M.J.; Tsai, P. and Olefsky, J.M. (1981):** Insulin binding, internalization and insulin receptor regulation in fibroblasts from type II NIDDM diabetic subjects. *Diabetes*, 30:596.
- Prockop, D.J.; Kivirikko, K.I.; Tuderman, L. and Guzman, N.A. (1979):** The biosynthesis of collagen and its disorders. *The New Eng. J. of Med.*, 301:13.
- Proctor, R.A.; Prendergast, E. and Mosher, D.F. (1982):** FN mediates attachment of staphylococcus aureus to human neutrophils. *Blood*, 59:681.
- Proctor, R.A. (1987-a):** FN: A brief overview of its structure, function and physiology. *Rev. of Infectious Diseases*, 9, Suppl. 4:S317.
- Proctor, R.A. (1987-b):** The staphylococcal FN receptor: evidence for its importance in invasive infections. *Rev. Infect. Dis.*, 9 Suppl. S335.

Proietto, J. (1990): Pathophysiology and genetics of non-insulin dependent diabetes mellitus. *Medicographa*, 12(2):18.

Pugh, J.A. (1989): The epidemiology of diabetic nephropathy. *Diabetes Metab. Rev.*, 5:531.

Puistola, V.; Risteli, L.; Kauppila, A.; Knip, M. and Risteli, J. (1993): Markers of type I and type III collagen synthesis in serum as indicators of tissue growth during pregnancy. *J. Clin. Endocr. Metab.* 77(1):178.

R

Rajaraman, R.; MacSeveen, J.M. and Murdock, C.A. (1983): Exogenous FN requirement for adhesion by neoplastic human cells. *Exp. Cell. Biol.*, 51:9.

Ramsden, J.J. (1993): Calcium-dependence of laminin binding to phospholipid membranes. *Biopolymers*, 33(3):475.

Rauterberg, J.; Voss, B. and Pott, G. (1981): Connective tissue components of the normal and fibrotic liver. *Klin. Wocheschr.*, 59:767.

Reaven, P.D.; Barrett-Connor, E.L. and Browner, D.K. (1990): Abnormal glucose tolerance and hypertension. *Diabetes Care.*, 13:199.

Repesh, L.A.; Fitzgerald, T.D. and Furcht, L.T. (1982): FN involvement in granulation tissue and wound healing in rabbits. *J. Histochem. Cytochem.*, 30:351.

- Rhodeo, R.K.; Miller, E.J. and Simon, P. (1978):** Physiochemical characterization and molecular organization of the collagen A and B chain. *Biochem. Biophys. Res. Comm.*, 80:866.
- Rhodin, J.A. (1962):** Fine structure of vascular walls in mammals with special reference to smooth muscle components. *Physiol. Re.*, 5 Suppl., 48.
- Risteli, J.; Drager, K.e.; Reqitz, G. and Neubauer, H.P. (1982):** Increase in circulating BM antigens in diabetic rats and effects of insulin treatment. *Diabetologia*, 23:266.
- Risteli, J. and Risteli, A. (1986):** Radio-immunoassays for monitoring connective tissue metabolism. *Rheumatology*, 10:216.
- Ristel, J., Niemi, S.; Trivedi, P.; Maentausta, O. and Risteli, A. (1988):** Rapid radioimmunoassay for the aminoterminal propeptide of human type III procollagen. *Clin. Chem.*, 34(1):715.
- Rochlitz, C.H.; Hasslacher, C.H.; Brocks, D.G. and Herrmarr, R. (1987):** Serum concentration of laminin and course of the disease in patients with various malignancies. *J. Clin. Oncol.*, 9:1424.
- Rohrbach, D.H.; Hassell, J.R.; Kleinman, H.K. and Martin, G.R. (1982):** Alteration in the basement membrane heparan sulfate proteoglycan in diabetic mice. *Diabetes*, 31:185.
- Rojkind, M. and Dunn, M.K. (1979):** Hepatic fibrosis. *Gastroenterology*, 76:849.
- Rossini, A.A.; Mardes, J.P. and Handler, E. (1988):** Perspectives in diabetes. Speculations on etiology of diabetes mellitus. Tumbler hypothesis. *Diabetes*, 37:257.

- Rowe, D.W.; Starman, B.J.; Fujimoto, W.Y. and Williams, R.H. (1977):** Abnormalities in proliferation and protein synthesis in skin fibroblast cultures from patients with diabetes mellitus. *Diabetes*, 26:248.
- Roy, M.S. and McCulloch, J.C. (1982):** Peripheral retinal haemorrhages in long duration insulin dependent diabetes with minimal background retinopathy. *Br. J. Ophth.*, 66:382.
- Ruoslahti, E. and Vaheri, A. (1974):** Novel human serum protein from fibroblast plasma membrane. *Nature*, 248:790.
- Ruoslahti, E.; Engvall, E. and Hyman, E.G. (1981):** FN: Current concepts of its structure and functions. *Cell Res.*, 1:95.

S

- Saba, T.M. (1970):** Physiology and physiopathology of the reticuloendothelial system. *Arch. Intern. Med.*, 126:1031.
- Saba, T.M. and Jaffe, E. (1980):** Plasma FN; its synthesis by vascular endothelial cells and role in cardiopulmonary integrity after trauma as related to reticuloendothelial function. *Am. J. Med.*, 68:577.
- Saba, T.M.; Blumenstock, F. and Shob, D. (1986):** Reversal of opsonic deficiency in surgical trauma and burn patients by infusion of purified human plasma FN. *A.M.J. Med.*, 80(2):229.
- Sank, A.; Wei, D.; Reid, J.; Ertl, D.; Nimni, M.; Weaver, F.; Yellin, A. and Twan, T.L. (1994):** Human endothelial cells are defective in diabetic vascular disease. *J. Surg. Res.*, 57(6):647.

- Sasaki, M.; Kleinman, H.K.; Huber, H.; Deutzmann, R. and Yamada, Y. (1988):** Laminin, a multidomain protein. The A chain has a unique globular domain and homology with the BM proteoglycan and the laminin B chains. *J. Biol. Chem.*, 263:16536.
- Sato, S. (1992):** 75 in patients with chronic liver disease with special reference to histological findings. *Nippon-Shokakibyō-Gakkai-Zasshi*, 89(9):2007.
- Scarpa, S.; Dorazi, G.; Ragano-Caracciolo, M.; Cardelli, P.; Masuelli, L. and Modesti, A. (1992):** Modulation of laminin synthesis in human neuroblastoma cells during retinoic acid induced differentiation. *Cancer*, 64(1):31.
- Scheiffarth, O.F.; Kampi, K.A.; Gunthern, H. and Vonder-Mark, K. (1988):** Proteins of the extracellular matrix in vitreoretinal membranes. *Grafe's Arch. Clin. Exp. Ophthalmol.*, 226:357.
- Schiffrin, A. (1986):** Reversibility of diabetic complications. *Recent advances in diabetes*, 11:145.
- Schirakami, A.; Shigekiyo, T. and Hvawi Y. (1986):** Plasma FN deficient in eight members of one family. *Lancet*, (847):437.
- Schmid, S.H. (1981):** Blood rheology and physiology of microcirculation. *LaRicerca Clin. Lab.* 11 (Suppl. 1):13.
- Schuppan, D.; Hahn, E.G. and Riecken, E.O. (1988):** Connective tissue proteins in the liver: collagenes and glycoproteins. In: *Cholestasis in neonates*. Waldschmidt, (Ed.), Zuchschwerdt, Verlag, Munchen, p. 97-115.
- Seaquist, E.R.; Goeto, F.C. and Rich, S. (1989):** Familial clustering of diabetic renal disease. *N. Engl. J. Med.*, 320:1161.

- Seeling, R.; Pott, G.; Seeling, H.P.; Licher, H.; Metzger, P. and Waldherr, R. (1984):** Virus binding activity of FN masking of hepatitis A virus. *J. Virol. Methods*, 8:335.
- Seghieri, G.; Bartolomei, G. and DeGiorgia, L.A. (1986):** Plasma FN in diabetic retinopathy and macroangiopathy. *Diabetes Metab. Aug.*, 12(4):186.
- Selby, J.V.; Fitz-Simmons, S.C.; Newman, J.M.; Hatz, P.P.; Sepe, S. and Showstack, J. (1990):** The natural history and epidemiology of diabetic nephropathy. *JAMA*, 263(14):1954.
- Sephel, G.C.; Tashiro, K.I.; Sasaki, M.; Grestorex, D.; Martin, G.R.; Tamada, Y. and Kleiman, H.K. (1989):** Laminin A chain synthetic peptide which supports neurite outgrowth. *Biochem. Biophys. Res. Commun.*, 162:821.
- Shen, T.F. and Strecker, H.J. (1975):** Synthesis of proline and hydroxyproline in human lung fibroblasts. *J. Biochem.*, 150:453.
- Shimiza M.; Vmeda, K.; Sugihara, N.; Yoshio, H.; Takedo, R. and Okada, Y. (1993):** Collagen remodelling in myocardia of patients with diabetes. *J. Clin. Pathol.*, 46(1):32.
- Shimuzu, K.; Kobayash, Y. and Muraoko, K. (1981):** Midperipheral fundus involvement in diabetic retinopathy. *Ophthalmology* 82:601.
- Simpson, W.A.; Courtney, H.S. and Ofek, I. (1987):** Interactions of FN with Streptococci: The role of FN as a receptor for streptococcus pyogenes. *Rev. Infect. Dis.*, 9(Suppl.4):S351.
- Singer, D.E.; Nothan, D.M.; Fogil, H.A. and Schanchat, A.P. (1992):** Screening for diabetic retinopathy. *Ann. of Int. Med.*, 116(8):660.

- Singer, I.I. (1979):** The fibronexus: a transmembrane association of FN containing fibers and bundles of 5 nm microfilaments in hamster and human fibroblasts. *Cell*, 16:67775.
- Skrha, J., Vockova, I., Skvasnicka, J., Stibor, V., Stolba, P., Richeter, H. and Hormann, H. (1990):** Plasma free N-terminal FN 30-KDa domain as a marker of endothelial dysfunction in type I diabetes mellitus. *Eur. J. Clin. Invest. Apr.*, 200:171.
- Smith, E.L. and Hill, R.L. (1983):** Connective tissue. In: Principles of biochemistry mammalian biochemistry. Smith, E.L. and Hill, R.L. (Eds.), 7th ed., McGraw Hill Book Company, New York, St. Louis, U.S.A., p. 211.
- Smith, H.S.; Riggs, J.L. and Mosesson, M.W. (1979):** Production of FN by human epithelial cells in culture. *Cancer, Res.*, 39:4138.
- Smith, J.W. (1968):** Molecular pattern in native collagen. *Nature*, 219:157.
- Smith, J.W. and Silbert, A. (1981):** Fibronectin and collagen of cultured skin fibroblasts in diabetes mellitus. *Biochem. Biophys. Res. Commun.*, 100:275.
- Snow, D.M. and Letourneau, P.C. (1992):** Neurite outgrowth on a step gradient of chondroitin sulfate proteoglycan. *J. Neurobiol.*, 23(3):322.
- Solerte, S.B., Fioravanti, M. and Ferrari, E. (1987):** Plasma FN as an indicator of microvascular damage in diabetic patients. Rapid and sensitive evaluation by a radial immunodiffusion technique. *Ric. Clin. Lab.*, Jul-Sept.; 19(3):251.
- Song, C.W.; Payne, J.T. and Levitt, S.H. (1972):** Vascularity and blood flow in α -irradiated Walker carcinoma 256 of rats. *Radiology*, 104:693.

- Sosenko, J.M.; Breslow, J.L.; Miettinen, O.S. and Gabbay, K.H. (1979):** Prospective study of diabetic patients. *N. Eng. J. Med.*, 302:650.
- Sotaniemi, E.A.; Stenback, F.G.; Risteli, L.; Vuori, J.A.; Rautio, A.; Hupponen, R. and Risteli, J. (1992):** Effect of insulin on serum aminoterminal propeptide of type III procollagen in non-insulin dependent diabetes mellitus. *Diab. Res. Clin. Pract.*, 17(3):209.
- Springer, T.A. (1990):** Adhesion receptors of the immune system. *Nature*, 346:425.
- Stack, S.; Gray, R.D. and Pizzo, S.V. (1991):** Modulation of plasminogen activation and type IV collagenase activity by a synthetic peptide derived from the laminin A-chain. *Biochemistry*, 30(8):2073.
- Stallmach, A.; Schuppan, D.; Dax, J.; Hanski, C. and Biecken, E.O. (1990):** Identification of laminin binding proteins in cell membranes of a human colon adenocarcinoma cell-line. *Gut*, 31:70.
- Stanbridge, E.J.; Der, C.J. and Doersen, C.J. (1982):** Human cell hybrids: analysis of transformation and tumor genecity. *Science*, 1219:252.
- Stanley, J.R.; Foidart, J.M.; Murray, J.C.; Martin, G.R. and Katz, S.I. (1980):** The epidermal cell which selectively adheres to a collagen substrate in the basal cell. *Invest. Dermatol.* 74:54.
- Stathakis, N.E.; Fountas, A. and Teianoe, E. (1981):** Plasma FN in normal subjects and in various disease states. *J. Clin. Pathol.*, 34:504.
- Stenman, S. and Vaheri, H. (1978):** Distribution of a major connective tissue protein FN, in normal human tissues. *J. Exp. Med.*, 147:1045.

Stern, W.H.; Anderson, D.H.; Fisher, S.K.; Borgula, G.A. and Erikson, P.A. (1981): Anatomy recovery following retinal detachment clinicopathologic correlations. *Aust. N.Z.J. Ophthalmol.*, 9:143.

Sternberg M.; Cohen-Fortner, L. and Peyroux, J. (1985): Connective tissue in diabetes mellitus: biochemical alterations of the intercellular matrix with special reference to proteoglycans collagens and BMs. *Diabetes, Metab.*, 11:27.

Studer, R.K.; Graven, P.A. and DeRubertis, F.R. (1993): Role of protein kinase C in the mediation of increased FN accumulation by mesangial cells grown in high glucose medium. *Diabetes*, 42:118.



Takahashi, S. and Lee, M.T. (1987): Collagen metabolism in fibrotic liver. *Biochem. J.*, 241:49.

Tanzer, M.L., Chandrasekaran, S.; Dean, J.W. and Giniger, M.S. (1993): Role of laminin carbohydrates on cellular interactions. *Kidney, Int.*, 43(1):66.

Tarsio, J.F.; Reger, L.A. and Furcht, J.T. (1987): Decreased interaction of FN, type IV collagen and heparin due to non-enzymatic glycation. *Biochemistry, Feb.*, 24:26(4):1014.

Tarsio, J.F.; Reger, L.A. and Furcht, J.T. (1988): Molecular mechanisms in BM complications of diabetes, alterations in heparin, and type IV collagen association. *Diabetes*, 37(5):532.

Taylor, R. (1989): Aetiology of non-insulin dependent diabetes mellitus. *Br. Med. Bull.*, 45(1):73.

- Timpl, R.; Johansson, S.; Van Delden, V.; Oberbaumer, I. and Hooh, M (1983):** Characterization of protease resistant fragments of laminin mediating attachment and spreading of rat hepatocytes. *J. Biol. Chem.*, 258:8922.
- Tinder, P. (1969):** Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Ann. Clin. Biochem.* 6:24.
- Toda, A.; Takeuchi, B., Tokagi, M. and Isogai, S. (1987):** The effect of glycemic control on plasma soluble fibrin monomer complexes and FN in diabetic patients. *Tohoku J. Exp. Med.* Oct.; 153(2):145.
- Todd, J.A.; Bell, J.I. and McDevitt, H.O. (1987):** HLA-DQB gene contributes to suseptibility and resistance to insulin dependent diabetes mellitus. *Nature*, 329:599.
- Tomono, S.; Kawazu, S.; Kato, N.; Ohno, T.; Vtusgi, T. and Muraka, K. (1991):** Clinical complications of serum levels of BM components in diabetic patients with and without albuminuria. *J. Diabet. Complications*, 5(2-3):193.
- Torre, D.; Zeroli, C.; Ferrario, G.; Bonetta, G.; Fiori, G.F. and Martegani, R. (1993):** Cerebrospinal fluid concentration of FN in patients with HIV-1 infection and central nervous system disorders. *J. Clin. Pathol.*, 46:1039.
- Touraine, J.L.; Tourane, F. and Revilland, J.P. (1975):** T-lymphocytes and serum inhibitors of cell-mediated immunity in renal insufficiency. *Nephron*, 14:195.
- Triolo, G.; Giardina, E.; Zarccone, M.R.; Seddiog, G. and Bompiani, G.D. (1989):** Serum levels of type III procollagen peptide in diabetes mellitus. *Horm. Metab. Res.*, 21(2):77.

Tronchin, G.; Bouchara, J.P.; Larcher, G.; Lissitzky, J.C. and Chabasse, D. (1993): Interaction between *Aspergillus fumigatus* and basement membrane laminin: binding and substrate degradation. *Biol. Cell.*, 77(2):201.

U

Uracko, R. (1974): Basal lamina scaffold-anatomy and significance of orderly tissue structure. *Am. J. Path.*, 77:314.

V

Vaheri, A.; Ruaslahti, E.; Westermarck, B. and Ponten, J. (1976): A common cell-type specific surface antigen in cultured human glial cells and fibroblasts: loss of malignant cells. *J. Exp. Med.*, 143:64.

Vaheri, A.; Salonem, E.M. and Vartio, T. (1985): FN in formation and degradation of the pericellular matrix in fibrosis. *Ciba Foundation Symposium (114):* Pitman, London, p. 111-120.

Varley, H.; Gowenlock, A.H. and Bell, M. (1980): Non-protein nitrogen, urea, urate, creatine and creatinine in: *Practical Clinical Biochemistry*, Varley, H.; Gowenlock, A.H. and Bell, M. (Eds.), William Heinemann Medical Books, London, Vol. 1, p. 456.

Vartio, T., Vaheri, A.; Von Essen, R.; Isomak, H. and Stenman, S. (1981): FN in synovial fluid and tissue in rheumatoid arthritis. *Eur. J. Clin. Invest.*, 11:207.

- Veijola, R.; Kni, M.; Risteli, L.; Risteli, J.; Kaar, M.L.; Puukka, R. and Ilonen, J. (1993):** Clinical characteristics and circulating collagen and laminin metabolites in IDDM diabetic children with joint and skin manifestations. *Pediatr. Res.*, 33(5):501.
- Velez, R.L. and Henrich, W.L. (1984):** The vasculature. In: The systemic consequences of renal failure. E. Knoyan, G. and Knochel, J. P. (Eds.), Grune & Stratton Inc., Orlando, Fl. 93.
- Vishwanath, V.K., Frank, C.A.; Elmets, P.J. and Manner, V.M. (1986):** Glycation of skin collagen in type I diabetes mellitus, correlation with long-term complications. *Diabetes*, 35:916.
- Voss, B.; Allam, S.; Rauterberg, J.; Ulrick, K. Gieselmann, V. and Von Figura, K. (1979):** Primary cultures of rat hepatocytes synthesize FN. *Biochem. Biophys. Res. Commun.*, 90:1348.
- 
- Wagner, D.D. and Hynes, R.O. (1979):** Domain structure of FN and its relation to function. Disulfide and sulphhydryl groups. *J. Biol. Chem.*, 254:6746.
- Waite, S. and Beetham, T. (1980):** Vascular lesions in type II diabetes mellitus. *N. Engl. J. Med.*, 212:367.
- Weidmann, P. and Weller, M. (1988):** The pathophysiology of proliferative vitreoretinopathy. *Acta Ophthalmol.*, 66: Suppl. 189.
- Weistener, M.; Krieg, J. and Horlein, D. (1977):** Inhibiting effect of procollagen peptide on collagen biosynthesis in fibroblast culture. *J. Biol. Chem.*, 57:47.

- Weller, M.; Weidmann, P.; Bresgen, M. and Heimann, K. (1991):** Vitronectin and proliferative interocular disorders. II- Expression of cell surface receptors for FN and vitronectin in periretinal membranes. *International Ophthalmology*, 15:103.
- Wiklund, T.A.; Blomqvist, C.P.; Risteli, L.; Risteli, J. and Eloma, I. (1993):** Effect of chemotherapy on collagen metabolism. A study of serum PIIINP in advanced sarcomas. *J. Cancer Res. Clin. Oncol.*, 119(3):160.
- Williamson, J.R. and Kilo, C. (1976):** Basement membrane thickening and diabetic microangiopathy. *Diabetes*, 25: Suppl. 2., 925.
- Wolff, S.P. (1992):** Blood glucose concentrations and progression of diabetic retinopathy. *B.M.J.* 304:505.
- Wright, S.D. and Meyer, B.C. (1985):** FN receptor of human macrophages recognizes the sequences Arg.-Gly.-Asp.-Ser. *J. Exp. Med.*, 162:762.



- Yang, C.W.; Vlassara, H.; Peten, E.P.; He, C.J.; Shiker, G.E. and Shiker, L.J. (1994):** Advanced glycation end products up-regulate gene expression found in diabetic glomerular disease. *Proc.-Natl.-Acad. Sci., U.S.A.* 91(20):9436.
- Yanoff, M. (1966):** Diabetic retinopathy. *N. Eng. J. Med.*, 274:1344.
- Yanoff, M.; Fine, B. and Berkow, J. (1970):** Vacuolation of the pigment epithelium of the iris: a histologic finding considered pathognomonic of diabetes. *Am. J. Ophth.*, 96:201.

- Yamada, K.M. and Weston, J.A. (1974):** Isolation of major cell surface glycoprotein from fibroblasts. *Proc. Natl. Acad. Sci. U.S.A.*, 71:34.
- Yamada, K.M.; Yamada, S.S. and Pastan, I. (1976):** Cell surface protein partially restores morphology adhesiveness and contact inhibition of movement to transformed fibroblasts. *Proc. Natl. Acad. Sci., U.S.A.*, 73:1217.
- Yamada, K.W. and Yamada, S.S. (1983):** Cell surface interaction with extracellular materials. *Ann. Rev. Biochem.*, 52:761.
- Yue, D.K.; McLennan, S.; Handelsman, D.J.; Delbridge, L.; Reeve, T. and Turtle, J. (1985):** The effects of cyclo-oxygenase and lipo-oxygenase inhibitors on the collagen abnormalities of diabetic rats. *Diabetes*, 34:74.
- Yurchenco, P.D.; Tsilibary, E.C.; Charonis, A.S. and Furthmayer, H. (1985):** Laminin polymerization in vitro, evidence for a two-steps assembly with domain specificity. *J. Biol. Chem.*, 260:7636.



- Zhu, G.G.; Risteli, J.; Puistala, V.; Kauppila, A. and Risteli, L. (1993):** Progressive ovarian carcinoma induces synthesis of type I and type III procollagens in the tumour tissue and peritoneal cavity. *Cancer, Res.*, 53(20):5028.
- Ziadeh, F.N.; Snipes, E.R.; Watanabe, M.; Alvarez, R.J.; Goldfarbs and Hoverty, T.P. (1990):** High glucose induces cell hypertrophy and stimulates collagen gene transcription in proximal tubule. *Am. J. Physiol.*, 259:F704.

- Ziayadeh, F.N. (1993):** Renal tubular basement membrane and collagen type IV in diabetes mellitus. *Kidney Int.*, 43(1):114.
- Zimmet, P.; Dowse, G. and Bennett, P. (1991):** Hyperinsulinaemia is a predictor of non-insulin dependent diabetes mellitus. *Diabetes, Metab.*, 17(1):101.
- Zucker, M.B.; Mosesson, M.W.; Broekman, M.J. and Kaplan, K.L. (1979):** Release of platelet FN from granules induced by thrombin or collagen: lack of requirement for plasma FN in ADP-induced platelet aggregation. *Blood*, 54:8.
- Zweifach, B.W. (1980):** Integrity of vascular endothelium, in: *Advances in microcirculation*, Vol. 9, p. 206-225, (Karger, Basel).