

REFERENCES

*Abramwicz D, Schandene L, and Goldman M .

Release of tumour necrosis factor interleukin-2 and gamma-interferon in serum after injection of OKT3 monoclonal antibody in kidney transplant recipients .

Transplantation ,47,606-8,1989.

*Auffray C, and Strominger JL.

Molecular genetics of the human major histocompatibility complex .
Adv Human Genet ,15,197-247,1985.

*Bailey DW.

Four approaches to estimating the number of H -loci

Transplant Proc ; 2, 32-8, 1970.

*Baldwin WM, Claas FHJ, and Paul LC.

All monocyte antigens are not expressed on renal endothelium.

Tissue Antigens , 21:254, 1983.

*Battye, F.L.

A fast cell sampler for flow cytometry.

Cytometry 6 (5) :492-494,1985.

*Bjorkman PJ, Saper MA, Samraoui B, Bennet WS, Strominger JL, and Wiley DC.

Structure of the human class I histocompatibility antigen, HLA-A2.

Nature ;329,506-12,1987.

*Blom ,J.

A comparison of two fluorescent activated cell sorters, the FACS IV (Laser) and FACS Tm (mercury Lamp), as research analyzers for the quantitation of T and B subsets in human peripheral blood .

J.Immunol. Methods 95 (1) :39-46 ,1986.

*Blume, P.A, and Stovel,L.A.

A spectrophotometric method for determining the stream sample core diameter of a flow cytometry.

Cytometry 10 (3) : 351-353 ,1989.

*Borst J,Alexander S,Elder J,and Terhorst C.

The T3 complex on human T lymphocytes involves four structurally distinct glycoproteins .

J.Biol Chem ;258, 5135-41,1983.

*Brasile L,Kremer JM,Clarke JI, and Cerilli J.

Identification of an autoantibody to vascular endothelial cell specific antigens in patients with systemic vasculitis .

Am.J.;87,74-80,1989

*BrennanDC, Jevnikar AM, Takei F,and Reubin Kelley VE.

Mesangial cell accessory functions:mediation by intercellular adhesion molecule-1.

Kidney International;38,1039-46,1990

*Brenner MB, Strominger JL, and Krangel MS.

The T cell receptor .

Adv Immunol ;43, 133-92,1988.

*Brewer Y, Bewick M, Palmer A, Severn A, Welch K, Taube D.

Prevention of renal allograft rejection by perfusion with antileukocyte common monoclonal antibodies is dependent on good uptake of antibody by interstitial dendritic cells .

Transplantation Proc;21,1172-3,1989.

*Brodsky FM, and Guagliardi LE.

The molecular immunology of transplantation and graft rejection .

Immunol Allergy Clin N America ;9,1-19,1989.

*Brown JH, Jardetzky T, Saper MA, Samraoui B, Bjorkman PJ, and Wiely DC.

A hypothetical model of the foreign antigen binding site of class II histocompatibility molecules .

Nature ; 332,845-50,1988.

*Calabi F, and Milstein C.

A novel family of human major histocompatibility -related genes not mapping to chromosome 6

Nature ,323,540-3,1986

*Cerilli J, Brasile L, Clarke J, and Galouzis T.

The vascular endothelial cell- specific antigen system:three years' experience in monocyte crossmatching.

Transplant Proc, 17:567,1985.

*Cerilli J,Holliday J,Wilson C and Sharma H.

Clinical significance of the 1-hour biopsy in renal transplantation.

Transplantation: 26:91,1978.

*Ceryl,JG.

Significance of vascular endothelial cell antigen.

Transplant Proceedings : 6,4468-70,1987.

*Class FHJ, Paul LC, and Van Rood JJ.

Antibodies against donor antigens on endothelial cells and monocytes in elutes of rejected kidney allografts.

Tissue antigens :15:19,1980.

*Clayberger C, Parham P, Rothbard J, Ludwig DS, Schoolnik GK,and Krensky AM.

HLA-A2 peptides can regulate cytotoxicity by human allogenic T lymphocytes .

Nature ;330,763-5,1987.

*Clift R,and R.Storb.

Histo compatibility bone marrow transplants in humans .

Annual Review of Immunology ;5 ,43-64,1987.

*Coppage M.

Further observation on the significance of non HLA antigens transplantation .

Human Immunol, vol 23, suppl 1, pg 6, 1991.

*Daar AS,Fuggle SV,Fabre JW,Ting A,and Morris PJ.

The detailed distribution of HLA-A,B,C antigens in normal human organs .

Transplantation ;38,287-94,1984.

*Davis MM.

T cell receptor gene diversity and selection .

Ann Rev Biochem;59,475-96 , 1990.

*dela Hera A, Muller U,Olsson C, Isaaz S,and Tunacliffe A.

Structure of the Tcell antigen receptor (TCR) :two CD3 subunits in a functional TCR/CD3 complex.

J Exp Med ;173,7-17,1991.

*Dolittle DP,Davisson MT, Roderick TH,and Hiallyard AL.

The locus map of the mouse .

Mouse genome ;87, 14-27,1990.

*Doyle C, and Strominger JL.

Interaction between CD4 and class II MHC molecules mediates cell adhesion .

Nature ; 330,256-9,1987

*Falk K, Rotzschke O, Stevanovic S, Jung G, and Rammensee H-J.

Allele-specific motifs revealed by sequencing of self peptides eluted from MHC molecules .

Nature ;351,290-6,1991.

*Ferrara JLM, and Deeg HJ.

Graft versus host disease.

New Eng J Med ;324,667-74 ,1991.

*Fraser JD, Goldsmith MA, and Weiss A.

Ligand induced association between the T cell antigen receptor and two glycoproteins.

Proc Natl Acad Sci (USA); 86,7133-7,1989.

*Garovoy MR.

Prospective flow cytometry crossmatching appears to improve graft survival in high PRA and regrant recipients.

Human Immunol, vol23, suppl 1, pg 69,1991.

*Gillitzer R, Pilarski L M.

In situ localization of CD45 isoforms in the human thymus indicates a medullary location for the thymic generative lineage .

J Immunol; 144,66-74,1990.

*Graff RJ, and Brown DH.

Estimates of histocompatibility differences between inbred mice strains .

Immunogenetics ;7,367-73, 1987.

*Groh V, porcelli S, and Fabbi M.

Human lymphocytes bearing T cell receptor are phenotypically diverse and eventually distributed throughout the lymphoid system .

J Exp Med ;169,1277-94,1989

*Halloran PF, Cockfield SM ,and Madrenas J.

The molecular immunology of transplantation and graft rejection .

Immunol Allergy Clin N America ;9,1-19,1989.

*Hedrick SM.

Specificity of the T cell receptor for antigen .

Adv Immunol;43,193-234,1988.

*Jardetzky TS, Lane WS ,Robioson RA,Madden DR,and Wiley DC.

Identification of self peptides bound to purified HLA-B27.

Nature ;353,326-9,1991.

*Jevnikar AM,Wuthrich RP, Takei F,Xu HW,Brennan DC, Glimcher LH,and
Rubin-Kelley VE.

Differing regulation and function of ICAM-1 and class II antigens on
renal tubular cells .

Kidney International ;38,417-25,1990.

*Jordan SC,Yap HK,and Alfonso P.

Hyperacute allograft rejection mediated by antivascular endothelial
cell antibodies with a negative monocyte crossmatch.

Transplantation: 46:585,1988.

*Joyce S, Flye MW, and Mohanakumar T.

Characterization of kidney cell specific, non-major
histocompatibility complex alloantigen using antibodies eluted from
rejected human renal allografts .

Transplantation ;46,362-9,1988.

*Kamentsky,L.A.

Cytology Automation.

Adv. Biol Med Phys 14:93 , 1973.

*Kelly AP,Monaco JJ,Cho S, and Trowsdale J.

A new human HLA class II-related locus .

Nature ,353,571-3,1991.

*Kisielow P, Bluthmann H, Staerz UD, Steinmetz M, and Von Boehmer H.
Tolerance in T cell receptor transgenic mice involves deletion of
nonmature CD4+8+ thymocytes .

Nature. ;333, 742-6, 1988.

*Klein J.

Natural History Of The Major Histocompatibility complex .
New York :John Wiley and Sons ,1986 .

*Koning F.

Lymphocyte antigen receptors: a common design?
Immunol Today;12,100-1,1991.

*Kourilsky P , and Claverie JM.

MHC-antigen interaction : What does the T cell receptor see?
Adv Immunol ;45,107-93,1989.

*Krangel MS.

Secretion of HLA-A and B antigens via an alternative RNA splicing
pathway.
J Exp Med ;163,1173 -90,1986.

*Krensky AM, A Weiss, G Crabtree , M M Davis , and P Parham.

T-lymphocyte -antigen interactions in transplant rejection .
New Eng J of Med ;322,510--517,1990.

*Lafferty, K.J, S.J.Prawse, R.J. Simeonovic , and H.S.Warren .

Immunobiology of transplantation.

Ann Rev Of Immunol; 1:43-173,1983.

*Lechler RI, Batchelor JR.

Restoration of immunogenicity to passenger cell depleted kidney allografts by the addition of donor strain dendritic cells .

J Exp Med ;155,31-41,1982 .

*Lechler RI, Lombardi G, Batchelor JR,Reinsmoen N,and Bach FH.

The molecular basis of alloreactivity .

Immunol Today; 11, 83-8,1990.

*Lillian W. Gabber, A. Osama. Gabber, Elizabeth A. Tolley, and Donna K.Hathaway.

Prediction by postvascularizaion biopsies of cadaveric kidney allografts of rejection, graft loss, and preservation nephropathy.

Transplantation , vol.53, 1219-1225, No 6, 1992.

*Little CC.

In Snell GD (ed). *The Biology of the Laboratory Mouse*.

New York:Dover ,1941

* Madden DR,Gorga JC,Strominger JL, and Wiely DC.

The structure of HLA-B27 reveals nonamer self peptides bound in an extended conformation .

Nature ;353,321-5, 1991.

*Male D,Champion B, Cook A,and Owen M.

Leukocyte surface molecules .

In Advanced Immunology (2nd ed). London :Gower Medical Press,1991

*Marrack P,and Kappler J.

The staphylococcal enterotoxins and their relatives .

Science ;248,705-11,1990.

*Marti, G.E.

Application of lymphocyte immunophenotyping in selected diseases .

Pathol,Immunopathol.Res. 7 (5) :319-337,1989.

*Mason , D.W. and P.J.Morris .

Effector mechanisms in allograft rejection.

Ann Rev Of Immunol;4:119-145,1986.b

*Metlay JP, Pure E,and Steinman RM.

Control of the immune response at the level of antigen -presenting cells : a comparison of the function of dendritic cells and B lymphocytes .

Adv Immunol ;47,45-116,1989.

*Moraes, J.N and Stastny, P.

Eight groups of human endothelial cell alloantigens.

Tissue Antigen 8, 273-276, 1979.

*Paul LC,Baldwin WM, and Van Es La .

Vascular endothelial alloantigens in renal transplantation.

Transplantation ;40,117-23,1985.

*Paul LC, VanES LA, and Fleuren G.

Demonstration of transplantation antigens on the endothelium of peritubular capillaries in renal allografts by the immunoperoxidase method .

Transplantation: 28:72,1979.

*Peterson PA,Cunningham BA,Berggard I and Edelman GM.

2-microglobulin - α free immunoglobulin domain.

Proc Natl Acad Sci (USA) ,69,1697-1701,1972.

*Ploegh HL,Orr Ht,and Strominger JL.

Major histocompatibility antigens :the human (HLA-A,-B,-C) and murine (H-2K,H-2D) class I molecules .

Cell ;24,287-90,1981.

*Poher LC, Collins T, Gimbron MA, Libby P, and Reiss CJ.

Inducible expression of class II major histocompatibility complex antigens and the immunogenicity of vascular endothelium.

Transplantation, 41:141, 1986.

*Poher JS,Guild BC,and Strominger JL.

Phosphorylation in vivo and vitro of human histocompatibility antigens (HLA-Aand ALA-B) in the carboxy -terminal intracellular domain.

Proc Natl Acad Sci (USA) ;75,6002-6,1978.

*Poher,JS,and Cotran.

The role of endothelial cells in inflammation.

Transplantation : 50,537-544,1990.

*Posselt AM,Barker CF,Tomaszewski JE,Markmann JF,Choti MA,and Naji A.
Induction of donor specific unresponsivness by intrathymic islet transplantation .

Science ;249, 1293-5,1990.

*Prowse SJ,Lafferty KJ.

The normal immune response. In Cerilli Gj.

Organ Transplantation and Replacement .

Philadelphia :J.B. Lippincott CO., 1988.

*Rojo S,Lopez D, Calvo V, and Lopez de Castro JA.

Conservation and alteration of HLA-B27-specific T cell epitopes on mouse cells .

J Immunol; 146, 634-42,1991 .

*Rudensky AY, Preston-Hurlburt P, Hong S-c, Barlow A and Janeway CA.
Sequence analysis of peptides bound to MHC class II molecules .
Nature ;353,622-7,1991.

*Russell TJ, Schultes LM, and Kubar DJ.
Histocompatibility (HAL) antigens associated with psoriasis.
New Engl J Med ;287,738-40,1972.

*Salter RD, Benjamin RJ, and Wesley PK.
A binding site for the T-cell co-receptor CD8 on the alpha 3 domain
of the HLA-A2.
Nature ;345,41-6,1990.

*Sedmak, and Orosz.
The role of vascular endothelial cells in transplantation.
Arch Pathol Lab Med : 115,260-265,1991.

*Shapiro, H.M.
The little laser that could :Applications of low power lasers in
clinical flow cytometry.
Ann.N.Y. Acad.Sc., 468:18-27,1984

*Simister NE, and Mostov KE.
An Fc receptor structurally related to MHC class I antigens .
Nature;337, 184-7,1989.

*Simpson E, Chandler P, Goulmy E, Distèche CM, Ferguson-Smith MA, and Page D.

Separation of the genetic loci for the H-Y antigen and for testis determination on human Y chromosome .

Nature ;, 326,876 -8 ,1987 .

*Simpson E.

Immunology of H-Y antigen and its role in sex determination .

Proc R Soc London Ser B; 220,31-46,1983 .

*Snell GD.

Studies in histocompatibility.

Science, 213,172-8,1981.

*Suciu-Foca N, Reed E, and D'Agati VD.

Soluble HLA antigens , anti-HLA antibodies and antiidiotypic antibodies in the circulation of renal transplant recipients .

Transplantation;51,593-601,1991 .

*Tiwari JI, and Terasaki PI.

HLA and Disease .

New York:Springer-Verlag, 1985.

*Todd I, Hammond LJ, James RF, Feldman M, and Botazzo GF.

Epidermal growth factor and transforming growth factor -alpha suppress HLA class II induction in human thyroid epithelial cells .

*Zuniga MC, Malissen B, and McMillan M.

Expression and function of transplantation antigens with altered or deleted cytoplasmic domains .

Cell ,34,535-44,1983