

Ahmed Zadek , N.; Shingu , M. and Nobunaga, M.(1990):

The Effect of Recombinant Tumour Necrosis Factor Alpha on Superoxide and Metalloproteinase production by Synovial cells and Chondrocytes .
Clin. Exp.Rheumatol. , 8 : 387.

Aitchenson , C. T. ; Peebles , C. ; Joplin , F. and Tan , E. M. (1980):

Characteristic of antinuclear antibody in rheumatoid arthritis .
Arth. Rheumat. , 23 : 528.

Akashi, M.; Loussararian,A.H.; Adelman, D.C.; Saito , M. and Koeffler, H.P.(1990):

Role of Lymphotoxin in expression of interleukin- 6 in human fibroblasts : stimulation and regulation

J. Clin. Invest., 85 : 121.

Alam, R.; Bodenbunrg , Y.; Forsythe , P.A.; Lett- Brown , M.A. and Grant, J.A.(1991):

Agonistic - antagonistic property of interleukin-8 on basophils : identification of IL-8 as a potent inhibitor of cytokine - induced histamine release .

J. Allergy Clin. Immunol. , 87:24.

Alderson , M.R.; Sassenfeld, H.M. and Widmer , M.B.(1990):

Interleukin - 7 enhances cytolytic T lymphocyte generation and induces Lymphokine - activated killer cells from human peripheral blood .

J. Exp . Med., 172 : 577.

Allen , G.C. and Kunkle , H.G.(1968):

Hidden rheumatoid factors with specificity for native gamma-globulin .
Arthritis Rheum., 9:758.

Allison, K.C.; Strober, W.and Harriman G.R.(1991):

Induction of IL-5 receptors on normal B cells by cross- linking surface Ig with anti - Ig dextran .

J. Immunol., 146 : 4197.

Altman, A.; Coggeshall, K.M. and Mustelin , T.(1990):

Amos, R.S.; Crockson, R.A. and Crockson, A. P.(1978):

Rheumatoid arthritis : C. reactive protein and erythrocyte sedimentation rate during initial treatment .

Br . Med. J. ,1:1396.

Anderson , K.C.; Morimoto, C.; Paul, S.R.; Chauhan , D.; Williams, D.; Cochran, M. and Barut , B.A.(1990):

Interleukin - II promotes accessory cell- dependent B cell differentiation in humans .

Blood, 80 : 2797.

Andus , T.; Geiger, T.; Hirano , T.and Einrich, P.C (1987):

Recombinant human B-cell stimulatory factor 32 (BSF - 2 IFN γ) regulates b fibrinogen and albumin mRNA levels in Fao-9cells.FEBS. Lett., 221:18.

Ansell , B.M.and Loewi, G.(1977):

Rheumatoid arthritis : general features .

Arthritis Rheum. , 3 : 385.

Arend , W.P. and Dayer , I.M.(1990):

Cytokines and Cytokines Inhibitors or Antagonists in Rheumatoid Arthritis.

Arthritis and Rheumatism , 33: 305.

Arend , W.P.;Joslin ,F.G. and Massoni , R.G.(1985) :

Effects of immune complexes on production by human monocytes of interleukin - 1 or interleukin - 1 inhibitor .

J . Immunol., 134 : 3868.

Armitage , R.;Namen, A.; Sassenfeld , H. and Grabstein , K.(1990) :

Regulation of human T cell proliferation by IL- 7 .

J. Immunol., 144: 938.

Arner, E.C. and Pratta, M.A.(1989) :

Independent effects of IL- I on proteoglycan breakdown , Proteoglycan synthesis and prostaglandin E2 release from cartilage in organ culture .

Arth . Rheum ., 32:288.

Arnett , F.C.; Edworthy , S.M.; and Block, D.A.(1988) :

American Rheumatism Association - 1987 revised criteria for the classification of rheumatoid arthritis .

Arthritis Rheum ., 31 : 315.

Asano , S.; Okano , A.; Ozano,K.; Nakahata, T.; Ishibashi ,T.; Koike , K. and Akigana , Y.(1990) :

In vitro effects of recombinant human IL- 6 primates : stimulation production of platelets .
Blood, 75: 1602.

Atkins , E.(1960):

Pathogenesis of fever .
Physiol. Rev., 40:580.

Aversa , G. ; Punnonen , J. ; Cocks , B. G. and De Vries , J. E.(1993):

An Interlukin-4(IL-4) mutant protein inhibits both IL-4 or IL-13 induced human IgG4 and IgE synthesis and B cell proliferation : Support for a common component shared by IL-4 and IL-13 receptors .
J. Exp. Med. , 178 : 2213.

Bacon , K.B. and Camp , R.D.R.(1990) :

Interleukin - 8 induced in vitro human lymphocyte migration is inhibited by cholera and pertussis toxins and inhibitors of protein kinase C.
Biochem. Biophys. Res. Commun ., 169 : 1099.

Bacon , P. A.(1982) :

Assessment of rheumatic diseases . In : Scientific Basis of Rheumatology , Panay: , G.S ., Churchill , Livingstone , Edinburg , London , Melbourne , New York , p 242.

Balavoine , J.F.; De Rochemonteix , B. ; Williamson, K. ; Sechinger, P. ; Cruchaud, A. and Dayer, J.M.(1986) :

Prostaglandin E₂ and collagenase production by fibroblasts and synovial cells is regulated by urine derived human interleukin-1 and inhibitor (s) .
J. Clin . Invest., 78 : 1120.

Barade , F. A. ; Brien , W. and Horwitz , A.(1981) :

Scientific Basis of Rheumatology . edited by Panayi , G. S. Churchill , Livingstone , Edinburgh , London , New York . , 70.

Barnes, C. G. and Mason, M.(1975) :

Rheumatoid Arthritis . In : An Introduction to Clinical Rheumatology, 2 nd ed. Mason, M., Currey, H. L. Eds . Pitman medical, P. 24.

Barnes, R. ; Hart, C. ; Alexander, L. and Steen, S.(1983) :

Immunological evaluation of renal transplant patients: Changes of B₂ - M , immunoglobulin and complement component during graft rejection .
J. Clin. Lab. Immunol., 12 : 17.

Bastos, M.G. ; Pankewycz, O. ; Rubinkelly, V.E. ; Murphy, J.R. and Strom, T.B.(1990):

Concomitant administration of hapten and IL-2 toxin (DAB₄₈₆- IL-2) results in specific deletion of antigen-activated T cell clones .
J. Immunol. , 145 : 3535.

Baum, J. and Ziff, M.(1979) :

Laboratory finding in rheumatoid arthritis . In : Arthritis and Allied Conditions . A textbook of Reumatology, 9th ed Mc Carty, D. J. , Ed. Lea & Febiger Philade & Phia, P. 491.

Baum , J. and Ziff , M. (1985):

Laboratory findings in rheumatoid arthritis .In : Arthritis and allied conditions . ed. by McCarty , D. J. and Holland , J. L. , 10th ed . Lea. , Febiger , Philadelphia , 40 : 643.

Baumann, H. ; Held, W. A. and Berger, F. G.(1984) :

The acut phase response of mouse live . Genetic analysis of the major acute phase reactants .
J. Biol. chem., 259 : 556.

Baumann, H. and Schendel, P.(1991) :

Interleukin-11 regulates the hepatic expression of the same plasma proteines as interleukin-6
J. Biol. Chem., 266 : 20424.

Beckham, J.C.; Caldwell, D.S.; Peterson, B. L.; Pippen, A.M. and Currie, M.S.(1992):

Disease Severity in Rheumatoid Arthritis : Relationships of Plasma Tumour Necrosis Factor Alpha, Soluble IL-2 receptor, Soluble CD4/CD8 Ratio, Neopterin, and Fibrin Dimer to Traditional Severity and Functional Measures, J. Clin. Immunol., 12 : 353.

Belosevic, M. ; Davis , C.E. ; Meltzer, M.S. and Nacy, C.A(1988):

Regulation of activated macrophage antimicrobial activities ; identification of lymphokines that cooperate with interferon gamma for induction of macrophage resitance to infection .
J. Immunol., 141 : 890.

Belosevic , M. ; Finbloom, D.S. ; Meltzer, M.S. and Nacy, C.A.(1990):

IL-2;a cofactor for induction of activated macrophage resistance to infection .
J. Immunol., 145 : 831.

Benjamin, D. ; Wormsley, S. and Dower, S.K.(1990) :

Heterogenicity in interleukin IL-1 receptors expressed on human B cell lines .

Differences in molecular properties of IL-1 α and IL-1 β binding sites . J. Biol. Chem., 265:9943.

Ben-Sassor, S.Z.; LeGros, G.; Conrad, D.H. ; Finkelman, E.D. and Paul, W.E.(1990):

IL-4 production by T cells from native donors; IL - 2 is required for IL-4 production .

J. Immunol., 145 : 1127.

Berggard, B. ; Bjorck, L ; Cigen, R. and Logdberg, L. (1980):

B₂ - Microglobulin .

Scand. J. Clin. Lab. Invest., 40 : 13.

Bernard, M. ; Vyskocil, A. and Lauwerys, P.(1981) :

Determination of B₂ - M in human urine and serum by latex immunoassay . Clin. Chem., 27 : 832.

Berggard, L. and Bearn, A.G.(1968) :

Isolation and Properties of low molecular weight B₂-globulin occurring in human biological fluids .

J. Biol. Chem., 243 : 4095.

Bessey, O.A.; Lowry, O.H. and Brock, M.J.(1946):

A method for the rapid determination of alkaline phosphatase with five cubic millimeters of serum .

J. Biol. Chem., 164 : 321.

Beutler, B. and Cerami, A.(1987):

Cachectin : More than a TNF .

Negl. J. Med., 316 : 379.

Bevilacqua, M.P.; Pober J.S.; Majeau, G.R.; Cotran R.S . and Gimbrone , M.A. Jr.(1984):

Interleukin-1 (IL-1) induces biosynthesis and cell surface expression of procoagulant activity in human vascular endothelial cells .

J . Exp . Med., 160 : 618.

Bevilacqua M.P.; Stengelin, S.; Gimbrone, M.A.; Bearn, B.(1989):

Endothelial leucocyte adhesion molecule I; an inducible receptor for neutrophils related to complement regulatory proteins and lectins .

Science , 243: 1160.

Bhardwaj, N., Santhanam, U.; Law, L.L.; Taffer, S.B.; Chrayeb, J. and Rivelis, M.(1989):

IL6/INF β 2 in synovial effusions of patients with rheumatoid arthritis and other arthritides . Identification of several isoforms and studies of cellular sources .

J. Immunol., 14:2153.

Bienenstak, H.(1975):

Rheumatoid Planter synovial cysts .

Ann. Rheum. Dis., 39:98.

Bignold, L.P.; Rogers, S.D.; Siaw, T.M. and Bahnisch, J.(1992).:

Inhibition of chemotaxis of neutrophil leukocytes to interleukin - 8 by endotoxins of various bacteria .

Infect. Immunol., 59:4255.

Bischoff, S.C.; Brunner, T.; Deweck, A.L. and Dahinden, C.(1990):

Interleukin-5 modifies histamine release and leukotriene generation by human basophils in response to diverse agonists .

J. Exp. Med., 172:1577, 1990 .

Bjerrum, W. ; Bjerrum, J. and Borregaard, N.(1987):

B₂ - M in neutrophils : An intragranular protein .

J. Immunol., 138 : 3931.

Black, D.R. and Bacon, P.A.(1981):

Ferritin reticuloendothelial cell dysfunction and systemic rheumatoid arthritis .

Ann. Rheum. Dis., 39:368.

Bland, J.H.(1974):

Rheumatoid arthritis of the cervical spines .

J. Rheum., 1: 319.

Bochner, B.S.; Mckelvey, A.A; Klunk, D.; Macglashan, D. and schleimer, R.P.(1989):

Function and expression of CD¹⁸ glycoproteins during IgE- dependent and independent activation of human basophils .

FASEB. J., 3:323.

Bomsztyk, K.; Sims, I.E.; Stanton, T.H.; Slack, J.; McMahan, C.J. ; valentine, M.A. and Dower, S.K.(1989):

Evidence different interleukin-I receptors in murine B- and T- cell lines .

Proc. Natl. Acad. Sci. USA, 86:8034.

Bonvini, E.; Ruscetti, F.W.; Ponzoni, M. and Farrar, W.(1987):

Interleukin-2 rapidly stimulates synthesis and breakdown of polyphosphoionsitides in interleukin-2 dependent, murine T cell lines .
J. Biol . chem., 262 : 4160.

Boom , W.H.; Liano , D. and Abbas, A.K.(1988):

Heterogeneity of helper/ inducer T lymphocytes. Effects of interleukin - 4 and interleukin- 2 producing T cell clones on resting B Lymphocytes .
J. Exp. Med., 167:1350.

Bowers, L.D. and Wang, E.T.(1980):

Kinetic serum creatinine assay.
Clin.Chem., 26:555.

Boyce, B.F.; Aufdemrte , T.B.; Carrett, L.R. and Mundy , G.R.(1989):

Effects of interleukin - 1 on bone turnover in normal mice .
Endocrinology, 125 : 1142.

Bozicevich, J.; Bunim , J.J.; Freund , J. and Ward , S.B.(1958):

Bentonite flocculation test for rheumatoid arthritis. '
Proc. Soc. Exp . Biol . Med ., 97:180.

Breen, E.; Rezai , A.R.; Nakajima, K. and Martinez - Maza, O.(1990):

Infection with HIV is associated with elevated IL-6 levels and production.
J. Immunol., 144:480.

Brennan , F.M.; Chantry, D. and Feldmann, M.(1989):

Inhibitory effect of TNF antibodies on synovial cell IL-1 production in rheumatoid arthritis . Lancet, II : 244.

Brennan , F.M.; Chanty, D.; Turner, M. and Feldman, M.(1990):

Detection of IL-8(m-RNA & protein) in rheumatoid arthritis.
Br.J. Rheumatol., 20 :165.

Broide , D.H.(1991):

Inflammtory cells ;structure and function. In : Stites , D.P. and Terr , A.L. (eds) : Basic and clinical Immunology . 7th edition . Appleton and Lange London, Sydney , Toronto, Mexico, New Delhi . Chapter 12, p141, 1991 .

Broudy, V.C.; Kaushansk, K.; Harlan and Adamson , J.W.(1987):

Interleukin - 1 stimulates human endothelial cells to produce granulocyte - macrophage colony stimulating factors and granlocyte colony stimulating factors .
J. Immunol., 139:464.

Brozik ,M.; Rosztoczg, I.; Mereteg, K. and Balint,G.(1992):

Interleukin- 6 levels in synovial fluids of patients with different arthritides: correlation with local IgM rheumatoid factor and systemic acute phase protein production .

J . Rheumatol., 19:63.

Bruce, A.G.; Hoggatt, I.H . and Rose, T.M.(1992):

Oncostatin M is a differentiation factor for myeloid leukemia cells .

J. Immunol ., 146:121.

Buchanan, W.W.(1978):

Clinical feature of rheumatoid arthritis . In: Copemans Textbook of the rheumatoid Diseases, 5 th . ed . Scott, J.J., .Ed. Livingstone , London , NewYork , p. 318.

Buchanan , W.W. and Kean, W.F.(1986):

Articular and systemic manifestations of rheumatoid arthritis from Copeman's textbook of the Rheumatic Diseases. Edited by Scott. J.T.6 th edition volumel.Churchill, Livingstone, Edinburgh, London, and NewYork ,26:653.

Bucknall, R. C . , Davis , P . ; Bacon , P. A. and Verrier-Jones , J.(1982):

Neutropenia in rheumatoid arthritis: Studies on possible contributing factors .

Ann. Rheum. Dis. , 41 : 242.

Burmester , G.R.; John, B.; Rohwer, P.; Zacher, J.; Winchester. R.J. and Kalden , J.R.(1987):

Differential expression of Ia antigens by rheumatoid synovial lining cells .

J. Clin.Invest., 80:595.

Burstein, H.J.; Tepper , R.I.; Leder, P. and Abbas, A.K.(1991):

Humoral immune functions in IL-4 transgenic mice .

J. Immunol. , 147:2950.

Campen, D.H.; Howitz, D.A. and Ouismoris , F.P. Jr. (1988):

Serum level of IL-2 Receptors and Activity of Rheumatic Active Diseases Characterized by Immune System Activation.

Arthritis Rheum., 31:1358.

Canalis , E.(1986):

Interleukin -1 and its biologically related cytokines .

Adv . Immunol. , 44:153.

Carding , S.R. and Bottomly, K.(1988):

IL-4 (B cell stimulatory factor) exhibits thymocyte growth factor activity in the presence of IL-2 .

J. Immunol ., 140:1519.

Carpenter, A.B.; Eisenbeis, C.H.; Carrabis, S. and Brown, M.C.(1990):

Soluble Interleukin - 2 receptor : Elevated Levels in serum and synovial fluid of patients with rheumatoid arthritis.

J. Clin . Lab. Anal., 4 : 130.

Carpenter, D.F. ; Golden, A. and Robert, W.V.(1967):

Quadrivalvular rheumatoid heart disease associated with left bundle branch block .

Am. J. of med . , 43:922.

Carson , A. A.(1985):

Diagnostic tests : Rheumatoid factor. In: Textbook of Rheumatology . edited by Kelley. M. N.; Harris. E. D. ; Ruddy S. and Sledge C. B. , 2 nd edition : W. B. Saunders Comping. Philadelphia, London, Toronto, Mexico city, Riodejaneiro, Sydney and Tokyo, 44 : 664.

Carson , O.A.; Lawrance , S. and Catalano , M.A.(1977):

Radioimmunoassay of Ig G and IgM rheumatoid factors reacting with human IgG .

J. immunol ., 119 : 295.

Cartwright , G.L. and Lee, G.R.(1971):

The anaemia of chronic disorders .

Br . J. Haemat., 21:174.

Case , J.P.; Lafyatis, R.; Remmers , E.F. ; Kumkumian and Wilder , R.L.(1990):

IL-1 regulation of Transin-Stromelysin transcription in rheumatoid synovial fibroblasts appears to involve two antagonistic transduction pathways; an inhibitory prostaglandin dependent pathway mediated by CAMP , and a stimulatory protein Kinase - c- dependent pathway.

J. Immunol., 145 :3755.

Castell, J.V. ; Geiger, T. ; Gross, V. ; Andus. T. ; Walter, E. ; Hirano, T. ; Kishimoto, T. and Heinerich, P.C.(1988):

Plasma clearance organ distribution and target cells of interleukin-6/hepatocyte-stimulating factor in the rat .

Eur. J. Biochem., 177: 357.

- Castell, J.V.; Gomez-Lechon, M.J.; David, M.; Andus, T.; Geiger, T.; Trul-lenque, R.; Fabra, R. and Heinrich, P.C.(1989):*
Interleukin-6 is the major regulator of acute phase protein synthesis in adult human hepatocytes .
FEBS., 242:237.
- Castenfors , H. ; Hulman , E. and Lovgren , O.(1964):*
The bromosulphthalein test (BSP) as measure of rheumatoid arthritis activity .
Acta. Rheumat. Scand. , 10 : 128.
- Chang, T.L.; Shea, C.M.; Urioste, S.; Thompson, R.C.; Boom, W.H. and Abbas, A.K.(1990):*
Heterogeneity of helper/inducer T lymphocytes . Responses of IL-2 and IL-4 producing (Th1 and Th2) clones to antigens presented by different accessory cells .
J. Immunol., 145 : 1803.
- Chen, W.F. and Zlotnik, A.(1991):*
IL-10; a novel cytotoxic T cell differentiation factor .
J. Immunol., 147 : 528.
- Chizzonite, R.; Truitt, T.; Kilian, P.L.; Stern, A.S.; Nunes, P.; Parker, K.P.; Kaffka, K.L.; Chua, A.O.; Lugg, D.K. and Gubler, U.(1989) :*
Two high-affinity interleukin 1 receptors represent separate gene products .
Proc. Natl. Acad. Sci . USA, 86:8029.
- Coeffier, E.; Joseph, D.; and Vargaftig, B.B.(1991):*
Activation of guinea pig eosinophils by human recombinant IL-5; selective antagonists .
J. Immunol., 147 : 2595.
- Coffman, R.L.; Seymour, B.; Hudak, S. and Rennisk, D.(1989):*
Antibody to interleukin-5 inhibits helminth-induced eosinophilia in mice .
Science, 245 : 308.
- Coffman, R.L.; Seymour, B.W.P.; Lebman, D.A.; Hiraki, D.D.; Christiansen, J.A.; Shrader, B.; Cherwinski, H.M.; Savelkoul, H.F.J.; Finkelman, E.D.; Bond, M.W. and Mosmann, T.R.(1988):*
The role of helper T cell products in mouse B cell differentiation and isotype regulation .
Immunol. Rev., 102 : 5.
- Colditz, I.; Znahlen, R.; Dewald, B. and Baggiolini, M.(1989):*
In vivo inflammatory activity of neutrophil-activating factor, a novel chemotactic peptide derived from human monocytes .
Am.J. Pathol., 134 : 755.

Colley, C.; Fleck, A.; Goode, A.; Muller B. and Myers, M.(1983) :
Early time course of the acute phase protein response in man .
J. Clin. Path., 36 : 203.

Cone , R. O. and Resnick , D.(1985):
Radiographic evaluation of articular disorders .In: Textbook of Rheumatology edited by Kelly , W. M. ; Harris , E. D. ; Ruddy , S. and Sledge , C. B. 2nd ed .
Saunders Company . Philadelphia, London , Toronto, Mexico city, Riodejaneiro , Sydney and Tokyo . 39 :569.

Conrad, D.; Ben-Sasson, S.; LeGros, G. and Paul, W.(1990):
Infection with *Nippostrongylus brasiliensis* or infection of anti-IgD antibodies markedly enhance Fc-receptor-mediated interleukin-4 production by non B, non T cells .
J. Exp. Med., 171 : 1497.

Constable, T.; Crockson, R.; Crockson, A. and McConkey, B.(1975) :
Drug treatment of rheumatoid arthritis .
Lancet, 1 : 1176.

Conti, P.; Real, M.; Fiore, S.; Cancell, A.; Angeletti, P.U. and Dinarello, C.A.(1988) :
Recombinant interleukin-1 and tumor necrosis factor acting in synergy to release thromboxane 6 keto-PGF1 α and PGF2 by human neutrophils .
Scand. J. Rheumatol., 75 : 318.

Cooper, E.H. and Ward, A.M.(1979):
Acute phase reactant proteins as aids to monitoring disease .
Invest. cell. Path., 2 : 293.

Cox, G.; Chattopadhyay, U.; Oppenheim, J. and Varesio., L.(1992):
IL-4 inhibits the costimulatory activity of IL-2 or picolinic acid but not of lipopolysaccharide on IFN- γ treated macrophages .
J. Immunol., 147 : 3809.

Creswell, P. ; Springer , T. ; Strominger , L. and Kubo, T.(1974):
Immunological identity of the small subunit of HLA antigens and B₂ - M and its turn over on the cell membrane .
Proc. Natl. Acad. Sci., 71 : 2123.

Crilly, A. and Madhok, R.(1992):
Relationship between IL-6 levels and platelet counts in systemic juvenile rheumatoid arthritis .
Arthritis and Rheumatism, 35 : 840.

Curtis, B.M.; Widmer, M.B.; deRoos, P. and Qwanrstrom, E.E.(1990):
IL-1 and its receptor are translocated to the nucleus .
J. Immunol., 144 : 1295.

*Darbonne, W.C.; Rice, G.C. Mohler, M.A.; Apple, T.; Hebert, G.A.;
Valente, A.J. and Baker, J.B.(1992):*
Red blood cells are a sink for interleukin-8:a leukocyte chemotaxin .
J. Clin .Invest., 88 : 1362.

Darby, P. W.(1953):
Liver function test in rheumatoid arthritis .
J. Clin. Path. , 6 : 331.

Darby, P.W.(1956):
Liver function tests in rheumatoid arthritis .
J. Clin . Path ., 9:153.

Darlington G.J. Wislon D.R. and Lachman , L.B.(1986):
Monocyte conditioned medium , interleukin - I, and tumor necrosis factor
stimulate the acute phase response in human hepatoma cells in vitro .
J. Cell Biol.,103: 787.

Dasgupta B., Corkill, M.; Kirkham , B. and Panayi, G.(1992):
Serial edtimation of interleukin-6 as a measure of systemic disease in
rheumatoid arthritis .
J. Rheumatol., 19 : 22.

Davenport , R.; Strieter , R.; Standiford, T. and Kunkel , S.L.(1991):
Interleukin-8 production in red blood cell incompatibility.
Blood , 76:2439.

Davis , J.A.; Coben , A.H.; Weisbart, R. and Paulus , H.E.(1979):
Glomerulonephritis in rheumatoid arthritis .
Arthritis Rheum ., 22:1018.

Davis , P. and Hughes , G. R. V.(1974) :
Significance of eosinophilia during gold therapy .
Arth. Rheumatol . , 17 : 964.

*Dayer, J.M.; Rochemonteix , B.O.; Burrus, B.; Demczuk, S. and
Dinarelo , O.A.(1986):*
Human Recombinant IL1 Stimulates Collagenase and PGE 2
Production by Human Synovial Cells.
J. Clin ., 77:645.

Dayer , L . M. and Demezuk,S.(1984):

Cytokines and Other Mediators in Rheumatoid Arthritis .
Springer Semin. Immunopathol., 17:387.

De Benedetti, F.;Massa, M .; Robbiori, P. and Martiri, A.(1991):

Correlation of serum interleukin-6 levels with joint involvement and thrombocytosis in systemic Juvenile rheumatoid arthritis.
Arthritis and Rheumatism, 34:1158.

Defrance , T . ; Carayon , P . ; Minty , A. and Ferrara , P.(1994):

IL-13 is a B-cell stimulating factor .
J. Exp. Med. , 179 : 135.

De Jana, E.; Brevario, F; Erroi, A. and Mantovani ,A.(1987):

Interleukin - 1 and its biologically related cytokines.
Blood, 69 : 695.

Depper, J.M.; Leomard, W.J. and Creene , W.C.(1985):

Interleukin-2 (IL-2) augments transcription of the IL-2 receptor gene .
Proc. Natl. Acad . Sci.USA, 82: 4230.

De-Waal-Malefyt, R.H.; Spits, Il. and Kastelein, R.(1992):

Interleukin- 10 and viral IL-10 strongly reduce antigen specific human T cell proliferation by diminishing the antigen-presenting capacity of monocytes via down - regulation of class II major histocompatibility complex expression .
J. Exp. Med., 174:915.

Dewhirst, F.; Stashenko, P.; Mole , J. and Tsurumachi, T.(1985):

Purification and partial sequence of human osteoclast- activating facotr : Identity interleukin beta .
J. Immunol., 135:2562.

Dextar, T.M.(1989):

Haemopoietic growth factors .
Br. Med . Bulletin , 45:337.

Dick , W. C. and Goodacre , J.(1985):

Rheumatoid arthritis. In : Introduction of Clinical Rheumatology. 2nd edition English language book . Society / Churchill Livingstone , 4 : 46.

Dieppe, P.A.; Doherty, Macjarlane , D., and Maddison , P.(1985):

Rheumatological Medicine. Churchill Livingstone, Edinburgh, London, Mebourne and New York. 4 : 41 and 26 : 479.

Dinarello , C.A.(1987):

The biology of interleukin-1 and comparison to tumor necrosis factor .
Immunol . Lect.,16:227.

Dinarello, C.A.(1989):

Interleukin -1 and its biologically related cytokines.
Adv. Immunol .44:153.

Dinarello , C.A.(1991):

Interleukin-1 and interleukin - 1 antagonism.
Blood 77: 1627.

Dinarello., C.A.; Cannon , J.G. and Wolff, S.M.(1988):

New concepts on the pathogenesis of fever .
Rev. Infect. Dis . , 10:168.

Dinarello C.A.; Rosenwasser L.J., and Wolff S.M.(1982):

Interleukin- 1 and its biologically related cytokines .
J. Immunol., 127:2517.

Dixon , A.St.J.(1971) :

The rheumatic foot in Modern trends in Rheumatology . edited by Hill.
A.G.and Butterworths . London. 11:158.

Dixon, J.S.; Bird, H.A. and Wright, V.(1981):

A Comparison of serum biochemistry in ankylosing spondylitis
seronegative and seropositive rheumatoid arthritis .
Ann. Rheum . Dis. , 40:404.

Dixon , J.; Pickup, M.; Lowe, J.; Lee, M. and Wright , V.(1980):

Discriminatory indices of response of patients with RA treated with D-
Penicillamine .
Ann. Rheum . Dis., 39:301.

Doherty , T. M. ; Kastelein , R. ; Menon , S. and Andrade , S.(1993):

Modulation of murine macrophages function by IL-13 .
J. Immunol. , 151 :7151.

Dower, S.K.; McMahan, C.; Flack, J.; Lupton, S. and Sins , G.E.(1990):

Molecular characterization of two types of IL- 1 receptor coding peptides .
J. Leukoc. Biol., 1:103.

Dresner, E. and Trombly, P.(1959):

The latex fixation reaction in non - rheumatoid diseases .
N. Engl. J. Med., 261: 981.

Du, X.X.; Neben, T.; Goldman, S. and Williams, D.A.(1993):

Effects of recombinant human interleukin - 11 on hematopoietic reconstitution in transplant mice ; acceleration of recovery of peripheral blood neutrophils and platelets .

Blood, 81:27.

Duncan . M.R. and Berman ,B.(1985):

Interferon is the Lymphokine and interferon is the monokine responsible for inhibition of fibroblast collagen production and late but not early fibroblast proliferation .

J. Exp. Med., 162:516.

Dustin , M.L.; Stauton , D.E. and Springer , T.A(1988):.

Supergene families meet at the immune system .

Immunology Today , 2:213.

Eastgate, J.A.; Symons , J.A.; Wood, N.C.; Grinlinton , F.M.; Digiovine, E.S. and Duff, C.W(1988):.

Correlation of plasma Interleukin-1 levels with disease activity in rheumatoid arthritis.

Lancet , 11: 706.

Edward, D. and Harris , J.R.(1981):

Rheumatoid arthritis:the clinical spectrum.In:Textbook of Rheumatology , vol . 1 Keue, M.W.; Harris, E.D.; Ruddy , S . and Sledge, C.B. (Eds). Saunders company . P, 930.

Ellis , H. A. and Felix-Davis , D.(1959):

Serum complement, rheumatoid factor and serum proteins in rheumatoid disease and systemic lupus erythematosus .

Ann. Rheum. Dis. , 18 : 215.

El- Shaer, O.S.; Zidan, M. and Ibrahim, A.A.(1992):

Serum and urinary B₂ - Microglobulin in patients with rheumatoid arthritis.

The New Egyptian Journal of Medicine , 7 :1.

Ennen, J.; Ersnt, M. and Flad H.D.(1991):

Activation of human monocytes by interleukin -2; role of T lymphocytes .

J. Biolumin. chemilumin. , 6: 3.

Esperson, G.T.; Vestergaard, M.; Ernst, E. and Grunnet , N.(1991):

Tumour necrosis factor alpha and Interleukin -2 in plasma from rheumatoid arthritis patients in relation to disease activity.

Clin. Rheumatol., 10 : 374.

Espinoza- Delgado, I .; Longo , D.; Gusella , G. and varesio , L.(1990):
IL-2 enhances c-fms expression in human monocytes.
J. Immunol ., 145: 1137.

Evrin, E., Peterson , A. and Widel, L.(1971):
Radioimmunoassay of B₂ - M in human biological fluids .
Scand. J. Clin. Lab. Invest., 28 : 439.

Fanslow, W.C.; Clifford KN.; park, L.S.; Rubin, A.S.; voice, R.F.; Beckmann, M.P. and Widmer, M.B.(1991):
Regulation of alloreactivity in vivo by IL-4 and the soluble IL- 4 receptor .
J. Immunol., 147:535.

Fryling, C.; Dombalagian, M.; Burgess, W.; Hollander, N. and Schreiber, B.B.(1989):
Purification and characterization of tumor inhibitory factor -2 : Its identity to interleukin-1 .
Cancer Res., 49:3333.

Farr, M.; Scott, D.; Constable, T.; Kawkins, C. and Staurt, J .(1983):
Thrombocytosis of active rheumatoid disease .
Ann . Rheun . Dis ., 42 : 545.

Feldmann, M.; Bernnan , F.M . and Maini, R.N.(1990):
Cytokine production in rheumatoid joint implications for treatment .
Ann . Rheum. Dis., 49 : 480.

Feldman , C.M. and Finbloom , D.S.(1990):
Induction and regulation of IL- 4 receptor expression on murine macrophage cell lines and bone marrow derived macrophages by IFN- γ .
J. Immunol. , 145 : 854 .

Fernandez - Botran, R.; Sanders , V.M. and Vitetta , E.S.(1989):
Interactions between receptor for interleukin-2 and interleukin-4 on lines of helper T cells (HT-2) and B lymphoma cells (BCL-1) .
J. Exp. Med., 169: 379.

Ferrua , B. ; Vincent C. and Revillard , P.(1980):
A sandwich method of enzyme immunoassay , assay for human B₂ -M compared with radioimmunoassay .
J. Immunol. Methods , 36 : 149.

Fine J.M. ; Lanbin , P. and Desjobert, H .(1988):
Determination of serun B2- microglobulin .
Acta Med . Scand ., 224:179.

Fiorentino, D. F.; Bond, M.A. and Mosmann, T.R.(1989):

Two types of T helper cell IV, Th-2 clones secrete a factor that inhibits cytokine production by TH-1 clones.

J. Exp. Med., 170:2081.

Fiorentino, D.; Zlotnik, A.; Mosmann, T. and Garra, A.(1992):

Interleukin-10 inhibits cytokine production by activated macrophage.

J. Immunol., 147: 3815.

Firestein, C.S. and Zvaifler, N.J.(1987):

Peripheral blood and synovial fluid monocyte activation in inflammatory arthritis, low levels of synovial fluid and synovial tissue interferon suggest that interferon is not the primary macrophage activating factor.

Rheum., 30 : 864.

Fleming, A.; Crown, J.M. and Corbett, M.(1976):

Early rheumatic disease onset.

Ann. Rheum. Dis., 35 : 357.

Forman, D.; Finn, W. and Mandel, S.(1983):

Comparison of enzyme immunoassay for serum B₂ - M with a radioimmunoassay technique.

Clin. Chem., 29 : 1245.

Frank, O. and Klemmayer, K. Z.(1979):

Rheumatol., 27 :142,1968. Quoted from Arthritis and Allied conditions :

A textbook of Rheumatology, McCarty, D.L.;

Ed. Lea, Febiger, Philadelphia.

Freeman, C.J.; Freedman, A.S.; Rabinowe, S.N.; Segil, J.M.;

Horowitz, J.; Rosen, K.; Whitman, J.F. and Nadler, L.M.(1989):

Interleukin - 6 gene expression in normal and neoplastic B cells.

J. Clin. Invest., 83:1512.

Fuji, M.; Sugamura, K.; Nakai, S.I.; Tanaka, Y.; Tozawa, H. and Hinuma, Y.(1986):

High and low affinity interleukin-2 receptors; distinctive effects of monoclonal antibodies.

J. Immunol., 137 : 1552.

Furukawa, S. and Matsubara, T.(1992):

Kawasaki disease with regards tumor necrosis factor and IL-6.

Eur. J. Pediatr. 11 : 44.

Gabay, C.; Silacci, P.; Genin, B. and Guerne, P.A.(1995):

Soluble IL- 6 receptor strongly increase the production of acute phase protein by hepatoma cells but exerts minimal changes on human primary hepatocyte .

Eur . J. immunol., 25: 2378.

Galizzi, J.; Zuber, C.; Cabrillat, H. and Banchereau, J.(1989):

Internalization of human interleukin 4 and transient down -regulation of its receptor in the CD23 inducible cells .

J.Biol. Chem., 264:6984.

Gallay, P.; Mach, J.P. and Carrel, S.(1991):

Characterization and detection of naturally occurring antibodies against IL-1 α and IL-1 β in normal human plasma .

Eur . Cytokine Netw., 2 :329.

Gallis, B.; Prickett, K.S.; Jackson, J.; Schooley, K.; Sims, J.E. and Dower, S.K.(1989):

IL-1 induces rapid phosphorylation of the IL-1 receptor.

J. Immunol., 143 : 3235.

Gallo, P.; Frei, K.; Rordorf, C.; Lazdins, J.; Tavitato, B. and Fontana, A.(1989):

Human immunodeficiency virus type I (HIV-I) infection of the central nervous system; an elevation of cytokines in cerebrospinal fluid .

J. Neuroimmunol., 23:109.

Gately, M.K.; Wolizky, A.G.; Quinn, P.M. and Chizzonite, R.(1992):

Regulation of human cytolytic lymphocyte responses by interleukin - 12.

Cell Immunol., 143 : 127.

Gatter, R. A. and Richmond, J. D.(1975):

Predominance of synovial fluid lymphocytes in early rheumatoid arthritis .

J. Rheumatol. , 2 : 340.

Gauldie, J.; Richard, C.; Harnish, D. and Baumann, H.(1987):

Interferon B₂/B-cell stimulatory factor type 2 shares identity with monocyte-derived hepatocyte stimulating factor and regulate the major acute phase protein response in liver cells .

Proc. Natl. Acad. Sci. USA, 84 : 7251.

Geiger, T.; Andus, T.; Klapproth, J.; Hirrano, T.; Kishimoto, T. and Heinrich, P.(1988):

Induction of rat acute-phase proteins by interleukin-6 in vivo .

Eur. J. Immunol., 18:717.

Gerondakis, S.(1990):

Structure and expression of murine germline immunoglobulin heavy chain transcripts induced by interleukin-4 .

Proc. Natl. Acad. Sci. USA, 87 : 1581.

Gewurz, H.; Mold, C.; Siegel, J. and Fiedel, B.(1982):

C-reactive protein and the acute phase response .

Adv. Internal. Medicine, 27 : 345.

Ghezzi, P. and Dinarello, C.A.(1989):

Interleukin-1 and its biologically related cytokines .

J. Immunol., 140 : 4238.

Gimbrone, M.; Obin, M.; Luis, E.; Hass, R.; Hebert, C.; Yip, Y.; Leung, D.; Lowe, D.; Kohr, W.; Darbonne, W. and Baker, J.B.(1989):

Endothelial interleukin-8; a novel inhibitor of leukocyte endothelial interactions .

Science, 246 : 1601.

Glynn, L.E.(1972):

Pathology, pathogenesis and aetiology of rheumatoid arthritis .

Ann. Rheum. Dis., 13 : 412.

Goddard, D.H.; Kirk, A.P.; Kerwan, J.R.; Johnson, G.D. and Holborow, E.J.(1984):

Impaired polymorphonuclear leucocytes chemotaxis in R.A.

Ann. Rheum. Dis., 43 : 151.

Goetz, E.J.(1976):

Defective responsiveness of ascorbic acid of neutrophil random and chemotactic migration in Felty's syndrome and SLE.

Ann. Rheum. Dis., 35 : 510.

Goitsuka, R.; Ohashi, T.; Ono, K.; Yasukawa, Y.; Koishibara, Y.; Fukui, H.; Ohsugi, Y. and Hasegawa, A.(1990):

IL-6 activity in infectious peritonitis .

J. Immunol., 144 : 2599.

Golde, D.W. and Glaspy, J.(1990):

Therapeutic use of granulocyte and monocyte colony-stimulating factors .

IN: Williams, W.J., Beutler, E.; Erlser, A.J. and Lichtman, M.A. : Haematology . McGraw-Hill, Inc, New York, St Louis, San Francisco, Colorado, Springs, Auckland . Chapter 27, p 273.

Golding, D.N.(1982):

Synopsis Rheumatic Disease, 4th ed. Bristol wright .
J., Sons, LTD.P. 63 .

Goldring, M.; Brikhead, J.; Sandell, L. and Krane, S.(1988):

IL-1 suppresses expression of cartilage specific type II and IX collagen and increases type I & Type III collagens in human chondrocytes .
J. Clin. Invest., 82: 2026.

Goldstein, H.; Koerholz, D.; Chesky, L. and Ambrus, J.L.(1990):

Divergent activities of protein kinases in IL-6 induced differentiation of a human B cell line .
J. Immunol., 145 : 952.

Good, R.A.; Venters, H.; Page, A.R. and Good, T.A.(1961):

Diffuse connective tissue diseases in childhood, with a special comment on connective tissue diseases in patients with gammaglobulinemia .
J. Lancet, 81 : 192.

Goodwin, R.G.; Friend, D.; Ziegler, S.F.; Jerzy, R.; Falk, B.A.; Gimpel, S.; Cosman, D.; Dower, S.K. and Namen, A.E.(1990):

Cloning of the human and murine interleukin-7 receptors; demonstration of a soluble form and homology to a new receptor superfamily .
Cell, 60 : 941 .

Granstein, R.; Murphy, G.; Margolis, R. and Amento, E.(1987):

γ -Interferon Inhibits Collagen Synthesis in vivo in the mouse.
J. Clin. Invest., 79: 1254 .

Greco, C.; Ameglio, F.; Alvino, S.; Cianciulli, A. and Lopez, M.(1994):

Selection of patients with monoclonal gammopathy of undermined significance is mandatory for a reliable use of Interleukin-6 and other non specific multiple myeloma serum markers .
Acta Haemat., 92: 1 .

Gross, V.; Andus, J.; Caesar, L. and Scholmerich, J.(1992):

IL-6 production in Grahm's disease.
Castroenterology, 102 : 514 .

Guerne, R.; Zuraw, B.; Vaughan, J.; Carson, D. and Lotz., M.(1989):

Synovium as a source of IL-6 in vitro contribution to local and systemic manifestation of arthritis.
J. Clin. Invest., 83 : 858 .

Haatagia, K., and Kalliomaki, J.L.(1978):

Laboratory scale for evaluating the activity of R.A.
Rhum. and Rehabilitation,17:83.

Hadden, J.W.; Hadden, E.M. and Coffey, R.G.(1987):

Interleukin-2 increases cyclic GMP levels in immature thymocytes and mitogen-primed T lymphocytes .
Int. J. Immunopharmacol., 9 : 851.

Hamerman, D.(1970):

Views on the pathogenesis of rheumatoid arthritis .
Med. Clin. Noth. Am., 1:322.

Hanolon, S.M.; Panay, G.S. and Laurent, R.(1980):

Defective polymorphnuclear leucocyte chemotaxis in rheumatoid arthritis associated with a serum inhibitor .
Ann. Rheum. Dis., 39:68.

Harada, N.; Takahashi, T.; Matsumoto, M. and Takatsu, K.(1987):

Production of a monoclonal antibody useful in the molecular characterization of murine T cell replacing factor/B cell growth factor II .
Proc. Natl. Acad. Sci. USA, 84 : 4581.

Harris, E.D.(1981):

Pathogenesis of rheumatoid arthritis in "Textbook of Rheumatology" edited by Kelley. W.N.; Harris, E.D.; Ruddy, S. and Sledge, C.B. Saunders Company . Philadelphia. Vol. 1 :896.

Harris , E. D.(1985):

Rheumatoid Arthritis : The clinical spectrum.In: Textbook of Rheumatology edited by : Kelley . W. N. ; Harris , E. S. ; Ruddy , S. and Sledge , C. B. 2 nd edition . W. B, Saunders Company. Philadelphia , London , Toronto , Mexico city, Riodejanerio, Sydney and Tokyo . , 60 : 950.

Harris, R.I.; Stone,P.C. and Stuart,J. (1984):

C-reactive protein rapid assay techniques for monitoring resolution of infection in immunosuppressed patients.
J.Clin. Path.,37:821.

Hart, P.H.; Vitti, G.F.; Burgess, D.R.; Whitty, G.A.; Piccoli, D.S. and Hamilton, J.H.(1989):

Potential anti-inflammatory effects of interleukin 4; suppression of human monocyte tumor necrosis factor- α , Interleukin-1 and prostaglandin E .
Proc. Natl. Acad. Sci. USA, 86 : 3803.

Haslock, I.(1983):

Report of the combined annual provincial meeting of the British Association for Rheumatology and Rehabilitation . The Herberden Society . The Royal Society of Medicine Section of Rheumatology and Rehabilitation and Irish Society of Rheumatology and Rehabilitation .
British J. of Rheum. , 22 : 57.

Hays, G. S. and Stinson, I. N.(1976):

Erythrocyte sedimentation rate and age .
Arch. Ophthalmol. , 94 : 939.

Hebert, C.A.; Luscinskas, F.W.; Kiely, J.M. and Baker, J.B.(1990):

Endothelial and leukocyte forms of IL-8; conversion by thrombin and interactions with neutrophils .
J. Immunol., 145 : 3033.

Hechtmant, D.; Cybulsky, M.; Baker, J. and Gimbrone, M. (1990):

Intravenous endothelial interleukin-8 reduces neutrophil accumulation at intradermal sites of inflammation .
FASEB.J., 4 : 890.

Heinzel, F.; Sadick, M.; Holaday, B. and Locksley, R.(1989):

Reciprocal expression of interferon- γ or interleukin-4 during the resolution or progression of murine leishmaniasis .
J. Exp. Med., 169 : 59.

Helfgott, D.; May, L.; Sthoeger, Z.; Tamm, I. and Sehgal, P. (1987):

Bacterial lipopolysaccharide (endotoxin) enhances expression and secretion of $\beta 2$ interferon by human fibroblasts .
J. Exp. Med., 166 : 1300.

Helin, H.; Korpela, M.; Mustonen, J. and Pasternack, A.(1986):

Mild mesangial glomerulopathy with hematuria or proteinuria .
Nephron, 42:224.

Helle, M.; Boeije, L. and Aarden, L. (1991):

ELISA for IL-6 in serum and synovial fluid.
J.Immunol. Methods, 138:47.

Helle, M.; Boeije, L.; Pascual-Salcedo, D. and Aarden, L.(1991):

Differential induction of interleukin-6 production by monocytes, endothelial cells and smooth muscle cells .
Prog. Clin. Biol. Res., 367 : 61.

Helle, M.; Brakenhoff, J.; De-Groot, E. and Aarden, L.(1988):

Interleukin-6 is involved in interleukin-1-induced activities .

Eur J Immunol., 18:957.

Hellsing , K. and Enstrom , H.(1977):

Pre-treatment of serum samples for immuno-nephelometric analysis by precipitation with polyethylene glycol .

Scand. J. Clin. Lab. Invest. , 37 : 529 .

Henderson, B. and Pettipher, E.R.(1989):

Actions of Recombinant IL-1 and TNF- α in the rabbit evidence for synergistic interacting between cytokines in vivo .

Clin. Exp. Immunol., 75: 306.

Heptinstall, R.H.(1983):

Pathology of the kidney .

Little Brown and Company, Boston , 1.

Hermann, G.G.; Geertsen, P.F.; Vonder,A. and Zeuthen, J.(1992):

Recombinant Interleukin-2 and lymphokine activated killer cell treatment of advanced bladder cancer; clinical results and immunological effects .

Cancer Res., 52:726.

Herzog, C.; Herzog, A.; Aeschlimann, A. and Muller, W.(1990):

SIL-2R in chronic polyarthritis .

Z-Rheumatol., 49 : 8.

Hetchman, D.H.; Cybulsky, M.I. and Gimbrone, M.A.(1991):

Inhibitor of polymorphonuclear leukocytes accumulation at sites of acute inflammation .

J. Immunol., 147:883.

Hill,A.B.(1979):

Statistical analysis ;In : A Short Textbook of Medical Statistics ; Hill , A.B., (10th) ed .; 1.

Hill, D.B.; Marsxano, L.; Cohe, D and McClain, C.J.(1992):

Increased plasma IL-6 in alcoholic hepatitis .

J. Lab. Clin. Med., 119 :547.

Hill, R.J.; Warren, M.K. and Levin, J.(1990):

Stimulation of thrombopoiesis in mice by human recombinant IL-6 .

Clin. Invest., 85:1242.

Hirano, T.(1992):

Interleukin-6 and its relation to inflammation and disease .
Clin. Immunol. Immunopathol., 62:60, 1992 .

Hirano, T. and Kishimoto, T.(1988):

Interleukin-6 (IL-6), in Sporn M.B. and Roberts A.B. Handbook of Experimental Pharmacology "Peptide Growth Factors and Their Receptors" Springer-Verlag, Berlin.

Hoekman , K.; Nieuwkoop, J. and Willemze , R.(1985):

The significance of B₂ - M in clinical medicine .
Neth. J. Med. , 28 : 551.

Hokanen, E.; Tornroth, T.; Pettersson, E. and Skifvars, B.(1987):

Membranous glomerulonephritis in rheumatoid arthritis not related to gold or D-penicillamine therapy : a report four cases and review of the literature
Clin. Nephrology, 27:87.

Holborow , E.(1978):

R.Fs and antinuclear antibodies.In:Copeman's Textbook of The Rheumatic Diseases . edited by Scott , J. T. Livingstone , London . 5th ed. , 167.

Holbrook, S.; Ohls, R.; Schibler, K. and Christensen, R.(1991):

Effect of interleukin-9 on clonogenic maturation and cell cycle status of fetal and adult hematopoietic progenitors .
Blood, 77:2129.

Hollander, A.; Atkins, R.; Eastwood, D. and Elson C.(1991):

Human cartilage is degraded by RA synovial fluid but not by recombinant cytokines in vitro.
Clin.Exp.Immunol., 83 : 52.

Holmes, W.E.; Lee, J.; Kuang, W.J.; Rice, G.C. and Wood, W.I.(1991):

Structure and functional expression of a human interleukin-8 receptor .
Science, 253:1278.

Homma,T.; Tsujinaka; Kido.Y.; Yasuda,T. and Mori, T.(1992):

Secrtion of B₂-M from human hepato-plastoma cells and hepatoma cells on stimulation with IL- 6 .
Eur. Surg. Res., 24 :204.

Honda, M.; Kjtamura, K.; Mizutani, Y. and Tokunaga, T.(1991):

Quantitative analysis of serum IL-6 and its correlation with increased levels of serum IL-2R in HIV induced diseases .
J. Immunol., 145:4059.

Hooper, B.; Whittingham, S.; Mathews, J.D.; Mackay, I.R. and Curnow, D.H.(1972):

Autoimmunity in rural community .
Clin. Exp. Immunol., 12:79.

Horii, Y.; Muraguchi, A.; Suematsu, S. and Kishimoto, T.(1988):

Regulation of BSF-2/IL-6 production by human mononuclear cells;
macrophage-dependent synthesis of BSF-2/IL-6 by T cells .
J. Immunol., 141:1529.

Horuk, R.; Huang, J.J.; Covington, M. and Newton, R.C.(1987):

A biochemical and kinetic analysis of the interleukin-1 receptor .
Evidence for differences in molecular properties of IL-1 receptors .
J. Biol. Chem., 262:16275.

Horwitz, D.A.; Linger-Israeli, M. and Gary, J.D.(1987):

Interleukin-1 and its biologically related cytokines .
J. Immunol. Immunopharmacol., 7:43.

Houssian, F.A.; Devogelaer, J.P. and Van Snick, J.(1988):

IL-6 in synovial fluid and serum of patients with R.A. and other arthritidis .
Arthritis rheum., 31-784.

Howard, M.; Farrar, J.; Hilfiker, M.; Johnson, B.; Takatsu, K.; Hamaoka, T. and Paul, W.E.(1982):

Identification of a T cell derived B cell growth factor distinct from IL-2 .
J. Exp. Med., 155:914.

Howe, G.B.; Fordham, J.N.; Brown K.A. and Curry, H.L.F.(1981):

Polymorphnuclear cell function in rheumatoid arthritis and Felty's syndrome .
Ann. Rheum. Dis., 40:370.

Huang, S. ; Xiao, H.; Paciotti, G. and Lawrence, M.(1995):

IL-13 expression at the sites of allergen challenge in patients with asthma .
J. Immunol. , 155 : 2688.

Huber, A.R.; Kunkel, S.L.; Tood, R.F. and Weiss, S.J.(1991):

Regulation of transendothelial Neutrophil Migration by Endogenous IL-8 .
Science: 254:99.

Hughes, G. R. V. and Currey, H. L. F.(1982) :

Laboratory investigation In:"An introduction of clinical rheumatology".
edited by Mason, M. Pilman Medical. P. 348 .

Hultner, L.; Druez, C.; Moeller, J.; Uyttenhove, C.; Schmitt, E.; Ruede, E.; Dormer, P.A. and Van Snick, J.(1990):

Mast cell growth enhancing activity (MEA) is structurally related and functionally identical to the novel mouse T cell growth factor P40/TCGFII (Interleukin-9) .

Eur. J. Immunol., 20:1413.

Hultner, L. and Moeller, J.(1990):

Mast cell growth enhancing activity stimulates interleukin-6 production in a mouse bone marrow derived mast cell line and a malignant subline .

Exp. Haematol., 18:873.

Hurd, E. R. and Cheatum, D. E.(1976):

Decreased spleen size and increased neutrophils in patients with Felty's syndrome . Effects of gold sodium thiomalate therapy .

JAMA. , 235 : 2215.

Iida, Y.; Miyata, T.; Inagi, R.; Sugiyama, S. and Maeda, K.(1994):

B₂- microglobulin modified with advanced glycation end products induces interleukin-6 from human macrophages, role in the pathogenesis of hemo dialysis-associated amyloidosis .

Biochem. Biophys. Res. Commun., 201:1235.

Ishibashi, T.; Kimwra, H. and Shikama, Y.(1989):

Interleukin-6 is a potent thrombopoietic in vivo in mice .

Blood, 74:1241.

Ishimi, Y.; Miyaura, C.; Jin, C.H.; Akatsu, T. and Suda, T.(1990):

IL-6 is produced by osteoblasts and induces bone resorption .

J. Immunol., 145:327.

Ishizaka, T.; Saito, H.; Hatoke, K. and Ishizaka, K.(1989):

Preferential differentiation of inflammatory cells by recombinant human interleukins .

Int. Arch. Allergy Appl. Immunol., 88:46.

Ito, H.; Kishikawa, T. and Toda, T.(1983):

Serum acute phase reactants in pediatric patients especially in neonates .

Japan. J. Surg., 13 :506 .

Jablons, D.M.; Mule, J.J.; McIntosh, J.K.; Sehgal, P.B.; May, L.T.;

Huang, C.M.; Rosenberg, S.A. and Lotze, M.T.(1989):

IL-6/IFN γ -2 as a circulating hormone . Induction by cytokine administration in humans .

J. Immunol., 142:1542.

Jaffe, H.S. and Sherwin, S.A.(1994):

Immunomodulators . IN: Stites, D.P. and Terr, A.I.(eds): Basic and Clinical Immunology . Appleton and Lange, London, Sydney, Toronto, Mexico, New Delhi . 7th edition . Chapter 59, p 781.

Jelinke, D.F. and Lipsky, P.E.(1988):

Inhibitory influence of IL-4 on human B cell responsiveness .
J. Immunol., 141:164.

Johanson, A.M.(1982):

Plasma protein assays. In: Clinical Diagnosis and Management . Beckman Sci., Bulletin, 6215, California.

Johanson, S. V. and Strandberg, P. O.(1972):

Haem biosynthesis studied in patients with rheumatoid arthritis .
J. Clin. Path. , 25 : 159.

Johnson, K.W. and Smith, K.A.(1990):

CAMP regulation of IL-2 receptor expression; selective modulation of the p75 subunit .
J. Immunol., 145:1144.

Joost, J. O. ; Francis, W.R. and Connie, F.(1994):

Cytokines. In : Basic and Clinical Immunology. Daniel, P. S. and John, D.S.edition . 8th eds.

Kakumus, F.; Askinagawa and Kusakabe, A.(1992):

Localisation of intrahepatic IL-6 in patients with acute and chronic liver diseases .
J. Clin. Pathol. 45:408.

Kampschmidt, R.F.(1981):

Leukocytic endogenous mediator/endogenous pyrogen, in Powanda MC, Canonico PG (eds) Leukocytic Endogenous Mediator/Endogenous Pyrogen . Amsterdam, the Nether, Elsevier/North Holland, p 55.

Kapp, A. and Zeck-Kapp, G.(1991):

Interleukin-5-induced granulocyte activation in atopic patients .
Br. J. Dermatol., 125:108.

Karasuyama, H.; Melchers, F. and Rolink, A.(1988):

Recombinant Interleukin-2 or 5 but not 3 or 4, induces maturation of resting mouse B lymphocytes and propagates proliferation of activated B cell blasts .
J. Exp. Med., 167:1377.

Karch, J.; Halbert, S.P.; Klima, E. and Steinberg, A.D.(1980):

Quantitative determination of RF by an enzymatic-labelled immunoassay .
J. Immunol. methods, 32:115.

Karlsson , A. ; Groth , T. and Peterson , A.(1980):

Turn over in humans of B₂ - M the constant chain of HLA antigen .
Eur. J. Clin. Invest . , 10 : 293.

Kato, K.; Yokoi, T.; Takano, N. and Taniguchi, N.(1990):

Detection by in situ hybridization and phenotypic characterization of cell expressing IL-6 mRNA in human stimulated blood .
J. Immunol., 144: 1317.

Katona, I.M.; Urban, J.F.; Kang, S.S. and Finkelman, F.D.(1991):

IL-4 requirements for the generation of secondary in vivo IgE responses .
J. Immunol., 146:4215.

Kawano, M.; Hirano, T.; Matsuda, T.; Taga, T.; Horri, Y.; Iwato, K.; Asaoku, H.; Tang, B. and Kishimoto, T.(1988):

Autocrine generation and requirement of BSF-2/IL-6 for human multiple myelomas .
Nature, 332:83.

Kawano, M.; Yamamoto, I.; Iwato, K.; Tanaka, H.; Asoku, H.; Tanabe, O.; Ishikawa, H.; Nobuyoshi, M.; Ohmoto, Y. and Kuramoto, A.(1989):

Interleukin 1- β rather than lymphotoxin as the major bone resorbing activity in human multiple myeloma .
Blood, 73:1646.

Kawashima, I.; Ohsumi, J.; Miyadai, K. and Takiguchi, Y.(1992):

Function, molecular structure, and gene expression of interleukin-11 (IL-11/AGIF) .
Nippon Rinsho, 50 :1833.

Kean, W. F. ; Dwosh, I. L.; Ford , P. M. and Kelley , H. G.(1980):

The toxicity pattern of D-penicillamine therapy : A guide to its use in rheumatoid arthritis .
Arth. Rheumatol. , 23 : 158.

Kelleher, K.; Bean K.; Clark, S.C. and Yang, Y.C.(1991):

Human interleukin-9; genomic sequence, chromosomal location, and sequences essential for its expression in human T cell leukemia virus (HTLV) I, transformed human T cells .
Blood, 77 :1436.

Kendall, M.G.(1976):

Biochemical studies in the assesment of rheumatic disease .
Clin. Rheum. Dis., 2:253.

Kennedy, A.C. and McGavin, D.D.M.(1975):

Rheumatoid scleritis producing exophthalmos .
Br. J. Clin. Practice, 29:73.

Kerr, L.D.; Oashaw, N.E. and Matrisian, I.M.(1988):

Transforming growth factor beta-1 and cAMP inhibit transcription of
epidermal growth factor and oncogene-induced transin RNA .
J. Biol. Chem., 263:16999.

Kierat, J.(1973):

Liver disease in R.A. and Sjogren's syndrome .
Rheumatologia, 11:11.

Kirkham, B.; Chikanza, I; Pitzalis, C. and Panayi, G.S.(1991):

Monoclonal antibody therapy in rheumatoid arthritis .A clinical and
immunological effects of a CD7 monoclonal antibody .
Br. J. Rheum., 30:459.

Kishimoto, T.(1989):

The biology of interleukin-6 .
Blood, 74:1.

Kiyokawa, T.; Shirono, K.; Haltori, T. and Takatsuki, K.(1989):

Cytotoxicity of IL-2 toxin toward lymphocytes from patients with adult T
cell leukemia .
Cancer Res. , 49:4042.

Kobayashi, S.; Teramura, M.; Sugawara, I. and Mizoguchi, H.(1993):

IL-11 acts as an autocrine growth factor for megakaryoblastic cell lines .
Blood, 81 :889.

Koj, A.(1985):

Definition and classification of acute phase proteins in the acute phase
response to injury and infection . Ed. A. H. Gordion and A. Koj.
Amsterdam: Elsevier . 10:139.

Kortiz, T.N.; Gurner, B.W.; Chalmers, D.G. and Coombs, R.R.(1980):

Measurement of rheumatoid factor in human sera by passive
haemagglutination of human erythrocytes carrying immunoglobulin linked
by chromic chloride .
J. Immunol. Meth., 32:1.

Kosaka, S. and Tozawa, M.(1976):

Alpha-1-antichemotrypsin in rheumatoid arthritis .
J. Exp. Med., 119:369.

Krane, S.M.; Conca, W.; Stephenson, M.L. and Goldring, M.B.(1990):

Mechanisms of matrix degradation in rheumatoid arthritis .
Ann. NY. Acad. Sci., 580:340.

Krane, S.M.; Dayer, J.M.; Simon, L.E. and Byrne, M.S.(1985):

Mononuclear cell conditioned medium containing mononuclear cell factor (MCF), homologous with IL-1, stimulates collagen and fibronectin synthesis by adherent rheumatoid synovial cells.
Coll. Relat. Res., 5:99.

Kreitman, R.J.; Chaudhary, V.K.; Kozak, R.W.; Fitzgerald, D.J.; Waldmann, T.A. and Pastan, I.(1992):

Recombinant toxins containing the variable domains of the anti-Tac monoclonal antibody to the interleukin-2 receptor kill malignant cells from patients with chronic lymphocytic leukemia .
Blood, 80 :2344.

Kuna, P.; Reddigari, S.R.; Kornfeld, D. and Kaplan, A.P.(1991):

IL-8 inhibits histamine release from human basophils induced by histamine releasing factors, connective tissue activating peptide III and IL-3 .
J. Immunol. 147:1920.

Kunkel, S.L.; Strieter, R.M.; Chensue, S.W.; Basha, M.; Standiford, T.; Ham, J. and Remick, D.G.(1990):

Tumor necrosis factor-alpha, interleukin-8, and chemotactic cytokine.
Prog. Clin. Biol. Res., 349:433.

Kupper, T.S.; Horowitz, M.; Leo, F.; Robb, R. and Flood, P.M.(1987):

Autocrine growth of T cells independent of interleukin-2; identification of interleukin-4 (IL-4, BSF-1) as an autocrine growth factor for a cloned antigen specific helper T cell .
J.Immunol., 138:4280.

Kurimoto, Y.; DeWeck, A.L. and Dahinden, C.A.(1989):

Interleukin-3 dependent mediator release in basophils.
J. Exp. Med., 170:467.

Kurt-Jones, E.A.; Beller, D.I.; Mizel, S.B. and Unanune, E.R.(1985):

Identification of a membrane associated interleukin-1 in macrophages .
Proc. Natl. Acad. Sci. USA, 82:1204.

Kurzrock, R.; Redman, J.; Cabanillas, F.; Jones, D; Rothberg, J. and Tallbaz, M.(1993):

Serum IL-6 levels are elevated in lymphoma patients and Correlate with survival in advanced Hodgkin's disease and with B symptoms .
Cancer-resp., 53 :2118.

Kushner, I.(1981):

The acute phase reactant and the E.S.R. In "Textbook of Rheumatology"
Editted by Kelley, W.N.; Harris, E.D.; Ruddy, S. and Sledge . C.B.
Saunder company, Philadilphia .Volume 1:672.

Kushner, I.; Gewarz, H. and Benson, M.D.(1981):

CRP and the acute phase response . Biologic reactivities and functions of
CRP .

J. Lab. Clin. Med., 67 :739.

Lacey, D.L. and Erdmann, J.M.(1990):

IL-1 and IL-4 modulate IL-1 receptor expression in a murine T cell line .
J. Immunol., 145:4145.

Lai, K.N.; Leung, J.C.; Li, RK.; Larche, M. and Ritter, M.A.(1991):

The effects of blockade of interleukin-2 receptors and interleukin-4
receptors on cytokine production .
ARMIS, 99:434.

Latte, D.; Weiss, J.B. and Jayson, M.I.V.(1981):

B2-M levels in serum and urine of R.A. patients on gold therapy .
Ann. Rheum. dis., 40:157.

Lefkovits , A. M. and Farrow , I. G.(1955) :

The liver in rheumatoid arthritis .
Ann. Rheum. Dis. , 14 : 162 .

Liao, Z.; Grimshaw, R.S. and Rosenstreich, D.L.(1989):

Interleukin-1 and its biologically related cytokines .
J. Exp. Med., 159:126 .

Libby, P.; Ordovas, J.M.; Auger, K.R.; Robbins, A.H.; Birinyi, I.K. and Dinarello, C.A.(1986):

Interleukin-1 and its biologically related cytokines .
J. Clin. Invest., 78:1432 .

Lipsky, P.E.(1985):

Interleukin-1 and its biologically related cytokines .
Contemp. Top. Mol. Immunol., 10:195.

Lipsky, P.E.(1989):

The control of antibody production by immunomodulatory molecules .
Arthr. Rheum., 32:194.

Lise, R.B.; Messner, R.P. and Troup, G.M.(1972):

Histocompatibility antigen and rheumatoid arthritis .
Arthr. Rheum., 15:524.

Lissoni, P.; Tisi, E.; Brivio, F.; Barni, S.; Rovelli, F.; Perego, M. and Tancini, G.(1991):

Increase in soluble interleukin-2 receptor and neopterin serum levels during immunotherapy of cancer with interleukin-2 .
Eur. J. Cancer, 27 :1014.

Liu, J.; Mondrell, B.; Aruffo, A.; Marken, J.S.; Taga, T.; Yasukawa, K.; Murakami, M.; Kishimoto, T. and Shoyab, M.(1992):

Interleukin-6 signal transducer gp 130 mediates oncostatin M signaling .
J. Biol. Chem., 267 :16763.

Loegdberg, L. and Shevach, E.M.(1985):

Interleukin-I and its biologically related cytokines.
Eur. J. Immunol., 15:1007.

Loegdberg, L., Wassmer, P. and Shevach, E.M.(1985):

Interleukin-I and its biologically related cytokines.
Cell Immunol., 94:299.

London, M. G. ; Huirten, K. D. and Hewit, J. V.(1963):

Serum cholesterol in rheumatoid arthritis .
Br. Med. J. , 1 : 1380.

Lowenthal, J.; Ransom, J.; Howard, M. and ziotnik, Z.(1988):

Upregulation of interleukin- 4 receptors on immature Lyt 2⁺
L 3T4⁺ thymocytes.
J.Immunol., 140: 474.

Lotz, M. and Carson, D.A.(1989):

Transforming growth factor and cellular immune responses in synovial fluid.
Arthr. Rheum., 32 : 542.

Lotz, M.; Jirik, F. and Kabowridis, P. (1988):

B-CSF-2/IL-6 is a co-stimulant for human thymocyte and T lymphocyte.
J.Exp.Med., 167 :1253.

Loughnan, M.S.; Sanderson, C.I. and Nassal, C.J.Y. (1988):

Soluble IL-2 receptors are released from the cell surface of normal murine B lymphocytes stimulated with Interleukin-5.
Proc. Natl . Acad. Sci. (USA) , 3115:4119.

Lowenthal, J.W. and MacDonald, H.R. (1986):

Binding and internalization of interleukin-1 by T-cells . Direct evidence from high and low affinity classes of interleukin-1 receptors .
J. Exp. Med., 164:1060.

MacNeil, I.; Suda, T.; Moore, K.; Mosmann, T. and Zlotnik, A. (1991):

IL-10, a novel growth cofactor for mature and Immature T cells .
J. Immunol., 145:4167.

Madhok, R.; Crilly, A.; Watson, J. and Capell, H.A. (1993):

Interleukin-6 levels in rheumatoid arthritis : Correlation with clinical and laboratory indices of disease activity .
Rheum. Dis., 52:232.

Makrynika, V.; Kabral, A. and Bradstock, K. (1991):

Effects of interleukin-7 on the growth of clonogenic cells in T cell acute lymphoblastic leukemia .
Leuk. Res., 15:879.

Malkovsky, M.; Loveland, B.; Gao, L.; Ward, P. and Friers, W. (1987):

Recombinant IL-2 directly augments the cytotoxicity of human monocytes .
Nature, 325: 262.

Malmquist, E. and Richard, H. (1962):

Serum ornithine carbamoyl transferase and transaminase activity in rheumatic diseases .
Acta Rheumatol. Scand. , 8 : 170.

Mancilla, J.; Ikejima, I.; Clark, B.D. and Dinarello, G. A. (1989):

Lectin binding suggest different glycosylation patterns of IL-1 receptors on different cells .
Cytokine, 1:95.

Manicourt, D.; Brauman, H. and Orloff, S. (1978):

Plasma and urinary levels of B₂-M in RA.
Ann. Rheum. Dis., 37:328.

Manicourt, D.; Brauma, H. and Orloff, S.(1980):

Synovial fluid B₂- microglobulin and hydroxy proline fractions in rheumatoid arthritis and non-autoimmune arthropathies .
Ann. Rheum. Dis., 39:207.

Marinkovic, S.; Jahreis, G.P.; Wong, G.G. and Baumann, H.(1989):

IL-6 modulates the synthesis of a specific set of acute phase plasma proteins in vivo .
J. Immunol., 142:808.

Marucha, P.T.; Zeff, R.A. and Kreutzer, D.L.(1990):

Cytokine regulation of IL-1 β gene expression in the human polymorph nuclear leucocyte .
J. Immunol., 145:2932.

Matsuda, T.; Hirano, T.; Nagasawa, S. and Kishimoto, T.(1989):

Identification of α_2 -macroglobulin as a carrier protein for IL-6 .
J. Immunol., 142:148.

Matsushima, K.; Yamada, M.; Furutani, Y. and Oppenheim, J.(1985):

Properties of specific IL-1 receptor on human Epstein Barr virus transformed B lymphocytes; identity of the receptor for IL-1 α and IL-1 β .
J. Immunol., 136:4496.

McBride, W.H.; Economou, J.S.; Comora, S. and Essner, R.(1990):

Influences of interleukin-2 and 4 on tumor necrosis factor production by murine mononuclear phagocytes .
Cancer Res., 50:2949.

McCarty, D.J.(1979):

Arthritis and Allied Conditions . A Textbook of rheumatology, 9 th ed .
Lea & Febiger Philadelpha, P. 400.

McCarty , D. J.(1985):

Synovial fluid In : Arthritis and allied conditions edited by McCarty , D. J. and Hollander , J. L. 10th ed .
Lea. Febiger , Philadelphia , 4 : 54.

McConkey, B.; Davis, P.; Crockson, R.A. and Amos, R.S.(1973):

The effect of some anti-inflammatory drugs on the acute phase protein in rheumatoid arthritis .
Quarterly J. Med., 42:785.

McConkey, B.; Davies, P.; Crockson, R. and Amos, R.(1979):

Effect of gold, dapsone and prednisone on serum C reactive protein and haptoglobin and erythrocyte sedimentation rate in rheumatoid arthritis .
Ann. Rheum. Dis., 38: 141.

McCormick , J. N.(1963):

An immunofluorescence study of rheumatoid factor .
Ann. Rheum. Dis. , 22 : 1.

McKinely, D.; Wu, Q.; Yang-Feng, T. and Yang, Y.C.(1992):

Genomic sequence and chromosomal location of human IL-11 gene .
Genomics, 13 :814.

Means, R.T. and Krantz, S.B.(1992):

Progress in understanding the pathogenesis of anaemia of chronic disease .
Blood, 80:1636.

Meikle,M.; Akinson, S.; Ward, R.; Murphy ,G. and Reunolds, J.(1989):

Gingival fibroblasts degrade I type I collagen films when stimulated with tumor necrosis factor and interleukin-I; evidence that breakdown is mediated by metalloproteinases .
J. periodont. Res, 24: 207.

Merriman , C.R.; puliam, L.A. and Kampsahimdt, R.F .(1977):

Comparison of leukocytic pyrogen and leucocytic endogenous mediator .
Proc. Sci. Exp. Biol. Med., 15:224.

Merz ,H.; Houssiau, F.A.; Orscheshek, K. and van Snick ,J.(1991):

Interleukin-9 expression in human malignant lymphomas : unique association with Hodgkin's disease and large cell anaplastic lymphoma .
Blood, 78 :1311.

Messner , P. (1984):

B₂ - M: An old molecule assumes a new look .
J. Lab. Clin. Med. , 104 : 141.

Metcalf,D.(1989):

Haemopoietic growth factor I.
Lancet , 1: 825.

Metcalf , D.(1992):

Introduction to the biology and pharmacology of interleukin-3 . IN:
perspective on Interleukin- 3 Symposium .
New Connaught Rooms , Grand Hall, 1:2.

Mikkelson, M.W. ; Dodge, H.J.; Duff, I.F. and Kad, H. (1967):

Estimation of the prevalence of rheumatic diseases in the population of Tecumseh, Michigan 1950-60.

J. Chronic Dis., 20: 315.

Miller, C.L.; Hooton, J.W.L.; Cillis, S. and Paetkau, V. (1990):

IL-4 potentiates the IL-2 dependent proliferation of mouse cytotoxic T cells.

J. Immunol., 144:1331.

Minty, A. P. ; Chalon, J. M. ; Derocq, X. and Caput, D. (1993):

IL-13 is a new human lymphokine regulating inflammatory and immune responses.

Nature, 362 : 248.

Mintz, G. (1985):

The acute phase reactants and the erythrocyte sedimentation rate. In: Rheumatoid Arthritis : Etiology, Diagnosis and Management., Utsinger P.O.; Zvaiflex, N. J. and Ehrlich, G.F. Eds., J.B. Lippincott company Philadelphia, London, Mexico City, New York, St Louis, Sao Paulo, Sydney, P.20.

Mitchell, L.C.; Davis, L.S. and Lipsky, P.E. (1989):

Promotion of human T lymphocyte proliferation by IL-4.

J. Immunol., 142 : 1548.

Miyasaka, N.; Sato.; Hashimoto, J. and Matsuda, T. (1989):

Constitutive production of IL-6/B cell stimulating factor 2 from inflammatory synovium.

Clin. Immunol. Immunopathol., 54:238.

Miyaura, C.; Jin, C.H.; Hirano, T.; Kishimoto, T. and Suda, T. (1989):

Production of IL-6 and its relation to the macrophage differentiation of mouse myeloid leukemia cells (M1) treated with differentiation inducing factor and $1\alpha, 25$ dihydroxy vitamin D₃.

Biochem. Biophys. Res. Commun., 158: 660.

Mizuguchi, J.; Beaven, M.A.; Ohara, J. and Paul, W.E. (1986):

BSF-1 action on resting B cells does not require elevation of inositol phospholipid metabolism or increased (CA²⁺).

J. Immunol., 137:2215.

Mizutani, H.; May, L.T., Sehgal, P.B. and Kupper, T.S. (1991):

Conversion of the inactive cytokine IL-1B, to biologically active species by human mast cell chymase.

J. Immunol., 147 : 3830.

Mock, B.A.; Nordan, R.P.; Justice, M.J. and Rudikoff, S.(1989):
The murine IL-6 gene maps to the proximal region of chromosome 5 .
J. Immunol., 142:1372.

Mongan, E. S. (1969):

Laboratory finding in rheumatoid arthritis "serum complement" . In : Arthritis and allied conditions edited by McCarty ,D. J. and Hollander , J. L. 10th edition .
Lea. , Febiger , Philadelphia P. 650.

Moore, F.D. Jr.; Schoof, D.D.; Rodrick, M. and Eberlein, T.J.(1991):

The systemic complement activation caused by interleukin - 2/ Lymphokine activated killer cell therapy of cancer causes minimal systemic neutrophil activation .
Int . J. cancer, 49 : 504.

Moore, M.A. and Warren, D.J.(1987):

Synergy of Interleukin -1 and granulocyte colony - stimulating factor : In vivo stimulation of stimulation of stem cell recovery and hematopoietic regeneration following 5- fluorouracil treatment of mice .
Proc. Natl . Acad . Sci . USA , 84 : 7134.

Mori, S.; Coto, F.; Coto, K.; Ohkawara, S. and Yoshinaga, M. (1988):

Cloning and sequence analysis of a cDNA for lymphocyte proliferation potentiating factor of rabbit neutrophils : Identification as rabbit Interleukin- 1B.
Biochem . Biophys . Res . Comm ., 150:1237.

Mosmann, T.R. and Coffman, R.L.(1989) :

TH-1 and TH-2 cells ; different patterns of lymphokine secretion lead to different functional properties .
Ann.Rev . Immunol ., 7:145.

Movitt, E.R. and Davis, A.E.(1953):

Liver biopsy in rheumatoid arthritis.
Am. J. Med. Sci., 226: 516.

Mowat, A. G. and Baum, J.(1971):

Chemotaxis of polymorphnuclear leucocytes from patients with RA.
J. clin. Invest., 50 : 2541.

Moxey- Mims, M.; Simms , H.; Frank , M. and Gaither, T.(1991):

The effects of IL-1, IL-2, and tumour necrosis factor on polymorphonuclear leucocyte Fc γ receptor mediated phagocytosis. J. Immunol., 147: 1823.

Muegge, K.; Williams, T.M.; Kant, J.; Karin, M.; Chin, R.; Schmidt, A.; Siebenlist, U.; Young, H.A. and Durum, S.K.(1989):
Interleukin-1 costimulatory activity on the interleukin-2 promoter via Ap-1.
Science, 246.

Mukaida, N.; Hishinuma, A. and Matsushima, K.(1991):
Regulation of human interleukin-8 expression and binding of several other members of the intercrine family to receptors for interleukin -8 .
Adv. Exp. Med. Biol., 305 : 31.

Murata, Y.; Takaki, S.; Migita ; Kikuchi, Y. and Takatsu, K.(1992):
Molecular cloning and expression of the human interleukin-5 receptor.
J.Exp. Med., 175: 341.

Nachman, R.L.; Hajjar, K.A. and Dinarello, C.A.(1986):
Interleukin -1 and its biologically related cytokines .
J.Exp. Med., 163:1544, 1986 .

Nakahama H.; Okada, M.; Meyazoki, M. and Kubori, S.(1992):
IL-6 and other laboratory parameters to treatment patient with poly arteritis nodosa .
Angiology, 43:512.

Nakanishi, K.; Matsui, K.; Hirose, S. and Higashino, K.(1990):
Lymphokine regulated differential expression mRNA for p75 KDa-IL-2E and p55 Kda IL-2R in a cloned B lymphoma line (BLC-CL- 3 cells).
J.Immunol., 145:1423.

Nakase, K.; Kita, K.; Ostuji, A. and Tsutani, H.(1992):
Diagnostic and clinical importance of interleukin- 2 receptor alpha chain expression on non - T cell acute leukaemia cells .
Br. J.Haematol., 80: 317.

Naume, B. and Espevik, T.(1992):
Effects of IL-7 and IL-2 on highly enriched CD56, natural killer cells .
J. Immunol., 147: 2203.

Nettlebladt, E.(1962):
Rheumatoid arthritis and hepatocellular injury .
Acl. Rhum. Scand., 6 : 256.

Nikaido, T.; Shimizu, A.; Ishida, N. and Honjo, T.(1984):
Molecular cloning of cDNA encoding human interleukin-2 receptor.
Nature, 311: 631.

Nilsson, L. A. (1968):

Determination of serum C-reactive protein.
Acta Path . Microbiol . Scand ., 73 : 129.

Nishida, T.; Nishino, N.; Takano, M.; Akai, S. and Hirai, Y. (1987):

cDNA cloning of IL- 1 alpha and IL-1 beta from mRNA of U937 cell line .
Biochem . Biophys . Res . Commun ., 143:345.

Numerol , R.P.; Dinarello, C.A.; Endres , S. and Mier, J. (1988):

Interleukin- 1 and its biologically related cytokines .
J. Immunol ., 144: 964.

Ogra, P.L.; Chiba, Y.; Ogra , S.S. and Hard, J. K . (1975):

Rubella - virus infection in juvenile rheumatoid arthritis .
Lancet, 1: 1157.

Ohta , K. ; Seno , A. ; Shintani , N. ; Kato , E. and Yachie , A. (1993):

Increased levels of urinary interleukin-6 in Kawasaki disease .
Eur. J. Pediat. , 152 : 647.

Ohzato, H.; Yoshizaki, K.; Nishimoto, N. and Mori, T. (1992):

IL-6 as a new indicator of inflammatory status.
Surgery, 111: 201.

Old, L.J. (1987):

TNF polypeptide mediator network .
Nature , 326:330.

Oliveira , E.B.; Gotschpich, E.C. and Lin, T. (1979):

Primary structure of human C- reactive protein
J. Biol. Chem., 254:489.

Onji, M. ; Yamaguchi, S.; Masumoto, T. and Ohta , Y. (1991):

Induction of the interleukin-2 receptor (p55) by interferons .
J. Interferon Res ., 11 : 237.

Oppenheim, J.J.; Rusctti, F.W. and Connie , F. (1994):

Cytokines . IN: Stites , D.P. and Terr, A.I. (eds) : Basic and clinical Immunogy . 7 th edition . Appleton and Lange , London , Sydney , Toronto, Mexico , New Delhi . Chapter 9.p 105.

Oppenheim, J.J.; Rusctti, F.W. and Fallynek, C. (1991):

Cytokines. In: Basic and Clinical Immunology textbook. (7th eddition),
Chapter 7 : Page 78 . edited by Daniel P. Stites and Abba I. Tem .
Published by Appleton and lange .

Ortaldo, J.; Masoon, A.; Smyth, M.; Falk, L. and Ruscetti, F. (1991):
 Mechanistic studies of transforming growth factor- β inhibition of IL-2
 dependent activation of CD 3⁺ large granular lymphocyte functions;
 regulation of IL-2 R β (P75) signal transduction .
 J. Immunol ., 146: 3791.

O' sborn, L.; Hession, C.; Tizard, R.; Vassallo, and Lobb, R. (1989):
 Direct expression cloning of vascular cell adhesion molecule 1, a cytokine
 induced endothelial protein that binds to lymphocytes .
 Cell , 59 : 1203.

Oshimi, K, and Hoshino, S. (1992) :
 Function and molecular structure of IL- 12.
 Nippon . Rinsho ., 50 : 1840.

Osmand, A.P.; Friedenson, B. and Gewurz, H. (1977):
 Characterization of C- reactive protein and the complement subcomponent
 C1q as homologous proteins displaying cyclic pentameric symmetric
 (pentraxins) .
 Proc . Natl . Acad . sci . USA , 74 : 739.

Osmand, A.P. ; Mertenser, R.F.; Siegle, J. and Gewurz, H. (1975):
 Interaction of C- reactive protein with the complement system .
 Complement - dependent passive haemolysis initiated by CRP.
 J. Exp . Med ., 142 : 1065.

Ota, K. ; Kitano, U. and Miyasaka, N. (1989):
 Clinical and pathophysiological aspects of dialysis - associated
 amyloidosis .
 Nephrol. Dial. Transplant., 4 : 7.

Ozaki, Y.; ohashi, T, and Kume, S. (1987):
 Potentiation of neutrophil function by recombinant DNA produced
 Interleukin- 1 α .
 J. Leucocyte Biol. , 42 : 621.

Ozaki, Y.; Ohashi, T.; Niwa, Y. and Kume, S. (1988):
 Effect of recombinant DNA produced tumor necrosis factor on various
 parameters of neutrophil function inflammation .
 Leuk . Res. , 12 : 297.

Panayi, G.S. (1982):
 Scientific Basis of Rheumatology, Churchill Livingstone ,
 Edinburgh, London , Melbourne , New York, p. 61.

Panuch , S.R.(1982):

Principle of Rheumatic Diseases , Awiley Medical Puplication . John Wiley Sons ,New York , Chichester , Brisbane , Toronto , P. 25.

Panush , R.S.; Franco , A.E. and Schur, P.H.(1986):

Rheumatoid arthritis associated with easinophilia .

Annals of Internal Medicine , 75 : 199.

Parekh , R.B.; Dwek R.A. and Radenmacher , T.(1992):

Glycosylation of interleukin-6 purified from normal human blood mononuclear cells .

Eur . J. Biochem ., 203 : 135.

Parker ,K.P.; Benjamin, W.R.; Kaffke , K.L. and Kilian , P.L.(1989):

Presence of IL-1 receptors on human and murine neutrophils relevance to IL-1 mediated effects in inflammation .

J.Immunol., 142 : 537.

Paul , S.R.; Barut , B.; Bennett , F. and Anderson , K.(1992):

Lack of a role of interleukin - 11 in the growth of multiple myeloma .

Leuk. Res ., 16 : 247.

Paul, S. R. and Schendel , P. (1992):

The cloning and biological characterization of recombinant human IL-11 .

Int . J. Cell Cloning , 10 : 135 .

Pavlidis , N.A.; Manoussakis , M.N. and Moutsopoulos , H.M.(1992):

Serum soluble IL-2 receptors in B cell lymphoproliferative malignancies .

Med . Pediatr . Oncol ., 20 : 26.

Pechumer. H.; Leinisch, E. and Ziegler- Heitbrock , H.(1991):

Recovery of monocytes after bone marrow transplantation,rapid reappearance of tumor necrosis factor alpha and IL- 6 production .

Transplantation, 52 : 698.

Pelliniemi , T.T.; Irjala,K.; Pulkki and Lahtinen, R.(1995):

Immunoreactive interleukin 6- and acute phase proteins as prognostic factors in multiple myeloma .

Blood, 85 : 765.

Pepys , M.B .(1982):

C- reactive protein and the acute phase reponse .

Nature , 296 : 12.

Peterson , A. ; Evrin , E. and Berggard , I.(1969):

Differentiation of glomerular , tubular and removal proteinuria .
Determination of urinary excretion of B₂ - M, albumin and total protein .
J. Clin. Invest. , 48 : 1189.

Phadke , K.(1987):

Interleukin - 1 and its biologically related cytokines .
Biochem . Biophys. Res. commun., 142 : 448.

Pickup , M. E. ; Dixon , J. S. and Wright , V.(1982):

Plasma viscosity : a new appraisal of its use as an index of disease activity
in rheumatoid arthritis .
Ann. Rheum. Dis. , 40 : 272.

Plaut , M.; Pierce , J.; Watson , C.; Hanley , J. and Paul , W.(1989):

Mast cell lines produce lymphokines in response to cross - linkage of
FcεR1 or to calcium ionophores .
Nature , 339 : 64.

Plesner,T.; Karle , H.; Rubin , B. and Thosmson , M .(1978):

Evidence for a change in the expression of B₂- M associated membrane
structure on leukaemic human cell .
Clin . Exp . Immunol ., 31:269.

Podlaski , F.J.; Nanduri , V.B. and Stern , A.S.(1992):

Molecular characterization of interleukin-12.
Arch. Biochem. Biophys., 294 : 230.

Postlethwaite , A.; Raghow, R.; Strickin , G. and Haug , A.(1988):

Modulation of fibroblast functions by Interleukin- 1 : increased steady state
accumulation of type I procollagen RNAs and stimulation of other
functions but not chemotaxis by human recombinant IL- I α and B .
J. Cell Biol., 106: 311.

Puri, R.Y.; Finbloom , D.S.; Leland, P. and Siegel J.(1990):

Expression of high affinity IL-4 receptors on murine tumor infiltrating
lymphocytes and their upregulation by IL- 1 .
Cell Immunol. , 97 : 110.

Quesniaux , V.F.; Clark , S.C.; Turner , K. and Fagg , B.(1992):

Interleukin-11 stimulates multiple phases erythropoiesis in vitro .
Blood, 80 :1218.

Qwarnstrom, E.E.; Page, R.C.; Gillis, S. and Dower, S.K.(1988):

Binding, internalization, and intracellular localization of interleukin-1 beta in human diploid fibroblasts .

J. Biol. Chem., 263:8261.

Raines, E.W.; Dower, S.K. and Ross R.(1989) :

IL-1 mitogenic activity for fibroblasts is due to PDGF-A.

Science 243:393.

Ralph, P.; Nakcinz, I. and Rennick, D.(1988):

Role of interleukin-2, interleukin-4, and α , β , and γ interferon in stimulating macrophage-antibody dependent tumoricidal activity .

J. Exp. Med., 167:712.

Reitman, S. and Frankle, S.(1957):

Determination of serum aminotransaminases .

Am. J. Clin. Path., 28: 57.

Renauld, J.C.; Goethals, A.; Houssiau, F. and Van Snick, J.(1990):

Cloning and expression of a cDNA for the human homolog of mouse T cell and mast cell growth factor p40 .

Cytokine, 2: 9.

Renz, H.; Domenico, J. and Gelfand, E.(1991):

IL-4 dependent up-regulation of IL-4 receptor expression in murine T and B cells .

J. Immunol., 146:3049.

Richards, C.D.; Brown, T.; Shoyab, M. and Gauldie, J.(1992):

Recombinant oncostatin M stimulates the production of acute phase proteins in Hep G2 cells and rat primary hepatocytes in vitro .

J. Immunol., 148: 1731.

Ritchie, R.T.(1979):

Specific protein. In : Henery, J.B. (ed) Clinical Diagnosis and Management by Laboratory Methods, 16th ed . W.B. Saunders Co. Philadelphia , London, Toronto, 243.

Roper, R.L.; Conread , D.H.; Brown , D.M. and Phipps , R.P.(1990):

Prostaglandin E, promotes IL- 4 induced IgE and IgG-1 synthesis .

J . Immunol., 145 : 2644.

Ropes , M. W. ; Bennett , G. A. and Jessar , R. A.(1959):

Diagnostic criteria for rheumatoid arthritis .

Ann. Rheum. Dis. , 18 : 49.

Rose , H.M.; Regan, C.; Perace,E . and Lipnan, M.O.(1949):

Differential agglutination of normal and sensitized sheep erythrocytes by sera of patients with rheumatoid arthritis .

Proceed Soc. Exp . Biol . Med ., 68 : 1.

Rose, P.E.; Johnson, S.A. and Mack, J.(1989):

Study of C - reactive protein during infection in leukemia .

J. Clin. Pathol ., 39 : 263 .

Rose , T.M . and Bruce , A.G.(1991):

Oncostatin M is a member of a cytokine family that includes leukemia - inhibitory factor , granulocyte colony - stimulating factor, and Interleukin-6.

Proc. Natl . Acad . Sci . USA ,88 : 8641.

Rosenberg , S.; Lotze, M.; Muul . Chang , A. and White , D .(1987):

A progress report on the treatment of 157 patients with advanced cancer using lymphokine - activated killer cells and Interleukin-2 or high dose Interleukin 2 alone .

N. Engl . J. Med., 316 : 889 .

Rosenwasser , L.M.; Dinarello , C.A . and Rosenthal , A.(1979):

Adherent cell function in murine T - lymphocyte antigen recognition . IV. Enhancement of murine T - cell antigen recognition by human leukocytic pyrogen .

J. Exp. Med., 150 : 709.

Roux - Lombard , P.; Modonix , C. and Dayer , J.M.(1988):

Inhibitoes of IL-1and TNF - α activities in synovial fluids and cultured synovial fluid cell supernatants. .

Calcif. Tissue Int., 42 : 47.

Rubin , L.; Snow , K.; Kurman , C. and Keystone , E(1990).:

Serial levels of soluble IL-2 receptor in peripheral blood of patients with rheumatoid arthritis : Correlation with disease activity .

J. Rheumatol., 17 : 597.

Saeland , S.; Duvert, V.; Pandrau, D.; Caux , C.; Durand , I.; Widerman, J.; Lee, F. and Banchereau , J.(1991):

Interleukin -7 induces the proliferation of normal human B cell precursors .

Blood , 78 : 2229 .

Saiki, O.; Tanaka , T.; Doi , S. and Kishimoto,S.(1988):

Cellular interaction in the humoral immune response .

J.Immunol. , 140 : 853 .

Saklatvala, J.; Sarfield, S.J. and Townsend, Y. (1985):

Pig interleukin purification of two immunologically different leukocyte proteins that cause cartilage resorption, lymphocyte activation and fever
J. Exp. Med., 162: 1208.

Sampson, H.A.; Broadbent, K.R. and Bernhisel-Broadbent, J. (1989):

Spontaneous release of histamine from basophils and histamine-releasing factor in patients with atopic dermatitis and food hypersensitivity.
N. Engl. J. Med., 321:228.

Scapigliati, G.; Ghiara, P.; Bartalini, A. and Boraschi, D. (1989):

Differential binding of IL-1 α and IL-1 β to receptors on B and T cells.
FEBS Lett., 243 : 398.

Schaller, J.; Beckwith, B. and Wedg Wood, R.J. (1970):

Hepatic involvement in juvenile rheumatoid arthritis
J. Pediat., 77 : 203.

Schardijn, C. and Statius, Van E. W. (1987):

B₂ - M. Its significance in the evaluation of renal function .
Kidney Int., 32 : 635 , 1987 .

Schibler, K.R.; Yang, Y.C. and Christensen, R.D. (1992):

Effect of interleukin-11 on cycling status and clonogenic maturation of fetal and adult hematopoietic progenitors .
Blood, 80 : 900.

Schmitt, E.; Beuscher, H.U.; Huels, C.; Montevne, P.; Van Brandwi, J.K. R.; Van Snick, J. and Ruede, E. (1992):

IL-1 serves as a secondary signal for IL-9 expression .
J. Immunol., 147:3848.

Schmitt, E.; Van Brandwi, J.K.R.; Van Snick, J. and Ruede, E. (1990):

TCGF III / P 40 is produced by native murine CD4⁺T cells but is not a general T cell growth factor .
Eur. J. Immunol., 19: 2167.

Schnittman, S.M.; Singer, K.H.; Greenhouse, J.J.; Stanley, S.K.; Whichard, L.P.; Haynes, B.F. and Fauci, A.S. (1992):

Thymic microenvironment induces HIV expression; physiologic secretion of IL-6 by thymic epithelial cells upregulates virus expression in chronically infected cells .
J. Immunol., 147: 2553.

Schonhaut, D.S.; Chua, A.O.; Wolitzky, A.G.; Quinn, P.M.; Dwyer, C.M.; McComas, W. and Gubler, U.(1992):

Cloning and expression of IL-12 .

J. Immunol., 148: 343.

Schroder, J.M.; Mrowietz, J. and Christophers, E.(1988):

Purification and partial biological characterization of a human lymphocyte derived peptide with potent neutrophil stimulating activity .

J. Immunol., 140 : 3534.

Schroeder, T.J.; Helling, T. and Jeffrey, J.(1992):

A multicenter study to evaluate a novel assay for quantitation of soluble interleukin-2 receptor in renal transplantation .

Transplantation, 53 : 34.

Schuster, J. ; Gold , P. and Pouli, K.(1976):

B₂ - M levels in cancerous and other disease states .

Clin. Chem. Acta , 67 : 307.

Seckinger, P.; Klein-Nulend, J.; Alander, C. and Raisz, L.(1990):

Natural and recombinant human IL-1 receptor antagonists block the effects of IL-1 on bone resorption and prostaglandin production .

J. Immunol., 145:4181.

Sehgal, P.; Zilberstein, A.; Ruggieri, R.; May, L. and Ruddle, F.(1987):

Human chromosome 7 carries the β_2 interferon gene .

Proc. Natl. Acad. Sci. USA, 84:3633.

Sehgal, P.B.; Helfgott, D.C.; Clarick, R.H. and May, L.T.(1988):

Regulation of the acute phase and immune responses in viral disease .

J. Exp. Med., 167:1951.

Seitz, M.; Dewald, B.; Geber, N. and Baggiolini, M.(1991):

Preferential production of IL-8 in RA relationship for rheumatoid factor in vivo and inhibition by Gold Sodium Thiomalate and Methotrexate in vitro .

Br. J. Rheum., 30 :85.

Sharon, M.; Gnarr, J.R.; Baniyashi, M. and Leonard, W.J.(1988):

Possible association between IL-2R and class I HLA molecules on T cells.

J. Immunol., 141: 3512.

Sher, A.; Fiorentino, D.; Caspar, P. and Mosmann, T.(1992):

Production of IL-10 by CD4⁺ T lymphocytes correlates with down-regulation of TH-1 cytokine synthesis in helminth infection .

J. Immunol., 147:2713.

Sherry, B. and Cerami, A.(1988):

Cachectin/Tumor necrosis factor exerts endocrine, paracrine, and autocrine control of inflammatory responses .
J.Cell Biol., 107:1269.

Shore, A.; Jaglal, S. and Keystone, E.C.(1986):

Enhanced IL-1 generation by monocytes in vitro is temporarily linked to an early event in the onset or exacerbation of rheumatoid arthritis .
Clin. Exp. Immunol., 65:293.

Sieff, C.A.(1987):

Hemopoietic growth factors .
J. Clin. Inv., 79:1549.

Siegel, J.P.; Sharon, M.; Smith, P.L. and Leonard, W.J.(1987):

The IL-2 receptor- β chain p(70); role in mediating signals for LAK, NK, and proliferative activities .
Science, 238:75.

Silberstein, D.S.; Schoof, D.D.; Rodrick, M.L. and Eberlein, T.(1989):

Activation of eosinophils in cancer patients treated with IL-2 and IL-2 generated lymphokine activated killer cells .
J. Immunol., 142:2162.

Silverman, E.D.; Laxer, R.M.; Nelson, D.L. and Rubin, L.A.(1991):

Soluble Interleukin-2 receptors in Juvenile Rheumatoid Arthritis .
J. Rheumatol 18: 1398 .

Simic, M.M. and Stosic-Grujicic, S.(1985):

Interleukin-1 and its biologically related cytokines .
Folia Biol., 31:410.

Sims, H. and Miller, L.(1984):

B₂ - M in laboratory diagnosis .
Lab. Med. , 15 : 803 .

Singer, J.M. and Plotz, C.M.(1956):

The latex fixation test . Application to the serologic diagnosis of RA.
Am . J. Med ., 21: 888 .

Skjonsberg, C.; Erikstein, B.; Smeland, E.B. and Blomhoff, H.(1991):

Interleukin-7 differentiates a subgroup of acute lymphoblastic leukemias .
Blood, 77 : 2445.

Smith, K.A.(1988):

The interleukin-2 receptor .
Adv. Immunol., 42:165.

Smith, K.A.(1989):

Receptors of interleukin-2 .
Ann. Rev. Cell Biol., 5:397.

Smith, K.A. and Cantrell, D.(1985):

Interleukin-2 regulates its own receptors .
Proc. Natl. Acad. Sci. USA, 82:864.

Smith, J.; Urba, W.; Steis, R.; Janik, J. and Longo, D.(1990):

IL-1 alpha: Results of a phase I toxicity and immunomodulatory trial.
Am. Soc. Clin . Oncol., 9:717.

Smyth, M.J.; Ortado, J.R.; Bere, W. and Young, H.A.(1990 a):

IL-2 and IL-6 synergize to augment the pore-forming protein gene expression and cytotoxic potential of human peripheral blood T cells .
J. Immunol., 145:1159.

Smyth, M.; Ortaldo, J.R.; Nakata, M. and Young, H.A.(1990 b):

Regulation of pore-forming protein gene expression in human peripheral blood leukocytes .
J. Exp. Med., 171:1269.

Smyth, M.J.; Zachariae, C.; Norihisa, Y. and Matsushima, K.(1991):

IL-8 gene expression and production in human peripheral blood lymphocytes subsets .
J. Immunol., 146:3815.

Solari, R.; Smithers, N.; Page, K. and Champion, B.R.(1990):

IL-1 responsiveness and receptor expression by murine Th1 and Th2 clones .
Cytokine 2:129.

Soulillou, J.; Cantarovich, D.; Hourmant, M. and Jacques, Y.(1990):

Randomized controlled trial of a monoclonal antibody against the interleukin-2 receptor as compared with rabbit anti-allografts .
N. Engl. J. Med., 322:1175.

Spence, C.; Nachman, M.; Gately, M.K. and Bailon, P.(1993):

Affinity purification and characterization of anti-Tac (FV)-c3-PE38KDEL; a highly potent cytotoxic agent specific to cells bearing IL-2 receptors .
Bioconjug Chem., 4:63.

Spits, H.; Yassel, H.; Palird, X.; Figdor, C. and De Vries, J.E.(1988):
IL-4 inhibits IL-2 mediated induction of human lymphokine-activated killer cells, but not the generation of antigen-specific cytotoxic T lymphocytes in mixed leucocyte cultures .
J. Immunol., 141: 29.

Standiford, T.J.; Kunkel, S.L.; Rolfe, M.W. and Strieter, R.M.(1991):
IL-8 gene expression from human alveolar macrophage; the role of adherence .
J. Resp. Cell Mol. Biol, 5:579.

Standiford, T.J.; Strieter, R.M.; Kasahara, K. and Kunel, S.L.(1990):
IL-4 inhibits the expression of IL-8 from stimulated human monocytes .
J. Immunol., 145:1435.

Steinberg, A.C. and Wilson, J.A.(1963):
Hereditary globulin factors and immune tolerance in man .
Science, 140:303.

Steins, S.H. and Phipps, R.P.(1992):
Antigen-specific IgG_{2a} production in response to prostaglandin E₂, immune complexes and IFN- γ .
J. Immunol., 147:2500.

Stephanson, M.L.; Krane, S.M.; Aments, E.P. and Byrne, M.(1985):
Immune interferon inhibits collagen synthesis by rheumatoid synovial cells associated with decreased levels of the procollagen m.RNAs.
FEBS Lett. ,180:43.

Stork, L.C.; Barczuk, L.; Kissinger, M. and Robinson, W.A.(1989):
IL-1 accelerates murine granulocyte recovery following treatment with cyclophamide .
Blood, 73:938.

Stork, L.C.; Peterson, U.M.; Rundus, C.H. and Robinson, W.A.(1988):
Interleukin-1 enhances murine granulopoiesis in vivo .
Exp. Haematol., 16:163.

Stotter, H.; Custer, M.C.; Bolton, E.S. and Lotze, M.T.(1991):
IL-7 induces human lymphokine-activated killer cell activity and is regulated by IL-4 .
J. Immunol., 146:150.

Stuart, P.M.; Zlotnik, A. and Woodward, J.G.(1988):
Induction of class I and class II MHC antigen expression on murine bone

marrow derived macrophages by IL-4 (β cell stimulatory factor 1) .
J. Immunol., 140:1542.

Sturk, C.(1992):

IL-6 and acute phase reactant in patient with acute myocardial infarction .
J. Lab. Clinic. Med., 114:148.

Suda, T.; Murray, R.; Fischer, M.; Yokota, T. and Zlotnik, A.(1990):

Tumor necrosis factor alpha and p40 induce day 15 murine fetal thymocyte proliferation in combination with IL-2 .
J. Immunol., 144: 1783.

Suda, T. and Zlotnik, A.(1991):

IL-7 maintains the T cell precursor potential of CD3⁺ CD4⁺ CD8⁺ thymocytes.
J. Immunol., 146:3068.

Sullivan, G.W.; Carpet, H.T.; Sullivan, J. and Manell, G.(1989):

Both recombinant interleukin-1 (beta) and purified human monocyte interleukin-1 prime human neutrophils for increased oxidative activity and promote neutrophil spreading .
J. Leucocyte Biol., 45:389.

Svenson, M.; Hansen, M.B.; Kayser, L.; Rasmussen, A.K.; Reimert, C.M. and Bendtzen, K.(1992):

Effects of human anti-IL-1-alpha autoantibodies on receptor binding and biological activities of IL-1 .
Cytokine, 4 : 125.

Swain, S.L.; Huston, G.; Tonkonogy, S. and Weinberg, A.(1991):

Transforming growth factor- β and IL-4 cause helper T cell precursors to develop into distinct effector helper cells that differ in lymphokine secretion pattern and cell surface phenotype .
J. Immunol., 147:2991.

Swanson, A. ; Tracy , P. ; Katzmman , A. and Young , S.(1982):

B₂ - M determination by radioimmunoassay with monoclonal antibody .
Clin. Chem. , 28 : 2033 .

Swezey, R.L.(1967):

Myocardial infraction due to rheumatoid arthritis .
JAMA., 191.

Symann, M.(1992):

Interleukin-3 and peripheral blood stem cell harvesting . IN:

Perspectives on Interleukin-3 symposium . New Connaught Rooms, Grand Hall, London, 1: 18.

Tada, S.; Kawabata, H.; Kitani, H. and Tanizaki, K.(1992):
The Significance of SIL-2R in RA with Interstitial Pneumonia .
Arerugi: 41, 428 .

Talal, N.; Grey, H.M.; Zvaifler, N. and Daniels, T.E.(1975):
Elevated Salivary and Synovial fluid B₂-M in Sjogren's Syndrome and RA.
Science, 187:1196.

Talmadge, J.E.; Pennington, R.; Bowersox, O. and Tribble, H.(1989):
Myelostimulatory activity of recombinant human interleukin-2 in mice .
Blood, 73 :1458.

Tanaka, T.; Saiki, O.; Suemura, M. and Kishimoto, S.(1988):
Cellular interaction in the humoral immune response .
J. Clin. Invest., 82:316.

Tang , E. and Van Den Bergh , W.(1981) :
The tubular handling of B₂ - M. in sickle cell nephropathy .
Nephron , 29 : 138.

Tanner, J.E. and Tosato, G.(1992):
Regulation of B cell growth and Ig gene transcription by Interleukin-6 .
Blood, 79 :452.

Tavernier, J.; Devors, R.; Cornelis, S. and Plaetinck, G.(1991):
A human high affinity interleukin-5 receptor (IL-5R) is composed of an IL-5 specific alpha chain and a beta shared with the receptor for GM-CSF .
Cell, 66:1175.

Teramura, M.; Kobayashi, S.; Hoshino, S. and Mizoguchi, H.(1992):
Interleukin-11 enhances human megakaryocytopoiesis in vitro .
Blood, 79 :327.

Terkeltaub, R.; Zachari, C.; Artin, J. and Matsushima, K.(1991):
Monocyte derived neutrophil chemotactic factor/IL-8 is a potential mediator of cystal induced inflammation .
Arth. Rheum., 34:894.

Tetta, C.; Camussi, G.; Modena, N. and Baglioni, C.(1990):
Tumour necrosis factor in serum and synovial fluid of patients with active and severe RA .
Ann. Rheum. Dis., 49 : 665.

Te-Velde, A.A.; Huijbens, R.J.; Heije, K. and Figdor, C.(1990):

Interleukin-4 (IL-4) inhibits secretion of IL-1 beta, tumor necrosis factor alpha, and IL-6 by human monocytes .
Blood, 76 :1392.

Tewari, A.; Buhles, W.C. and Starnes, H.F.(1990):

Preliminary report: Effects of interleukin-1 on platelet counts .
Lancet, 336 : 712.

Thompson, H.L.; Burbelo, P.D.; Yamada, Y. and Metcalf, D.D.(1989):

Laminin promotes mast cell attachment .
J.Immunol., 143:3323.

Tigges, M.A.; Casey, L.S. and Koshland, M.E.(1989):

Mechanism of interleukin-2 signaling; mediation of different outcomes by a single receptor and transduction pathway .
Science, 243:781.

Touw, I.; Pouwels, K.; Van-Agthosven, T. and Lowenberg, B.(1990):

IL-7 is a growth factor of precursor B and T acute lymphoblastic leukemia.
Blood, 75:2097, 1990 .

Tsuda, M.; Kitamura, F. and Miyasaka, M.(1989):

Characterization of the interleukin-2 receptor β -chain using three distinct monoclonal antibodies .
Proc. Natl. Acad. Sci. USA, 86: 1982.

Tsukada, K. ; Katoh, H. ; Suzuki, T. ; and Nagamachi, Y.(1993):

Correlations of peritoneal interleukin-6 , serum B₂ - M and urinary B₂ - M after elective abdominal surgery .
Ampis , 101 : 409 .

Tulkens, M.(1986):

Experimental studies of nephrotoxicity of aminoglycosides at low doses .
Am. J. Med. , 80 : 105.

Uhl, J.; Newton, C.R.; Giri, J.G.; Sandlin, G. and Horuk, R.(1989):

Identification of IL-1 receptors on human monocytes .
J. Immunol., 142 : 1576.

Utsinger, P.D.; Zvaifler, N.J. and Ehrlich, G.E.(1985):

Rheumatoid Arthritis: Etiology, Diagnosis, Management, J.B. Lippincott Company Philadelphia, London, Mexico City, New York, St Louis, Sao Pule, Sydney, P.209.

Vaisberg, M. and Scheinberg, M.A.(1990):

Simultaneous evaluation of membrane bound and soluble IL-2 receptor expression in the blood and synovial fluid of patients with R.A.
Clin. Exp. Rheumatol., 8 :579.

Valent, P. and Bettelheim, P.(1990):

The human basophil .
CRC. Crit. Rev. Oncol. Haematol, 73:1763.

Valiante, N.M.; Rengaraju, M. and Trinchieri, G.(1992):

Role of the production of natural killer cell stimulator factor (NKSF/IL-12) in the ability of B cell lines to stimulate T and NL cell proliferation .
Cell Immunol., 145 : 187.

Van-Damme, J. and Opdenakker, G.(1991):

Interaction of interferons with reactive cytokines; from IL-1 to IL-8 .
J. Invest. Dermatol., 95 : 90.

Van Gameren, M.M.; Willemse, P.H.B. and Mulder, N.H.(1994):

Effect of recombinant human interleukin-6 in cancer patients .
Blood, 84:1434.

Van Leeuwen, M.; van der Heijde, D. and Van Rijswijk, M.H.(1994):

Inter-relationship of outcome measures and process variables in early rheumatoid arthritis .
J. Rheumatol., 21:425.

Van Leeuwen, M.A.; Westra, J. and Van Rijswijk, M.H.(1995 a):

Interleukin-6 in relation to other proinflammatory cytokines, chemotactic activity and neutrophil activation in rheumatoid synovial fluid .
Ann. Rheum. Dis., 54:33.

Van Leeuwen, M.; Westra, J. and Van Rijswijk, M.(1995 b):

Clinical significant of IL-6 in early R.A.:relation with laboratory,clinical and radiological progression in a three year prospective study .
Ann. Rheum. Dis., 54:674.

Van Snick, J.(1990):

Interleukin-6 an overview .
Ann.Rev. Immunol., 8:253.

Villiger, P.; Cronin, M.; Amenomori, T. and Lotz, M.(1991):

IL-6 production by human T lymphocytes; expression in HTLV-1 infected but not in normal T cells .
J. Immunol., 146:550.

Vincent, C. ; Pozet, N. and Revillard, J.(1980):

Serum levels and urinary excretion of B₂ - M in patients under haemodialysis or after renal transplantation .

Acta Clin. Belg. , 35 : 31 .

Vincent, C. and Revillard, P.(1976) :

Comparison of radioimmunoassay and lymphotoxicity inhibition technique for determination of B₂ - M.

J. Immunol. Methods , 10 : 253 .

Vreugdenhil, G.; Lindemans, J.; Van-Eijk, H. and Swaak, A.(1992):

Elevated serum transcobalamin levels in anaemia of rheumatoid arthritis : Correlation with disease activity but not with serum TNF and IL-6.

J. Intern. Med., 231:547.

Waler, E.(1940):

On the occurrence of a factor in human serum activating the specific agglutination of sheep blood corpuscles .

Acta. Path. Microb. Sand., 17:172.

Wahl, S.M.; McCartney-Francis, N.; Hunt, A.; Smith, P.D.; Wahl, L.M. and Katona, I.M.(1988):

Monocyte interleukin-2 receptor gene expression and interleukin-2 augmentation of microbicidal activity .

J. Immunol., 139:1342.

Waller, M.(1970):

Determination of serum rheumatoid factor .

Reviews in Medical Sci., 173.

Walsh, L.; Davis, P. and Macomkey, B.(1979):

Relationship between erythrocyte sedimentation rate and serum C-reactive protein in rheumatoid arthritis .

Ann. Rheum. Dis., 38:362.

Walz, G.; Aniffo, A.; Kolnaus, W. and Seed, B.(1990):

Recognition by ELAM-1 of the Sialyl-Le determinant on myeloid and tumor cells .

Science, 250:1132.

Wang, J.M.; Sica, A.; Peri, G.; Walter, S.; Padura, I.M.; Libby, P.; Ceska, M.; Lindley, L.; Colotta, F. and Mantovani, A.(1991):

Expression of monocyte chemotactic protein and interleukin-8 by cytokine activated human vascular smooth muscle cells .

Arterioscler Thromb., 11:1166.

Wasserman, S.I.(1991):

Mediators of inflammation . IN: Stites, D.P. and Terr, A.I. (eds): Basic and Clinical Immunology . 7th edition . Appleton and Lange, London, Sydney, Toronto, Mexico, New Delhi, Chapter 13, p 154.

Wegelius, O.; Wafin,F.; Flack, H. and Tomroth, T.(1979):

Follow up study of amyloidosis secondary to rheumatic disease . IN : Proceedings of the third international symposium on amyloidosis, Glenner, G.G.; Costa, P.P.; Freitas, A.F., Eds. Excerpta Medica Amesterdam, P. 1639.

Weinstein , A. ; Peters , S. ; Bronn , S. and Blueston , R.(1972):

Metabolism of the third component of complement(C_3)in patients with RA. Arthritis Rheumat. , 15 : 49.

White, M.; Yoshimura, T.; Hook, W. and Leonard, E.(1989):

Neutrophil attractant / activation protien (NAP-1) causes human basophil histamine release . Immunol. Lett., 22:151.

Wibell , L.(1976):

Studies on B_2 - M in patients and normal subjects . Acta. Clin. Belg. , 31 : 14 .

Wickremasinghe, R.G.; Mire-Sluis, A.R. and Hoffbrand, A.B.(1987):

Interleukin-2 binding to activated human T lymphocytes triggers generation of cAMP but not inositol phosphates . FEES Lett., 220:52.

Williams, D.E. and Broxmeyer, H.E.(1988):

IL-1 alpha enhances the in vitro survival of purified murine granulocyte-macrophage progenitor cells in the absence of colony stimulating factors . Blood, 72: 1608.

Williams, D.; Morrissey, P. and Broxmeyer, H.E.(1990):

T cell growth factor p40 promotes the proliferation of myeloid cell lines and enhances erythroid burst formation by normal murine bone marrow cells in vitro . Blood, 76:906.

Wirth, J.J.; Kierszenbaum, F. and Zlotnik A.(1989):

Effects of IL-4 on macrophage functions; increased uptake and killing of a protozoan parasite (*Trypanosoma cruzi*) . Immunol., 66:296.

Wood, N.C.; Symons, J.A. and Duff, G.W.(1988):

Serum soluble Interleukin-2 receptor in rheumatoid arthritis: A prognostic indicator of disease activity .

J. Autoimmun., 1:353.

Wu, C.Y.; Fargeas, C.; Nakajima, T. and Delespesse, G.(1991):

Glucocorticoids suppress the production of interleukin-4 by human lymphocytes .

Eur. J. Immunol., 21 :2645.

Yamaguchi, Y.; Suda, T.; Shiozaki, H.; Hitoshi, Y.; Tominaga, A.; Takatsu, K. and Kasahara, T.(1990):

Role of IL-5 in IL-2 induced eosinophilia; in vivo and in vitro expression of IL-5 mRNA by IL-2 .

J. Immunol., 145: 873.

Yamasaki, K.; Taga, T.; Hirata, Y.; Ywata, H.; Kawanishi, Y.; Seed, B.; Taniguchi, T.; Hirano, T. and Kishimoto, T.(1988):

Cloning and expression of the human interleukin-6 (BSF-2/IFN β) receptor. Science, 241:825.

Yang, Y.C.; Ricciardi, S.; Ciarletta, A.; Calvetti, J.; Kelleher, K. and Clark, S.C.(1989):

Expression cloning of cDNA encoding a novel human hematopoietic growth factor; human homologue of murine T cell growth factor p40 .

Blood, 74:1880.

Yaron, I.; Meyer, F.A.; Dayer, J.M.; Bleiberg, I. and Yaron, M. (1989):

Some recombinant human cytokines stimulate glycosaminoglycan synthesis in human synovial fibroblasts cultures and inhibit it in human articular cartilage cultures .

Arthritis Rheum., 32:173.

Yin, T.; Miyazawa, K. and Yang, Y.C.(1992):

Characterization of interleukin-11 receptor and protein tyrosine phosphorylation induced by interleukin-11 in mouse 3T2-L1 cells .

J. Biol. Chem., 267 :8347.

Yocun, D.E.; Esparza, L.; Dubry, S. and Scuderi, P.(1989):

Characteristics of TNF production in RA.

Cell Immunol., 122:131.

Yoshimura, T.; Robinson, E.A.; Appela, E.; Matsubama, K.; Showalter, S.D.; Skeel, A. and Leonard, E.J.(1989):

Three forms of monocyte derived neutrophil chemotactic factor (MDNCF) distinguished by different lengths of amino-terminal sequence .

Mol. Immunol., 26:87.

Yoshizawa, H.; Chang, A.E. and Shu, S.(1991):

Specific adaptive immunotherapy mediated by tumor-draining lymph node cells sequentially activated with anti-CD3 and IL-2 .

J. Immunol., 147:729.

Young, J.C.; Gishizky, M.L. and White, O.N.(1991):

Hyperexpression of interleukin-7 is not necessary or sufficient for transformation of a pre-B lymphoid cell line .

Mol. cell Biol., 11:854.

Zacharaski, L.R. and Kyle, R.A.(1967):

Significance of extensive evaluation of ESR .

JAMA., 202:264.

Zhang, Y.; Lin, J.X. and Vicek, J.(1988):

Synthesis of interleukin-6 (interferon- γ /B-cell stimulatory factor 2) in human fibroblasts is triggered by increase in intracellular cyclic AMP .

J. Biol. Chem., 263:6177.

Zvaifler, N.J.(1988):

New perspectives on the pathogenesis of rheumatoid arthritis .

Am. J. Med. 85A:12.

Zvaifler, N.J. and Schur, P.(1968):

Reactions of aggregated mercaptoethanol treated gammaglobulin with rheumatoid factor-precipitin and complement fixation studies .

Arthritis Rheum., 11:523.

Zubiaga, A.N.; Munoz, E. and Huber, B.T.(1991):

Production of IL-1 α by activated TH type 2 cells . Its role as an autocrine growth factor .

J. Immunol., 146: 3849.

Zurawski, G. and De Vries, J. E. (1994):

IL-13 : an IL-4 like cytokine that acts on monocytes and B cells , but not on T. cells .

Immunol. Today , 15 : 19 .

Table (25) : Clinical and Hematological data of the studied subjects :

No.	Age(y)	Dis.dur.	ttt	SC nod.	Hb(g/dl)	RBCs(mil)	WBCs	Plat(1000)	ESR(mm)
1	47	4	NSAI	-	11.0	3.50	6900	310	55
2	34	4	CS	-	11.0	3.60	6500	250	30
3	66	5	CS	+	8.3	2.90	8500	380	70
4	27	4	CS	-	11.9	3.90	7150	290	30
5	50	3	NSAI	+	10.6	3.20	7200	320	55
6	36	5	CS	-	10.5	3.40	8650	260	40
7	45	6	NSAI	-	9.3	3.20	7100	240	35
8	50	7	NSAI	-	10.6	3.10	9100	400	70
9	35	6	NSAI	-	8.6	3.00	8200	420	60
10	32	5	NSAI	-	9.3	2.50	6000	290	20
11	50	6	NSAI	-	10.2	3.60	6250	250	20
12	32	6	NSAI	-	10.3	3.60	6000	260	22
13	45	4	IS	-	10.8	3.90	6500	260	25
14	29	5	IS	-	12.0	4.10	7100	240	28
15	32	5	IS	-	11.2	3.80	6200	260	20
16	37	7	NSAI	-	10.4	3.20	7100	300	30
17	41	5	IS	-	11.0	3.30	6900	250	25
18	50	6	IS	-	9.5	3.90	8100	400	60
19	41	6	NSAI	+	10.8	3.60	9000	390	50
20	48	7	IS	-	10.5	3.40	8600	380	45
21	32	5	NSAI	-	11.2	4.00	7100	240	20
22	40	11	NSAI	-	11.3	3.80	6100	250	20
23	42	7	NSAI	-	11.5	3.90	5750	260	25
24	40	8	NSAI	-	9.5	3.11	6750	300	30
25	38	7	CS	-	10.5	3.20	6200	240	28
26	42	4	NSAI	-	10.0	3.00	7000	290	20
27	39	7	NSAI	-	10.8	3.60	6500	280	28
28	47	3	IS	-	11.0	4.10	8750	280	50
29	52	5	NSAI	-	11.0	3.50	5100	270	25
30	38	6	CS	-	10.5	3.20	10000	410	45
31	40	5	CS	+	11.0	3.10	9250	250	40
32	35	6	CS	+	10.3	3.30	8600	350	50
33	30	3	NSAI	-	11.4	3.30	8600	200	25
34	49	13	NSAI	-	11.5	3.60	8000	220	30
35	30	3	CS	-	12.7	4.10	6100	260	20
36	40	11	NSAI	-	10.0	3.10	5600	230	25
37	32	12	NSAI	-	10.5	3.70	5200	260	25
38	38	7	CS	-	10.0	3.10	6850	250	20
39	40	8	NSAI	-	10.8	3.30	6000	270	28
40	51	5	IS	-	9.5	3.10	6750	260	25
41	30	00	-	-	14.0	4.50	5600	200	10
42	31	00	-	-	13.2	4.10	4500	190	12
43	29	00	-	-	13.0	4.00	5100	175	11
44	35	00	-	-	12.5	3.90	6500	210	15
45	40	00	-	-	13.5	4.10	4100	190	8
46	51	00	-	-	13.7	4.20	5200	185	9
47	43	00	-	-	12.7	3.98	4600	220	7
48	47	00	-	-	11.8	3.50	5250	175	10
49	66	00	-	-	12.9	3.90	4950	195	12
50	32	00	-	-	13.5	4.20	5650	160	15
51	45	00	-	-	14.2	4.60	5150	210	12
52	28	00	-	-	13.5	4.10	6100	240	13
53	37	00	-	-	13.0	3.90	4950	180	15
54	41	00	-	-	12.8	3.50	4800	198	10
55	50	00	-	-	12.5	3.45	5150	170	15
56	60	00	-	-	13.8	4.10	5000	250	10
57	61	00	-	-	13.0	4.00	4100	240	7
58	63	00	-	-	12.9	3.95	5050	200	8
59	42	00	-	-	12.8	3.80	5250	175	12
60	30	00	-	-	13.0	4.20	5900	185	15

Table (26) : Biochemical and Hematological data of the studied subjects:

No.	St.	Seg.	Eos.	Bas.	Lym.	Mono.	SGOT	SGPT	Alk.ph.	Creat.
1	2	64	2	0	30	2	18	16	3.0	0.8
2	2	62	0	0	36	0	9	7	0.8	0.9
3	4	58	2	0	36	0	30	28	3.2	0.6
4	2	68	0	0	26	0	22	25	3.2	0.9
5	2	64	0	0	32	2	25	23	2.2	0.7
6	2	68	0	0	24	2	18	16	1.2	0.8
7	4	70	2	0	26	0	17	15	2.0	1.5
8	2	62	2	0	32	2	20	21	2.8	0.5
9	4	62	0	1	33	0	25	22	2.5	0.5
10	0	72	2	0	24	2	20	22	1.3	0.7
11	2	52	2	0	40	4	10	8	2.0	0.6
12	2	60	0	0	36	2	17	18	2.1	0.7
13	4	50	2	0	44	0	25	22	1.9	1.1
14	2	62	2	0	32	2	20	22	2.8	1.3
15	2	70	0	0	28	0	18	20	2.5	1.2
16	4	68	0	0	26	2	44	30	2.8	1.5
17	0	66	2	0	30	2	35	25	3.2	1.2
18	2	56	0	0	40	2	10	12	1.5	1.2
19	4	70	2	0	24	0	11	13	1.2	1.1
20	2	58	0	0	38	2	10	15	2.3	1.2
21	2	58	0	0	38	2	22	20	1.8	1.3
22	4	70	2	0	24	0	18	14	2.6	1.1
23	2	56	0	0	40	2	12	14	1.5	1.2
24	0	66	2	0	30	2	18	16	1.9	1.2
25	4	28	0	0	26	2	15	17	2.0	0.9
26	2	70	0	0	28	0	32	40	3.1	1.3
27	2	62	2	0	32	2	11	10	1.7	0.8
28	4	50	2	0	44	0	42	38	3.2	1.3
29	2	60	0	0	36	2	12	10	2.9	0.9
30	2	52	2	0	40	4	20	18	1.2	0.6
31	0	72	2	0	24	2	19	18	1.4	0.7
32	4	63	0	1	32	0	19	17	1.9	1.1
33	2	62	2	0	32	2	20	17	2.8	1.0
34	4	70	2	0	26	0	23	16	1.5	0.7
35	0	64	2	0	34	0	24	18	2.0	1.5
36	2	48	0	0	48	2	18	16	1.2	0.9
37	2	64	2	0	30	2	29	25	2.3	0.8
38	2	62	0	0	36	0	20	29	1.5	1.0
39	4	58	2	0	34	2	29	27	1.9	2.1
40	2	26	0	0	26	0	18	26	2.3	1.1
41	2	60	2	0	34	2	10	12	1.0	0.9
42	4	58	0	0	38	0	8	9	0.8	0.7
43	6	50	4	0	40	0	11	10	0.9	0.5
44	2	66	2	0	26	4	13	11	0.7	0.6
45	2	58	0	0	38	2	18	20	1.1	0.9
46	0	56	4	0	36	4	7	9	1.0	1.1
47	2	60	2	0	34	2	9	8	0.8	0.5
48	6	50	4	0	40	0	11	13	0.7	0.6
49	2	58	4	0	34	2	13	15	0.7	0.9
50	2	60	2	0	36	0	10	12	0.6	0.8
51	6	50	4	0	40	0	11	10	1.1	0.7
52	2	70	0	0	26	2	9	8	1.3	0.5
53	4	48	2	0	42	4	10	9	0.7	1.0
54	4	62	4	0	28	2	11	7	1.0	0.5
55	2	54	2	1	37	4	9	7	0.8	0.5
56	4	56	0	0	40	0	10	8	0.6	0.4
57	4	60	4	0	30	2	10	9	0.7	0.6
58	2	56	2	0	36	4	13	11	1.0	0.4
59	6	50	4	0	40	0	11	13	0.6	0.5
60	2	54	4	0	36	4	8	10	0.7	0.4

No.	B ₂ -M	Activity	RF titer	CRP	IL-6	ANA
1	5.1	Active	72	42	412.00	+
2	1.5	Inactive	00	30	305.00	-
3	4.2	Active	96	60	721.00	-
4	2.3	Inactive	48	36	482.00	-
5	3.3	Active	84	42	546.00	-
6	3.8	Inactive	60	30	400.00	-
7	1.9	Inactive	48	24	395.00	-
8	4.3	Active	96	48	750.00	-
9	4.5	Active	96	48	780.00	-
10	1.5	Inactive	48	24	480.00	-
11	2.0	Inactive	36	24	200.00	-
12	1.8	Inactive	00	18	150.00	-
13	1.4	Inactive	48	24	489.00	-
14	1.8	Inactive	60	24	295.00	-
15	1.6	Inactive	48	18	480.00	-
16	3.0	Inactive	00	24	220.00	-
17	2.8	Inactive	36	18	160.00	-
18	4.0	Active	96	48	790.00	+
19	3.8	Active	96	48	800.00	-
20	2.9	Active	84	42	750.00	+
21	1.9	Inactive	48	24	36.00	-
22	2.1	Inactive	42	18	130.00	-
23	1.9	Inactive	42	24	200.00	-
24	1.5	Inactive	00	24	100.00	-
25	2.0	Inactive	00	24	160.00	-
26	2.2	Inactive	36	18	481.00	-
27	1.5	Inactive	36	24	110.00	-
28	3.7	Active	60	36	220.00	-
29	1.6	Inactive	00	18	95.00	-
30	3.8	Active	96	48	802.00	-
31	3.5	Inactive	84	42	220.00	-
32	3.2	Active	84	48	624.00	-
33	2.5	Inactive	48	24	105.00	-
34	2.0	Inactive	60	30	483.00	-
35	1.5	Inactive	00	12	200.00	-
36	2.0	Inactive	36	18	210.00	-
37	2.3	Inactive	36	24	250.00	-
38	3.1	Inactive	36	18	200.00	-
39	3.4	Inactive	36	24	300.00	-
40	2.9	Inactive	48	24	105.00	-
41	0.6	--	00	00	14.12	-
42	1.0	--	36	00	23.50	-
43	1.3	--	00	12	35.12	-
44	0.8	--	00	00	15.10	-
45	1.2	--	00	00	13.65	-
46	0.7	--	00	00	13.19	-
47	2.0	--	36	00	24.11	-
48	0.9	--	00	00	34.35	-
49	1.5	--	00	12	15.13	-
50	1.8	--	00	00	25.19	-
51	0.9	--	00	00	44.69	-
52	0.7	--	00	18	13.11	-
53	1.1	--	00	00	7.21	-
54	0.6	--	00	00	14.16	-
55	1.0	--	00	00	14.95	-
56	0.7	--	00	00	23.02	-
57	0.8	--	00	12	34.65	-
58	1.2	--	00	00	13.19	-
59	0.9	--	00	18	13.65	-
60	0.6	--	00	18	23.48	-