




REFERENCES

References

Abbas, A.K. , Lichtman, A.H. and Pober, J.S. (1994):-

Immunity to viruses. In:Cellular and molecular immunology (2nd edition).Saunders ,W.B.(ed.);Saunders Company ; Philadelphia; London ; Toranto ; Montreal ; Sydney ;Tokyo ;1994: pp. 273-329

Abd el-Hadi A.M. (1987):

A histopathological and immunohistochemical study of collagen in schistosomal hepatic fibrosis before and after chemotherapy. PH.D. Thesis, Al-Azhar Univ., PP. 37-48.

Abd el-Kader S., Amin M., Hamdy H. (1997):

Causes of minimal hepatic fibrosis present in Egypt. J. Egypt Soc. Parasitol., 27 (3): 919-928.

Abdel Wahab, M.F. , Zakaria, S. , and Kamal, M (1994):

High seroprevalence of hepatitis C infection among risk groups in Egypt. Am. J. Trop. Med. Hyg: 51(5):563-7.

Adami H., Hsing A. and Mclaughlin J. (1992):

Alcoholism and liver cirrhosis in the aetiology of primary liver cancer. Int. J. cancer, 51: 898-902.

Ahmed E., Daffala A., Christensen N. (1996):

Pattern of infection and transmission of human schistosomiasis. Ann. Trop. Parasitol. 90 (2) : 173-181.

Alberti, A. (1998):

Recent advances in understanding HCV. International Conference on Hepatitis C. Monte Carlo, 9-11 January, (Abstr.).

Alberti, A. , Morsica, G. and Chemello, L. (1992) :

Hepatitis C viremia and liver disease in symptom free individuals. *Lancet*; , 340:697-698.

Alexander, G.J.M. (1990) :

Immunology of hepatitis B virus infection. *British and Liver Disease Medical Bulletin* ;1990 – 46:345-367.

Alter, H.J (1989):

Chronic consequences of non-A, non-B hepatitis. In: *Current perspectives in Hepatology*. Seel, L.B. and Lewis, J.H. (eds.), Plenum Publishing, New York; pp. 83-97.

Alter, H.J. (1990):

Clinical, virological and epidemiological basis for the treatment of chronic non-A, non-B hepatitis. *Journal of Hepatology*; 11 (suppl.1):S 19-S25.

Alter, M.J. (1997) :

Hepatitis C, clinical spectrum of disease. National Institute of health conference on hepatitis C. Bethesda Maryland, p.24.

Andrade Z. A., Santana S. and Rubin E. (1962):

Hepatic changes in advanced schistosomiasis. *Gastroenterology*. 42, 393.

Angeli P., Gatta A., Caregaro L. (1990):

Tubular site of renal sodium retention in ascitic liver cirrhosis evaluated by Lithium clearance. *Eur. J. Clin. Invest.*; 20:117.

Angelico M., Renganathan E., Gandin C. (1997):

Chronic liver disease in the Alexandria governorate, Egypt: Contribution of schistosomiasis and hepatitis virus infections. *Hepatology*, 26: 236-243.

Arakaki_Arakaki N, Kawakami S, Nakamura O, Ohnishi T, Miyazaki H, Ishii T, Tsubouchi H. (1995):

Evidence for the presence of an inactive precursor of human hepatocyte growth factor in plasma and sera of patients with liver diseases. *Hepatology* Dec ; 22(6) : 1728-34.

Arroyo V., Gines P., Alexander L., Dudmeyer F.J. and Gentilini P. (1996):

Definition and diagnostic criteria of refractory ascites and hepatorenal syndrome in cirrhosis. *Hepatology*, 23 (1): 164.

Arroyo V., Gines P., Jimenez W. and Rodes J. (1999):

Renal dysfunction in cirrhosis. In: *Oxford Textbook Of Clinical Hepatology*. Bircher J., Behamou J.P., McIntyre N., Rizzeto M. and Rodes J. (eds), 2nd edition, Oxford: Oxford University press. Volume I, p: 734.

Arroyo V., and Rodis J.A. (1975):

A rational approach to the treatment of ascitis. *Postgrad. Med. J.*; 51:558.

Arroyo V., and Rodis J. (1976):

Prognostic value of spontaneous hyponatraemia in cirrhosis with ascitis. *Am.J.Dig.Dis.*; 21:249.

Aufiro B. and Schneider R.J. (1990):

The hepatitis B virus X-gene product transactivates both RNA polymerase II and III promoters EMBO J. 9:497-504.

Bach N., Kahn H., Thung S.N. (1991):

Hepatocellular carcinoma in a long-term survivor of intrahepatic biliary duct hypoplasia. Am.J. gastroenterol, 86: 1527-1530.

Badalamenti S., Graziani G., Salerno F. and Fonticelli C. (1993):

Hepatorenal syndrome new perspectives in pathogenesis and treatment. Arch. Intern. Med., 153 (Sep. 13): 1957.

Baddour N., and Yonis L. (1996):

Causal relationship between inflammation and hepatocytes regeneration: An immunohistochemical study using PCNA. J. Hep. Gastroenterol. Infect. Dis. JHGID 4 (2): PP. 27:35.

Badizadegan K., Atayde P., Jonas M. (1998):

Histopathology in liver in children with chronic hepatitis C viral infection. Hepatology, 28:1416-1423.

Bahr, M. , Bell, D.R. and Walz, A. (1987) :

Viral hepatitis. In : Manson's tropical disease ; 14th edition ; English Book Society ; London, 170-8.

Ballaridini G., Groff P., Zoli M. (1994):

Increased risk of hepatocellular carcinoma development in patients with cirrhosis and high hepatocellular proliferation. J. Hepatol.; 20:218-222.

Ballistreri F. and Rej R. (1996):

Liver function. In: Tietz fundamentals of clinical chemistry. Burtic A and Ashwood R (eds), 4th edition, Philadelphia: W.B. Saunders Company, chapter 31, p 539.

Barrera, J. , Brancis, B. , Ercilla, G. and Pontisso, P. (1995):

Improved detection of anti-HCV in post-transfusion hepatitis by a third-generation ELISA. Vox. Sanguinis, 68:15-8.

Bartolozzi, C. (2000):

Diagnosis and staging of hepatocellular carcinoma. In: Omar, S.; El-Mawla, N.; El-Zawahry, H. (2000) In European School of Oncology. Educational Course on Liver Tumors, P: 52.

Bassily, S. , Farid, Z. and Higashi, G. (1979):

Chronic hepatitis Bs antigenaemia in patients with hepatosplenic chistosomiasis. J. Trop. Med. Hyg. ; 82:248-251

Bazarbashi, S. (2000):

Hepatocellular carcinoma; Epidemiology and Etiology. In: Omar, S. El-Mawla, N.; El-Zawahry, H. (eds.) European School of Oncology Educational courses on Liver Tumors, P: 29.

Beasley, R.P. , Hwang, L.Y. and Lin, C.C. (1981):

Hepatitis B immune globulin (HBIG) efficacy in the interruption of perinatal transmission of hepatitis B virus carrier state. Initial report of a randomized double-blind placebo-controlled trial. J.Infec. Dis. ; 2:388-393.

Becker M.D., Schewer P.J., Baptista A. and Sherlock S. (1970):

Prognosis of chronic persistent hepatitis. Lancet: 53-56.

Benhamou J.L. and Lebrec D. (1985):

Non-cirrhotic intrahepatic portal hypertension in adults. Clin. Gastroenterol., 14:21.

Bennet, J.C. , Plum, F. , Jill, G.N. , Kokko, J.P. , Mandell, G.L. , Ockner, R.K. and Smith, J.W. (1996) :

Chronic hepatitis ; Cecil Textbook of Medicine ;20th edition ; W.B. Saunders Company : Philadelphia ; 776-780.

Bernardi M., Di Marco C., Trevisani F. (1993):

Renal sodium retention during upright posture in preascitic cirrhosis. Gastroenterology; 105-188.

Bernardi M., Trevisame F., Santini C. (1983):

Aldosterone related blood volume expansion in cirrhotic before and after the early phase of ascitis formation. Gut; 24:761.

Bianchi, L. (1993):

Glycogen storage disease and hepatocellular tumours. Eur J Pediatr 152: 62-70.

Bilezikci B., Haberal A. N. and Demirhan B. (2001):

Hepatocyte growth factor in patients with three different stages of chronic liver disease including hepatocellular carcinoma, cirrhosis and chronic hepatitis: an immunohistochemical study. Can. J. Gastroenterol, 15 (3): 159-65.

Bird G., Lau Y., Koskinasa J. (1991):

Insulin and glucagon infusion in acute alcoholic hepatitis. A prospective randomized controlled trial. Hepatology; 14(6): 1097-1101.

Block E., Addel-Wahab, M. and Warron K. (1972):

In vivo microscopic observation of the pathogenesis and pathophysiology of hepatosplenic schistosomiasis in mouse liver. *Am.J. Trop. Med. Hyg.* 21:546.

Borzio M., Trere D., Borzio F. (1998):

Hepatocyte proliferation rate is a powerful parameter for predicting hepatocellular carcinoma development in liver cirrhosis. *Mol. Pathol.*; 51 (2) : 96-101.

Bosch, F.X. (1997):

Global epidemiology of hepatocellular carcinoma in: Okuda K and Edward T. *Liver cancer* 1st edition. Ch. 2. P: 13-28 Churchill Livingstone.

Bottaro D.P., Rubin J.S., Faletto D.L. (1991):

Identification of the hepatocyte growth factor receptor as the C-met protooncogene product, *science* 251:802-804.

Bradbear R.A., Bain C., Siskind V. (1985):

Cohort study of internal malignancy in genetic hemochromatosis and other chronic nonalcoholic liver diseases. *J Natl Cancer Inst* 75:81-84.

Brechot C., Pourcol C., Louis A. (1980):

Presence of integrated hepatitis B virus DNA sequences in cellular DNA in human hepatocellular carcinoma. *Nature*, 533-535.

Brunetto, M.R. , Oliveri, F. and Rocca, G. (1989):

Natural course and response to interferon in chronic type B hepatitis, accompanied by antibody to hepatitis Be antigen. *Hepatology*; 20:198-202

Bucher N. (1991):

Liver regeneration: an overview. *J. Gastroenterol Hepatol*; 6 (6): 615-624.

Buendia M. (1998):

Hepatitis B virus and carcinogenesis. *Biomed pharma cother*; 52 (1): 34-43.

Busby W.F.Jr. and Wogan G.N. (1984):

Aflatoxins. In Searle CE (ed.): *Chemical carcinogens 2nd Ed. Vol. 12.* American Chemical Society Washington, Dc, PP: 945-1136.

BWahlefeld, A.W. (1978) :

Optimization of methods for aspartate aminotransferase and alanine aminotransferase. *Clin. Chem.*, 24/1:58-73

Casey C., Camacho K., Tuma D. (1992):

The effects of chronic ethanol administration on the rates of internalization of various ligands during hepatic endocytosis *Biochim, Biophys Acta*; 1134 (2): 96-104.

Cazaja A., and carpenter H. (1997):

Histological finding in chronic hepatitis C with autoimmune features *Hepatology*, 26: 459-466.

Cerny, A. , Fowler, C. , Brothers, M.A. , Bradley, D.W. and Miller, R.H.(1995) :

Induction in vitro of a primary human **antiviral** cytotoxic T cell response. *Eur. J. ImmunoL*, 25:627-630.

Cerny, A., McHutchinson, J.G. , Pasquinelli, C. , Brothers, M.A. and Folwler, C. (1995b) :-

Cytotoxic T lymphocyte response to hepatitis C virus-derived peptides containing the HLA-A2.1 binding motif. *J. Clin. Invest.*, 95:521-530.

Chan, C.Y. , Lee, S.D. , Hwang, S.J. , McOmish, F. and Holmes, E.C. (1995) :-

Quantitative branched DNA assay and genotyping for hepatitis C virus in Chinese patients with acute and chronic hepatitis C. *J. Infect. Dis.*, 171:443-446.

Chang, K.M. , Reherrmann, B. and McHutchinson, J.G. (1997) :-

Cytotoxic T lymphocyte epitope variants in patients chronically infected with the hepatitis C virus. 4th International meeting on hepatitis C and related viruses, Kyoto, Japan (Abstr.). *Hepatology*; 22:218A

Chau, K.H. , Dawson, G.J. , Mushahwar, I.K. , Gutierrez, R.A. , Johnson, R.G. , Lesniewski, R. , Mattson, L. and Weiland, O. (1991) :

IgM antibody response to hepatitis C virus antigens in acute and chronic post-transfusion NANB hepatitis. *J. Virol. Methods*; 35(3):343-352.

Chen, D.S. (1993):

From hepatitis to hepatoma. Lessons from type B viral hepatitis science 262: 269-270.

Chen D.S., Hsu N.H.M., Sung J.L. (1987):

A mass vaccination program in Taiwan against hepatitis B virus infection in infants of Hepatitis B surface antigen carrier mothers. *JAMA* 257: 2597-2608.

Chen S.H., Hu C.P. and Chang C. (1992):

Hepatitis B virus replication in well differentiated mouse hepatocyte cell lines transformed by plasmid DNA. *Cancer Res*, 52: 1329-1335.

Chen S.H., Hu C., Lee C.K. and Chang C. (1993):

Immune reactions against hepatitis B viral antigens lead to rejection of hepatocellular carcinoma in BALB/c mice. *Cancer Res*, 53: 4648-4651.

Cheng W.S., Govindarajan S. and Redeker A.G. (1992):

Hepatocellular carcinoma in a case of Wilson's disease. *Liver* 12: 42-45.

Chu, C.M. , Karayiannis, P. , Craxi, A. , Montano, L. , Goodall, A. , Thomas, H.C. and Fowler, M.J. (1985) :-

Natural history of chronic hepatitis B virus infection in Taiwan : Studies of HBV DNA in serum. *Hepatology*; 5:431-434

Colombo M., DeFranchis R., Del Ninna . (1991):

Hepatocellular carcinoma in italian patient with cirrhosis *N Engl. J. Med.* 325; 675-680.

Colombo M., Rumi M.G., Damato M.F. (1991):

Hepatitis C antibody in patient with chronic liver disease and hepatocellular carcinoma. *J. Med. Viral.* 43: 125-128.

Colucci G., Colombo M. and Ninno E. (1983):

In situ characterization by monoclonal antibodies of the mononuclear cell infiltrate in chronic active hepatitis *Gastroenterology*; 85-237.

Conn H.O. (1973):

A rational approach to the hepatorenal syndrome. *Gastroenterol.*, 65:321.

DeGrotte J., Desmet V.J., Gedigk, P. (1968):

A clasification of chronic hepatitis *Lancet*; 48-66.

Delhaye M., Louis H., Degraef C. (1996):

Relationship between hepatocyte proliferative activity and liver functional reserve in human cirrhosis. *Hepatology*; 23: 1003-1011.

Derenzini M., (2000):

The Ag NORs. *Micron* 31 (2): 117-20.

Dervan P., Magee H., Buchley C. (1992):

Proliferating cell nuclear antigen counts in formalin fixed paraffinembedded tissue correlate with Ki-67 in fresh tissue. *Am.J.Clin. Pathol.* 97 (Supp): S21-S28.

Desmet, V.J. , Gerber, M. , Hoofnagle, J.M. , Manns, M. and Scheuer, P.J.(994) :

Classification of chronic hepatitis : Diagnosis, grading and staging. *Hepatology*, 19(6): 1513-1520.

Desmet V.J., Sciote R., and Van Eyken P. (1990):

Differential diagnosis and prognosis of cirrhosis: role of liver biopsy: *Acta Gastroenterology*; Belg.L. III: 198-208.

Deuginer Y., Guyader D., Crantock, I. (1993):

Primary liver cancer in genetic hemochromatosis: a clinical pathological and pathogenetic study of 54 cases. *Gastroenterology*, 104: 228-234.

Diehl A. and Rai R. (1999):

Liver regeneration. In: Schiff E., Sorvell M. Maddry W. (eds): Schiff's diseases of the liver "8th edition" lippincott- Raven publishers, philadelphia, 39-52.

Dienes, H.P. , Popper, H. and Arnold, W. (1982):

Histological observations in human nori-A, non-B hepatitis. Hepatology, 2:552.

Doria, M., Klein, N., Lucito, R. (1995):

The hepatitis B virus HBX protein is a dual specificity cytoplasmic activator of Ras and nuclear activator of transcription factors EMBO J. 14: 4797-4757.

Doughty B., Ottesen E. and Nash T. (1984):

Delayed hypersensitivity granuloma formation around S. mansoni egg in vitro granuloma formation and modulation in human S. Mansoni. J. of Immunol.; 133:993.

Dusheiko, G.M. (1992):

Hepatitis C virus. In: Recent advances in gastroenterology 9 ; Pounder, R. ed. ; Churchill Livingstone ; New York ; 195:216.

Eguchi S., Kamlot A., Ljubimova J. (1996):

Fulminant hepatic failure in rats: Survival and effect on blood chemistry and liver regeneration. Hepatology; 24 (6): 1452-1459.

Eisenmenger W.J., Blondheim S.H., Bongiovanni A. M. (1950):

Electrolyte studies on patients with cirrhosis of the liver. J. Clin. Invest.; 29: 1491.

Fagan, E.A. and Williams, R.C. (1990) :

Fulminant viral hepatitis. British Medical Bulletin; 46:462-480. Failla, C. ; Tomei, L. and De Francesco, R. (1994) : Both NS3 and NS4 are required for proteolytic processing of hepatitis C virus nonstructural proteins. J. ViroL, 68:3753-3760.

Farghaly A. and Barakat R. (1993):

Prevalence impact and risk factors of hepatitis C infection. J. Egypt Public Health Assoc.; 68 (1-2): 63-75.

Fargion S., Mandelli C., Piperno A. (1992):

Survival and prognostic factors in 212 Italian patients with genetic hemochromatosis. Hepatology, 15: 655-639.

Fausto N., Laird A.D., Webber E.m. (1995):

Liver regeneration. 2. Role of growth factors and cytokines in hepatic regeneration. Faseb. J. 9 (15): 1527-1536.

Feitelson M.A., Zhu, M., Duan L.X. and London W.T. (1993):

Hepatitis B X antigen and p53 associated in vitro and in liver tissues from patients with primary hepatocellular carcinoma. Oncogene, 8: 1109-1117.

Flicher MS., Packham D.A., Pryor J.p. and Yates A.J. (1981):

Hepatic dysfunction in renal carcinoma. Br. J. Urol., 53:536.

Foschi D., Castoldi L., Lesma A. (1993):

Effects of ischemia and reperfusion on liver regeneration in rats. Eur. J. Surg.; 159 (8): 393-398.

Francavilla A., Barone. M., Van thiel DH. (1991):

Further steps of hepatic stimulatory substance purification Dig Dis Sci. 36:674-680.

French S.W. (1981):

Nature, pathogenesis and significance of the Mallory body. Semin. Liver Dis., 1:217.

Fujimoto J. (2000):

Gene therapy for liver cirrhosis. J. Gastroenterol. Hepatol.: 15 Suppl: D33-6.

Ghaffar Y., Kamel M., Abdel Wahab M. (1990):

Hepatitis B vaccination in children infected with schistosoma Mansoni: Correlation with ultrasonographic data. Am. J. Trop. Med. Hyg.; 43: 516-519.

Gines P., Arroyo V. and Rodes J. (1996):

Disorders of renal function in cirrhosis. In: Hepatology. A Textbook Of Liver Disease. 3rd edition, Zakim D and Boyer TD (eds), Philadelphia: W.B. Saunders company, p: 650.

Gohda E., Tsubouchi H., Nakayama H. (1988):

Purification and partial characterization of hepatocyte growth factor from plasma of a patient with fulminant hepatic failure, J. Clin. Invest. 81: 414-419.

Gohda E., Yamaski T., Tsubouchi H. (1990):

Biological and immunological properties of human hepatocyte growth factor from plasma of fulminant hepatic failure. Biochim. Biophys. 1053:21-6.

Goldstein N., Blue D., Hankin R. (1999):

Serum alpha feto-protein levels in patients with chronic hepatitis C. Relationships with serum alanine transferase values, histologic activity index, and hepatocyte MIB-1 scores. *Am. J. Clin. Pathol.*, 111 (6), 811-6.

Goodman Z., and Ishak K. (1995):

Histopathology of hepatitis C virus infection. *Sem. Liv. Dis.*; 15: 70-81.

Grimaud J. and Borojevic R. (1976):

Schistosomal pigments in human and experimental infection with *S. mansoni*. *Trans. Roy. Soc. Trop. Med. Hg.*, 70 (1): 73-88.

Grimaud J. and Borojevic R. (1986):

Portal fibrosis: intrahepatic portal vein pathology in chronic human *S. mansoni*. *J. submicrosc. Cytol.*, 18 (4): 783-794.

Groopman J.D., Donahue P.R., Zhu J.Q. (1985):

Alfatoxin metabolism in humans: detection of metabolites and nucleic acid adducts in urine by affinity chromatography. *Proc. Natl. Acad. Sci, USA*, 82: 6492-6496.

Gretch, D. , Corey, L. and Wilson, J. (1994) :-

Assessment of hepatitis C virus RNA by quantitative competitive RNA polymerase chain reaction: high-titer viremia correlates with advanced stage of disease. *J. Infec. Dis.*, 169:1219-1225.

Gretch, D. , Dela Rosa, D. , Corey, L. and Wilson, J. (1996) :

Assessment of hepatitis C viremia using molecular amplification technologies. *Viral hepatitis Reviews*, 2:85-96.

Hananoue M., Kawaida K., Takao S., Shimizu H., Noji S. and Matsumoto K. (1992):

Rapid and marked induction of hepatocyte-growth factor during liver regeneration after ischemic or cruch injury *Hepatology* 16, 1485-1492.

Harold O.C. and Colin E.A. (1993):

Cirrhosis. In: Leon Schiff and Eugene R Schiff (eds.). *Diseases of the liver*, 7th (ed.), vol. 2, J. B. Lippincott Company, Philadelphia, PP 875.

Haque S., Chandra B., Gerber M.A. (1996):

Iron overload in patients with chronic hepatitis C: a clinicopathologic study. *Hum. Pathol.*; 27: 1277-1291.

Helal T., Danial M. and Ahmed H. (1998):

The relationship between hepatitis C. virus and schistosomiasis: Histopathological evaluation of liver biopsy specimens. *Hum. Pathol.* 29 (7) 743-760.

Helwing F.C. and Schutz C.B. (1932):

A liver kidney syndrome. Clinical, pathological and Experimental studies. *Surg. Gyecol. Obstet.*, 55:570.

Higashio K., Shina N., Goto M. (1990):

Identity of a tumor cytotoxic factor from human fibroblasts and hepatocyte growth factor, *Biochem Biophys Res. Commun.* 170:397-404.

Hioki O., Watanabe- A., Minemura -M. and Tsuchida- T. (1993):

TI Clinical significance of serum hepatocyte growth factor levels in *J-Med.* VOL: 24 (1), P: 35-46.

Hirano T, Fujimoto J, Ueki T. (1998) :

Persistent gene expression in rat liver *in vivo* by retroviral transfections using HIV-liposome. *Gene Ther*; 5: 459-64

Ho R.t., Liew C.T. and Lai KN. (1999):

The expression of hepatocyte growth factor (HGF) and interleukin 6 (IL-6) in damaged human liver and kidney tissues. *Hepatogastroenterology*; 46 (27): 1904-9.

Hohne, M., Schaefer, S., Seifer, M. (1990):

Malignant transformation of immortalized transgenic hepatocytes after transfection with hepatitis B virus DNA. *EMBO J* 9: 1134-1145.

Ikeda K., Saitoh S., Koida I. (1993):

A multivariate analysis of risk factors for hepatocellular carcinogenesis: a prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology*, 18: 47-53.

Ilan Y., Roy-Chowdhury N., Prakash R. (1997):

Massive repopulation of rat liver by transplantation of hepatocytes into specific lobes of the liver and ligation of portal vein branches to other lobes *Transplantation*; 64 (1): 8-13.

Ishak K.G., Baptista A. Bianchi. (1995):

Histological grading and staging of chronic hepatitis. *J. Hepatol*; 22: 696-699.

Ishi T., Sato M., Sudo K., Suguki M., Nakai H., Hishida T., Niwa T., Umezu K. and Yuasa S. (1995):

Hepatocyte growth factor stimulates liver regeneration and elevates blood protein level in normal and partially hepatectomized rats (Bibliographic Citation): *J-Biochem-Tokyo*.

Ishiki Y., Ohnishi H., Muto Y., Matsumoto K. and Nakamura T. (1992):

Direct evidence that hepatocyte growth factor is a hepatotrophic factor for liver regeneration and has a potent antihepatitis effect in vivo *Hepatology* 16, 1227:1235.

Jameson, C. (1989):

Primary hepatocellular carcinoma in hereditary haemorrhagic telangiectasia. *Histopathology*, 15: 550-552.

Jeffers, L.J. , Dailey, P.J. and Coelho-little, E. (1993) :

Correlation of HCV-RNA quantitation in sera and liver tissue of patients with chronic hepatitis C. *Gastroenterology*; 106:A866.

Joly A., Desjardins J., Fremond B., (1997):

Survival, proliferation and functions of porcine hepatocytes encapsulated in coated alginate beads: a step toward a reliable bioartificial liver. *Transplantation*; 63 (6): 795-803.

Jones H., Clark N., Barrass N., (1993):

Phenobarbital-induced hepatocellular proliferation: anti-bromodeoxyuridine and anti-proliferating cell nuclear antigen immunocytochemistry. *J. histochem Cytochem*, 41: 21-27.

Kage M. Shimamtu K., Nakashima E. (1997):

Long-term evolution of fibrosis from chronic hepatitis to cirrhosis in patients with chronic hepatitis C: Morphometric analysis of repeated biopsies. *Hepatology*, 25: 1028-1031.

Kakumu, S. , Yoshioka, K. , Wakita, T. , Ishikawa, T. , Takayanagi, M. and ., Higashi, Y. (1989) :

A pilot study of ribavirin and interferon beta for the treatment of chronic hepatitis C. *Gastroenterology*; 105:505-512.

Kan .M, Huan J , Mansson P , Yasumitsu H . (1989):

Heparin – binding growth factor type1 (acidic fibroblast growth factor) : a potential biphasic autocrine and paracrine regulator of hepatocyte regeneration . Proc. Nat. Acad . Sci. USA ; 86 : 7432-7436.

Kan M., Zhang G., Zarnegar R. (1991):

Hepatocyte growth factor-hepatopoietin A stimulates the growth of rat kidney proximal tubular epithelial cells (RPTE), rat non parenchymal liver cells, human melanoma cells, mouse keratinocytes and stimulates anchorage independent growth of SV-40 transformed RPTE, Biochem Biophys Res. Commun 174: 331-337.

Kane, M. (1992) :

Control of hepatitis B virus infection through universal infant mmunization with hepatitis B vaccine. In : Viral hepatitis management Standards for the future. Cannes, France, p. 14.

Kaplan M.M. (1986):

Serum alkaline phosphatase: another piece is added to the puzzle. Hepatology, 6: 526.

Kawakita N., Seki S., Sakaguchi H. (1992):

Analysis of proliferating hepatocytes using monoclonal antibody against proliferating cell nuclear antigen/cyclin in embedded tissues for various liver diseases fixed in formaldehyde. Am. J. Patho., 140, 513-520.

Keeffe E., Pinson C., Ragsdale J. and Zonana J. (1993):

Hepatocellular carcinoma in arteriohepatic dysplasia. Am. J. Gastroenterol, 88: 1446-1448.

Kew M., Song E., Mohamed A. and Hodgkinson J. (1990):

Contraceptive steroids as a risk factor for hepatocellular carcinoma: a case control study in South African black woman. *Hepatology*, 11: 298-302.

Kinoshita, T., Hirao, S., Matsumoto, K. (1991):

Possible endocrine control by hepatocyte growth factor of liver regeneration after partial hepatectomy, *Biochem Biophys Res. Commun.* 177: 330-335.

Kinoshita T., Tashiro K., Nakamura T. (1989):

Marked increase of HGF mRNA in non parenchymal liver cells of rats treated with hepatotoxins, *Biochem Biophys Res. Commun* 105: 1229-1234.

Kirk A.P., Dooley J.S. and Hunt R.H. (1980):

Peptic ulceration in patients with chronic liver disease. *Digestive Diseases and Sciences*; 25: 756-760.

Kitamura T., Watanabe S., Sato N. (1998):

Liver regeneration, liver cancer and cyclins. *J. gastroenterol hepatol.*, 13 (Suppl.), S96-S99.

Kiyosawa, K. , Akahane, Y. and Nagata, A. (1984) :

Hepatocellular carcinoma after non-A, non-B transfusion hepatitis. *American Journal of Gastroenterology*; 79:777-781.

Kiyosawa, K. , Sodeyama, T. and Tanaka, E. (1990) :

Interrelationship of blood transfusion, non-A, non-B hepatitis and hepatocellular carcinoma : analysis by detection of antibody to hepatitis C virus. *Hepatology*; 12:671.

Koziel, M.J. (1997) :

The role of immune responses in the pathogenesis of hepatitis C virus infection. *J. Virol. Hepatitis*, 4(Suppl.2):31-41.

Koike K., Moriya, K., Lino, S. et al. (1994):

High level expression of hepatitis B virus HBx gene and hepatocarcinogenesis in transgenic mice *Hepatology*, 19: 810-819.

Kondo F., Ebara M., Surgiura N., (1990):

Histological features and clinical course of large regenerative nodules: evaluation of their precancerous potential, *Hepatology*; 12: 592.

Kren B.T., Trembley J.H., Krajewski S. (1996):

Modulation of apoptosis-associated genes bcl-2, bcl-x, and bax during rat liver regeneration. *Cell growth differ*; 7: 1633-1642.

Kunio Matsumoto and Toshikazu Nakamura (2001):

Hepatocyte growth factor : Renotropic role and potential therapeutics for renal diseases . *Kidney international* ; 59:2023-2038.

Lai K.N., Lui S.F., Li PKT Au T.C., Tam J.S., Tong K.L. and Lai F.M. (1991):

Membranous nephropathy related to hepatitis B virus in adult. *N. Engl. J. Med.*, 324:1457.

Lawrence S. (1996):

Chronic hepatitis. In: Lawrence M., Stephen J. and Maxine A. (eds): *Current medical diagnosis and treatment* "35th edition". Lange Medical publication 584-586.

Lencioni R., Bartolozzi, Caramella D. (1995):

Treatment of small hepatocellular carcinoma with percutaneous ethanol injection analysis of prognostic factors in 105 western patients. *Cancer* 76: 1737-1776.

Liaw Y.f., Tai D.Y., Chu C.M. (1986):

Early detection of hepatocellular carcinoma in patient with chronic type B hepatitis a prospective study. *Gastroenterology* 90: 263-267.

Liaw Y.F., Tsai S.L., Chang J.J. (1994):

Displacement of hepatitis B virus by hepatitis C virus as the cause of continuing chronic hepatitis. *Gastroenterology*, 106: 1048-1053.

Lim H.W. and Mascaro J.M. (1995):

The porphyrias and hepatocellular carcinoma. *Dermatol, Clin.* 13: 135-142.

Lindroos P.M., Zarnegar R., Michalopoulos G.K. (1991):

Hepatic growth factor (hepatopoietin A) rapidly increases in plasma before DNA synthesis and liver regeneration stimulated by partial hepatectomy and carbon tetrachloride administration, *Hepatology*, 13: 743-750.

Litofsky N., Mix. T., Baker S., (1998):

Ki-67 (Clone MIB-1) in recurrent glial neoplasms. No prognostic significance. *Surg. Neurol*; 50 (6): 579-585.

Liu KW, Kati Y, Yamazaki M, Nakamura T .,(1993)

Decrease in hepatic clearance of hepatocyte growth factor in carbon tetrachloride – intoxicated rats . *Hepatology* 17:65 -660 .

Liu KX, Kato Y., Narukawa M., Kim DC, (1922):

The importance of liver in the plasma clear of hepatocyte growth factor (HGF) in ratd . Am Jphysiol ;3:27-54.]

Lohr J.W., Lee T.P., Farooqui M. and Mookerjee B.K. (2000):

Increased levels of serum hepatocyte growth factor in patients with end-stage renal disease. J. Med. 31 (3-4): 131-41.

Magnia A., Vallari D., DiBisceglie A.M. (1994):

Use of confirmatory tests for hepatitis C viral infection in patient with hepatocellular carcinoma. J. Med. Virol, 43: 125-128.

Malatino L.S., Mallamaci F., Benedetto F.A., Bellanuova I., Cataliotti A., Tripepi G and Zoccali C. (2000):

Hepatocyte growth factor predicts survival and relates to inflammation and intima media thickness in end-stage renal disease. Am J Kidney Dis. 36 (5): 945-52.

Mandelli C., Fraquelli M., Fargion S. (1994):

Comparable frequency of hepatocellular carcinoma in cirrhosis of different aetiology. Eur. J. Gastroenterol, Hepatol, 6: 1129-1134.

Maroto A., Gines P., Arroyo V. and Gines A. (1994):

Brachial and femoral artery blood flow in cirrhosis: Relationship to kidney dysfunction. Hepatology, 17: 768.

Martinot, P.M. ; Marcellin, P. ; Pouteau, M. ; Pouteau, U. and Benhamou, J.P. (1995) :

Pretreatment serum hepatitis C virus RNA levels and hepatitis C virus genotypes are the main and independent prognostic factors of sustained response to interferon alpha therapy in chronic Yieipa

Masuhara M., Yasunaga M., Tanigawa K. (1996):

Expression of hepatocytes growth factor, transforming growth factor alpha and transforming growth factor beta 1 mRNA in various human liver diseases and correlation with hepatocytes proliferation. *Hepatology*; 24 (2): 323-329.

Matsuda-Y, Matsumoto-K., Yamada-A. (1997):

Preventive and therapeutic effects in rats of hepatocyte growth factor infusion on liver fibrosis/cirrhosis. *Hepatology*, 26 (1): 81-89.

Matsumoto K. and Nakamura T. (1992):

Hepatocyte growth factor molecular structure, roles in liver regeneration and other biological functions. *Crit Rev Oncog*; 3, 27:54.

Matsumoto K., Mizuno S., Nakamura T. (2000):

Hepatocyte growth factor in renal regeneration, renal disease and potential therapeutics. *Curr. Opin. Nephrol. Hypertens.* 9 (4): 395-402.

Matsumoto K., Nakamura T. (2001):

Hepatocyte growth factor: renotropic role and potential therapeutics for renal diseases. *Kidney Int*: 59 (6): 2023-38.

Matsumoto K., Tujima I.I., Hamaque M. (1992):

Identification and characterization of "injurin" an inducer of expression of the gene for hepatocyte growth factor; *Proc. Natl. Acad. Sci USA* 89: 3800-3804.

Matsunaga T., Bohdo E., Tahoba T., Wu Y.L., Twao M., Kataoka I.I. and Yamamoto T. (1994):

Expression of hepatocyte growth factor is regulated through activation of c AMP-mediated pathway. *EXP CELL RES* 210 (2): 325-35.

Mazinque C., Stadler S., Quantanneus B. (1986):

Schistosome derived inhibitory factor: an immunosuppressive agent preferentially active on T lymphocytes. *Int. Archs. Allergy Appl. Immuno.*, 60, 347.

McCullough A.J., Mullen K.D., and Kalhan S.C. (1991):

Measurement of total body and extracellular water in cirrhotic patients with or without ascites. *Hepatology*; 14: 1103.

McFarlane I.G., Smith H.M., Johnson P.J. (1990):

Hepatitis C virus antibody in patients with chronic active hepatitis pathogenetic factor or false positive result. *Lancet* 335: 754-757.

Mcintyre, N. (1990) :

Clinical presentation of acute viral hepatitis. *British Medical ulletin*; 46:533-547.

Meki T., Kaneda Y., Tsutsu H. (1999):

Hepatocyte growth factor gene therapy of liver cirrhosis in rats. *Nat. Med.*, 5 (2) 226-230.

Michalopolous G.K. (1984):

Control of hepatocyte replication by two serum factors, cancer *Res* 44: 4414-19.

Michalopoulos G. (1994):

Control mechanism of liver regeneration. *J. gastroenterol*, 29 (Suppl 7): 23-29.

Michalopoulos G. and Defrances M. (1997):

Liver regeneration. *Science*; 276 (5309): 60-66.

Michalopolous G.K., zarnegar R. (1992):

Hepatocyte growth factor. *Hepatology*, 15: 149-155.

Mikkelsen WL., Edmondson H.A., Peters R.L., Redeker A.G. and Reynolds T.B. (1965):

Extra- and intrahepatic portal hypertension without cirrhosis (hepatoportal sclerosis). *Ann. Surg.*, 162:602.

Miyazawa K., kitamura A., Naka D. (1991):

An altenativity processed mRNA generated from human hepatocyte growth factor gene, *Eur J. Biochem*, 197: 15-22.

Monis A. (2000):

Management of hepatocellular carcinoma update in: Omar S, El-Mawla, N.; El-Zawahry, H. (eds.): In European School of Oncology, Educational course on Liver Tumors, PP: 17-19.

Montesano R. : (1991)

Identification of a fibroblsast – derived epithelial morphogen as hepatocyte growth factor .67:901-908.

Moriya K., Yotsuyanogi H., Shintani Y. (1997):

Hepatitis C virus protein induces hepatic steatosis in transgenic mice. *J. Gen. Virol.*; 78: 1527-1531.

Morris J.D.H., Eddelstone A.L.W. and Crook, W. (1995):

Viral infection and cancer. *Lancet* 346: 754-758.

Moses H., Proper J., Volkemant M. and Swartzendruber D. (1980):

Growth arrest of AKR-28 cells maintained in the presence of epidermal growth factor 12, 12-o- tetradecomolyphorbol-13-acetate:

evidence for two separate G1 arresting points. *J. Cell. Physiol.*, 102:367.

Mosnier J-F, Degot C., Marcellin P. (1993):

The intraportal lymphoid nodule and its environment in chronic active hepatitis C: an immunohistochemical study. *Hepatology*; 17: 366-371.

Muto Y, Nouri – Aria KT , Meager A . (1988):

Enhanced tumor necrosis factor and interleukin – in fulminant hepatic failure. *Lancet* :2:72-74.

Naccarato R. and Farinati, F. (1991):

Hepatocellular carcinoma, alcohol, and cirrhosis; facts and hypotheses. *Dig. Dis. Sci*, 36: 1137-1142.

Nagaike M., Hirao S., Tajima H. (1991):

Renotropic functions of hepatocyte growth factor in renal regeneration after unilateral nephrectomy, *J. Biol. Chem*, 226: 22781-22784.

Nakajima T., Kagawa K., Meda K., (1994):

Evaluation of hepatocytic proliferative activity in chronic liver diseases and hepatocellular carcinomas by proliferating cell nuclear antigen (PCNA): Immunohistochemical staining of methanol-fixed tissues. *J. Gastroenterol* 29: 450-454.

Nakamura T., Hayama M., Sokas T. (1993):

Proliferative activity of hepatocytes in chronic viral hepatitis as revealed by immunohistochemistry for proliferating cell nuclear antigen. *Hum. Pathol.* 24: 750-753.

Nakamura T. (1986) :

Inhibitory effect of transforming growth factor [beta] on DNA synthesis of adult rat hepatocytes in primary culture. *Biochem. Biophys. Res. Commun.* 133:

Nakamura T., Kitazawa T., Ichihara A. (1986):

Partial purification and characterization of masking protein for B-type transforming growth factor rat platelets, *Biochem. Biophys Res. Commun* 141: 176-184.

Nakamura T., Nishizawa T., Hagiya M. (1989):

Molecular cloning and expression of human hepatocytes growth factor, *Nature* 342: 440-443.

Nakayama N. (1995):

Expression of hepatocyte growth factor (HGF) in LEC rats at various phases of hepatitis and hepatoma *Hokkaido-Igaku-Zasshi* Vol: 70 (4), p: 609-21, ISSN: 0367-6102.

Nakayama N., Kashiwazaki H., Kobayashi N. (1996):

Hepatocyte growth factor and C-met expression in long-Evans Cinnamon rats with spontaneous hepatitis and hepatoma. *Hepatology*, 24 (3): 596-602.

Nakayama M., Okamoto Y., Morita T. (1990):

Promoting effect of citrulline in hepatocarcinogenesis: possible mechanism in hypercitrullinemia. *Hepatology*, 11: 819-832.

Naldini L., Vigna E., Narsimhan R.P. (1991):

Hepatocyte growth factor (HGF) stimulates the tyrosine kinase activity of the receptor encoded by the proto-oncogene C-Met, *Oncogene* 6.

Napoli, J. , Bishop, G.A. , McGuinness, P.H. , Meyer, D.M. , Dustin, M.L. and Canion, C.P. (1996) :

Progressive liver injury in chronic hepatitis C infection correlates with increased intrahepatic expression of Thi-associated cytokines. *Hepatology*, 24:759-765.

Natoli G., Avantaggiati M. L., Chirillo P. (1994):

Induction of the DNA binding activity of C-Jun/C. fos heterodimers by the hepatitis B virus. *Transactivars. Mol. Cell. Biol.* 14: 989-998.

Ng I., Lai E., Fan S., (1994):

Prognostic significance of proliferating cell nuclear antigen expression in hepatocellular carcinoma. *Cancer*; 73: 2268-2274.

Nooman Z., Hassan A., Mishriky A., (1995):

The use and limitation of ultrasonography in the diagnosis of liver morbidity attributable to schistosoma mansoni infection in community – based surveys. *Mem. Inst., Oswaldo Cruz.*, 90: 147-154.

Okuda K. (1992):

Hepatocellular carcinoma; recent progress. *Hepatology*; 15: 948-963.

Okuda K. (1998):

Hepatitis C virus and hepatocellular carcinoma. In Okuda K and Tabor E (eds.). *Liver cancer*, 4: 39-45.

Okuda H., Obata H. (1995):

Epidemiological and clinical features of Budd-Chiari syndrome in Japan, *J. Hepaol.* 22: 1-9.

Oshita M., Hayashi N., Kasahara A. (1994):

Increased serum hepatitis C virus RNA levels among alcoholic patients with chronic hepatitis C. *Hepatology*; 20: 1115-1120.

Papper S., and Rosenbaum J.D. (1952):

Abnormalities in the excretion of water and sodium in compensated cirrhosis of the liver. *J.lab.Clin.Med.*; 40:523.

Patel T. (2000):

Apoptosis in hepatic pathophysiology: In Herrera J. (eds) *North America clinics of Liver disease. Vol. 4* philadelphia: W.B. Saunders Co. 295-317.

Pateron D., Ganne N., Trichet J.C. (1994):

Prospective screening for hepatocellular carcinoma in caucasian patients with cirrhosis. *J. Hepatol* 20; 65-71.

Poley J.R. (1994):

Malignant liver disease in alpha I antitrypsin deficiency *Acta paediat. Suppl.* 393-32.

Pontisso P., Ruvoletto M., Fatovich G. (1993):

Clinical and virological profiles in patients with multiple hepatitis viral infections. *Gastroenterology*, 105: 1529-1540.

Poper H. (1966):

What are major types of hepatic cirrhosis? In: Ingelfinger F; Pelman A; Finland M (eds.). *Controversy in Internal Medicine*, Philadelphia Saunders, 233-243.

Powell C.J. (1994):

Copper-overload causes cancer. The LEC rate a model for human hepatitis, liver cancer and much more. *Hum. Exp. toxicol*, 13: 910-912.

Prata A. (1982):

Infection with *S. mansoni*: In "Schistosomiasis" (ed). By Jordon P. and Webb G. William Heinman Medical Book. London; 5408-5412.

Pratt DS. And Kaplan M. (1999):

Evaluation of the liver. In: Schiff R., Sorrellaf and Maddrey WC (eds) Schiff's Diseases Of The Liver, 8th edition, Philadelphia: Lippincott Raven Publishers, p: 205.

Propst T., Propst A., Dietze O. (1994):

Prevalence of hepatocellular carcinoma in alpha antitrypsin deficiency-J. Hepatol 21: 1006-10111.

Punukollo R.C. and Gopalswamy (1990):

The hepatorenal syndrome Renal disease. *Med. Clin. Nor. Amer.*, 74 (4): 933.

Puoti G., Magrini A. Stati T. (1997):

Clinical, histological and virological features of hepatitis C virus carries with persistently normal or abnormal alanine transaminase levels. *Hepatology*; 26: 1393-1398.

Quiroga J., Beloqui O. and Castilla A (1992):

Cirrhosis. In Prieto J, Rodes J, Shafrit Z (eds.). *Hepatobiliary Diseases*. Springer-verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, Hong Kong, Barcelona, Budapest, pp 323.

Rabinovitz M., Gavalier J., Kelly R. (1992):

Lack of increases in heterozygous alpha antitrypsin deficiency among patients with hepatocellular and bile duct carcinoma. *Hepatology* 15: 407-410.

Raia S., Mies S. and Macedo A.L. (1985):

Portal hypertension in schistosomiasis. *Clin. Gastroent.*, 14:57.

Rosenthal P. (1996):

Living – related liver transplantation for adults? The time has come! *Pediatric Gastroenterol Nutr.*, 23 (3): 335-336.

Rubin R.A., Chan A. M. L., Bottaro D.P. (1991):

A broad. Spectrum human lung fibroblast-derived mitogen is a variant of hepatocyte growth factor, *Biochemistry* 88: 415-419.

Russell, W.E., McGowan, J.A., Bucher, NLR. (1984):

Biological properties of the hepatocyte growth factor from rat platelets, *J cell Physiol* 119: 193-197.

Ryder S., Koskinas J., Rizzi P. (1995):

Hepatocellular carcinoma complicating autoimmune hepatitis: role of hepatitis C virus. *Hepatology*, 22: 718-722.

Sakiyama P., Moris S., Ichimura M. (1996):

Comparative studies on the time course of proliferating cell nuclear antigen and bromodeoxyuridine staining pattern in regenerating rat liver cells. *Rinsho Byori.*, 44 (1): 61-66.

Sathar A., Chitra S, Nabeen C. (1997):

Hepatocytic proliferation in chronic liver disease: A study of liver biopsies using immunohistochemical localization of proliferating cell nuclear antigen. *Annals of Saudi Medicine*; 17 (3): 363-367.

Scheuer P.J. (1991):

Pathology of cirrhosis. In: McIntyer N., and Benhamou J.P. (eds). *Oxford Textbook of Clinical Hepatology*: Oxford University Press, Vol. 1, pp: 383.

Schewer, P.J. (1988):

Chronic hepatitis. In: *Liver biopsy interpretation*. London, Balliere Tindall, 113-130.

Schluger, L.K. , Sheiner, P.A. and Thung, S.N. (1996) :

Severe recurrent cholestatic hepatitis C following orthotopic liver transplantation. *Hepatology*;347-1643.

schmid, M. , Pirovino, M. , Altorfer, J. , Baldi, M. and Abate, M. (1982) :

Acute hepatitis Non-A, Non-B are there any specific light microscopic features . *Liver*, 2:61-67.

Schuster V. L. and Seldin D.W. (1992):

Renal Clearance. In: *The Kidney: Physiology And Pathophysiology*. Seldin DW, Giebisch G (eds). 2nd edition. New York: Raven, p: 943.

Sciot R., Staessen D., Van Damme B., van Steenberg W., Fevery J., De Groote J. and Desmet V.J. (1988):

Incomplete septal cirrhosis: histopathological aspects. *Histopathology*, 13: 593.

See G. and Marchal G., Odievere M.(1984):

Hepatocarcinoma chezun adulte suspect dune intolerance hereditaire au fructose. Ann Pediciency. Acta Paediatr Suppl 393: 27-32.

Seki S., Sakaguchi H., Kawakita N. (1990):

Identification and fine structure of proliferating hepatocytes in malignant and non malignant liver disease by use of a monochonal antibody against DNA polymerase alpha. Hum. Pathol.; 21, 1020:1030.

Senaldi G., Portmann B., Mowat A.P. (1992):

Immuno hisochemical features of the portal tract mononuclear cell infiltrate in chronic aggressive hepatitis Arch. Dis. Childh.

Seto E., Mitchell P.J. and Ten T.S.B. (1990):

Transactivation by the hepatitis B virus X protein depend on AP-2 and other transcriptional factors. Nature, 44: 72-74.

S. Girodano et al.,(1989)

Medical Oncogene 4, 1383 .

Shahin MN. Schuppan D., Waldherr R., (1992):

Serum procollagen peptides and collagen type III for the assessment of activity and degree of hepatic fibrosis and alcoholic liver disease. Hepatology, 15: 637-644.

Shamaa S., El-Bedeway A., Aita M. (2000):

Analysis of 192 Egyptian patients with Hepatocellular carcinoma: Etiology, clinical data, radiology, markers, pathology, and management in Omar, S.; El-Mawla, N.; El-Zawahry, H. In: Educational Course on Liver Tumors, PP: 86-93.

Shearman D.J.C., Niall F., Camilleri M., and Carter D. (1997):

Cirrhosis; Diseases of the gastrointestinal and liver. Third edition, pp. 915-982.

Sheen I.S., and Liaw Y.F. (1989):

The prevalence and incidence of cholecystolithiasis in patients with chronic liver disease: a prospective study. *Hepatology*; 9: 583.

Sherlock S. and Dooley J. (1997):

Diagnostic criteria of the hepatorenal syndrome, *Diseases of the liver and biliary system*, 129-134.

Sherlock S., Feldman CA., Moran B. and Scheuer P.J. (1966):

Partial nodular transformation of the liver with portal hypertension. *Am. J. Med.*, 40:195.

Shimamatsu K., Kage M., Nakashima O., (1994):

Pathomorphological study of HCV antibody positive liver cirrhosis. *J. Gastroenterol Hepatol*, 9: 624-630.

Shiota G., kamo J., Kawasaki H., Rawamotot., and Nakamusa T. (1995).

Serum Hepatocyte growth factor Levels: liver diseases: clinical implication *Hepatology* 21: 106-112.

Shiratori Y., Shiina S., Imamura M. (1995):

Characteristic difference of hepatocellular carcinoma between hepatitis B. and C-viral infection in Japan. *Hepatology*; 22: 1027:1033.

Simon M.A., Diez J., and Prieto J. (1991):

Abnormal sympathetic and renal response to sodium restriction in compensated cirrhosis. *Gastroenterology*; 101:1354.

Skullman S., Ihse I. And Larson J. (1990):

Influence of malnutrition on regeneration and composition of the liver in rats. *Acta Chir Scand*; 156 (10): 717-722.

Stagle B.L., Lee T.H. and Butel J.S. (1992):

Hepatitis B Virus and hepatocellular carcinoma. *Prog. Med. Viral*, 39:167-203.

Stoker M, Gherardi E , Perryman M . (1987) :

Scatter factor is a fibroblast – derived modulator of epithelial cell mobility. *Nature* 327: 239-242 .

Sugimura T. (1992):

Multistep carcinogenesis: a 1992 prospective *Science* 258: 663-667.

Tajima H , Matusmoto K Nakamura T . (1991)

Hepatocyte growth factor has potent anti – proliferative activity in various tumor cell lines . *FEBS Lett .*, 291 : 229-232 .

Takehara T., Hayashi N., Mita E. (1998):

Delayed fasmediated hepatocyte apoptosis during liver regeneration in mice: Hepatoprotective role of TNF alpha. *Hepatology*; 27: 1643-1651.

Tariq M.A. and Presis I.L. (1994):

Fluctuations in Fe, Cu, Zn, Br, As, Se, and Rb concentrations in C57L/J mice bearing BW7756 murine hepatoma using radioisotopic-induced X-ray fluorescence. *Biol. Trac Elem. Res*, 42: 97-114.

Tashiro K.(1990):

Deduced primary structure of rat hepatocyte growth factor and expression of the mRNA in rat tissues. Proc Natl Acad Sci U.S.A ,87:3200-3204.

Teixeira C.C. and Andrade Z.A. (1985):

Hepatic regeneration following partial hepatectomy in mice infected with schistosoma mansoni. Brazilian J. Med. Biolo. Res.; 18: 201-213.

Theocharis S., Skopetitou A., Margeli A., (1994):

Proliferating cell nuclear antigen (PCNA) expression in regenerating rat liver after partial hepatectomy Dig. Dis. Sci.; 39: 245-252.

Thomas D. (1991):

Exogenous steroid hormones and hepatocellularcarcinoma. In Tabor, E.; DiBisceglie, A.; Purcell R (eds.): etiology, Pathology and Treatment of Hepatocellular Carcinoma in North america. Portfolio, The Woodlands, Texas, pp: 67-89.

Thorgeirsson S.S. (1996):

Hepatic stem cells in liver regeneration. Faseb. J., 10 (11): 1249-1256.

Threre D. (2000):

Ag NOR staining and quantification. Micron, 31 (2): 127-131.

Tomiya T., Nagoshi and S., Fujiwara K.(1992):

Significance of serum hepatocyte growth factor levels in patients with hepatic failure. Hepatology: 15: 1-4.

Tsau Y.K., Lu M. Y. and Ni YH. (2001):

Nephromegaly and elevated plasma hepatocyte growth factor-transforming growth factor-beta 1 ratio in infants with fulminant hepatitis or biliary atresia. *Am J Kidney Dis*: 38 (2): 27-85.

Tsubouchi H., Gohda E., Strain AJ. And Daikuhara Y. (1993):

The role of HGF-SF in animal and human hepatic physiology and pathology. *EXS*. 65:251-74.

Tsubouchi H., Niitani Y., Hhirono S., Nakayama H., Gohda E., Arakaki N., Sakiyama O., (1991):

Levels of the human hepatocyte growth factor in serum of patients with various liver diseases determined by an enzyme-linked immunosorbent assay. *Hepatology*: 13: 1-5.

Tsukuma H., Hiyama T., Tanaka S. (1993):

Risk factors for hepatocellular carcinoma among patients with chronic liver disease. *N. Engl. J. Med*. 328: 1797-1801.

Tsutsumi Y., Horis S., Onoda N., (1990):

DNA polymerase alpha: an immunohistochemical marker for proliferating cells in normal and neoplastic tumour tissues. *Am.J. Clin. Pathol*. 93: 643-665.

Ueki T., Kaneda Y., Tsutsui H., Nakanishi K., Sawa Y., Morishita R., Matsumoto K., Nakamura T., Takahashi H., Okamoto E., Fujimoto J. (1999):

Hepatocyte growth factor gene therapy of liver cirrhosis in rats. *Nat Med*: 5 (2): 226-30.

Vaamonde C.A. (1988):

Renal water handling in liver disease. In: Epstein M. ed. *The kidney in liver disease*, 3rd ed. Baltimore, William & Wikins.: 31-72.

Van R.S., Cook M.P., Van S.D. (1985):

Hepatocellular carcinoma and dietary aflatoxin in Mozambique and Transkei, Br. J. Cancer 51: 713-726.

Van Ness M., and Dichl., A.M. (1989):

Liver biopsy useful in the evaluation of patients with clinical elevated liver enzymes? Ann. Intern. Med. 111: 473-478.

Van Spronsen F., Thomasse Y., Smit, G.P.A. (1994):

Hereditary tyrosinemia type 1: a new clinical classification with difference in prognosis on dietary treatment. Hepatology 20: 1187-1191.

VanThial D.H., gavalier J.S., and Schade R.R. (1985):

Liver disease and hypothalamic pituitary gonadal axis. Seminars in liver disease; 5: 35-45.

Wang X.W., Gibson. M.K., Vermeulen W. (1995):

Abrogation of p53 induced apoptosis by the hepatitis B virus X gene cancer Res, 55: 6012-6016.

Webber E., Godowski P., Fausto N. (1994):

In vivo response of hepatocytes to growth factors requires an initial priming stimulus. Hepatology; 19 (2): 489-497.

Weidner M., Arakaki N., Hartmann G. (1991):

Evidence for the identity of human scatter and human hepatocyte growth factor, Proc. Natl Acad Sci. USA 88: 7001-7005.

Weinberg A., Mize C. and Worthen H. (1976):

The occurrence of hepatoma in the chronic form of hereditary tyrosinaemia. J pediater 88: 434-438.

Welton A, Watson A.J. and Rock R.C. (1996):

Nitrogen metabolites and renal function. In: Tietz fundamentals Of Clinical Chemistry. Burits CA and Ashwood ER (eds). 4th edition. Philadelphia: W.B. Saunders Company. P: 569.

Wilkinson S.P., Partman B., Cochran A.M.G. (1978):

Clinical course of chronic lobular hepatitis report of five cases. J. Med., 188: 412-429.

Wolf H.K., Zarnegar, Michalopoulos G.K., (1991):

Localization of hepatocyte growth factor in human and rat tissues: Hepatology; 14: 488-494.

Wood I.J., Massie D., McLean A.J., (1988):

Renal sodium retention in cirrhosis: tubular site and relation to hepatic dysfunction. Hepatology; 8: 831.

Wu P., Lau J., Lau T. (1993):

Relationship between intrahepatic expression of hepatitis B viral antigens & histology in Chinese patients with chronic hepatitis B infection. Am. J. Clin. Pathol.; 100: 648-653.

Yamada Y., Kirillova I., Peschon J. (1997):

Initiation of liver growth by tumour necrosis factor: deficient liver regeneration in mice lacking type I tumour necrosis factor receptor. Proc. Natl. Acad. Sci. USA, 94 (4): 1441-1446.

Yamagami H, Moriyama M, Tanaka N, Arakawa Y. (2001):

Detection of serum and intrahepatic human hepatocyte growth factor in patients with type C liver diseases. Intervirology: 44 (1): 36-42.

Yamamoto T., Kajina K., Ogawa M. (1998):

Hepatocellular carcinoma infected with the novel DNA virus lack viral integration Biochem. Biophys. Res. Commun, 1: 389-343.

Yamane Y., Sakai K. and Umeda, T. (1984):

Suppressive effect of cupric acetate on DNA alkylation, DNA synthesis and tumorigenesis in the liver of dimethylnitrosamine treated rats. Jpn. J. Cancer Res. (Gann) 75: 1062-1069.

Yen T.S.B. (1996):

Hepadnaviral X protein, review of recent progress J. Biomed. Sci, 3: 20-30.

Yoshida Y., Komatsu M., Ozeki A. (1997):

Ethanol represses thymidylate synthase and thymidine kinase at mRNA level in regenerating rat liver after partial hepatectomy (in process citation). Biochim Biophys Acta, 1336 (2): 180-186.

Zakim D., and Boyer T.D. (1996):

Hepatic fibrosis and cirrhosis, Hepatology; A textbook of Liver disease, vol. I, pp: 506-526.

Zarnegar R., Defrances M.C., Kost D.P. (1991):

Expression of hepatocyte growth factor mRNA in regenerating rat liver after partial hepatectomy, Biochem Biophys Res. Commun, 177: 559-565.

Zarnegar R., Michalopolous G. (1989):

Purification and biological characterization of human hepatopoietin A, a polypeptide growth factor for hepatocytes, m cancer Res. 49: 3314 -3320.

Zarnegar R., Muga S., Rahija R. (1990):

Tissue distribution of hepatopoietin-A: a heparin binding polypeptide growth factor for hepatocytes *Pro Natl Acad. Sci USA* 87: 1252: 1256.

Zeldin G., Yang S.Q., Yin M. (1996):

Alcohol and cytokine- inducible transcription factors. *Alcohol Clin. Exp. Res.* 20 (9): 1639-1645.

Zucoloto S., Silva F., Junior O. (1990):

Hepatic regeneration after hepatectomy in schistosomatic fibrosis. *Int. J. Exp. Path.* 71: 465-479.