




REFERENCES

Al-Dujaili, A.H. and Harris, D.M. (1974):

Evaluation of commercially available antisera for serotyping of Pseudomonas aeruginosa.

J. Clin. Path. 27, 559.

Allen - K.D.; Bartzokos - C.A.; Graham-R; Gibson M.F;

Gilbertson - A.A. (1987):

Acquisition of endemic Pseudomonas aeruginosa on an intensive therapy unit.

J. Hosp. Infect. Sep, Vol: 10, P: 156 - 64.

Anderson, T. (1970):

Testing susceptibility to antimicrobial agents and assay of antimicrobial agents in body fluids in (Manual of Clinical Microbiology) eds. Blain et al., the American Society for Microbiology Bethesda MD.

Asensio C, Penez - Díez TC, Martínez MC, Baquero F. (1976):

A new family of low molecular weight antibiotics
from Enterobacteria.

Biochem Biophys Res Commun 69: 7 - 14.

Ayliffe, G.A.J.; Barry, D.R.; Lowbury, E.J.L. Roper-Hall,
M.J. and Martin Walker, W. (1966):

Lancet I, 1113 cited from Clarke and Richmond 1975.

Baltch - A - L; Griffin - P-E; Hammer - M. (1979):

Pseudomonas aeruginosa bacteraemia: relationship
of bacterial enzyme production and pyocin types
with clinical prognosis in 100 patients.

J. Lab. Clin. Med. Apr. Vol. 93.

Baquero F. Moreno F. (1984):

The microcins FEMS Microbiol Lett 23: 117 - 124.

Barder, J., and Waterworth, P.M. (1966):

Activity of gentamicin against Pseudomonas and
hospital Staphylococci.

Brit. Med. J.L.: 2030.

Bergan. T. (1968):

Typing of Pseudomonas aeruginosa by pyocin production.

Acta. Pathol. Microbiol. Scand. 72: 401 - 411.

Bergan. T. (1972):

Comparison of numerical procedures for grouping

Pseudomonas bacteriophages according to lytic sepectra.

Acta. Pathol. Microbiol. Scand. Sect. B 80: 55 - 70.

Bergan, T., (1973):

Epidemiological markers for Pseudomonas aeruginosa

III. Comparison of bacteriophage typing, serogrouping and pyocin type.

Acta. Pathol. Microbiol. Scand. Sect. B 81: 91 - 101.

Bergan, T; Kozaczek. W. (1979):

Epidemiological markers for Pseudomonas aeruginosa

VII: Nosocomial occurance in urological comparison of two phase typing sets, pyocin typing and serotyping.

Arch. Immunol. Ther. Exp. Vol: 27, P : 447 - 58.

Bergey's Manual of systemic Bacteriology (1984):

Vol. I - 141 - 199 The Williams and Wilkin's Co.

Bobo, R.A., E.J. Newton, L.F. Jones. L.H. Farmer and J.J.

Farmer III (1973):

Nursery outbreaks of Pseudomonas aeruginosa

Epidemiological conclusion from five different
typing methods.

Appl. Microbiol. 25: 414 - 420.

Bodey, G.P. Whetecar. J.R.J.P. Middlemen, E. and Rodrig-

uez. V. (1971):

Carbenicillin therapy for Pseudomonas infection

J. Amer. Med. Ass. 218: 62 - 66.

Botterell, E. & Magner, D. (1945):

Meningitis due to Ps. aeruginosa, penetrating
wounds of head.

Lancet I : 112.

Brokopp. C.D.; R. Gomez. Lus and J.J. Farmer III (1977):

Serological typing of Pseudomonas aeruginosa: Use
of commercial antisera and live antigen.

J. Clin. Microbiol. 5: 640 - 649.

Brokopp, C.D; J.J. Farmer (1979):

Typing methods for Pseudomonas aeruginosa.

P. 89 - 133. In R.G. Dogget (ed). Pseudomonas aeruginosa. Academic Press. New York.

Brown, M.R.W., Faster, J.H.S. and Clamp, J.R. (1969):

Composition of Ps. aeruginosa slime.

Biochem. J. 112: 521 - 525.

Brown, M.R.W., Anwar H, Lambert P A (1984):

Evidence that mucoid Pseudomonas aeruginosa in the cystic fibrosis lung grows under iron restricted condition.

F E M S Microbiol. Lett 21: 113 - 117.

Bruun, J.N., G.J. Mc Garrity, W.S. Blakemore and L.L.

Coriell (1976):

Epidemiology of Pseudomonas aeruginosa infection determination by pyocin typing.

J. Clin. Microbiol 3: 264 - 271.

Bryan. L. (1979):

Resistance to antimicrobial agents: the general nature of the problems and the basis of resistance, Ps. aeruginosa. Clinical manifestation of infection and current therapy. Edited by R.G. Doggett. New York.

Bryan, C.S.; Reynolds, K.L.; and Brenner, E.R. (1983):

Analysis of I, 186 episodes of gram negative bacteraemia in non university hospitals: the effect of antimicrobiol therapy.

Rev. Infect. Dis. 5: 629 - 638.

Cervantes - Vega, C; Chavez, J; Rodriguez, M.G. (1986):

Antibiotic susceptibility of clinical isolates of Pseudomonas aeruginosa.

Antonie - Van - Leeuwenhoek, Vol: 52, P: 319 - 24.

Chitkara, Y.K; Feierabend, T.C. (1981):

Endogenous and exogenous infection with Pseudomonas aeruginosa in a burn unit.

Int - Surg. Jul-Sep. Vol: 66 P: 237 - 40.

Clarke, P.H. & Richmond, M.H. (1975):

The Biochemistry and genetics of Ps. aeruginosa

P. 37 - 65 eds. John. Wiley London.

Colwell, R. (1964):

A study of features used in the diagnosis of Ps. aeruginosa.

Can. J. Microbiol. 37; 181.

Conroy - J.V., Baltch, A.L, Smith, R.P, Hammer M.C.;

Griffin, P.E. (1983):

Bacteraemia due to Pseudomonas aeruginosa:

Use of a combined typing system in an eight year study.

J. Infect. Dis. Sep. Vol.: 148, P: 603.

Cox, C.D. (1985):

Iron transport and serum resistance in Pseudomonas aeruginosa.

Antibiotic Chemother 36. 1 - 12.

Cruickshank, R.; J.P. Duguid; B. P. Marmion & R.H.A. Swain,
(1975):

Medical Microbiology Vol. 2, 12th edn P. 444, 448 eds.

The English Language Book Society & Churchill Livingstone.

Darrell, J.H. and Wahba, A.H. (1964):

Pyocin typing of hospital strains of Ps. pyocyanea
J. Clin. Pathol 17; 236.

Darrell, J. & Waterworth, P. (1967):

Dosage of gentamicin for Ps. aeruginosa infection
Brit. Med. J. 2; 535.

Daschner, F.; Langaack, H. and Wiedermann, B. (1978):

Antibiotic resistance in intensive care unit areas.
Infect. Control 4: 382 - 387.

Del - Piano, M, La - Palombara - P, Picci - A, Nicosia-
R (1986):

Serological and pyocin typing and antibiotic sen-
sitivity of Pseudomonas aeruginosa strains.
Microbiologica Apr, Vol.: 9, P: 235 - 8.

Doring G.; Obernsser HJ; Botzenhart K, et al. (1983):

Proteases of Pseudomonas aeruginosa in patients
with cystic fibrosis.
J. Infect. Dis. 147: 744 - 70.

Drasar, F.A.; Farrell, W.; Maskell, J. and Williams J.D.,

(1976):

Tobramycin, amikacin, sisomicin and gentamicin
resistant gram negative rods.

Brit. Med. J., 2, 1284.

Edmonds, P., R.R. Suskind, B.G. Mac Millan, and I.A.

Holder (1972):

Epidemiology of Pseudomonas aeruginosa in a burns
hospital. Evaluation of serological, bacteriophage
and pyocin typing methods.

Appl. Microbiol 24: 213 - 218.

Eykyn, S. and Phillips, I. (1975):

Gentamicin resistant Pseudomonas aeruginosa

Lancet. I. 861.

Falkiner, F.R. and Kean. C. (1977):

Epidemiological information from active and passive
pyocin typing of Pg. aeruginosa.

J. Med. Microbiol. 10; 447.

Falkiner, F.R.; Jacoby, G.A., Keane, C.T.; McCann S.R.,

(1982):

Amikacin, gentamicin and tobramycin resistant

Pseudomonas aeruginosa in a leukaemic ward.

Epidemiology and genetic studies.

J. Hosp. Infect. Sep, Vol.: 3, P: 253 - 61.

Farmer, J.J., III and L.G. Herman (1969):

Epidemiological fingerprinting of Pseudomonas

aeruginosa by the production of and sensitivity

to pyocin and bacteriophage.

Appl. Microbiol. 18 - 760 - 765.

Farmer, J. & Herman, L. (1974):

Pyocin typing of Ps. aeruginosa. J. Infect. Dis.

130; 543.

Finland (1960):

Treatment of Pneumonia and other serious infections.

New Eng. J. Med. 263; 207.

Funk, E. (1901):

Amer. Med. 2; 738 - 773.

Cited from J. Infect. Dis. 5; 542 (1908).

Fyfe J A M, Harris G M, Govan J R W. (1984):

Revised typing method of Pseudomonas aeruginosa.

J. Clin. Microbiol 20: 47 - 59.

Gadde, H.B., Jahagirdar, V.L. and Sengupta. S.R. (1980):

Aeruginocin typing of Ps. aeruginosa as an epidemiological marker.

Indian. J. Microbiol 20: 265.

Garrod, L.P. and Waterworth, P.M. (1969):

Susceptibility of Ps. aeruginosa to tobramycin and gentamicin alone and in combination with carbenicillin.

J. Clin. Pathol 22; 534.

Garrod, L.P., Lambert, H.P. and O. Grady, F. (1981):

"Antibiotic and Chemotherapy" 5th. ed.

Churchill Livingstone (London)

Gessard (1982):

Comp. Rend. Acad. Sci., Paris 94; 536.

Cited from Bergey's Manual of Determinative Bacteriology. The Williams and Wilkin's Co. P.99 (1957).

Gillies, R.R. and Govan, J.R.W. (1966):

Typing of Pseudomonas Pyocyanea by pyocin production.

J. Pathol. Bact. 91, 339.

Gordon, J. & McLead, J. (1928):

The principal application of the direct oxidose reaction - bacteriology.

J. Pathol. Bacterial. 31; Vol. I, 185.

Govan, J.R. W and Gillies, R.R. (1969):

Further studies in the pyocin typing of Pseudomonas Pyocyanea.

J. Med. Microbial 2, 17.

Govan, J.R.W. (1974):

Studies on the pyocins of Pseudomonas aeruginosa.

Production of contractile and flexuous pyocins in Pseudomonas aeruginosa.

J. Gen. Microbial. 80: 17 - 39.

Govan, J.R.W. (1974 a):

Studies on the pyocins of Pseudomonas aeruginosa;
morphology and mode of action of contractile pyocins.
J. Gen. Microbiol. 80, 1.

Govan, J.R.W. (1978):

Pyocin typing of Pseudomonas aeruginosa.
In "Methods in Microbiology"
Academic press ed. Bergan & Norris P. 61 - 91 London.

Govan, J.R.W. (1986):

In vivo significance of bacteriocins and bacteriocin
receptors.
Scand. J. Infect. Dis. (Suppl). Vol, 49 P ; 31 - 7.

Graber, C.L. Tumbusch, W.T.I. and Vogel E.H.J. (1961):

Invitro-sensitivity of Pseudomonas from burned
patients to colistin sulphate antibiotic.
Ann. 7: 77 cited from J. Clin. Inverst. 40:
2064 - 2075.

Granstrom M, Ericsson A, Strandvik B, et al., (1984):

Relationship between antibody response to Pseudomonas
aeruginosa exoproteins and colonic infection in
patients with cystic fibrosis.
Act. Paediatr. Scand, 73: 772 - 777.

Hall, J.H., Calway, J.L. Tindall, J.P. and Graham, S.J.,
(1968):

Archives of Dermatology 97, 312.

Cited from Clarke and Richmond (1975).

Harris, R.C., Buxbaum, L.V., and Applebaum, E., (1946):

Secondary Bacillus pyocyaneus infection in meningitis following intrathecal penicillin therapy.

J. Lab. Clin. Med. 31: 1113 - 1120.

Heckman, M.G.; Babcock, J.; Rose, H. (1972):

Pyocin typing of Pseudomonas aeruginosa:

Clinical and epidemiological aspect.

Am. J. Clin. Pathol. 57; 35.

Henderson, A. Machaurin, J. and Scott, J.M. (1969):

Pseudomonas in a Glasgow baby unit,

Lancet, 2, 316.

Higerd, T.B., Baechler, C.A. and Berk, R.S. (1967):

In vitro and in vivo characterisation of pyocin.

J. Bact. 93, 1976.

Hojyo - Tomoka, M.T., and Richard, R.M. (1973):

Pseudomonas infection in superhydrated skin.

Arch. Dermatol. 107: 723 - 727.

Holloway, B.W. (1960):

Grouping *Pseudomonas aeruginosa* by Isogenicity and pyocinogenicity.

J. Pathol. Bacterial., 80, 448.

Howe T R, Wretlind B, Iglewski BH. (1983):

Comparison of two methods of genetic exchange in determination of the genetic locus of the structural gene for *Pseudomonas aeruginosa* elastase,

J. Bacterial; 156: 58 - 61.

Hubbard, J.D., King, H. and Beemer, P.R. (1957):

Pathogenicity of *Pg. aeruginosa*.

Amer. J. Clin. Path, 28: 243 - 257.

Hunter, C.A. & R.R. Ensign (1947):

Amer. J. Pub. Health 37; 1160.

Cited from Topley and Wilson's principles of Bacteriology and Immunology 5th. edn. Vol. 1
P.: 636 - 644.

Baltimore Williams and Wilkins.

Iglewski BH, Kabat D. (1975):

NAD dependent inhibition of protein synthesis by

Pseudomonas aeruginosa toxin.

Proc Natl. Acad. Sci. U.S.A; 72: 2284.

Iglewski BH, Sadoff J C, Bjorn MJ, et al., (1978):

Pseudomonas aeruginosa exoenzyme S: an adenosine
diphosphate ribosyl transferase distinct from toxin A.

Proc. Natl. Acad. Sci U.S.A; 75: 3211 - 3215.

Ito, S. and Kageyama, M. (1970):

J. Gen Appl. Microbiol (Tokyo) 16, 231 - 240.

Cited in Govan (1978).

Ito, S.; Kageyama, M. and Egami, F. (1970):

Isolation and characterisation of pyocins from
several strains of Pseudomonas aeruginosa.

J. Gen. Appl. Microbial (Tokyo), 16, 205.

Jacoby, G.A. (1974):

Properties of R. plasmid in Pseudomonas aeruginosa
producing amikacin (BB - KB), butirosin, kanamycin,
tebramycin and sisomicin resistance.

Antimicrob. Agents Chemother., 6, 807.

Jacop, F. (1954):

Biosynthese induit par mode d, action d'une pyocine
antibiotique de Pseudomonas aeruginosa

Ann. Inst. Pasteur 86: 149 - 160.

Jagger K S; Robinson D L; Franz M N, et al., (1982):

Detection by enzyme - linked immunosorbent of
antibody specific for Pseudomonas protease and
exotoxin A in sera from cystic fibrosis patients.

J. Clin. Microbiol, 15: 1054 - 1058.

Jawetz, E. (1952):

Antibiotic synergism and antagonism

Review of experimental evidence.

Arch. Intern. Med. 90, 301.

Jawetz, E.; Melnick, J.L. Adelberg, E.A. (1984):

Review of Medical Microbiology

16th. ed, P.: 243 - 248.

Joklick, W.K.; H.P. Willett and Amos, D.B. (1980):

17th. edition Zinsser Microbiology (des)

Appleton - Century - Crofts, New York, P. 273 - 274.

Jones, L.F., J.P. Zakanycz, E.T. Thomas, and J.J. Farmer

III (1974):

Pyocin typing of Pseudomonas aeruginosa:

A simplified method.

Appl. Microbiol 27: 400 - 406.

Kageyama, M. (1964):

Studies of a pyocin. I. physical and chemical pr-

operties J. Biochem, Tokyo 55, 49.

Kageyama, M. (1970 a):

J. Gen. App. Microbiol, (Tokyo) 16, 523 - 530

Cited in Govan (1978).

Kageyama, M. (1970 b):

J. Gen. Appl. Microbiol. (Tokyo) 16, 531 - 535

Cited in Govan (1978).

Kageyama, M. (1975):

Bacteriocins and bacteriophage in Pseudomonas aeruginosa. P: 291 - 305. In S. Mitsuhashi and H. Hashimoto (ed). Microbiol drug resistance. University of Tokyo Press. Tokyo.

Kapur, R; Shriniwas (1986):

Prevalance of Protease and elastase production by clinical isolates of Pseudomonas aeruginosa in relation to aeruginosa typing pattern.

Eur - J- Epidemiol. Sep., Vol.: 2, P: 182 - 5.

Kass, E.H.(1955):

Symposium on newer aspects of chemotherapeutic and antibiotic drugs in management of infections of urinary tract.

Amer. J. Med. 18: 764.

Kaziro, Y. and M. Tanaka (1965):

Studies on the mode of action of pyocin I. inhibition of macromolecular synthesis in sensitive cells.
J. Biochem 57. 689 - 695.

Kelly, N.M.; Fitzgerald, M.X. Tenpany, E.D Boyle, C; Falk-
iner, F.R; Kean, C.T. (1982):

Does Pseudomonas Cross infection occur between
cystic fibrosis patients.

Lancet, Sep. 25, Vol. 2, P.: 688 - 90.

King, E.O.; Ward, M.B. & Raney, D. (1954):

Two simple media for the demonstration of pyocyanin
and fluorescin.

Journal of Laboratory and Clinical Medicine 44, 301.

Klastersky, J.C.; Hensgens, C. & Daneau D. (1975):

Therapy of Staphylococcal infections: comparative
study of cephaloridine and gentamicin.

Am. J. Med. Sci. 269; 201.

Klinger J.D., Straus D C, Hilton C B, et al. (1978):

Antibiotic to proteases and exotoxin A of Pseudomonas aeruginosa in patients with cystic fibrosis:

demonstration by radio immunoassay.

J. Infect. Dis. 138: 49 - 58.

Kominos, S.D.; Copeland, C.E. and Grosiak, B. (1972):

Mode of transmission of Pseudomonas aeruginosa in a burn unit and an intensive care unit in a general hospital.

Appl. Microbiol, 23, 309.

Komiyama, K.; Habbick, B, F.; Martin, T., Tumber, S.K.,
(1987):

Characterization by pyocin typing and serotyping of oral and sputum strains of Pseudomonas aeruginosa isolated from cystic fibrosis patients Can-

J. Microbiol, Mar. Vol.: 33, P.: 221 - 5.

Kovac's N. (1956):

Identification of Ps. pyocyanea by the oxidase reaction. Nature (London) 178, 703.

Kumazaki - T - Ishii - S. (1982):

Comparative study on fibers isolated from four -
R type pyocins, phage tail like bacteriocins of
Pseudomonas aeruginosa.

J. Biochem (Tokyo) No. Vol.: 92 P.: 1559 - 66.

Kuroda K., Kageyama R. (1983):

Biochemical relationship among three F - type
Pyocins, pyocin F₁, F₂ and F₃ and phage KFi₁

J. Biochem 94: 1429 - 1441.

Lanyi, B.; T. Bergan (1978):

Serological characterisation of Pseudomonas
aeruginosa.

P. 93 - 168. In T. Bergan and J.R. Norris (ed)

Methods in microbiology, Vol. 10.

Academic press. Inc. London.

Liu, P.V., Abe, Y. and Bates J.L. (1961):

The role of various fraction of Pg. aeruginosa
in it's pathogenesis.

J. Infect. Dis. 108; 218.

Liu PV. (1979):

Toxins of *Ps. aeruginosa*. In Dogget R G. ed.

Ps. aeruginosa. Clinical manifestation of infection and current therapy.

New York. Academic Press.

Margartten, W.; Nakai, H.; and Landing, B.H. (1961):

Significance of selective vasculites and the "bone marrow" syndrome in Pseudomonal septicaemia.

New. Eng. J. Med. 265; 773.

Mc Millan, M., Price, T.M.L., Mac Laren, D.M., and Scott,

G.W.(1962):

Ps. pyocyanea infections treated with colistin methane sulphonate.

Lancet 2: 737.

Merrikin, D.T., and C.S. Terry (1972):

Variability of pyocin type and pyocin sensitivity in some strains of *Pseudomonas aeruginosa*.

J. Appl. Bacterial. 35, 667 - 672.

Migula (1895):

In Engler & Prantl, Die natural Pflnzenfam. I.

La 29. Cited from Bergey's Manual of Determinative
Bacteriology. 7th. edn.

Baltimore, Williams and Wilkin's Co. P. 99.

Migula (1900):

Syst. D. Bakt. 2. 884.

Cited from Bergey's Manual of Determinative
Bacteriology 7th. edn.

Baltimore Williams and Wilkin's Co. P. 99 (1957).

Moody, M.; Young, V.; Schimpff, S. and Kenton, D. (1970):

Relationship of Pseudomonas aeruginosa immunotypes
to antibiotic susceptibility patterns.

Antimicrob. Ag. Chemother. P. 249.

Murphy, G. & McDonald, D. (1961):

Proteus, Pseudomonas, Achromobacter, Septicaemia
as complication of chronic infection in anatomically
predisposed urinary tract.

J. urol. (Baltimore) 85; 672.

Newton, B.A. (1955):

The properties and mode of action of the polymyxin
Bac. Rev. 20: 14 - 27.

Nicas TI, Frank D W, Strenzel P, et al. (1985):

Role of exoenzyme S in chronic Pseudomonas aerug-
inosa lung infection.

Eur. J. Clin - Microbiol.; 4: 175 - 179.

Ogle. J.W, Vasil ML. (1985):

The epidemiology of Pseudomonas aeruginosa (PA)
in cystic fibrosis as defined by a cloned D N A
Probe derived from P A exotoxin (E T A) Abstracts
of the 25th. Interscience conference on Antimicro-
biol Agents and Chemotherapy: 122.

Ohkawa, I. Kageyama, M and Egami, F. (1973):

Purification and properties of pyocin S₂
J. Biochem, Tokyo 73, 281.

Ohkawa, I. Shiga S, Kageyama M. (1981):

Effect of iron concentration in the growth medium
on the sensitivity of Pseudomonas aeruginosa to
pyocin S₂.

J. Biochem 87: 323 - 331.

Ohman D E, Chakrabarty A M. (1982):

Utilization of human respiratory secretions by
mucoid Pseudomonas aeruginosa of cystic fibrosis
origin.

Infect. Immun.; 37: 662.

Ohman D E, Goldberg J.B., Flynn J A, et al. (1985):

Genetics of exopolysaccharide production by mucoid
Pseudomonas aeruginosa.

Antibiot Chemother; 36: 13 - 22.

Palleroni N J. Family I. (1984):

Pseudomonadaceae. In: Krieg NR, ed. Bergey's
Manual of Systemic Bacteriology.

Baltimore: Williams and Wilkins, 141.

Parker, M.T. (1972):

Characterisation and typing of Pseudomonas
aeruginosa.

J. Gen. Microbial, 73.

Penketh, A.; T. Pitt; D. Roberts; M.E. Hodson and J.C.

Batten (1983):

The relationship of phenotypic changes in Pseudo-
monas aeruginosa to the clinical condition of pa-
tients with cystic fibrosis.

Am. Rev. Respir. Dis. 177: 605 - 608.

Pericival, A., and Leigh, D.A. (1967):

Clinical and laboratory studies with carbenicillin
a new penicillin active against Ps. aeruginosa.

Lancet I: 1289 - 1293.

Pier C B. (1985):

Pulmonary disease associated with status of the
host bacterium interaction.

J. Infect. Dis.; 151: 575 - 580.

Pines, A; Raafat, H. and Plucinski, K. (1967):

Gentamicin and Colistin in chronic bronchial infections Brit. Med. J. 2: 535 - 537.

Pitt, T.L. (1980):

State of the art. typing Pseudomonas aeruginosa
J. Hosp. Infect. I: 193 - 199.

Pones, R. (1927):

Ann. Inst. Pasteur 41: 1338

Cited from Topley and Wilson's principles of Bacteriology and Immunology. 5th. edition Vol. I,
636 - 644.

Baltimore the Williams and Wilkin's Co.

Porebska, A.M. (1979):

Pyocin typing of Pseudomonas aeruginosa strains
isolated in Ahmadu Bello University Teaching Hospital, Zaria, Afr. J. Med - Sci - Mar. Jun P: 19 - 23.

Price, K.E.; Pursiano, T.A.; De Furia, M.D. & Wright, G.E.,
(1974):

Activity of amikacin against clinical isolates resistant to one or more aminoglycoside antibiotics.
Antimicrobiol Agents and Chemotherapy. 5, 143.

Rampling, A. Whit by, J.L and Widly, P. (1975):

Pyocin sensitivity testing as a method of typing

Pseudomonas aeruginosa: Use of phage free Preparations of pyocin.

J. Med. Microbial., 8, 531.

Reynolds H.H. (1985):

Phagocytic defense in the lung

Antibiot chemother, 36: 74.

Ringen L. & Drake C. (1952):

A study of the incidence of Ps. aeruginosa from various natural sources.

J. Bact. 64; 841.

Ruben, F.; Norden C. & Hruska E (1978):

Factors associated with acquisition of Ps. aeruginosa resistant to gentamicin

Am. T. Med. Sci. 275; 173 - 179.

Schable - B; Olson, D.R.; Smith, P.B. (1986):

Improved, computer generated system for pyocin
typing of Pseudomonas aeruginosa (Published err-
atum appears in J. Clin - Microbial (1987 Feb.,
25 P.: 466).

Schroeter (1872):

In Cohn. Bertrag. 2, Biologic I, Heft. 2, 126:

Cited from Bergey's Manual of Determinative
Bacteriology 7th. edn. P. 99.

The Williams and Wilkines Co. (1957).

Sengupta, S.R., Johagirdas, V.L. and Gadde, H.B. (1980):

Epidemiology of Pseudomonas infection in burnward.
Indian. J. Pathol Microbial 23: 241.

Shooter, R.A.; Cooke, E.M.; Gaya, H.; Kumar, P., Patel, N.;

Parker, M.T., Tham, B.T. and France, D.R. (1969):

Food and medicaments as possible source of hospital
strains of Pseudomonas aeruginosa. Lancet I, 1227.

Shriniwas (1974):

Aeruginocine typing of Ps. aeruginosa T. Clin,
Path, 27, 92.

Shriniwas (1976):

Further typing of Pseudomonas aeruginosa strains
not typable by aeruginocin typing method.
Indian I. Med. Res. 64: 1747.

Sidberg, H.D.; Sadoff, J.C. (1977):

Pyocin sensitivity of Neisseria gonorrhoea and its
feasibility as an epidemiological tool. Infect
Immun. Feb. Vol.: 15, P.: 628 - 37.

Smith A.L. (1985):

Principle of Microbiology, 10th ed., P. 25.
Times Mirror, Mosby, College Publishing.

Sonne, M. and Jawetz, E. (1969):

Combined action of carbenicillin and gentamycin
on Ps. aeruginosa infection in vitro.
Appl. Microbial. 17: 893 - 896.

Southern PM, Mays B B, Preece AK, et al. (1970):

Pulmonary clearance of Ps. aeruginosa.

J. Lab. Clin. Med. 76: 548.

Stanly, P.G., Shepherd, R.G., and White, H.J. (1947):

Pobymyxin, a new chemotherapeutic agent. Bull.

Johns. Hopkins. Hosp. 81: 43 - 54.

Stapleton MJ, Jagger KS, Warren RL. (1984):

Transposon mutagenesis of Ps. aeruginosa exoprotease genes.

J. Bacterial; 157: 7 - 12.

Stoodly, B. and Thomb, B. (1970):

J. Med. Microbial. 2; 367.

Cited from Clarke and Richmond (1975).

Sussman, M. and Stevens, J. (1960):

Ps. pyocyanea wound infection, an outbreak in an arthopedic unit.

Lancet 2: 734.

Takeya, K. Minamishima, Y., Ohnishi, Y. and Amako, K.
(1969):

Rod shaped pyocin 28. J. Gen. Virol., 145.

Tinne, J, E.; Gordon A.M. Batn W.H. and Mackey, W.A. (1967):

Cross infection by Pseudomonas aeruginosa as
a hazard of intensive surgery. Br. Med. J. 4, 313.

Topley and Wilson's (1975):

Principle of Bacteriology, Virology and Immunology
6th. ed. Vol. I : 802 - 826.

Baltimore the Williams and Wilkin's Company.

Topley and Wilson's (1984):

Principles of Bacteriology, Virology and Immunology
7th. ed. Vol. 3 Page, 246: 262. Eward Arnold.

Tripathy, G.S. and Chadurick P (1971):

The effect of mitomycin - C on the pyocin typing
patterns of hospital strains of Ps. aeruginosa.

Can. J. Microbial 17. 829.

Trivisan (1885):

Atti. Acad. Fis. Stat. Milano. Ser. 4, 311.

Cited from Bergey's Manual of Determinative Bacteriology 6th.ed. The Williams and Wilkin's Co.

P. 89 (1948).

Uratani, Y. and Kageyama (1977):

A fluorescent probe response to the interaction of pyocin R_1 with sensitive cells.

J. Biochem. 81: 333 - 341.

Uratani, Y. (1982):

Dansyl Chloride labeling of Pseudomonas aeruginosa treated with pyocin R_1 : Change in permeability of the cell envelop.

J. Bacterial, 149: 523 - 528.

Urbano & Boddì (1977):

A relational code for reporting patterns in bacteriocin and bacteriophage typing.

J. Med. Microbial 11, 63.

Wretling B, Pavlovskis OR (1981):

The role of Proteases and exotoxin A in the
pathogenicity of Pseudomonas aeruginosa infections
Scand. J. Infect. Dis.; 29 (Suppl: 13 - 19, 28):

Yow, M. (1952):

Development of proteus and Pseudomonas infections
during antibiotic therapy.

J. Amer. Med. Ass. 149; 1184.

Young, V.M, and M.R. Moody (1974):

Serotyping of Pseudomonas aeruginosa.

J. Infect. Dis. 130 (Suppl 47 - 52).

Zabransky, E.E. and Day, F.E. (1969):

Protein typing of clinical strains of Pseudomonas
aeruginosa.

Appl. Microbial 17, 293.

Zopf (1884):

Spaltzilfe, 2 API 83.

Cited from Bergey's Manual of Determinative
Bacteriology 6th. edn.

Baltimore. The Williams and Wilkin's Co. (1948).