

References

Ackerman, A.D. and Dean, J.M: Meningitis and infectious encephalitis. In: Rogers, M.C (ed). Textbook of pediatric intensive care, 1st edition P895 Williams and Wilkins. Baltimore, Sydney, 1989.

Adams, D.H. and Liloyd, A.R: Chemokines: leukocyte recruitment and activation cytokines. Lancet 349: 490, 1997.

Adams, D.O. and Hamilton, T.A: The cell biology of macrophage activation. Ann. Rev. of Immunol., 2: 283, 1984.

Akira, S., Taga, T. and Kishimoto, T: Interleukin 6 in biology and medicine. Adv. Immunol, 54:1, 1993.

Alam, R., stafford, S. and Forsythe, P: RANTES is a chemotactic and activating factor for human eosinophils. J. Immunol., 150: 3442, 1993.

Alcami, A. and Smith, G.L: cytokine receptors ecoded by pox viruses: a lesson in cytokine biology. Immunol Today, 16:474, 1995.

Allavena, P., Bianchi, G. and Zhou, D: Introduction of natural killer cell migration by monocyte chemotactic protein-1, protein-2 and protein-3. Eur. J. Immunol, 24: 3233, 1994.

Ameglio, F., Capobianchi, M.R. and Dolei, A: Differential effects of gamma interferon on expression of HLA class II molecules controlled by the DR and DC loci. Infect. Immunol., 42: 122, 1983.

Arai, K.F., Lee, A. and Miyajina, A.: Cytokines: Coordinator of immune and inflammatory responses. *Ann. Rev. Biochem.*, 59: 783, 1990.

Azuma, H., Tsuda, N. and Sasaki, K.: Clinical significance of cytokine measurement for detection of meningitis. *J. Pediatr.*, 131: 463, 1997.

Bacon, K.B. and Schall, T.J.: Chemokines as mediators of allergic inflammation. *Int. Arch. Allergy Immunol.*, 109: 97, 1996.

Bacon, K.B., Fincham, N.J. and Quinn, D.G.: Tissue-derived leukocyte attractant factors in inflammation; Biology of the chemotactic response in Armitage, J.P., Lackie JM (eds) P379. Cambridge, Cambridge University Press, 1990.

Bacon, K.B., Flores-Romo, L. and Life, P.F.: IL-8 induced signal transduction in T-lymphocytes involves receptor-mediated activation of phospholipases C and D. *J. Immunol.*, 154: 3654, 1995.

Baggiolini, M. and Dahindin, C: C-C chemokines in allergic inflammation. *Immunol Today*, 15:127,1994.

Baggiolini, M., Dewald, B. and Moser, B.: Interleukin 8 and related chemotactic cytokines – C-X-C and C-C chemokines. *Adv. Immunol.*, 55: 97, 1994.

Baggiolini, M., Walz, A., Kunkel, S.L.: Neutrophil-activating peptide-1/interleukin 8, a novel cytokine that activates neutrophils. *J. Clin. Invest.*, 84: 1045, 1989.

Balkwill, F.R. and Burke, F.: The cytokine network: Immunology Today, 10: 299, 1989.

Barker, J.N.W.N., Sarma, V. and Mitra, R.S: Marked synergism between tumor necrosis factor- α and interferon- γ in regulating of keratinocyte- derived adhesion molecules and chemotactic factors. J.Clin. Invest, 85:605,1990.

Bazzoni, F., Cassatella, M.A. and Rossi, F.: Phagocytosing neutrophils produce and release high amounts of the neutrophil-activating peptide 1/interleukin 8. J. Exp. Med., 173: 771, 1991.

Becker, S., Quary, J. and Koren, H.S.: Constitutive and stimulated Mcp-1, GRO α , B & γ expression in human air way epithelium and broncho alveolar macrophages. Am. J. Physiol, 266: 278, 1994.

Begg, Pepper, D.S. and Chesterman, C.N: Complete covalent structure of human beta-thromboglobulin. Biochemistry,17:1739,1978.

Beisel, W.B.: Mediators of fever and muscle proteolysis (editorial) N. Engl. J. Med., 308: 586, 1991.

Bell, W.E. and McCormic, W.F.: Bacterial meningitis: General concepts and managements, 2nd edition p21, Philadelphia. WB Saunders Co. 1991.

Besemer, Hujber, A. and Kuhn, B: specific binding internalization, and degradation of human neutrophil activating factor by human polymorphonuclear leukocytes. J.Biol. Chem. In press, 1989.

Beutler, B. and Cerami, A.: Cachectin: More than a tumour necrosis factor. *N. Engl. J. Med.*, 316: 379, 1991.

Beuvery, E.C., Van Rossum, F. and Lauwers, S.: Comparison of counter immunoelectrophoresis and ELISA for diagnosis of bacterial meningitis. *Lancet*, 1: 1154, 1991.

Bevilacqua, M.P.: IL-1 acts on cultured human vascular endothelium to increase the adhesion of polymorphnuclear leukocytes, monocytes and related leukocyte cell lines. *J. Clin. Invest.*, 76: 2003, 1987.

Bierer, B.E., Sleckman, B.P. and Ratnofsky, S.E.: The biologic roles of CD2, CD4 in T-cell activation. *Ann. Rev. Immunol.*, 7: 579, 1989.

Bjorkman, P.J., Saper, M.A. and Samraoul, B.: Structure of the human class I histocompatibility antigen, HLA-A2. *Nature*, 329: 506, 1987.

Bland, R.D., Lister, R.C. and Ries, J.P.: Cerebrospinal fluid lactic acid level and PH in meningitis. *Am.J. Dis. Child*, 128:151,1984.

Bochner, B.A., Luscinskas, F.W. and Gimbrone, M.A.Jr.: Adhesion of human basophils, eosinophils and neutrophils to interleukin-1-activated human vascular endothelial cells: contribution of endothelial cell adhesion molecules. *J. Exp. Med.*, 173: 1553, 1991.

Brandtzae, P.P., Kierulf, P. and Gauoted, A.: Plasma endotoxin as predictor of multiple organ failure and death in systemic meningococcal disease. *J. Infect. Dis.*, 45: 132, 1988.

Braun, R.K., Franchini, M. and Erard, F: Human peripheral blood eosinophils produce and release interleukin-8 on stimulation with calcium ionophore. *Eur.J.Immunol*, 23:956,1993.

Brezinschek. R.L., Lipsky, P.E. and Galea, P: Phenotypic characterization of CD4+ T cells that exhibit a transendothelial migratory capacity. *J.Immunol*, 154:3062,1995.

Brook, I., Brichmell, K.S. and Bverturf, G.D: Measurement of lactic acid in cerebrospinal fluid of patient with infections of the central nervous system. *J. infect. Dis.*, 137:384,1988.

Brown, Z., Gerristen, M.E. and Carley, W.W.: Chemokine gene expression and secretion by cytokine-activated human microvascular endothelial cells – Differential regulation of monocyte chemoattractant protein-1 and interleukin-8 in response to interferon- γ . *Am. J. Pathol.*, 145: 913, 1994.

Bruijnzeel, P.L.B., Kuijper, P.H.M. and Rihs, S.: Eosinophil migration in atopic dermatitis. 1: Increased migratory responses to N-formyl- methionyl- leucyl- phenylalanine, neutrophil- activating factor, platelet activating factor and platelet factor 4. *J. Invest. Dermatol*, 100: 137, 1993.

Buck, C., Bundschu, J. and Gallati, H.: IL-6: A sensitive parameter for the early diagnosis of neonatal bacterial infection. *Pediatrics*, 93: 54, 1994.

Burstein, L. and Breining stall, G.N: Movement disorders in bacterial meningitis. *J. Pediatr., Infect. Dis*, 109:250, 1986.

Capobianchi, M.R., DeMarco, F. and Marco, P.: Acid labile human interferon alpha production by peripheral blood mononuclear cells stimulated by HIV-infected cells. *Arch. Virol.*, 99: 9, 1988.

Carr, M.W., Roth, S.J. and Luther, E.: Monocyte chemoattractant protein 1 acts as a T-lymphocyte chemoattractant. *Proc. Natl. Acad. Sci. USA*, 91: 3652, 1994.

Cavaillon, J.M. and Haeffner-Cavaillon, N.: Cytokines and inflammation. *Revue du Practien*, 43: 547, 1993. (English abstract).

Cazzola, M.: Natural history of idiopathic refractory sideroblastic anaemia. *Blood*, 71: 305, 1988.

Ceska, M., Effenberger, F. and Peichl, P.: Purification and characterization of monoclonal and polyclonal antibodies to neutrophil activation peptide (NAP-1). The development of high sensitive ELISA-methods for determination of NAP-1 and anti-NAP-1 antibodies presented at the second International Workers on Cytokines. Society for Leukocyte Biology, Hilton Head, Sc., 41: 72, 1989.

Chavanet, P., Bonnotte, B. and Guiguet, M.: High concentration of intrathecal interleukin-6 in human bacterial and nonbacterial meningitis. *J. Infect. Dis.*, 166: 428, 1992.

Christie, A.B. : Infectious diseases: Epidemiology and clinical practice. 4th edition p852 churchill living stone, Edinburg, London, New york, 1987.

Clarke, D. and Cost, K: Use of serum C-reactive protein in differentiating septic from aseptic meningitis in children. *J. Pediatr.*, 102: 718, 1993.

Cohen, A.B., Mac Arthur, C. and Idell, S: A peptide from alveolar macrophages that releases neutrophil enzymes into the lungs in patients with the adult respiratory distress syndrome. *Am. Rev. Respir. Dis.*, 137:1151, 1988.

Colditz, I.G.: Desensitization mechanisms regulating the plasma leakage and neutrophil emigration; in Gordon JL (ed): *Vascular endothelium: Interactions with circulating cells* p 175 New York, Elsevier, 1991.

Collins, P.D., weg, V.B. and Faccioli, L.H: Eosinophil accumulation by human IL-8 in the guinea-pig in vivo. *Immunology*, 79:312, 1993.

Collins, T., Korman, A.J. and Wake, C.T.: Immune interferon activates associated in variant chain gene in human endothelial cells and dermal fibroblasts. *Proc. Natl. Acad. Sci. USA*, 81: 4917, 1984.

Controni, G., Rodrigue, Z.W.J. and Hicks, J.M: cerebrospinal fluid lactic acid levels in meningitis. *J. pediatr.*, 91:379, 1987.

Coonrod, J.D. and Rytel, M.W.: Determination of aetiology of bacterial meningitis by counter – Immunoelectrophoresis. *Lancet*, 2: 197, 1992.

Cooper, M.D.: B-lymphocytes. Normal development and function. *N. Engl. J. Med.*, 317: 1452, 1987.

Cserr, H.F.: Physiology of the choroid plexus. *physiol Rev*, 51:273, 1981.

D'Andrea, A., Aste-Amezaga, M. and Valiaite, N.M.: Interleukin-10 inhibits human lymphocyte interferon gamma-production by suppressing natural killer cell stimulatory factor/IL-12 synthesis in accessory cells. *J. Exp. Med.*, 178: 1041, 1993.

Dahinden, C.A. and Geiser, T., Brunner, T.: Monocyte chemotactic protein 3 is a most effective basophil and eosinophil-activating chemokine. *J. Exp. Med.*, 179: 751, 1994.

Davey, P.G., Grukshank, J.K. and McManus, I.C.: Bacterial meningitis ten years experience. *JHYG (London)*, 88: 383, 1987.

De Benedetti, F., Massa, M. and Pignatti, P: Serum soluble IL-6 receptor and IL-6/ soluble IL-6 receptor complex in systemic juvenile rheumatoid, arthritis. *J.clin. Invest*, 93:2114, 1994.

De Simoni, M.G., Sironi, M. and De Luigi, A.: Intracerebro-ventricular injection of IL-1 induces high circulating levels of IL-6. *J. Exp. Med.*, 171: 1773, 1990.

Defrance, T., Vanber Vliet, B. and Aubry, J.P.: B-cell growth promoting activity of recombinant human interleukin-4. *J. Immunol.* 139: 1135, 1987.

Del Rio, M., Chrane, D. and Shelton, S: Ceftriaxone versus Ampicillin and chloramphenicol for treatment of bacterial meningitis in children. *Lancet*, 1:1241,1991.

Deorari, A.K., Verma, I.C. and Maheshwari: Prognostic factors related to mortality in meningoccal disease. *Indian. J. Med.* 86: 212, 1992.

Dewald, Thelen, M. and Baggiolini: Two transduction sequence are necessary of neutrophil activation by receptor agonists. *J.Biol. Chem.* 263:16179,1988.

Dianzani, F. and Antonelli, G.: Mechanism of induction of alpha interferon. The interferon system. A current review to 1987 (II). *Science*, 4: 51, 1987.

Dianzani, F. and Capobianchi, M.R.: Mechanism of induction of alpha interferon. The interferon system. A current review to 1987 (I). *Science*, 4: 21, 1987.

Digiovine, F.S. and Duff, G.W.: Interleukin 1: The first interleukin. *Immunology Today*, 11: 13, 1990.

Dinarello, C.A. and Mier, J.W.: Current concepts lymphokines. N. Engl. J. Med., 317: 940, 1987.

Dinarello, C.A: Interleukin-1. Rev. Infect. Dis, 6:51, 1991

Dixit, Kunkel, S.L. and Sarma, V: Molecular cloning of an endothelial derived neutrophil chemotactic factor: Identity with monocyte derived factor. FASEB (Fed. Am. Soc. Exp. Biol.) J, 3: A305, 1989.

Dodge, P.R. and Feigin, R.D.: Bacterial meningitis: Newer concepts of pathophysiology and neurological sequelae. Pediatr. clin. North. Am, 23:541, 1986.

Dowton, S.B. and Colten, H.R.: Acute phase reactants in inflammation and infection. Semin Hematol, 25: 84, 1990.

Dulkerian, S.J., Kilpatrick, L. and Costarino, A.T.: Cytokine elevations in infants with bacterial and aseptic meningitis. J. Pediatr, 126: 872, 1995.

Ehrlich, L.C., Hu, S. and Sheng, W.S.: Cytokine regulation of human microglial cell IL-8 production. The Journal of Immunology, 160: 1944, 1998.

Elsasser-Beile, U. and Von Kleist, S.: Cytokines as therapeutic and diagnostic agents. Tumour Biology, 14: 69, 1993.

Feigin, R.D.: Bacterial meningitis beyond the neonatal period. In: Feigin RD, Cherry JD (eds.) Text book of pediatric infectious diseases, 2nd edition P439 WB saunders company, Philadelphia, London, 1987.

Feigin, R.D.: Bacterial meningitis beyond the new born period. . In: Feigin RD, Cherry JD (eds.). Text book of pediatric infectious diseases 1st edition p293 Philadelphia, WB saunders Co, 1981.

Feigin,R.D. and Dodge, P.R.: Bacterial meningitis: Newer concepts of pathophysiology and neurologic sequelae. *Pediatr. Clin. North. Am*, 23:541,1986.

Figari, I.S., Mori, N.A. and Palladino and M.A. Jr.: Regulation of neutrophil migration and superoxide production by recombinant tumour necrosis factor alpha and beta: Comparison to recombinant interferon-gamma and IL-1 alpha. *Blood* 70: 979, 1989.

Fikrig , S.M., Berkovich, S. and Emmett, S.M.: Nitroblue tetrazolium dye test and differential diagnosis of meningitis. *J. pediatr.*, 82:855,1983.

Fikry, M.E.: 30 fevers: cerebrospinal fever, 4th edition, p117 Alexandria Al Maaref Establishment, 1987.

Fisher, R.M., Lipinski, J.K. and Cremin, B.J.: Ultrasonic assessment of infectious meningitis. *Clin. Radiol.*, 35: 267, 1990.

Fishman, R.A., sligar, K. and Hake, R.B: Effect of leukocytes on brain metabolism in granulocytic brain edema. *Ann. Neurol.*, 2:89,1988.

Fishman, R.A.: Brain oedema. *N.Engl. J.Med*, 293:706,1985.

Fong, Y. and Lowry, S.F.: Tumour necrosis factor in the pathophysiology of infection and sepsis. *Clin. Immunol. Immunopathol*, 55: 157, 1990.

Forrest, M.J., Eiermann, G.J. and Meurer, R.: The role of CD 18 in IL-8 induced dermal and synovial inflammation. *Br.J. Pharmacol*, 106:287, 1992.

Frei, K., Piani, D. and Pfister, H.W.: Immune-mediated injury in bacterial meningitis. *Intern. Rev. Exp. Pathol.*, 34: 183, 1993.

Frie, K., Malipiero, U.V. and Leist, T.P.: On the cellular source and function of IL-6 produced in the central nervous system in viral diseases. *Eur. J. Immunol.*, 19: 689, 1989.

Friedman, R.: Effect of interferon on cell membrane. *Tex. Resp. Biol. Med*, 41: 313, 1982.

Fu, X.Y., Kessler, D.S. and Levy, D.E.: Purification and functional characterization of ISGF3, the positive regulator of IFN-stimulated gene transcription. *J. Interf. Res.*, 9: 123, 1989.

Furia, M.B. and Ramdolph, G.J.: Chemokines and tissue injury *Am.J.pathol*, 146:1287, 1995.

Gallagher, R.B.: A special issue: The biology of complement. *Immunology Today*, 12: 291, 1991.

Georas, S.N., Ebisawa, M. and Bochner, B.S.: Expression of a functional laminin receptor $\alpha 6 \beta 1$, very late activation antigen 6 on human eosinophils. *Blood*, 82:2872, 1993.

Gertis: Infectious meningitis and encephalitis in Abbasia fever hospital from 1966 to 1989. *Am. Trop. J. Med.*, 2:47, 1991.

Guidet, B., Staikowsky, F. and Offenstadt, G.: Interleukines and TNF in septic shock. *Revue du praticien*, 43: 13, 1993.(English abstract).

Gurusawaami, P. and Ramakrishnan, S.: Biochemical parameters in the differential diagnosis of meningitis. *Indian J. Med.*, 86: 226, 1990.

Hack, C.: Interleukin 8 in sepsis: relation to shock and inflammatory mediators. *Infect. Immun.*, 60: 2835, 1992.

Hack, C.E., De Groot, E.R. and Felt-Bersma, R.J.F.: Increased plasma levels of interleukin-6 in sepsis. *Blood*, 74/5: 1704, 1989.

Halstensen, A., Ceska, M. and Brandtzaeg, P.: IL-8 in serum and CSF from patients with meningococcal disease. *J. Infect. Dis.*, 167 (2): 471, 1993.

Hammami, A.A.: cerebrospinal fluid meningitis in the Eastern Mediterranean region, *Epidemiology review. J.Egypt Pub. Health Ass*, 44:28,1978.

Handa, S.: Concentrations of interleukin-1 beta, interleukin-6, interleukin-8 and TNF-alpha in cerebrospinal fluid from children with septic or aseptic meningitis. *Kurume Med. J.*, 39(4): 257, 1992.

Haris, M.C., Costarino, A.T. Jr. and Sallivan, J.S.: Cytokine elevations in critically ill infants with sepsis and necrotizing enterocolitis. *J. Pediatr.*, 124: 105, 1994.

Harlan, J.M., Harber, L.A. and striker, S.E.: Effect of lipopolysaccharide on human endothelial cells in culture. *Thromb.Res*, 29:15,1991.

Harold, V.: Practical clinical biochemistry, 4th edition, XXVI P 699
New York: William Heinmann 1969.

Herrmann, F.: Interleukin 1 stimulates T lymphocytes to produce granulocyte-monocyte colony-stimulating factor. *J.clin Invest*, 18:1415,1990.

Hibi,M., Murakami, M. and Saito,M.: Molecular cloning and expression of an IL-6 signal transducer, gp 130. *Cell*, 63:1149,1990.

Hickman, C.J., Grim, J.A. and Mostowshi, H.S.: Regulation of human cytotoxic T-lymphocyte development by IL-7. *J. Immunol*, 145: 2415, 1990.

Honda,M., Yamamoto, S. and Cheng, M.: Human soluble IL-6 receptor: Its detection and enhanced release by HIV infection.*J.Immunol* 148:2175,1992.

Hopkins, S.J. and Rothwell, N.J.: Cytokines and the nervous system 1. Expression and recognition. *Trends Neuro sci*, 18:83,1995.

Huber, A.R., kunkel, S.L. and Todd, R.F.: Regulation of transendothelial Neutrophil migration by endogenous interleukin-8. *Science*, 254:99,1991.

Hughes, J.H.: IL-1 inhibits insulin secretion from isolated rat pancreatic islets by a process that requires gene transcription and mRNA translation. *J. Clin. Invest.*, 86: 856, 1993.

Hurwitz, A.A., Lyman, W.D. and Berman, J.W.: Tumour necrosis factor- α and transforming growth factor β up-regulate astrocyte expression of monocyte chemoattractant protein-1. *J. Neuroimmunol*, 57: 193, 1995.

Igarashi, M., Gilmartin, R.C. and Gerlad, B.: cerebral arteries and bacterial meningitis *Arch.Neurol.*, 41:531,1990.

Isaac, A. and Lindenmann, J.: Virus interference: The interferon. *Proc. R. Soc. Lond. (Biol)*, 147: 258, 1957.

Issekutz, T.B., Stoltz, J.M. and Webster, D.M.: Role of interferon in lymphocyte recruitment into the skin. *Cell Immunol*,99:322,1986.

Issekutz, T.B.: Dual inhibition of VLA-4 and LFA-1 maximally inhibits cutaneous delayed type hypersensitivity induced inflammation. *Am. J. Pathol.*, 143: 1286, 1994.

Issekutz, T.B.: Inhibition of in vivo lymphocyte migration to inflammation and homing to lymphoid tissues by the TA-2 monoclonal antibody: A likely role for VLA-4 in vivo. *J. Immunol*, 147: 4178; 1991.

Jackson, M.A. and Nelson, J.D.: Etiology and management of acute suppurative bone and joint infections in pediatric patients. *J. Pediatr. Orthop*, 2: 313, 1992.

Kennedy, D.H. and Fallon, R.J.: Tuberculous meningitis. J.A.M.A., 241: 264, 1989.

Kishimoto, T.: The biology of interleukin-6. Blood, 74: 1,1991.

Klbrick, S.: Current status of coxsackie and Echo viruses in human disease. Prog. Med. Virol, 6:22, 1979.

Klein, B., wijdenes, J. and Zhang, X.G.: Murine anti-interleukin-6 monoclonal antibody therapy for a patient with plasma cell leukemia. Blood, 78:1198,1991.

Klein, J.O., Feigin, R.D. and McCracken, G.H.Jr: Report of the task force on diagnosis and management of meningitis. Pediatrics, 78(5):959,1986.

Klempner, M.S., Dinarello, C.A. and Gallin, J.J.: Human leukocyte pyrogen induces release of specific granule contents from human neutrophils. J. Clin. Invest., 77: 1330, 1988.

Kluger, M. J.: Fever: Role of pyrogens and cryogens. Physiol. Rev, 71:93,1991.

Kunkel, Streiter, R.M. and Chensue, S.W.: Regulation of tumor necrosis factor-alpha and neutrophil activating protein-1 gene expression: potential role of cytokine directed cell communication during multiple organ injury. In respiratory distress syndrome: molecular to Bringham , M.K. and Stahlmann, M. editors. Vander bilt press, Nashville, TN. In Press. 1989.

Kupper, T.S.: Immune and inflammatory processes in cutaneous tissues. Mechanism and speculations. *J. clin. Invest.*, 86:1783, 1990.

Kurihara, N., Bertolini, D. and Suda, T.: IL-6 stimulates osteoclast-like multinucleated cell formation in long term human marrow cultures by inducing IL-1 release. *J. Immunol.*, 144:4226, 1990.

Larsen, C.G., Anderson, A.O. and Appella, E.: The neutrophil-activating protein (NAP-1) is also chemotactic for T lymphocytes. *Science*, 243: 1464, 1989.

Laurenzi, M., Siden, A. and Persson, M.A.A.: Cerebrospinal fluid interleukin-6 activity in HIV infection and inflammatory and noninflammatory disease of the nervous system. *Clin. Immunol. Immunopathol.*, 57: 233, 1990.

Lazaar, A.L. and Pure, E.: CD44: A model for regulated adhesion function. *Immunologist*, 3: 19, 1995.

Le, J. and Vilcek, J.: Interleukin 6: A multifunctional cytokine regulating immune reaction and the acute phase protein response. *Lab. Invest.*, 61:588, 1989.

Le, J. and Vilcek, J.: Tumor necrosis factor and IL-1: Cytokines with multiple overlapping biological activities. *Lab. Invest.*, 56: 234, 1990.

Lee, E.C., Lotz, M.M. and Mercurio, A.M.: The integrin $\alpha 6 \beta 4$ is a laminin receptor. *J. cell Biol.*, 117:671, 1992.

Leinon, M. and Drow, D.L.: Commercial kits for bacteriological diagnosis of meningitis. *Lancet*, 1: 500, 1989.

Leist, T.P., Freik and Kam-Hasen, S.: Tumor necrosis factor α in cerebrospinal fluid during bacterial, but not viral meningitis. *J. Exp. Med.*, 167: 1743, 1993.

Lesley, J., Hyman, R. and Kincade, P.W.: CD44 & its interaction with extracellular matrix. *Adv. Immunol.* 54: 271, 1993.

Liebler, Kunkel, S.L. and Burdick, M.D.: Production of IL-8 and monocyte chemotactic peptide-1 by peripheral blood monocytes: disparate responses to phytohemagglutinin and lipopolysaccharide. *J. Immunol.* 152:241, 1994.

Lindely, Aschauer, H. and Seifert, J.M.: Synthesis and expression in *Escherichia coli* of the gene encoding monocyte-derived and recombinant neutrophil-activating factor. *Proc. Natl. Acad. Sci. USA*, 85:9199:1988.

Lindquist, L., Linne, T. and Hansson, L.O.: Value of cerebrospinal fluid analysis in the differential diagnosis of meningitis. *Eur. J. Clin. Microbiol. Infect. Dis.*, 7: 374, 1988.

Loda, F.A., Collier, A.M. and Clezen, W.P.: Occurrence of diplococcus pneumonia in the upper respiratory tract of children. *J. Pediatr.* 87:1087, 1985.

Loetscher, P., Seitz, M. and Clark-Lewis: Monocyte chemotactic proteins MCP-1, MCP-2, MCP-3 are major attractants for human CD4+ and CD8+ T- lymphocytes. *FASEB. J.*, 8: 1055, 1994.

Long, N.C., Otterness, I. and Kunkel, S.L.: The roles of IL-1 β and TNF in lipopolysaccharide fever in rats. *Am J. Physiol*, 259: R724, 1990.

López-Cortés, L.F., Cruz-Ruiz, M. and Gómez-Mateos, J.: Interleukin-8 in cerebrospinal fluid from patients with meningitis of different etiologies: Its possible role as neutrophil chemotactic factor. *J. of infectious diseases*, 172: 581, 1995.

Luheshi, G. and Rothwell, N.J.: cytokines and fever. *Int. Arch. Allergy. Immunol*, 109:301, 1996.

Luheshi, G., Hopkins, S.J. and Le Feuvre, R.A.: Importance of brain IL-1 type II receptors in fever and thermogenesis in the rat. *Am. J. Physiol.*, 265:E585, 1993.

Luscinskas, F.W., Kiely, J.M. and Ding, H.: In vitro inhibitory effect of IL-8 and other chemoattractants on neutrophil-endothelial adhesive interactions. *J. Immunol.*, 149: 2163, 1992.

Lyle, W.H.: Lymphocytic meningo-encephalitis with myalgia and rash. *Lancet*, 21:1042, 1976.

Macneil, J.A., Suda, T. and Moore, K.W.: IL-10, a novel growth cofactor for mature and immature T-cells. *J. Immunol*, 143: 1210, 1989.

Maghazachi, A.A., Al-Aoukaty, A. and Schall, T.J.: C-C chemokines induce the chemotaxis of natural killer and IL-2 activated natural killer cells: A role for G-proteins. *J. Immunol*, 153: 4969, 1994.

Mandal, B.K. and Mayoum-white, R.T.: Bacterial meningitis. In: *Lecture notes on the infectious disease*. 4th edition P107 Oxford: Blackwell scientific publication, 1989.

Maraskovsky, E., Chen, W.F. and Shortman, K.: Interleukin-2 and interferons are two necessary lymphokines in the development of cytolytic T-cells. *J. Immunol*, 143: 1210, 1989.

Martich, G.D., Danner, R.L. and Ceska, M.: Detection of IL-8 and tumor necrosis factor in normal human after intravenous endotoxin: The effect of anti-inflammatory agents. *J. Exp. Med.*, 173: 1021, 1991.

Matsushima, Morishita, K. and Yoshimura, T.: Molecular cloning of human monocyte-derived neutrophil chemotactic factor (MDNCF) and the induction of MDNCF mRNA by interleukin-1 and tumor necrosis factor-*J.Exp. Med*, 167:1883, 1988.

Matsuzono, Y., Narita, M. and Akutsu, Y.: IL-6 in cerebrospinal fluid of patients with central nervous system infections. *Acta Paediatr.*, 84(8): 879, 1995.

Maurice, H.L.: Immunity and disease: Basic components of the immune system. The Medicine Group (Journals) Ltd, 1: 7, 1993.

Maxon, E.R. and Ostrow, P.T.: *Haemophilus influenzae* meningitis in infant rats: Role of bacteremia in pathogenesis of age-

dependent inflammatory responses in cerebrospinal fluid. *J. infect. Dis.*, 135:303,1989.

Mc Menamin, J.B. and Volpe, J.J.: Bacterial meningitis in infancy: Effects on intracranial pressure and cerebral blood flow velocity. *Neurology*, 34:500,1989.

Menkes, J.H.: The cause for low spinal fluid sugar in bacterial meningitis: Author look. *Pediatrics*, 44: 1, 1980.

Merico, F., Bergui, L. and Gregoret, M.G.: Cytokines involved in the progression of multiple myeloma. *Clin. Exp. Immunol.*, 92: 27, 1993.

Meydani, S.N. and Dinarello, C.A.: Influence of dietary fatty acids on cytokine production and its clinical implications. *Nutr. Clin. Pract.*, 8: 65, 1993.

Michaels, R.H.: Increase in H.influenza meningitis. *N.Engl. J. Med.*, 285: 666,1986.

Mielke, V., Bauman, J.G.J. and sticherling: Detection of neutrophil activating peptide NAP/IL-8 and mRNA in human recombinant IL-1 α and human recombinant tumor necrosis factor- α stimulated human dermal fibroblasts. *J. Immunol*,144:153,1990.

Minty, A., Chalon, P. and Deroc, J.M.: IL-13 is a new human lymphokine regulating inflammatory and immune responses. *Nature*, 362: 248, 1993.

Miyamasu, M., Hirai, K. and Takahashi, Y.: Chemotactic agonists induce cytokine generation in eosinophils. *J. Immunol*, 154: 1339, 1995.

Miyaura, C., Kusano, K. and Masuzawa, T.: Endogenous bone-resorbing factors in estrogen deficiency: Cooperative effects of IL-1 and IL-6. *J. Bone Miner Res.*, 10:1365, 1995.

Möller, A., Lippert, V. and Lessmann, D.: Human mast cells produce IL-8. *J. Immunol*, 151:3261, 1993.

Movat, H.Z., Burrowes, C.E. and Cybulsky, M.I.: Acute inflammation and a shawartzman-like reaction induced by interleukin-1 and tumor necrosis factor. *Am.J. Pathol*, 129:463, 1989.

Murray, P.D., McKenzie, D.T. and Surain, S.L.: Interleukin-5 and interleukin-4 produced by Peyer's patch T cells selectively enhance immunoglobulin A expression. *J. Immunol*, 139: 2669, 1987.

Nachbour, D., Gratwohl, A. and Herold, M.: Cytokine serum levels during treatment with high dose recombinant human IL-3 in a patient with severe aplastic anaemia. *Ann. Hematol.*, 66: 71, 1993.

Nachum, R., Lipsey, A. and Siegel, R.E.: Rapid detection of gram negative bacterial meningitis by the limulus lysate test. *N.Engl. J. Med.*, 289:931, 1983.

Neches, W. and Platt, M.: Cerebrospinal fluid LDH in 287 children, including 53 cases of meningitis of bacterial and nonbacterial etiology. *Pediatrics*, 41: 1097, 1988.

Nelson, N., Ecg-olofsson, O. and Larsson, L.: The diagnostic and predictive value of cerebrospinal fluid lactate in children with meningitis. *Act. Pediatr. Scand*, 75:52,1986.

Niemöller , V.M. and Tauber, M.G.: Brain oedema and increased intracranial pressure in the pathophysiology of bacterial meningitis. *Eur.J. Clin. Microbiol.Infect. Dis.*, 8:109,1989.

Niho, Y., Akashi, K. and Hayashida, K.: Basis of cytokines-history and present status. *Japan. J. Clin. Pathol.*, 41: 355, 1993.

Nohidge V.A.: Pneumococcal meningitis in sickle cell disease in childhood *Am.J. Dis. Child*, 137:29,1986.

Odio, C.M., Faingezicht, I. and Paris, M.: The beneficial effects of early dexamethasone administration in infants and children with bacterial meningitis. *N. Engl. J. Med.*, 324: 1525, 1991.

Okada, M., Kitahara, N. and Kishimoto, S: IL-6/BSF-2 functions as a killer helper factor in the in vitro induction of cytotoxic T cells. *J.Immunol* , 141:1543,1988.

Oppenheim, J.J., Kovacs, E.J. and Matsushima, K.: There is more than one IL-1. *Immunol Today*, 7: 45, 1991.

Oppenheim, J.J., Zachariae, C.O.C. and Mukaida, N.: Properties of novel proinflammatory supergene "intercrine" cytokine family. *Annu. Rev. Immunol.*, 9: 617, 1991.

Pagon, R.A., Bird, T.D. and Detter, J.C.: Hereditary sideroblastic anaemia and ataxia: An X-linked recessive disorder. *J. Med. Genet*, 22: 267, 1985.

Panton, Marrack's: Clinical pathology. 6th edition. P447 Churchill-London JA: Churchill Ltd, 1985.

Paquet, Ph. and Piérard, G.E.: Interleukin-6 and the skin. *Int. Arch. Allergy Immunol.*, 109: 308, 1996.

Pelletier, L., Bellon, B. and Druet, P.: Cytokines and immune response. *Revue du praticien*, 43: 536, 1993. (English abstract).

Peltola, H. and Valmari, P.: C-reactive protein in meningitis. *Lancet*, 2: 791, 1994.

Pestka, S.: The human interferons. From protein purification and sequence to cloning and expression in bacteria. *Arch. Biochem. Biophys.*, 221: 1, 1983.

Peveri, P., Walz, A. and Dewald, B.: A novel neutrophil activating factor produced by human mononuclear phagocytes. *J. Exp. Med*, 167: 1547, 1988.

Picker, L.J. and Butcher, E.C.: Physiological and molecular mechanisms of lymphocyte homing. *Annu. Rev. Immunol.*, 10: 561, 1992.

Pitha, P.M., Billelo, J.A. and Riggin, C.H.: Effect of interferon on retrovirus replication. *Tex. Rep. Biol. Med*, 41: 603, 1982.

Poli, V., Balena, R. and Fattori, E.: IL-6 deficient mice are protected from bone loss caused by estrogen depletion. *E.M.B.O.*, 13:1189, 1994.

Raimondi, A.J. and Dirocco, C.: The physiopathogenetic basis for the angiographic diagnosis of bacterial infection of brain and its covering in children. *Neurosurgery*, 5:398, 1989.

Ramadori, G.: IL-6, the 3rd mediator of the acute phase reaction, modulates hepatic protein synthesis in human and mouse. Comparison with IL-1 β and T.N.F. *Eur. Immunol*, 18: 1259, 1990.

Ramilo, O., Mustafa, M.M. and Porter, J.: Detection of IL-1 beta in cerebrospinal fluid of children with aseptic meningitis. *Am. J. Dis. Child*, 144: 349, 1990.

Ramilo, O., Saez-Lorenz, X. and Mertsola, J.: Tumor Necrosis factor α / cachectin and interleukin 1 β initiate meningeal inflammation. *J. Exp. Med*, 172:497, 1990.

Ratner, H.: Diagnosis of meningococcal meningitis. *J.A.M.A.*, 257: 1474, 1993.

Rendauld, J.C., Goethals, A. and Houssian, F.: Human P40 / IL-9. *J. Immunol*, 144: 4235, 1990.

Ridgway, D., Borzy, M.S. and Bagby, G.C.: Granulocyte macrophage colony-stimulating activity production by cultured human thymic non-lymphoid cells is regulated by endogenous interleukin-1. *Blood*, 72:1230, 1990.

Robbins, J.B., Schnerson, R. and Argamann, M.: Haemophilus influenzae type b: disease and immunity in humans. *Ann. Intern. Med.*, 78:259,1986.

Rola, Pleszczynski, M. and Grouin, S.: Asbestos-induced lung inflammation: role of local macrophage-derived chemotactic factors in accumulation of neutrophils in the lungs. *Inflammation*, 8:53,1984.

Rolfe, M.W., Kunkel, S.L. and Standiford, T.J.: Pulmonary fibroblasts expression of interleukin-8: A model for alveolar macrophage-derived cytokine networking. *Am. J. Respir. Cell. Mol. Biol.*, 5: 493, 1991.

Rollins, B.J., Walz, A. and Baggiolini, M.: Recombinant human MCP-1/JE induces chemotaxis, calcium flux and the respiratory burst in human monocytes. *Blood*, 78: 1112, 1991.

Romson, J.L., Hook, B.G. and Kunkel, S.L.: Reduction of the extent of ischemic myocardial injury by neutrophil depletion in the dog. *Circulation*, 67:1016,1983.

Roos, K.L.: Dexamethasone and nonsteroidal anti-inflammatory agents in the treatment of bacterial meningitis. *J. Clin. Therap.*, 12(4): 290,1990.

Rose-John, S. and Heinrich, P.C.: soluble receptors for cytokines and growth factors: Generation and biological function. *Biochem. J.*, 300:281,1994.

Rot, A., Kreiger, M. and Brunner, T.: RANTES and macrophage inflammatory protein-1 α induce the migration and activation of human eosinophil granulocytes. *J. Exp. Med.*, 176: 1489, 1992.

Rot, A.: Binding of neutrophil attractant/ activating protein-1 (Interleukin-8) to resident dermal cells. *Cytokine*, 4:347,1992.

Rot, A.: Neutrophil attractant/activation protein-1(Interleukin-8) induced in vitro neutrophil migration by haptotactic mechanism. *Eur.J.Immunol*,23:303,1993.

Rotbart, H.A. and Glode, M.P.: Haemophilus influenzae type b septic arthritis in children. *Pediatrics*, 75: 257, 1993.

Rothenberg, M.E., Luster, A.D. and Lilly, C.M.: Constitutive and allergen-induced expression of eotaxin in the guinea pig lung. *J. Exp. Med.*, 181: 1211, 1995.

Rothwell, N.J: Functions and mechanisms of interleukin-1 in the brain. *Trends pharmacol. Sci*,12:430,1991.

Rothwell, N.J: Mechanisms of the pyrogenic actions of cytokines *Eur. Cytokine Netw*, 1:211,1990.

Rusconi, F., Parizzi, F. and Garlaschi, L.: IL-6 activity in infants and children with bacterial meningitis. *Pediatr. Infect. Dis. J.*, 10: 117, 1991.

Saez-Lioren, X., Ranilo, O. and Mustafa, M.M.: Molecular pathophysiology of bacterial meningitis: Current concepts and therapeutic implications. *J.Pediatr*, 116:671,1990.

Samanta, A.K., Oppenheim, J.J. and Matsushima, K.: IL-8 (MDNCF) dynamically regulates its own receptor expression on human neutrophils. *J. Biol. Chem.*, 265: 183, 1989.

Samanta, Oppenheim, J.J. and Matsushima, K.: Identification and characterization of specific receptors for monocyte derived neutrophil chemotactic factor (MDNCF) on human neutrophil. *J. Exp. Med.*, 169: 1185, 1989.

Samuel, C.E.: Interferon induction of the antiviral state: proteins induced by interferons and their possible roles in the antiviral mechanisms of the interferon action. 2nd edition P111 Boca Raton: CRC Press, 1987.

Sarkar, R., Sharma, S. and Natarajan, R.: sulphadiazine resistant group "A" meningococci isolated during the meningitis out break in Delhi. *Indian J. Med. Res.*, 85: 612, 1987.

Saukkonen, K., Sande, S. and Cioffe, C.: The role of cytokines in the generation of inflammation and tissue damage in experimental gram-positive meningitis. *J. Exp. Med.*, 171: 439, 1990.

Schall, T.J. and Bacon, K.B.: Chemokines, leukocyte trafficking and inflammation. *Curr. Opin. Immunol.*, 6: 865, 1994.

Schall, T.J., Bacon, K. and Camp, R.D.R.: Human macrophage inflammatory protein α (MIP-1 α) and MIP-1 β attract distinct populations of lymphocytes. *J. Exp. Med.*, 177: 1821, 1993.

Schall, T.J., Bacon, K. and Toy, K.: Selective attraction of monocytes and T lymphocytes of the memory phenotype by cytokine RANTES. *Nature*, 347: 669, 1990.

Schall, T.J.: Biology of the RANTE/SIS cytokine gene family. *Cytokine*, 3: 165, 1991.

Scheld, W.M., Long, W.J.Jr. and Drodeur, J.P.: Effect of experimental *E. coli* meningitis in the blood-brain barrier: In vitro and in vivo studies (Abstract) *Clin. Res.*, 30:877 A, 1990.

Scheld, W.M., Quagliarello, V.J. and Wisplevey, B.: Potential role of host cytokines in *Haemophilus influenzae* lipopolysaccharide-induced blood brain-permeability. *Pediatr. Infect. Dis. J.*, 8: 910, 1992.

Schleimer, R.P., Sterbinski, S.A. and Kaiser, J.: IL-4 induces adherence of human eosinophils and basophils but not neutrophils to endothelium: Association with expression of VCAM-1. *J. Immunol.*, 148: 1086, 1992.

Schmid, J. and Weissmann, C.: Induction of mRNA for a serine protease and a beta -thromoglobulin-like protein in mitogen stimulated human leukocytes. *J. Immunol.*, 139:250, 1987.

Schroeder, J.M. and Christophers, E.: Identification of C5 adhesion protein and an anionic neutrophil-activating peptide (ANAP) in psoriatic scales. *J. Invest. Dermatol.*, 87:53, 1986.

Schroeder, J.M. and Christophers: Secretion of novel and homologous neutrophil- activating peptides by LPS-stimulated human endothelial cells. *J. Immunol*, 142:244, 1989.

Schwartz, J.F.: Ataxia in bacterial meningitis. *Neurology*, 22:1071, 1992.

Schweizer, R.C., Welmers, B.A.C. and Raaijmakers, J.A.M.: RANTES and interleukin-8- induced responses in normal human eosinophils: Effects of priming with interleukin-5. *Blood*, 83:3697, 1994.

Sebok, K., Woodside, D. and Al-Aoukaty, A.: IL-8 induces the locomotion of IL-2-activated natural killer cells. *J. Immunol*, 150: 1524, 1993.

Sehgal, P.B., Greininger, G and Tosato, G.: Acute phase and Immune Responses: Interleukin-6. *Ann. NY. Acad. Sci.*, 557: 581, 1989.

Sehmi, R., Cromwell, O. and Wardlaw, A.J.: Interleukin-8 is a chemo-attractant for eosinophils purified from subjects with a blood eosinophilia but not from normal healthy subjects. *Clin. Exp. Allergy*, 23:1027, 1993.

Seki, T., Joh, K. and Oh-Ishi, T.: Augmented production of interleukin-8 in cerebrospinal fluid in bacterial meningitis. *Immunology*, 80: 333, 1993.

Sell, S.H.: long term sequelae of bacterial meningitis in children *pediatr. Infect. Dis*, 2: 91, 1993.

Shaw, J., Messier, J.M. and Mercurio, A.M.: The activation dependent adhesion of macrophages to laminin involves cytoskeletal anchoring and phosphorylation of the $\alpha 6 \beta 4$ integrin. *J.Cell Biol.*,110:2167,1990.

Sheldon, M.W.: Cecil text book of Medicine: Meningitis 16th edition P1473 Philadelphia, London, Toronto, Tokyo, Sydney: WB saunders company 1983.

Shock, M.G. and Walling, A.D.: Meningitis: Describing the community health problem. *AMJ. Public health* 75:550,1989.

Silverstein, S. and Unkeless, J (eds): Innate immunity: Current opinion in immunology. 1st edition. P 97. Philadelphia: WB Saunders Co. 1987.

Smith, C.W., Kishimoto, T.K. and Abbass, O.: Chemotactic factors regulate lectin adhesion molecule 1 (LECAM-1) dependent neutrophil adhesion to cytokine-stimulated endothelial cells in vitro. *J. Clin. Invest.*, 87: 609, 1991.

Smith, D., Burrows, L. and Westwick, J.: Chemotactic cytokines: Interleukin-8- mediator of inflammatory lung disease?, in west-wick, J., piper,P.J(eds) p119- New York, Plenum press1991.

Smith, K.A.: Interleukin-2. Inception, impact and implications. *Science*, 240: 1169, 1988.

Smith, W.B., Gamble, J.R. and Vadas, M.A.: Interleukin-8 induces neutrophil transendothelial migration. *Immunology*, 72: 65, 1991.

Spagnuolo, P.J.: Haemophilus influenzal meningitis. The spectrum of disease in adults. *Medicine*, 61:74,1989.

Spiro, R.G.: Glycoproteins, their biochemistry, biology and role in human disease. *N. Engl. J. Med.*, 281, 1969.

Spits, H., Yssel, H. and Takebe, Y.: Recombinant interleukin-4 promotes the growth of human T cells. *J. Immunol*, 139: 1142, 1987.

Springer, T.A.: Traffic singals for lymphocyte recirculation and leukocyte emigration: The multistep paradigm. *Cell*, 76: 301, 1994.

Stahl, N. and Yancopoulos, G.: The alphas, Betas and kinases of cytokine receptor complexes. *Cell*, 74:587,1993.

Stahl, N., Boulton, T. and Farruggella, T.: Association and activation of Jak-Tyk kinases by CNTF-LIF-OSM-IL-6 beta receptor components. *Science*, 263:92,1994.

Standiford, T.J., Kunkel, S.L. and Phan, S.H.: Alveolar macrophage derived cytokines induce monocyte chemoattractant protein-1 expression from human pulmonary type II. Like epithelial cells. *J. Biol. Chem.*, 266, 1991.

Stanier, R.Y., Doudoroff, M. and Adelberg, E.A.: The microbial world, 2nd edition p451, permission of Prentice-Hall, Inc, Englewood Cliff's, N.J., 1963.

Strieter, Phan, S.H. and Showell, H.J.: Monokine- induced neutrophil Chemotactic factor gene expression in human fibroblasts. *J. Biol. Chem.* In press 1989.

Strieter, R.M., kasahara, K. and Allen, R.: human neutrophils exhibit disparate chemotactic factor gene expression. *Biochem. Biophys-Res commun*, 173:725, 1990.

Stutman, H.R. and Marks, M.I.: Bacterial meningitis in children: Diagnosis and therapy. *Clin. Pediatr.* 26: 431, 1990.

Suzuki, H., Yasukawa, K. and Saito, T.: Serum soluble interleukin-6 receptor in MRL/lpr mice is elevated with age and mediates the interleukin-6 signal. *Eur. J. Immunol.* 23:1078, 1993.

Sworn, M.J. and Urich, H.: Acute encephalitis in infectious mononucleosis. *J. path.*, 100:201, 1980.

Syrogianopoulos, G.A., Hansen, E.J. and Erwin, A.L.: *Haemophilus influenzae* type b Lipooligosaccharide induces meningeal inflammation. *J. Infect. Dis.* 157: 237, 1988.

Tabib Zadeh, S., Kong, Q.F. and Babakina, A.: Progressive rise in the expression of interleukin-6 in human endometrium during menstrual cycle is initiated during the implantation window. *Mol. Hum. Reprod.* 1:2793, 1995.

Tabib Zadeh, S.S., Sanhanam, U. and Sehgal, P.B.: Cytokine-induced production of IFN- 2β / IL-6 by freshly explanted human endometrial stromal cells. Modulation by estradiol- 17β . *J. Immunol.* 142: 3134, 1989.

Taga, T., Hibi, M. and Hirata, Y.: Interleukin-6 triggers the association of its receptor with a possible signal transducer, gp130. *Cell*, 58:573, 1989.

Tamura, T., Udagawa, N. and Takahashi, N.: Soluble IL-6 receptor triggers osteoclast formation by IL-6. *Proc. Natl-Acad. Sci. USA*, 90: 11924, 1993.

Tauber, M.G.: Brain oedema, intracranial pressure and cerebral blood flow in bacterial meningitis. *Pediatr. Infect. Dis. J.* 8:915, 1991.

Teir, H., Rytomaa, T. and Cederberg, A.: Studies on the elimination of granulocytes in the intestinal tract in rat. *Acta pathol. Microbiol. scand.*, 59:311, 1983.

Teti, G., Mancuso, G. and Tornasello, F.: Production of tumor necrosis factor-alpha and interleukin-6 in mice infected with group B streptococci. *Circ. Shock*, 38: 138, 1992.

Thelen, Peveri, P. and Kernen, P.: Mechanism of neutrophil activation by NAF, a novel monocyte-derived peptide agonist. *FASED (Fed-Am.Soc.Exp.Biol)* J, 2:2702, 1988.

Trinchieri, G.: Biology of natural killer cells. *Advanc. Immunol.*, 47: 187, 1989.

Tsai, M.L., Chen, W.C. and Wang, Y.C.: Cerebrospinal fluid interleukin-6, interleukin-8 and tumor necrosis factor-alpha in children with central nervous system infections. *Chung Hua Min Kuo Hsiao Erh Ko I Hsueh Hui Tsa Chil*, 37(1): 16, 1996.

Tugwell, P., Greenwood, B.M. and Warrel, A.D.: Pneumococcal meningitis a clinical and laboratory study. *J. Med.*, 45: 583, 1986.

Tunkel, A.R., Wispelwey, B. and Scheld, W.M.: Pathogenesis and pathophysiology of meningitis. *Infect Dis. Clin. N. Am.*, 4(4):555, 1990.

Tunkel, A.R., Wisplewey, B. and Scheld, W.M.: Bacterial meningitis recent advances in pathophysiology and treatment. *Ann Intern. Med.*, 112: 610, 1990.

Tuomanen, E., Liu, H. and Hengstler, B.: The induction of meningeal inflammation by components of the pneumococcal cell wall. *J. Infect. Dis.*, 151: 859, 1988.

Tureen, J.H., Dworkin, R.J. and Kennedy, S.L.: Loss of cerebrovascular autoregulation in experimental meningitis in rabbits. *J. Clin. Invest.*, 85:577, 1988.

Turksen, K., Kupper, T. and Degenstein, L.: IL-6: In sight to its function is skin by over expression in transgenic mice. *Proc. Natl. Acad. Sci. USA*, 89:5068, 1992.

Underman, A.E., Overturf, G.D. and Leedom, J.M.: Bacterial meningitis. In: *Disease-a-month* volume 24th P16 Chicago; Year Book Medical publisher 1987.

Vaddi, K. and Newton, R.C.: Regulation of monocyte integrin expression by β -family chemokines. *J. Immunol.*, 153:4721, 1994.

Valmari, P., Luhtalak, O. and Peltola, H.: C-reactive protein as detector of organic complication during recovery from childhood purulent meningitis. *J. Pediatr.*, 104: 869, 1994.

Van Damme, J., Proost, P. and Put, W.: Induction of monocyte chemotactic protein Mcp-1 and Mcp-2 in human fibroblasts and leukocytes by cytokines and cytokine inducers. *J. Immunol.*, 152: 5495, 1994.

Van Damme, J.: Interleukin-8 and related molecules. In: Thomson AW, ed. *The cytokine handbook*. 4th edition London: P201 Academic Press 1991.

Van Furth, A.M., Roord, J.J. and Van Furth, R.: Roles of proinflammatory and anti-inflammatory cytokines in pathophysiology of bacterial meningitis and effect of adjunctive therapy. *Infect. Immun.*, 64: 4883, 1996.

Van Snick, J.: Interleukin-6: An overview. *Annu. Rev. Immunol.*, 8:253, 1990.

Vazquez, A., Auffredon, M.T. and Gerard, J.P.: Sequential effect of high molecular weight B cell growth factor and of interleukin 2 on activated human B cells. *J. Immunol.*, 139: 2344, 1987.

Velasco, S., Tarlow, M. and Olsen, K.: Temperature-dependent modulation of lipopolysaccharide-induced interleukin-1 beta and tumor necrosis factor alpha expression in cultured human astroglial cells by dexamethasone and endomethacin. *J. Clin. Invest.*, 87: 1674, 1991.

Waage, A., Brandtzaeg, P. and Halstensen, A.: The complex pattern of cytokines in serum from patients with meningococcal septic shock. *J. Exp. Med.*, 169: 333, 1989.

Waage, A., Halstensen, A. and Shalaby, R.: Local production of tumor necrosis factor α , interleukin-1 and interleukin-6 in meningococcal meningitis. Relation to the inflammatory response. *J. Exp. Med.*, 170: 1859, 1989.

Ware, S.J. and Mclauchlan, S.: Haemophilus meningitis in portsmouth. *Lancet*, 1: 208, 1986.

Warringa, R.A., Schweizer, R.C. and Maikoe, T.: Modulation of eosinophil chemotaxis by interleukin-5. *Am. J. Respir. Cell Mol. Biol.*, 7:631, 1992.

Warringa, R.A.J., Koenderman, L. and Kok, P.T.M.: Modulation and induction of eosinophil chemotaxis by granulocyte-macrophage colony-stimulating factor and IL-3. *Blood*, 77:2694, 1991.

Warringa, R.A.J., Mengelers, H.J.J. and Raaymakers, J.A.M.: Up regulation of formyl-peptide and IL-8 induced eosinophil chemotaxis in patients with allergic asthma. *J. Allergy Clin. Immunol.*, 91:1198, 1993.

Watson, M.L., Westwick, J. and Fincham, N.J.: Elevation of PMN cytosolic free calcium and locomotion stimulated by novel peptides from IL-1 treated human synovial cell cultures. *Biochem. Biophys. Res. Commun.*, 155:1154, 1988.

Webb, L.M.C., Ehrenguber, M.U. and Baggiolini, M.: Binding to heparan sulphate or heparin enhances neutrophil responses to IL-8. *Proc. Natl. Acad. Sci. U.S.A.*, 90: 7158, 1993.

Welch, P.A., Namen, A.E. and Goodwin, R.G.: Human IL-7 A novel T-cell growth factor. *J. Immunol.*, 143: 3562, 1989.

Wenger, J.D., Hightower, A.W. and Facklam, R.R.: The bacterial meningitis study group. Bacterial meningitis in the United States, 1986; report of a multistate surveillance study. *J. Infect. Dis.*, 162: 1316, 1990.

White, Yoshimura, T. and Hook, W.: Neutrophil attractant /activation protein 1 causes human basophil histamine release. *Immunol lett*, 22: 151, 1989.

WHO: Epidemiological surveillance and control of cerebrospinal meningitis in Africa, 27: 347, 1983.

Williams J.D. and Burnie, J.: Selected aspects of the pathogenesis and pathophysiology of bacterial meningitis In: bacterial meningitis. 2nd edition P1 London, New York, Austria, Sydney, Academic Press, 1989.

Witt, D.P. and Lander, A.D.: Differential binding of chemokines to glycosaminoglycan subpopulations. *Curr. Biol.*, 4: 394, 1994.

Wolpe, S.D. and Cerami, A.: Macrophage inflammatory proteins 1 and 2: members fo a novel super-family of cytokines. *FASED. J.*, 2: 2565, 1989.

Xu, L., Kelvin, D.J. and Ye, G.Q.: Modulation of IL-8 receptor expression on purified human T lymphocytes in association with changed chemotactic responses to IL-6. *J. Leukoc. Biol.*, 57: 335, 1995.

Yokoyama, T., Oda, M. and Ogura, S.: Relationship of IL-8 and colony-stimulating factors to neutrophil migration in aseptic meningitis. *Acta Pediatr.*, 85: 303, 1996.

Yoshimura, Robinson, E.A. and Appella, E.: Three forms of monocyte-derived neutrophil chemotactic factor (MDNCF) distinguished by different lengths of the amino-terminal sequence. *Mol. Immunol.*, 26:87,1989.

Yoshimura,T., Matsushima, K. and Oppenheim, J.: Neutrophil chemotacitic factor produced by Lipopolysaccharide (LPS) stimulated human blood mononuclear leukocytes: Partial characterization and separation from interleukin-1(IL-1) *J.Immunol.*, 139: 788, 1987.

Zucali, J.R., Effenbein, G.J. and Barth, K.C.: Effects of human IL-1 and human T.N.F. on human T lymphocyte colony formation. *J. Clin. Invest.*, 80: 772, 1991.
