



REFERENCES



REFERENCES

1. *Akhan SE, lyibozkurt AC, Turfanda A*: Unscarred uterine rupture after induction of labor with misoprostol: a case report. Clin Exp Obstet Gynecol 2001; 28 (2): 118 – 20.
2. *Allot, H.A. and Palmer, C.R. (1993)*: Sweebling the membranes. A valid procedure in stimulating the onset of labor. Br. J. Obstet. Gynecol., 100: 898.
3. *American College of Obstetricians and Gynecologist (1991)*: Induction and augmentation of labor. Technical Bulletin No. 157, Washington, DC, July.
4. *American College of Obstetricians and Gynecologists(1999)*. Induction of labor. Practice bulletin no. 10. Washington, D.C.:ACOG,
5. *Amico, J.A.; Seitchik, J. and Robinso, A.G. (1984)*: Studies of oxytocinin plasma of women during hypocontractile labor. J. Clin. Endocrinol., Metab., 58: 274 – 9.
6. *Anderson, M.M. (1965)*: The state of the cervix and surgical induction of labor. J. of Obstet. & Gynecol. Of the British Commonealth, 72: 711.
7. *Anon (1991)*: A death associated with mifepristone/ sulprostone Lancet, 337: 969 – 70.
8. *Armando Fuentes and Mark Williams (1995)*: Cervical assessment. Clin. Obstet. Gynecol., 38: 224 – 231.
9. *Arulkumaran, S. and Ratnam, S.S. (1989)*: Influence of parity and cervical score on obscure on obstetric autcome of induced labor. In : Belfort, P., Pionotti, J.A.; Eskes TKAB (eds): Pregnancy and albor. The Parthenon publishing group Inc. New jersey, P. 91.

10. **Bados, O. and Backstrom O. (1987):** *Induciton of labor: a prospective randomized study into amniotomy and oxytocin as induction methods in a total uselected population.* Acta Obstet. Scand., 66: 537 – 541.
11. **Bell, R.J.; Sutton, B. and Eddiel, W. (1989):** Relaxin level in antenatal patients in vitro fertilization fertile & Steril., 52: 85 – 87.
12. **Benirschke, K. and Kaufman, P. (1990):** Pathology of the human placenta. New York: Springer Verlag, 130.
13. **Bennett, P.R.; Rose, M.P.; Myatt, L. and Elder, M.G. (1987):** Preterm labor; stimulation of arachidonic acid metabolism in human amnion cells by oacterial productions. Am. J. obstet. Gynecol., 156: 649.
14. **Birlain M.** Comparison of low doses of misoprostol with the traditional use of oxytocin for effective cervical ripening and labor induction. Obstet Gynecol 2001; 97 (4 Suppl 1): 567.
15. **Bishop, E.H. (1964):** Pelvic scoring for elective induction. Obstet. Gynecol., 24: 266 – 268.
16. **Blumenthal, P.D. and Ramanauskas, R. (1990):** Randomized trial of Dilapan and laminaria as cervical ripening agents before induction of labor. Obstet. Gynecol., 5:365.
17. **Blust D, Carlan SJ.** Buccal versus vaginal misoprostol for cervical ripening labor induction. Obstet Gynecol 2001; 97 (4 Suppl 1): S5
18. **Breeme, K. and Nilsson, B. (1984):** Prediction of time to delivery interval from start of contraction in induced labor: A life table analysis approach. Int. J. Gynecol. Obstet,22: 225.
19. **Bricker L, Luckas M.** Amniotomy alone for induction of labour. Cochrane Database Syst Rev 2002; 2:CD002862.

20. **Buccellato CA, Stika CS, Frederiksen MC.** A randomized trial of misoprostol versus extra-amniotic sodium chloride infusion with oxytocin for induction of labor. *Am J Obstet Gynecol* 2000; 182: 1039 – 44.
21. **Burghardt, R.C.; Barhoumi, R. and Dookwah, H. (1993):** endocrine regulation of myometrial gap junctions and their role in parturition. *Seminars reprod. Endocrin.*, 11: 250.
22. **Burnett, J.E. (1966):** Preinduction scring: an objective approach to induction of labor. *Obstet. Gynecol.*, 28: 479.
23. **Busowski, J. and parsons, M.(1995):** Amniotomy to induce labor. *CLin. Obstet. Gynecol.*, 38: 246 – 258.
24. **Bygdemman, M.; Christnsen, N.J. and Green, K. (1983):** Terminationm of early pregnancy – future development. *Acta obstetrical et gynecologica scandnavica supplement*, 113: 125 – 129.
25. **Calder, A.A. (1980):** Pharmacological management of the unripe cervix in the human. In: Naftolin, F. and stubblefiedl, P.G. (eds): *Dilation of the uterine cervix*, New York, Raven press, P. 317.
26. **Calder, A.A. (1990):** control of parturition. Scientific and clinical aspects in uterine contractility. In: Garfield, R.E. (ed.) *serono symposi*. Norwalk, Massachusetts, PP., 283 – 293.
27. **Clader, A.A. and Embrey, M.P. (1973);** Prostaglandins and unfavorable cervix. *Lancet*, 2: 1322.
28. **Clder, A.A.; Empery, M.P. and Tait, T. (1977):** Ripening of the cervix with extra – amniotic prostaglandin E2 in viscous gel before induction of labor. *Br. J. Obstet. Gynecol.*, 84: 264 – 268.

28. *Calder, A.A. (1990):* control of parturition. Scientific and clinical aspects in uterine contractility. In: Garfield, R.E. (ed.) serono symposi. Norwalk, Massachusetts, PP., 283 – 293.
29. *Calder, A.A. and Embrey, M.P. (1973);* Prostaglandins and unfavorable cervix. *Lancet*, 2: 1322.
30. *Calder, A.A.; Embrey, M.P. and Tait, T. (1977):* Ripening of the cervix with extra – amniotic prostaglandin E2 in viscous gel before induction of labor. *Br. J. Obstet. Gynecol.*, 84: 264 – 268.
31. *Calderyo – Barcia, R. and Poseiro, J.J. (1960):* Physiology of the uterine contractions. *CLin. Obstet. Gynecol.*, 386 – 408.
32. *Calkins, L.A. (1930):* variation in the length of labor. *Am. J. Obstet. Gynecol.*, 22: 294 – 297. Cited by Armando, Fuentes and mark Williams: Cervical assessment. *Clin Obstet. Gynecol.*, 38, 1995; 244 – 231.
33. *Calkins, L.A. (1941):* On Predicting the length of labor. *Am. J. Obstet, Gynecol.*, 42: 802. Cited by Armando, Fuentes and Mark Williams: Cervical assessment. *Clin. Obstet. Gynecol.*, 38, 1995; 224 – 231.
34. *Calkins, L.A.; Litzenberg, J.C. nad plass, E.D. (1931):* the length of labor. *Am. J. Obstet. GFynecol.*, 22: 604 – 614. Cited by Armando, Fuentes and Mark Williams: Cervical assessment. *Clin. Obstet. Gynecol.*, 38, 1995: 224 – 231.
35. *carlan SJ, Bouldin S, Blust D, O'Brien WF.* Safety and efficacy of misoprostol orally and vaginally: a randomized trial. *Obstet Gynecol* 2001; 98(1): 107 – 12.
36. *Carsten, M.E. and Miller, J.D. (1987):* A new look at uterine muscle contraction. *Am. J. Obstet. Gynecol.*, 157: 1303.

37. *Casey, M.L. and Macdonald, P.C. (1986):* Initiation of labor in women. IN the physiology and Biochemistry of the uterus in pregnancy and labor (Huszar, G. ed). CRC Press. Inc. Boca, Raton, Florida, 155 – 161.
38. *Challis J.R.G. and Vaughan, M. (1987) :* Steroid synthetic and PG measuring activity present in different cell population from human fetal membranes and deciduas. Am. J. obstet. Gynecol. 1470 : 1474.
39. *Chard, T. and Gibbens, G.L.D. (1983):* Spurt release of oxytocin during surgical induction of labor in women. Am. J. Obstet. Gynecol., 147: 678 – 80.
40. *Chaye, B. Tejani, N. and Verma, U. (1986):* Induction of labor with an elective breast pump. J. Reprod. Med., 31: 116 – 118.
41. *Cocks, D.P. (1955):* Significance of initial condition of cervix uteri to subsequent course of labor. BMJ, 1: 327. Cited by Armando, Fuentes and Mark Williams: Cervical assessment. Clin. Obstet. Gynecol., 38, 1995: 224 – 231.
42. *Collins, P.W. (1990) :* Misoprostol discovery development and chemical application. Med. Res. Rev. 10 : 149-72.
43. *Costa, S.H. and Vessey, M.P. (1993):* Misoprostol and illegal abortion in Riode Janeiro, Brazil. Lancet, 341: 1258 – 61.
44. *Crane JM, Young DC, Butt KD, Bennett KA, Hutchens D.* Excessive uterine activity accompanying induced labor. Obstet GYNecol 2001; 97(6): 926 – 31.
45. *Cross, W.G. and Pitkin, R.M. (1978):* Laminaria as an adjunct in induction of labor. Obstet. Gynecol., 51: 606.
46. *Crump, W.J. (1987):* Bishop score and labor duration: A new look. South med. J., 80: 1294.

47. **Cummiskey, K.C. and Dawood, M.Y. (1990):** Induction of labor with pulsatile oxytocin. *Am. J. Obstet. Gynecol.*, 163: 1868 – 74.
48. **Cunningham FG, Gant NF, Leveno KJ, Gilstrap LC, Hatt JC, Wenstrom KD.** Induction and augmentation of labor. In: *Williams Obstetrics*. 21st ed. McGraw – Hill. Norwalk, San Mateo and California. 2001a; 2: 470 – 481.
49. **Cunningham FG, Gant NF, Leveno KJ, Gilstrap LC, Hatt JC, Wenstrom KD.** Postterm pregnancy. In : *Williams Obstetrics*. 21st ed. McGraw – Hill. Norwalk, San Mateo and California. 2001 b; 2: 730 - 742
50. **Curbelo V, Bejar R, Benivschke K. and Gluck L. (1981):** Premature labour, 1. prostaglandin precursors in human placental membranes. *Obstet. Gynecol.* 57 : 973.
51. **Dawood, M.Y. (1990):** Uterine stimulation with oxytocin. In: Garfield, R.E., ed. *Uterine contractility*. Serono Symposia. Norwell, Massachusetts.
52. **Dawood, M.Y.; Raghavan, K.S.; Pociask, C. and Fuchs, F.(1978a):** Oxytocin in human pregnancy and parturition. *Obstet. Gynecol.*, 51: 138 – 43.
53. **Dawood, M.Y.; Wang, C.F.; Gupta, R. and Fuchs, F. (1978b):** Fetal contribution to oxytocin in human labor. *Obstet. Gynecol.*, 52: 205 – 9.
54. **Dawood, M.Y.; Ylikorkola, O.; Trivedi, D. and Fuchs, F. (1979):** Oxytocin in maternal circulation and amniotic fluid during pregnancy. *J. Clin. Endocrinol., Metab.*, 49: 429 – 34.
55. **Dhall, K.; Grover, V. dn mahendru, S.K. (1986):** prelabor, status evaluation and course of labor Asia. Oceania. *J. Obstet. Gynecol.*, 12: 25.

56. *Dudley, D.J.; Hunter, C.; Mitchell, M.D. and Varner, M.W. (1994):* clinical value of amniotic fluid interleukin-6 determination in the management of preterm labour. Br. J. Obstet. GYnecol., 101: 592 – 597.
57. *Dutton, W.A.W. (1958):* The assessment of the cervix at surgical induction. Can. Med. Assoc. J., 79: 463. Cited by Armando Fuentes and Mark Williams: Cervical assessment. Clin. Obstet. Gynecol., 38, 1995; 224 – 231.
58. *Edwards PK, Richards DS.* Preinduction cervical assessment. Clin Obstet GYnecol 2000; 43: 440 - 6
59. *Ekman, G.; Forman, A.; Marsail, K. and ulmsten, U. (1983):* Intravaginal versus intracervical application of prostaglandin E₂ in viscous gel for Cervical priming and induction of labor at term in patients with an unfavorable cervical state. Am. J. Obstet. Gynecol., 1 47: 657 – 61.
60. *Ekman, G.; Uldbjerg, N. and Ulmsten, U. (1985):* Comparison of intravenous oxytocin and vaginal prostaglandin E₂ gel in women with unripe cervixes and premature rupture of the membranes. Obstet. Gynecol., 66: 307 – 10.
61. *El Refaey, H.; Calder, L.; Wheatley, D.N. and Templeton, A. (1994):* Cervical priming with PGE₁ analogues misoprostol and gemeprost. Lancet, 343: 1207 – 9.
62. *El Refaey, H and Templeton, A. (1994):* Early Abortion induction by a combination of mifepristone and oral misoprostol. A comparison between two dose regimens of misoprostol and their effect on blood pressure. Br. J. Obstet. Gynecol. 101 : 792-796.
63. *El Refaey, H O'Brien, P., Motafa, W., Walder, J. and Rodeck, C.H. (1997):* Use of oral misoprostol in the prevention of post partum haemorrhage. British Journal of Obstetrics and Gynaecology. 104 : 336-339.

64. *El Refaey, H, Rajasekar, D., Abdalla, M.; Calder L. and Templeton, A. (1995):* induction of abortion with mifepriston (RU 486) and oral or vaginal misoprostol. N. Engl. J. Med. 332 : 983-987.
65. *Embrey, M.P. (1979):* Inductin of labor. Br. Med. J., 2: 793.
66. *Evans, M.L.; Dougan, M.B.; Moawad, A.H.; Evans, W.J.; Bryant – Greenwook, G.D. and Greenwood, F.C. (1983):* Ripening of the human cervix with porcine ovarian relaxin. Am. J. Obstet. Gynecol., 147: 410 – 414.
67. *Fields, H. (1966):* Induction of labor. Am. J. Obstet. Gynecol., 95: 426.
68. *Fletcher, H.M.; Mitchell, S.; Simeon, D.; Frederick, J. and Brown, D. (1993):* Intravaginal misoprostol as a cervical ripening agent. Br. J. Obstet. Gynecol., 100: 461 – 644.
69. *Foong LC, Vanaja K, Tan G, GHua S.* Membrane sweeping in conjunction with labor induction . Obstet gynecol 2000; 96: 539 – 42.
70. *Friedman, E.A. and Sachtleben, M.R. (1962):* Dterminant role of initial cervical dilation on the course of labor. Am. J. Obstet. Gynecol., 84: 930 – 935. cited by Armando Fuentes and Mark Williams: Cervical assessment. Clin. Obstet. Gynecol., 38, 1995; 224 – 231.
71. *Friedman, E.A.; Niswander, K.R.; Bayonet – Rivera, N.P. and Sachtleben, M.R. (1966):* Relation of prelabor evaluation to inducibility and the course of labor. Obstet. GYnecol., 28: 495.
72. *Frydman, R.; Lelaidier, C.; Baton, Saint Mleux, C. et al., (1992):* Induction of labor in women at term with mefiprisone, RU (486): A double blind, Randomized Placebo controlled study. Obstet. Gynecol., 80: 972 – 975.

73. **Fuchs, A.R.; Fuchs, F.; Husslein, P. and Soloff, M.S. (1984):** Oxytocin receptors in the human uterus during pregnancy and parturition. *Am. J. Obstet. Gynecol.*, 150: 734 – 41.
74. **Garret, W.J. (1960):** Prognostic signs in surgical induction of labor. *Med. J. aust.*, 47: 929. Cited by Armando Fuentes and mark Willimas: Cervical assessment. *Clin. Obstet. Gynecol.*, 38, 1995; 224 – 231.
75. **Gaud, H.T. and Connors, K.A. (1992):** Misoprostol dehydration kinetics in aqueous solution in the presence of hydroxypropyl methyl cellulose. *J. Pham. Sci.* 302 : 275-289.
76. **Gerson Weiss, M.D. (1995):** Relaxin used to produce the cervical ripening. *Clin. Obstet. Gynecol.*, 38: 293 – 300.
77. **Giannopoulos, G.; Jackson, K.; Kredentster, J. and Tulchinsky, K. (1985):** Prostaglandin E₂ and F₂ Rceptors inhuman myometrium during the menstrual cycle an din pregnancy and labor. *Am. J. Obstet. Gynecol.*, 153: 904.
78. **Gibb, D.M.F.; Arulkumaran, S.; Heng, S.H. and Ratnam, S.S. (1985):** Characteristics of induced labor. *Asia Oceania, J. Obstet. Gynecol.*, 11: 27 – 31.
79. **Goldman JB, Wigton TR.** A randomized comparison of extra-amniotic saline infusion and intracervical dinoprostone gel for cervical ripening. *Obstet Gynecol* 1999; 93: 271 - 4
80. **Gordon, A. and calder, A.A. (1977):** Estradiol applied locally to ripen the unfavorable cervix *Lancet*, 2: 1319.
81. **Gordon – Wright, A.P. and Elder, M.G. (1979):** Prostaglandin E₂ Tablets used intravaginally for the induction of labor. *Br. J. Obstet. Gynecol.*, 86: 32 – 36.

82. *Greer, I.A. and Galder, A.A. (1989):* Preinduction cervical ripening with extra-amniotic and vaginal PGE₂. J. Obstet. Gynecol., 16: 18.
83. *Guinn DA, Goepfert AR, Christine M, Owen J, Hauth JC.* Extra-amniotic saline, laminaria, or prostaglandin E(2) gel for labor induction with unfavorable cervix; a randomized controlled trial. Obstet Gynecol 2000; 96: 106 - 12
84. *Gustavii, B. (1973):* Studies on the mode of action of intra – amniotically and extra – amniotically injected hypertonic saline in therapeutic abortion. Acta. Obstet. Gynecol., scand. (Suppl.), 25:5.
85. *Hall, R, Duarte – Gardea M, Harless F., (2002):* Oral versus vaginal misoprostol for labor induction. Obstet Gynecol. 2002, Jun; 99(6) 1044 – 8.
86. *Harrison, R.F.; Flynn, M. and Graft, I. (1977):* assessment of factors an inducibility profile. Obstet. Gynecol., 49: 270.
Hendricks, C.H.; Brenner, W.E. and Kraus, G. (1970): Norma, cervical dilatation pattern in late pregnancy and labor. Am. J. Obstet. GYnecol., 106: 1065.
87. *Hirst, J.J., Haluska, G.J., Cook, M.J. and Novy; M.J. (1993) :* Plasma oxytocin and nocturnal uterine activity maternal but not fetal concentration increase progressively during late pregnancy and delivery in rhesus monkeys. Am. J. Obstet. Gynecol. 169 : 415.
88. *Hoffmann RA, Anthony J, Fawcus S. Oral misoprostol vs. Placebo in the management of prelabor rupture of membranes at term. Int J Gynecol Obstet 2001; 72(3): 215 – 221.*
89. *Hofmeyr GJ, Gulmezoglu AM.* Vaginal misoprostol for cervical ripening and induction of labour. Cochrane Database Syst Rev 2002;2:CD000941.

90. *Hughey, M.J.; McE, T.W. and Bird, C.C. (1976):* An evaluation of preinduction scoring systems. *Obstet. Gynecol.*, 48: 635.
91. *Husslein, P.; Fuchs, A.R. and Fuchs, F. (1981):* Oxytocin and the initiation of human parturition & prostaglandin release during induction of labor by oxytocin. *Am. J. Obstet. Gynecol.*, 141: 688 – 93.
92. *Inman, W.H.W. (1991):* report on current PEM studies: drugs for peptic ulcer or reflux. *Preser event monitor news*, 7: 32 – 4.
93. *Jeeva, M.A. and Dommissie (1982):* Laminaria tents or vaginal PG for cervical ripening. A comparative study. *Safr. Med. J.*, 61: 402
94. *Jonasson, A. and Larsson, B. (1989):* Biochemidal changes in human cervix connective tissue during pretreatment with laminaria tents in legal first trimester abortion. *Int. J. Obstet. Gynecol.*, 28: 361 – 364.
95. *Kalra, P.A.; Litherland, D.; Sallomi, D.F.; Critchley, H.O.D.; Falconer, G.F. and Holmes, A.M (1989):* Cardiac standstill induced by prostaglandin pessaries. *Lancet*, i : 1460 – 61
96. *Karim, S.M.M. (1971) :* Action of PG in pregnant woman. *Am. YN. Acad. Sci.*, 180 : 483.
97. *Karim, S.M.M.; Trussel, R.R.; Patel, R.C. and Hillier, K. (1968):* Response of pregnant uterus to PGF_{2a} induction of labor. *BR. Med. J.*, 4: 621.
98. *Karim, A. (1987):* Antiulcer prostaglandin misoprostol : Single and Multiple dose pharma cokinetics profile. *Prostaglandins*. 33 (Suppl) : 40-50.

99. *Katz VL, Farmer RM, Dean CA, Carpenter ME, (2000):* Use of misoprostol for cervical ripening. *South Med J*; 93(9): 881 – 4.
100. *Kase, N.G. (1990):* the hormones of reproduction na dhow they work. In: Kase, N.G.; Wingold, A.B.; Gershenson, D.M. (eds.). *Principles and practice of clinical gynecology*. Second edition. Churchill livingstone, Inc. New York, 91 – 119.
101. *Kavanagh J, Kelly AJ, Thomas J. (2002):* Sexual intercourse for cervical ripening and induction of labour. *Cochrane database syst Rev*; 2:CD003093
102. *Kazzi, G.M.; Bottoms, S.T. and Rosen, M.G. (1982):* Efficacy and safety of laminaria digitate for preinduction ripening of the cervix, *Obstet. GYnecol.*, 60 : 440.
103. *Keirse, M.J.N.C. (1992):* Estrogen for cervical ripening. In: Chalmes, I.; Enkin, M.; Keirse, M.J.N.C. (eds): *Effective care in pregnancy and childbirth*. Oxford University press, P. 1057.
104. *Keirse, M.J.N.C. (1993):* Prostaglandins in predinduction cervical ripening: Meta – analysis of worldwide clinical experience. *J. Reprod. Med.*, 38 – 100.
105. *Keirse, M.J.N.C.; Thiery, M.; Parewijck, W. and Mitchell, M.D. (1983):* Chronic stimulation of uterine PG synthesis during cervical ripening before onset of labor. *Prostaglandin*, 25: 671.
106. *Kelly AJ, Kavanagh J, Thomas, J.(2002).* Castor oil, bath and / or enema for cervical priming and induction of labour. *Cochrane Database Syst Rev* 2: CD003099
107. *Kramer, R.L.; Gilson, G.J.; Morrison, D.S.; Martin, D.; Gonzales, J.L. and Qualls, C.R. (1997):* A randomized trial of misoprostol and oxytocin for induction of labor. Safety and efficacy. *Obstet. Gynaecol.*, 89: 387-91.

108. *Lange, A.P.; Secher, N.J.; Westergaard, J.G. and skovagard, I. (1982):* Prelabor evaluation of inducibility. *Obstet. Gynecol.*, 60: 137.
109. *Lazor, L.Z.; Philipson, E.H.; Ingardia, C.J., Kobetitsch, E.S. and Curry, S.L. (1993):* A randomized comparison of 15 – and 40 – minutes dosing protocols *Obstet. GYnecol.*, 82: 1009 – 12.
110. *Lappet, P.C. and Yusy (1991):* Three dimensional structures of uterine elastic fibres: scanning electron microscopic studies. *Connect tissue res.*, 26: 15 – 31.
111. *Lee, H.Y. (1997) :* Aradomized double blind study of vaginal misoprosotol VS dinoprostone for cervical ripening and labour induction in prolonged pregnancy. Singapore. *Med. J.* 38 (7) : 292-4.
112. *Leppert, P.C (1995) :* Anatomy and physiology of cervical ripening. *Clin. Obstet. Guyaecol*, 38 : 267-279.
113. *Liggins, G.C. (1978):* Ripening of the cervix. In: Oliver, T.K. and Kirschbaum, T.K. (eds.). *New York Sem. Perinatol.*, 2: 261.
114. *Lindmark, G.; Zador, G. and Nilsson, B.A. (1975):* The induction of labor with PGF_{2a} by intravenous infusion. *Acta Obstet. Gynecol., Scand. (Suppl.)*, 37: 17.
115. *Lopez Bernal, A.; Bryant – Greenwood, G.D.; Hansell, D.J.; Hicks, B.R.; Greenwood, F.C and turnbull, A.C. (1987):* Effect of relaxin on prostaglandin E production by human amnion: Changes in relation to the onset of labor. *Br. J. Obstet. Gynecol.*, 94: 1045.
116. *MacDonald, D. (1970):* Surgical induction of labor. *Am. J. Obstet. Gynecol.*, 107: 908 – 911.

117. *MacDonald, P.C. and Casey, M.L. (1993)* : The accumulation of PG in amniotic fluid is an after effect of labour and not indicative of a role for PGE2 or PGF2a in the initiation of human parturition. *J. Clin. Endocrinol. Metab.* 76 : 1332.
118. *Mackenzie, I.Z. (1994)*: Labor including pregnancy termination for fetal anomaly. In: James, D.K.; Steer, P.J.; Weinger, C.P. and Gonik, B. eds. *High risk pregnancy management options*. Philadelphia, WB, Saunders.
119. *Mackenzie, E.Z. and Embrey, M.P. (1977)*: Cervical ripening with intravaginal prostaglandin E2 gel. *Br. Med. J.*, 1381 – 1381.
120. *MacInnes, D.R.; Naftolin, F. and VanDe Rrest, M. (1980)*: Cervical changes in pregnant women. In Naftolin, F. and Stubblefield, P.G. (eds). *Dilatation of the Uterine cervix* New York, Raven Press, pp, 1811.
121. *MacLennan, A.H. (1991)*: The role of the hormone relaxin in human reproduction and pelvic girdle, relaxation. *Scand. J. Rheumatol.*, 88 : 7 – 15.
122. *Mac Vicar, J. (1971)*: Failed induction of labor. *J. Obstet. Gynecol. Br. Commonw.*, 78: 1007 – 9.
123. *Manabe, Y.; Sagawa, N. and Mori, T. (1987)*: Condition of the cervix for the increase of plasma levels of 13, 14-dihydro – 15-keto- prostaglandin F_{2α} after amniotomy at term. *Prostaglandins*, 33: 757-766.
124. *Margulies, M.; Perez, C.G. and Voto, L.S. (1992)*: Misoprostol to induce labor. *Lancet*, 339 – 64.
125. *Mariani - Neto, Leao, E.J.; Baretto, E.M.; Ken, J.G. and DeAquino, M.M. (1987)*: Use of misoprostol for labor induction in still birth. *Rev. Paul. Med.*, 105: 325 – 8.

126. *Massil, H.Y.; Baker, A.C. and O'Brien, P.M.S. (1988):* A Comparison of oral prostaglandin E2 tablets with intravenous oxytocin for stimulation of labor after premature rupture of membranes at term. *Acta Obstet. Gynecol. Scand*, 67: 703-9.
127. *Mastrogiannis, D.S. and Knuppel, R.A. (1995):* Labor induced using methods that do not involve oxytocin. *Clin. Obstet. Gynecol.*, 38: 259 – 266.
128. *Mercer, B.; Pilgrim, P. and Sibai, B. (1991):* Labor induction with continuous low dose oxytocin infusion: Radomized trial. *Obstet. Gynecol.*, 77: 659- 63.
129. *Mercer BM, McNanley T, O'Brien JM, Randal L, Sibai BM(1995).* Early Versus late amniotomy for labor induction: A randomized trial. *Am J Obstet Gynecol* ; 173: 1371.
130. *Michael, M.V.; Michael, S.G. and Michael, J. (1995):* Misoprostol, a gastric cytoprotective agent. In : Michael, M.V.; Michael, S.; Michael, J. (eds). *Handbook of Gastrointestinal Drug – therapy*. Second Edition. Little, Brown and Company, New York, 315-318.
131. *Miller, W.L.; Redfield, M.M. and Brumett, J.C.Jr. (1989):* Integrated cardiac, renal and endocrine actions of endothelin. *J. Clin. Invest.*, 83: 317- 20.
132. *Mitchell, M.D.; Dudley, D.J.; Edwin, S.S and Lundin – Schiller, S. (1991);* Interleukin –6 stimulates prostaglandin production by human amnion and decidual cells. *Eur. J. Pharmacol.*, 192: 189 – 191.
133. *Mitchell, M.D.; Flint, A.P.F.; Bibby, J. et al. (1977):* Rapid increases in plasma prostaglandin concentrations after vaginal examination and amniotomy. *BMJ.*, 2: 1183 – 1185.

134. *Mitchell, B.F.; Challis, J.R.G. and Lukash, L. (1987) : Progesterone synthesis by human amnion, chorion and deciduas at term. Am. J. Obstet. Gynecol. 157 : 349.*
135. *Mitchell, B.F.; Rogers, K. and Wong, S. (1993) : The dynamics of prostaglandin metabolism in human fetal membranes and deciduas around time of parturition. J. Clin. Endocrinol. Metab. 77: 759.*
136. *Mortimer, G.; Hunter, I.C.; Stimson, W.H. and Gavan, A.D.T. (1985): A role for amniotic epithelium in control of human parturition. Lancet, 1: 1074.*
137. *Mortimer, G.; MacDonald, D.J. and Smeeth, A. (1985): A pilot study of the frequency and significance of placental villitis. Br. J. Obstet and Gynecol. 92 : 629-33.*
138. *Nagi, S.W.; To, W.K.; Lao, T.; Ho, P.C. (1996): Cervical priming with oral misoprostol in pre-labor rupture of membranes at term. Obstet. Gynecol., 87: 923 – 926.*
139. *Nakla, S.; Skinner, K.; Mitchell, B.F. and Challis, J.R.G. (1986) : Changes in PG transfer across human fetal membranes obtained after spontaneous labour. Am. J. Obstet. Gynecol. 155 : 1337.*
140. *Neiger, R.; Greaves, P.C. (2001) : Comparison between vaginal misoprostol and cervical dinoprostone for cervical ripening and labor induction. Tenn Med, 94 (1) : 25-7.*
141. *Neilson JP, Mifepristone for induction of labour (2002): Cochrane Database Syst Rev 2002; 2:CD002865*
142. *Nicolaides, K.H.; Welch, C.C; Koullapis, E.N. and Filshie, G.M. (1983): Cervical dilatation by lamipril – studies on the mechanism of action. Br. J. Obstet. Gynecol., 90: 1060 – 1064.*

143. *Nixon, W.C.W. (1952):* Abnormal uterine action in labor. J. Obstet. Gynecol. Br. Emp., 59: 624. Cited by Armando Fuentes and Mark Williams : Cervical assessment. Clin. Obstet. Gynecol., 38, 1995; 224-231.
144. *Norman, J.E; thong, K.J. and Baird, D.T. (1991):* Uterine contractility and induction of abortion in early pregnancy by misoprostol and mifepristone. Lancet, 338: 1233- 6.
145. *Norwitz E, Robinson J, Repke J(2002).* Labor and delivery. In: Gabbe SG, Niebyl JR, Simpson JL, eds. Obstetrics: normal and problem pregnancies. 4th ed. New York: Churchill Livingstone, 353 - 94
146. *Nugteren, D.H.; Beerthuis, and VanDrop, D.A. (1966) :* The enzymatic conversion of all - Cis, .8 -11 - 14 eicosatrienoic acid into PGE1 Recueil des travaux chimiques des pays Bas. 85 : 405 - 419.
147. *Okazaki, Sagawa, N, Okita, J.R.; Bleasdale, J.E.; MacDonald, P.C. and Johnston, J.M. (1981):* Diacylglycerol metabolism and arachidonic acid relase in human. Fetal membranes and deciduas year. J. Biol. Chem. 256 : 7316.
148. *Olah, K.S (1994):* Magnetic resonance imaging in the assessment of the cervical hydration state. Br. J. Obstet. Gynecol., 101: 255-257.
149. *Olson, D.M. and Zakart, T. (1993) :* Intrauterine tissue prostaglandin synthesis : regulatory mechanisms seminars. Reprod Endocin. 11 : 234.
150. *Paterson-Brown, S.; Fisk, N.M.; Edmonds, D.K. and Rodeck, C.H. (1991):* Preinduction cervical assessment by Bishop's score and transvaginal ultrasound. Eur. J. Obstet. Gynecol. Repro. Biol., 40: 17.

151. **Poulsen, H.K.; Moller, L.K.; Westergaard, A.G.; Thomsen, S.g.; Giersson, R.T. and Argrimsson, R. (1991):** Open randomized comparison of prostaglandin E₂ given by intracervical gel or vagitory for preinduction cervical ripening and induction of labor. *Acta. Obstet. Gynecol. Scand.*, 70: 549-53.
152. **Prins, R.P., Botton, R.N.; Marck, Neilson, D.R. and Watson, P. (1983):** Cervical ripening with intravaginal PGE₂ gel. *Obstet. Gynecol.*, 61: 459.
153. **Rabe, T.; Basse, H., Thuro, H.; Kiesel, L. and Runnebanum, b. (1987):** Action of the PGE₁ methyl analogue misoprostol on the pregnant human uterus in the first trimester. *Geburtshilfe frauenheilked*, 47: 324- 31.
154. **Rall, T.W. and Scheifer, L.S. (1985):** Oxytocin, prostaglandins, ergot alkaloids, and other drugs, tocolytic agents. In : Gilman, A.G., Goodman, L.S., Rall, T.W.; Murad, F. (ed): *The pharmacological basis of therapeutics*, 7th edition. New York, Macmillan publishing.
155. **Rayburn, W.; Gosen, Ramadei, C; Woods, R. and Scott, J. (1988):** Outpatient cervical ripening with prostaglandim E₂ gel in uncomplicated post-date pregnancies. *Am. J. Obstet. Gynecol.*, 158: 1417 – 23.
156. **Rayburn, W.F. (1989):** PGE₂ gel for cervical ripening and induction of labor. A critical analysis. *Am. J. Obstet. Gynecol.* 160: 529.
157. **Robert, A.; Nezamis, J.E. and Phillips, J.P. (1967):** Inhibition of gastric secretion by prostaglandins. *Am. J. Dig. Dis.*, 12: 1073 – 6.
158. **Romano, W.M.; Lukash, L.A.; Challis, J.R.G. and Mitchell, B.F. (1986):** Substrate utilization for estrogen synthesis by human fetal membranes and deciduas. *Am. J. Obstet. Gynecol.*, 155: 1170 – 1175.

159. *Sanchez-Ramos L, Kaunitz AM (1993): A randomized trial of extra-amniotic fluid infusion with intra-cervical Foley catheter balloon versus PGE₂ vaginal gel for ripening of the cervix and inducing labor in patients with unfavorable cervix. Obstet. Gynecol., 82 : 290.*
160. *Sanchez – Ramos, L.; Kaunitz, A.M.; DelValle, G.O.; Delke, I.; Chroeder, P.A. and Brines, D.K. (1993): Labor induction with the prostaglandin E₁ methyl analogue misoprostol versus oxytocin: A randomized trial. Obstet. Gynecol., 81: 332-6, 172 : 87-90.*
161. *Sanchez – Ramos, L.; Farah, L.A.; Kaunitz, A.M.; Adair, C.D. Delvalle, G.O.; and Fuqua, P. (1995): Preinduction cervical ripening with commonly available prostaglandin E₂ gel : A randomized, double blind comparison with hospital-compounded preparation Am. J. obstet. Gynecol, 173 : 1079-4.*
162. *Sanchez – Ramos, L.; Kaunitz, A.M.; Wears, R.Z.; Dellke, I. and Gaudier, F.L. (1997): Misoprostol for cervical ripening and labor induction : A meta analysis: Obstet. Gynaecol., 89 : 633-42.*
163. *Sanchez-Ramos L, Gaudier FL, Kaunitz AM (2000). Cervical ripening and labor induction after previous cesarean delivery. Clin Obstet Gynecol. 43: 513-23.*
164. *Sanchez – Ramos, L.; Peterson, D.E.; Delke, I.; Goudier, F.L. and Kaunitz, A.M. (1998): Labour induction with prostaglandin E₁ misoprostol compared with dinoprostone vaginal insert a randomized trial. Obstet. Gynecol. 91 (3) : 401-5.*
165. *Sanchez-Ramos L and Kaunitz AM.(2000). Misoprostol for cervical ripening and labor induction: a systematic review of the literature. Clin Obstet Gynecol; 43: 475 - 88*

166. **Satin, A.J.; Hankins, G.D.V. and Yeomans, E.R. (1991):** A prospective study of two dosing regimens of oxytocin for the induction of labor in patients with unfavorable cervixes. *Am. J. Obstet. Gynecol.*, 165: 980 – 4.
167. **Satin, A.J.; Leveno, J.K.; Sherman, L. and McIntire, D.D. (1992):** Factors affecting the dose response to oxytocin for labor stimulation. *Am. J. Obstet. Gynecol.*, 166 : 1260 – 1.
168. **Schiff, E.; Ben-Baruch, G.; Peleg, E.; Rosenthal, T.; Alcalay, M.; Devir, M. and Mashlach, S. (1992):** Immunoreactive circulating endothelin-1 in normal and hypertensive pregnancies. *Am. J. Obstet. Gynecol.*, 166: 624 – 628.
169. **Schonhofer, P.S. (1991):** Brazil; misuse of misoprostol as an abortifacient may induce malformation. *Lancet*, 337 : 1534 – 2.
170. **Secher, N.J.; Lange, A.P.; Nielsen, F.H.; Pedersen, G.T. and Westergaard, J.G. (1981):** Induction of labor with and without primary amniotomy. *Acta. Obstet. Gynecol. Scand.*, 60 : 237 – 241.
171. **Shermn DJ, Frenkel E, Pansky M, Caspi E, Bukovsky I, Langer R (2001).** Ballon cervical ripening with extra – amniotic infudion of saline or prostaglandin E2: a double – blind, Randomized controlled study. *Obstet; Gynecol* 97: 375 – 80.
172. **Shepherd, J.H.; Sims, C. and Craft, I. (1976):** Extra-amniotic PGE₂ gel and unfavorable cervix. *Lancet*, 2: 709.
173. **Sherwood, O.P. (1988):** Relaxin IN : Knobil, E.; Neill, J. (eds) : The physiology a reproduction. New York; Raven Press, 585 – 673.
174. **Shirley, K.S. and William, F. (1995):** Outpatient cervical ripening. *Clin. Obstet. Gynecol.*, 38 : 301 – 309.

175. *Shyken, J. and Petrie, R. (1995):* Oxytocin to induce labor. Clin. Obstet. Gynecol., 38: 232 – 245.
176. *Smith CA, Crowther CA (2002).* acupuncture for induction of labour. Cochrane Database Syst Rev 2:CD002962
177. *Speroff, F.F; Robert, H.; Glass, Nathan, G. and Kase, (1989):* Clinical Gynecologic, Endocrinology and Infertility. 11 : 351-353.
178. *Speroff, L.; Glass, R. and Kase, N. (1994):* Prostaglandin. In : Clinical Gynecologic endocrinology and Infertility. Ed. 5. Baltimore, Williams & Wilkins, pp., 291 – 317.
179. *Steiner, A.L. and Creasy, P.K. (1983):* Methods of cervical priming. Clin. Obstet. Gynecol., 47 – 602.
180. *Stewart, P.; Kennedy, J.; Barlow, D.H. and Calder, A.A. (1981):* A Comparison of estradiol and PGE₂ for ripening the cervix. Br. J. Obstet. Gynecol., 88: 236.
181. *Stewart, P.; Kennedy, J.H.; Hillan, E. and Calder, A.A. (1983):* The unripe cervix – management with vaginal or extra-amniotic PGE₂. J. Obstet. Gynecol., 4: 90.
182. *Stubbs TM, Oxytocin for labor induction (2000).* Clin Obstet Gynecol 43: 489 - 94
183. *Szalay, S.; Husslein, P. and Grunberget, W. (1981):* Local application of prostaglandin E₂ and its influence on collagenolytic activity of cervical tissue. Singapore, J. Obstet. Gynecol., 12 : 15.
184. *Theiry, M.; Baines, C.J. Keirse, M.J.C. (1989):* The development of methods for inducing labor. In: Chalmers, I.; Enkin, M.; Keirse, M.J.N.C., Eds Effective care in pregnancy and childbirth Oxford: oxford University press, 969 – 980.

185. *Thornton, S.; Davison, J.M. and Baylis, P.H. (1989):* Amniotomy induced labor is not mediated by endogenous oxytocin. *Br. J. Obstet. Gynecol.*, 96: 974 – 978.
186. *Trofatter, K.F. (1993):* Endocervical prostaglandin E2 gel for preinductino cervical ripening; clinical trial results. *J. Reprod. Med.*, 38 : 78 – 82
187. *Turnbull, A.C. (1983):* Historic development of overview. In initiation of parturition; prevention of prematurity (Macdonald, P.C. & porter, J., eds.) *Reprot of the fourth ross conference on obstetric Research, Ross laboratories, Columbus. Ohio*, 87 – 94.
188. *Turnbull, A.C. and Anderson, A.B.M. (1967):* Induction of labor. Part I. Amniotomy *Journal of obstet. & Gynecol. Of British common Wealth*, 74: 849 – 54.
189. *Uldbjerg, N. (1989):* Cervical connective tissue in relation to pregnancy, labor and treatment with prostaglandin E₂. *Acta Obstet. Gynecol., Scand., Supplement*, 148:
190. *Uldbjerg, N.; Ekman, G.; Malmstorm, A.; Olsson, K. and Ulmsten, U. (1983a):* Ripening of the human uterine cervix related to changes in collagen, Gycosaminglycans and collagenolytic activity. *Am. J. Obstet. Gynecol.*, 147: 662.
191. *Ulmsten, U. (1988):* Interacervical ripening andor induction of labor. *Facultas Vienna. Facultar*, P42.
192. *Ulmsten, U.; Wingerup, L. and Ekman, G. (1983):* Local application of prostaglandin E2 for cervical ripening of induction of term labor. *Clin. Obstt. Gynecol.*, 26: 95 – 104.
193. *Ulmsten, U.; Wingerup, L.; Belfrage, P.; Ekman, G. and Wiqvist, N. (1982):* Intracervical gel for induction of term labor. *Obstet. GYnecol.*, 59: 336 – 339

194. *Ulmsten, U.; Wingerup, L.; Andersson, K.E. (1979):*
Comparison of prostaglandin, E₂ and intravenous oxytocin for induction of labor. *Obstet. Gynecol.*, 54: 581 – 584.

195. *Varaklis, K.; Gumina, R.; Stubblefield, P.G. (1995):*
Radomized controlled trial of vaginal misoprostol and intracervical prostaglandin E₂ Gel for induction of labour at term. *Obstet. Gynecol.* 86 : 541-544.

196. *Von Maillot, K.V.; Stuhlsatzh, W.; Mohama-radhakrishnan, V. and Greiling, H. (1979):* Changes in glycosaminoglycans distribution pattern in the human uterine cervix during pregnancy and labor. *Am. J. Obstet. GYnecol.*, 135: 503.

197. *Wein, P. (1989):* Efficacy of different starting doses of oxytocin for induction of labor. *Obstet. GYnecol.*, 74: 863 – 8.

198. *Willcourt, R.J.; Pager, D.; Wendel, J. and Hale, R.W. (1994):*
Induction of labor with pulsatile oxytocin by a computer – controlled pump. *Am. J. Obstet. Gynecol.*, 170: 603 – 8.

199. *Wing, D.A.; Jones, M.M.; Rahall, A.; Goodwin, T.M and Paul, R.H. (1995):* A comparison of misoprostol and prostaglandin E₂ gel for preinductin cervical ripening and labor induction. *Am. J. Obstet. Gynecol.*, 172: 1804 – 10.

200. *Wing, D.A.; Ortiz-Omphroy, G. and Paul R.H. (1997) :*
Acomparision of intermittent vaginal administration of misoprostol with continous dinoprostone for cervical ripening and labour induction. *Am. J. obstet. Gynecol.* 177 (3) : 612-8.

201. *Wingerup, L.; Anderson, K.E. and Ulmsten, U. (1978):*
Ripening of the uterine cervix and induction of labor at term with PGE₂ in viscous gel. *Acta. Obstet. GYnecol., Scand.*, 57: 403.

202. *Word, R.A.; Kamm, K.E. and Casey, M.L. (1992):* Contractile effects of prostaglandins oxytocin, and endothelin in human myometrium in vitro. Refractoriness of myometrial tissue of pregnant women to PGE₂ and F_{2a}. J. Clin. Endocrinol. Metab., 75: 1027.
203. *Xenakis, E.M.; Pipre, J.M.; Canway, D.L.; Langer, O. (1997):* Induction of labor in the Nineties : Conquering the unfavorable cervix. Obstet Gynaccol., 90 : 235-9.
204. *Yamada, K.M. (1983):* Cell surface uneractions with extracellular materials. Annu. Rev. Biochem., 52 : 761 – 799.
205. *Young, J.H. and James Hamilton (1963):* Obstetrician and controversialist. Med. Hist., 7 : 62 – 73.
206. *Zanini, A.; Ghidini, A.; Norchi, S.; Benetta, E.; Cortinovis, I. and Bottinos (1990):* Preinductin cervical ripening with PGE₂ gel intracervical versus intravaginal route. Obstet. Gynecol., 70: 681
207. *Zeeman GG, Khan-Dawook FS, MY (1997):* OXytocin and its receptor in pregnancy and parturition: current concepts and clinical implications. Obstet Gynecol 89(5 pt 2): 873 - 83