

References

- Abdui-Karim, R. and Assail, N.S. (1961):** Pressor response to angiotensin in pregnant and non-pregnant women. *Am J Obstet Gynecol*, 82:246.
- Abe Y, Hasegawa Y, Miyanwto K, Yarnagushi M, Andoh A, Ibuki Y, and Igarashi M. (1990):** High concentrations of plasma immunoreactive inhibin during normal pregnancy in women. *J. Endocrinol. Metab* 71(1): 133-137.
- AlaFossi SL, Maenpaa J, Blauer M, Aine R Tuohirnaa P, and Punnonen R (1993).** Inhibin-A and -B in pre and postmenopause. *Maturitas* 30(3):273-281.
- American College of Obstetricians and Gynecologists (1996):** Hypertension in Pregnancy. Technical Bulletin No. 219.
- Ananth CV, Bowes WA, Savitz DA, Luther ER (1997):** Relationship between pregnancy-induced hypertension and placenta previa: A population-based study. *Am J Obstet Gynecol* 177:997.
- Anawalt BD, Bebb RA, and Matsumoto AM.** Serum inhibin-B levels reflect Sertoli cell function in normal men, and men with testicular dysfunction *J. Gun Endocrinol Metab* 1996; 8 1:334 1- 3345.
- Anumba DOC, Ford GA, Boys Ri, Robson SC (1999):** Stimulated nitric oxide release and nitric oxide sensitivity in forearm arterial vasculature during normotensive and preeclamptic pregnancy. *Am I Obstet Oynecol* 181:1479.
- Arias F (ed) (1993):** Practical guide to high risk pregnancy. Moasby Co.
- Auger J, Kunstmann JM, Czyplik, F. and Jouannet P.** Decline in semen quality among fertile men in Paris during the past 20 years. *N.Engl. J. Med* 1995; 322:281-285.
- Badonnel Y, Barbe D, Legagneur H, Poncelet F. and Schweitzer M. (1994):** Inhibin as a marker of hydatidiform mole: a comparative study with the determination of intact human chonionic gonadotrophin and its free?-subunit. *Clin. Endocrinol* 41:155-162.
- Bagatell CJ, McLachlan RI, and DeKresler DM (1989):** A comparison of the suppressive effects of testosterone and of a new potent GnRH antagonist on gonnadotropin and inhibin levels in normal men. *JGlin. Endocrinol Metab*; 69: 43-48.
- Baird DT, and Smith KB (1993).** Inhibin and related peptides in the regulation of reproduction. *Oxford Rev. Reprod. Biol* 15:191 - 232.

- Baird HG, Fraser HM, Hillier SG, Illingworth PJ, McNeilly A 5, Reddi K, and Smith K (1990):** Production and secretion of ovarian inhibin in women. In: Neuroendocrine Regulation of Reproduction. S.S.C. Yen and W.W. Vale, Eds., Mass. Serono Symposia, USA; pp:195-205.
- Baird D (1969):** Combined Textbook of Obstetrics and Gynaecology for Students and Practitioners. Edinburgh, Livingston, p 631
- Baker PN, Cunningham FG (1999):** Platelet and coagulation abnormalities. In Lindheimer MD, Roberts JM, Cunningham FG (eds): Chesley's Hypertensive Disorders in Pregnancy. 2nd ed. Stamford CT. Appleton & Lange. p 349
- Baker PN, Krasnow I, Roberts JM, Yeo K(1995):** Elevated serum levels of vascular endothelial growth factor in patients with preeclampsia. Obstet Gynecol 86:815.
- Baker, P.N. and Hackett, G.A. (1994):** The use of urinary Albumin/creatinine ratios and Calcium-creatinine ratios as screening tests for pregnancy-induced hypertension. Obstet Gynecol 83:745-9.
- Baker, P.N. (1993):** Screening tests for pregnancy-induced hypertension. In: Progress in Obstetrics and Gynecology, Volume 10, edited by John Studd. p. 69.
- Baker, P.N. and Broughton-Pipkin (1993):** Screening tests for pregnancy induced hypertension. In Studd J. (ed.) Progress in Obstet Gynecol. Churchill Livingstone, Edinburgh, London, Vol(10):69-81.
- Bardeguez AD, McNeer R, Frieri M, Verma UL, Tejani N (1991):** Cellular immunity in preeclampsia: Alterations in Tlymphocyte subpopulations during early pregnancy. Obstet Gynecol 77:859
- Barton JR, Sibai BM, Whybrew WD, Mercer BM (1993):** Urinary endothelin-1: Not a useful marker for preeclampsia. Am J Obstet Gynecol 168:599.
- Beaufils, M., Uzan, S., Donisimoni, R. and Colau, J.C. (1985):** Prevention of preeclampsia by early antiplatelet therapy. Lancet, 1:840-842.
- Beer, A.E. (1975):** Quoted from Doland, I., (ed): Practical obstetric problems 5th (ed), Lloyed: Luke 284:1979.
- Belizan JM, Villar J (1980):** The relationship between calcium intake and edema-, proteinuria-, and hypertension-gestosis: An hypothesis. Am J Clin Nutr 33:2202.
- Benedetti, T.J., Kates, R. and Williams, V. (1985):** Hemodynamic observations in severe pre-eclampsia complicated by pulmonary edema. Am J Obstet Gynecol, 152:330-334.
- Benedetto C, Marozio U, Neri I, Giarola M, Volpe A, Facchinetti F (2000):** Increased

- L-citrulline/L-arginine plasma ratio in severe preeclampsia. *Obstet Gynecol* 96:395.
- Benyo DF, Adatisa P, Conrad KP (2000):** Levels of inflammatory cytokines in normal term and preeclamptic placentas and their regulation by oxygen. *I Soc Gynecol Investig (Abstr 894)*, 7(1) (Suppl).
- Berg CI, Atrash HK, Koonin UM, Tucker M (1996):** Pregnancy-related mortality in the United States. 1987—1990. *Obstet Gynecol* 88:161.
- Bergmann, F., Rotmensch, S., Rosenzweig, B. How, H. and Chediak, J. (1991):** The role of Von Willebrand factor in pre-eclampsia. *Thromb Hemost*, 66:525-8.
- Bernstein, L., Dupue, R.J., Ross, R.K., and Henderson, B.F. (1989):** Cigarette smoking in pregnancy results in marked decrease in maternal HCG and Oestradiol levels. *Br J Obstet Gynaecol*, 96:92-6.
- Bewley S, Cooper D, Campbell S (1991):** Doppler investigation of uteroplacental blood flow resistance in the second trimester: A screening study for preeclampsia and intra-uterine growth retardation. *Br J Obstet Gynaecol* 98:871, 1991
- Bick, R.K. (1978):** Disseminated intravascular coagulation and related syndromes: Etiology, pathophysiology, diagnosis and management. *Am J Hematol*, 5:265-282.
- Blech, J.J.F., Thorburn, J., Greer, I.A. (1985):** Intravenous prostacyclin in management of pregnancy complicated by severe hypertension. *Chin Exp Hypertens*, B4:75-86.
- Bower S, Bewley S, Campbell S (1993):** Improved prediction of preeclampsia by two-stage screening of uterine arteries using the early diastolic notch and the color Doppler imaging. *Obstet Gynecol* 82:78.
- Broughton-Pipkin, F., Morrison, R.A. and O'Brien, P.M.S. (1986):** The effects of dietary supplementation with histolinic acid on the pressor and biochemical response to exogenous angiotensin II in human pregnancy. *Prog Lipid Res*, 25:425-429.
- Brown, C.E.L., Purdy, P.D. and Cunningham, P.G. (1988a):** Head computed tomographic scans in women with eclampsia. *Am J Obstet Gynecol*, 159:915.
- Brown, M.A., Gallery, E.D.M., Ross, M.R. and Esber, R.P. (1988b):** Sodium excretion in normal and hypertensive pregnancy: A prospective study. *Am J Obstet Gynecol*, 159:297.
- Brown, C.E.L., Cunningham, P.G. and Pritchard, J.A. (1987):** Convulsions in hypertensive, proteinuric primiparas more than 24 hours after delivery: Eclampsia or some other cause? *J Reprod Med*, 32:499.
- Brubaker DB, Ross MG, Marinoff D (1992):** The function of elevated plasma fibronectin in preeclampsia. *Am J Obstet Gynecol* 166:526.

- Burch, J.W., Standford, P.W. and Majerus, P.W. (1978):** Inhibition of platelet prostaglandin synthetase by oral aspirin. *J Clin Invest*, 61:314-19.
- Brunner HR, Gavras H (1975):** Vascular damage in hypertension. *Hosp Pract* 10:97.
- Balash J, C'reus M, Fabregues F, Garmona F, Casamitiana R, Ascaso C, and Vanrehl IA (1996):** Inhibin, FSH and age as predictors of ovarian response in IVF cycles stimulated with GnRI-I agonist and gonadotropin treatment. *Am. J. Obstet. Gynecol* 175(5):1226-1230.
- Beckmanm RB, Ling WE, Buzansky MR, Butes GW, Herbert WN, Laube WP. and Smith RP, Eds. GTD (1996):** in: *Obstetrics and Gynecology*, 2nd edition. Williams and Wilkins, USA, pp: 467-473.
- Birdsall M, Ledger W, Groome N, Abdalla H. and Muttukrisliid S (1997):** Inhibin-A and activin-A in the first trimester of human pregnancy. *J. Clin. Endocrinol. Metab* 82(5):1557-1560.
- Blaakaer, J.; Bennett, P; Micic, S.; Totager, L.K.; Hording, U.; Bock, J.E. and Lebecli, P.E. (1993):** The post-operative gonadotropin level Prostaglandins 35, 55 ; 46.
- Blaakaer J, Hoydall cJC, Micic S, Toftager LK, Hording U, Bennett P.,and Hoc J. (1995):** Ovarian carcinoma serum markers and ovarian steroid activity-is there a link in ovarian cancer? A correlation of inhibin, tetraoestin and CA 125 to ovarian activity and the gonadotropin level. *Eur. J. Obstet. Gynecol. Reprod. Biol*, 59(1)13-56.
- Bogges J F, Soules MR, Goff BA, Greer BE, Cain JM. and Tamitni HK (1997):** Serum inhibin and disease status in women with ovarian granulosa cell tumors. *Gynecol. Oncol*, 64(1):64-69.
- Browne FI (1946):** Sensitization of the vascular system in preeclamptic toxemia and eclampsia. *Br J Obstet Gynaecol* 53:510
- Brubaker DB, Ross MG, Marinoff D (1992):** The function of elevated plasma fibronectin in preeclampsia. *Am J Obstet Gynecol* 166:526.
- Buckler HM, Evans CA, Hamtara H, Burger H G. and Anderson DC (1991) .** Gonadotropin, steroid and inhibin levels in women with incipient ovarian failure during anovulatory and ovulatory rebound cycles. *J. Endocrinol. Metab*; 72: 116-124.
- Burger HG (1993):** Clinical utility of inhibin measurements. *J. Clin. Endocrinol. Metab*; 76:1392-1396.
- Burger HG, Robertson DM, and Cahir N (1996).** Characterization of inhibin

immunoreactivity in post-menopausal women with ovarian tumors. *Clin. Endocrinol*; 44:413-418.

Burger HG, Tin SC, Bangal ML, and DeKrester DM (1990): HCG raises serum immunoreactive inhibin levels in men with hypogonadotrophic hypogonadism. *Reprod. Fertil*; 2:137- 144.

Burger HG, and Igarashi M (1988): Definition and nomenclature, including related substances. *Endocrinology*; 122: 1701-1702.

Cameron, H.A. and Ardlie, N.G. (1982): The facilitating effects of adrenaline on platelet aggregation. *Prostaglandins Leukot Med*, 9:117-128.

Campbell, D.M. and MacGillivray, I. (1985): Preeclampsia in second pregnancy. *Br J Obstet Gynaecol*, 92:13 1-40.

Caritis S, Sibai B, Hauth I, Uindheimer MD, Dkebanoff M, Thorn E, Van Dorsten P, Uandon M, Paul R, Miodovnik M, Meis P, Thurnau 6 (1998): Low-dose aspirin to prevent preeclampsia in women at high risk. National Institute of Child Health and Human Development Network of Maternal— Fetal Medicine Units. *N Engl J Med* 338:701.

Caron C, Goudemaud I. Marey A. Beague D. Ducroux G, Drouvin F (1991): Are haemostatic and fibrinolytic parameters predictors of preeclampsia in pregnancy-associated hypertension? *Thromb Haemost* 66:410.

Castle WM (1977): Statistics in small doses. 2nd ed. Churchill Livingstone, London.

Chang 1K, Roman C, Heymann MA (1992): Effect of endothelium-derived relaxing factor inhibition on the umbilical— placental circulation in fetal lambs in utero. *Am J Obstet Gynecol* 166:727.

Chappell LC, Seed PT, Briley AL, Kelly FL, Lee R, Hunt RI, Parmar K, Bewley SJ, Shennan AH, Steer P1, Poston U (1999): Effect of antioxidants on the occurrence of preeclampsia in women at increased risk: A randomized trial. *Lancet* 354:810.

Chesley LC (1985): Diagnosis of preeclampsia. *Obstet Gynecol* 65:423.

Chesley, L.C. (1974): A short history of eclampsia. *Obstet Gynecol*, 43:599.

Chesley, L.C. and Cooper, D.W. (1986): Genetics of hypertension in pregnancy; possible single gene control of preeclampsia and eclampsia in the descendants of eclamptic women. *Br J Obstet, Gynaecol*; 93:898-908.

Chesley, L.C. (1984): History and epidemiology of preeclampsia/eclampsia. *Chin Obstet Gynecol*, 27(4): 801.

- Clark BA, Halvorson U, Sachs B, Epstein FH: Plasma endothelin levels in preeclampsia (1992):** Elevation and correlation with uric acid levels and renal impairment. *Am J Obstet Gynecol* 166:962.
- Clark SU, Cotton DB, Hankin GDV, Phelan JP (1991):** *Critical Care Obstetrics* 3rd ed. Maiden, MA, Blackwell Science.
- Conde-Agudelo A, Belizan JM (2000):** Risk factors for preeclampsia in a large cohort of Latin American and Caribbean women. *Br J Obstet Gynaecol* 107:75.
- Conde-Agudelo A.; Belizan M.J.; Lede, R. and Bergel, E.F. (1993):** What does elevated mean arterial blood pressure in the second half of pregnancy predict—gestational by hypertension or preeclampsia? *Am.J. Obstet. Gynecol*, 169:509-514.
- Conrad KP, Vernier KA (1989):** Plasma level, urinary excretion and metabolic production of cGMP during gestation in rats. *Am J Physiol* 257:R847.
- Cooper DW, Liston WA (1979):** Genetic control of severe preeclampsia. *J Med Genet* 16:409.
- Coppleson M, Monaghan JM, Marrow Cp, Tattersall HN. (1992):** Epidemiology of the cancer of the ovary In *Gynecological Oncology, Fundamentals, Principles and clinical practice*. Volume I. Second edition. Churchill Living stone Edinburgh London, P: 38-44.
- Cotter A, Molloy A, Scott JM, Daly SF (2001):** Elevated plasma homocysteine in early pregnancy: A risk factor for the development of severe preeclampsia. Presented at the 21st Annual Meeting of the Society of Maternal—Fetal Medicine, February 5—10, 2001 held in Reno, Nevada (abstract)
- Cowles T, Saleh A, Cotton DB (1994):** Hypertensive disorders in pregnancy. In James DK, Steer P1, Weiner CP, Gonik B (eds): *High Risk Pregnancy: Management Options*. London, Saunders, p 253
- Cuckle H, Sehmi I, Jones R (1998):** Maternal serum inhibin A can predict pre-eclampsia. *Br J Obstet Gynaecol*;105: 1101-1103.
- Cuckle H, Sehmi I, Jones R (1988):** Maternal serum inhibin A can predict preeclampsia. *Br J Obstet Gynaecol* 105:1101.
- Cuckle H (1998):** Effect of maternal age curve on the predicted detection rate in maternal serum screening for Down syndrome. *Prenat. Diagn*; 18(11):1 127-1 130.
- Cunningham, F.G., MacDonald, P.C. and Gnat, N.F. (2001):** Hypertensive disorders in pregnancy. *Williams Obstetrics*. 21st ed. Norwalk, Connecticut: Appleton and

Lang, 658-72.

- Cunningham, F.G., MacDonald, P.C., Gnat, N.F., Leveno, K.J. and Gilstrap, I.I.I. (1993):** Hypertensive disorders in pregnancy. Williams Obstetrics. 19th ed. Norwalk, Connecticut: Appleton and Lang, 763-817.
- Davidge ST, Hubel CA, Braden RD, Capeless EC, McLaughlin MK (1992):** Sera antioxidant activity in uncomplicated and preeclamptic pregnancies. *Obstet Gynecol* 79:897.
- D'Antona D, Wallace EM, Shearing C, Ash by JP, and Groome NP (1998):** Inhibin-A and pro-alpha C inhibin in Down's syndrome and normal pregnancies. *Prenat. Diagn*; 18(11):1122-1126.
- Davey DA, MacGillivray I (1988):** The classification and definition of the hypertensive disorders of pregnancy. *Am J Obstet Gynecol*; 158:892-898.
- Davidson EJ; Riley S C, Stephen A. Roberts S A; Shearing C H; Groomed N P; Martina C W (2003):** Maternal serum activin, inhibin, human chorionic gonadotrophin and 0L-fetoprotein as second trimester predictors of pre-eclampsia. *Br J Obstet Gynecol*, 76: 46.
- Debieve F, Beerlandt 5, Hubinont C, and Thomas K (2000):** Gonadotropins, prolactin, inhibin-A inhibin-B and activin-A in human fetal serum from mid pregnancy and term pregnancy *J. Clin. Endocrinol. Metab* 85(1):270-274.
- Degani, S., Abinader, E., Eibschitz, I., Sharpiro, I., Sharf, M. and Oettinger, M. (1985):** Isometric exercise test for predicting gestational hypertension. *J Obstet Gynecol*, 65:652-54.
- Dekker GA, Sibai BM (1998):** Etiology and pathogenesis of preeclampsia: Current concepts. *Am J Obstet Gynecol* 179:1359.
- DeKrester DM, Foulds LM, Hancock M, and Robertson DM (1994).** Partial characterization of inhibin, activin and follistatin in the term human placenta. *J. Clin. Endocrinol. Metab*; 79 (2):502-507.
- Dekker, G.A., Makovitz, J.W. and Wallenburg, H.C.S. (1990):** Prediction of pregnancy-induced hypertensive disorders by angiotensin II sensitivity and supine pressor test. *Br J Obstet Gynecol*, 97:817-821.
- DeKrester, DM, and Robertson DM (1989).** The isolation and physiology of inhibin and related proteins. *Biology of Reproduction*; 40:33- 47.
- Demura R, Suzuki T, Tajima 5, Mitsunashi, 5, Odagiri E, Denura H, and Ling N (1993).** Human plasma free activin and inhibin levels during the menstrual cycle. *J. Clin. Endocrinol. Metab*; 76(4): 1080-1082.

- Deng, L, Bremine, K., Hansson, L.O. and Blombach, M. (1994):** Plasma levels of Vonendothelial damage in preeclampsia. *Obstet Gynecol*, 84:941-5.
- Denson, K.W.E. (1977):** The ratio of factor VLII-related antigen and factor VIII biological activity as an index of hypercoagulability and intravascular clotting. *Thromb Res*, 10:107-19.
- Desvaux D, Haddad B. (2003):** Prevention of preeclampsia. *Presse Med.* 2003 Oct 11;32(33):1559-65.(Abstract)
- Dieckmann WI, Michel HL (1937):** Vascular—renal effects of posterior pituitary extracts in pregnant women. *Am J Obstet Gynecol* 33:131.
- Douglas KA, Redman CWG (1994):** Eclampsia in the United Kingdom. *BMJ* 309:1395.
- Dudley DI, Hunter C. Mitchell MD. Varnier MW, Gately M (1996):** Elevations of serum interleukin-12 concentrations in women with severe preeclampsia and HELLP syndrome. *J Reprod Immunol* 31:97.
- Duffus, G. and MacGillivray, I. (1968):** The incidence of preeclamptic toxemia in smokers and non-smokers. *Lancet*, 1:994-5.
- Emma J. Davidson EJ; Rileya S C, Stephen A. Roberts S A; Shearing C H; Groomed N P; Martina C W(2003):** Maternal serum activin, inhibin, human chorionic gonadotrophin and 0L-fetoprotein as second trimester predictors of pre-eclampsia. *Br J Obstet Gynecol*, 76: 46.
- Eramaa M, Heikinheimo K, Tuuri T, Hilden K, and Ritvos O (1993).** Inhibin/activin subunit mRNA expression in human granulosa-lutein cells. *Mol. Cell. Endocrinol*; 92:R15-R20.
- Eriksen, H.O., Kern Hansen, P. Brock, V. and Jensen, B.A. (1987):** Plasma fibronectin concentration in normal pregnancy and pre-eclampsia. *Acta Obstet Gynecol Scand*, 66:25-28.
- Estelles, A., Gilabert, J., Aznar, J., Loskoroff, D.H. and Schlef, R.R. (1989):** Changes in the plasma levels of type 1 and type 2 plasminogen activator inhibitors in normal pregnancy and in patients with severe pre-eclampsia. *Blood*, 74:1332-1338.
- Everett RB, Worley RI. MacDonald PC. Gant NF (1978):** Effect of prostaglandin synthetase inhibitors on pressor response to angiotensin 11 in human pregnancy. *J Clin Endocrinol Metab* 46:1007.
- Faas MM. Schuiling GA. Linton EA. Sargent IL. Redman CWG (2000):** Activation of peripheral leukocytes in late pregnancy and experimental preeclampsia. *Am J Obstet Gynecol* 182:351.

- Farnworth PG, Robertson DM, DeKrestet DM, and BurgerHG (1988):** Effects of 3 1K-daltons bovine inhibin on FSH and LH in rat pituitary cells in vitro: actions under basal conditions. *Endocrinology*; 122:207-213.
- Feng ZM, Bardin CW, and Chen CLC (1989).** Characterization and regulation of testicular inhibin B-subunit mRNA. *Mol. Endocrinol*; 3:939-948.
- Ferrazzani S. Caruso A. De Carohis S. Martino IV. Mancuso S (1990):** Proteinuria and outcome of 444 pregnancies complicated by hypertension. *Am J Obstet Gynecol* 162:366,
- Ferris, T.F. (1988):** Prostanoids in normal and hypertensive pregnancy. In: Rubin, P.C. ed. *Handbook of hypertension*, Vol. 10: Hypertension in pregnancy. Amsterdam Elsevier, pp. 102-117.
- Findlay JK (1993):** An update on the role of inhibin, activin and follistatin as local regulators of folliculogenesis. *Biol. Reprod*; 48:15-23.
- Findlay JK, Tsonis CG, Doglton B, and Forage RG. (1989):** Immunization against His 1 -Arg 166 (N) of the α -subunit of inhibin impairs fertility in sheep. *Endocrinology* 1989; 124:3122-3124.
- Fisher ER. Pardo V. Paul R. Hayashi IT (1969):** Ultrastructural studies in hypertension. 4. Toxemia of pregnancy. *Am J Pathol* 55: 901.
- Fitzgerald, DJ., Entman, S.S., Mulloyk, K. and FitzGerald, G.A. (1987):** Decreased prostacyclin biosynthesis preceding the clinical manifestation of pregnancy-induced hypertension. *Circulation*, 75:956-963.
- Fleisher, A., Schulman, H., Farmakides, G. (1986):** Uterine artery Doppler velocimetry in pregnant women with hypertension. *Am J Obstet Gynecol*, 154:806-13.
- Fowler PA, Evans LW, Groome NP, Templeton A, and Knight, PG. (1998):** A longitudinal study of maternal serum inhibin-A, inhibin-B, activin-A., activin-AB, pro- α C and follistatin during pregnancy. *Hum. Reprod* 1998; 13(12):3 530-3536.
- France JT, Keelan I, Song L, Liddell H, Zandderigo A, and Knox B (1996):** Serum concentrations of hCG and immunoreactive inhibin in early pregnancy and recurrent miscarriage: a longitudinal study. *Aust. N.Z.J. Obstet. Gynecol*; 36(3):325-330.
- Franchimont P, and Poncelet E (1994):** Two-site enzyme-immunoassays of inhibin. *Ares Serono Symposia Series. Frontiers in Endocrinology*; 3: 45-54
- Fraser RF, Mcasey ME, and Coney P (1998):** Inhibin and pro- α -C are elevated in pre-eclamptic pregnancy and correlate with hCG.s *Am. J. Reprod. Immunol*; 40(1):37-42.

- Frias AE Jr, Belfort MA. (2003):** How should we be managing severe preeclampsia? *Curr Opin Obstet Gynecol.* Dec;15(6):489-95.
- Friedman SA, Lindheimer MD (1999):** Prediction and Differential Diagnosis. In Lindheimer MD, Roberts JM, Cunningham FG (eds): *Chesley's Hypertensive Disorders in Pregnancy*. 2nd ed. Stanford, CT, Appleton & Lange, p 201
- Friedman SA, Lindheimer MD: Prediction and Differential Diagnosis (1999):** In Lindheimer MD, Roberts JM, Cunningham FG (eds): *Chesley's Hypertensive Disorders in Pregnancy*. 2nd ed. Stanford, CT, Appleton & Lange, p 201
- Friedman EA, Neff RK: Pregnancy outcome as related to hypertension. edema, and proteinuria. In Lindheimer MD. Katz Al, Zuspan FP (eds) (1976):** Hypertension in pregnancy. New York, Wiley, p 13
- Gallery, E.D.M., Hunyor, S.N., Ross, M. and Gyory, A.A. (1977):** Predicting the development of pregnancy associated hypertension. The place standardised blood pressure measurement. *Lancet*, I: 1273.
- Gant NF. Chand S. Worley RI. Whalley P1. Crosby UD. MacDonald PC (1978):** A clinical test useful for predicting the development of acute hypertension in pregnancy. *Am J Obstet Gynecol* 120:1.
- Gant, N.F., Jimenez, J.M., Whalley, P.J., Chands, S. and MacDonald, P.C. (1977):** A prospective study of angiotensin II pressor responsiveness in pregnancies complicated by chronic essential hypertension. *Am J Obstet Gynecol* 127:369.
- Gant, N.F., Chand, S., Worley, R.J., Whalley, P.J., Grosby, U.D. and MacDonald, P.C. (1974):** A clinical test useful for predicting the development of acute hypertension in pregnancy. *Am J Obstet Gynecol*, 120:1-7.
- Gant, N.F., Daley, G.L., Chand, S., Whalley, P.J. and MacDonald P.C. (1973):** A study of angiotensin II pressor response throughout primigravid pregnancy. I. *Chin Invest*, 52:2682-89.
- Garner, P.R., D'Alton, M.E., Dudley, D.K., Huard, P. and Hardic M (1990):** Preeclampsia in diabetic pregnancies. *Am J Obstet Gynecol*, 163:505-8.
- Garvasi M-T. Chaiworapongsa T. Pacora P. Naccasha N. Yoon BH. Mavmon E. Romero R (2001):** Maternal systemic inflammation: A mechanism of disease in preeclampsia. Presented at the 21st Annual Meeting of the Society of Maternal—Fetal Medicine. February 5—10. held in Reno. Nevada (abstract).
- Goodnight, S.H. (1989):** Vascular effects of n-3 fatty acids. *Invest Dermatol*, 93:102S-106S.
- Greer, I.A., Walker, J.J., Forbes, C.D.(1986):** The low dose aspirin controversy solved

at last? *Br Med J*, 391:1277-1278.

- Grobman WA, Wang EY (2000):** Serum levels of activin A and inhibin A and the subsequent development of preeclampsia. *Obstet Gynecol*;96:390-394.
- Groome NP, Illingworth PJ, O'Brien M, (1995):** Quantification of inhibin pro- α C-containing forms in human serum by a new ultrasensitive two-site enzyme-linked immunosorbent assay. *J Clin Endocrinol Metab*;80:2926-2932.
- Groome NP, Illingworth PJ, O'Brien M, Cooke I, Ganesan TS, Baird DT, and McNeilly AS (1994):** Detection of dimeric inhibin throughout the human menstrual cycle by two-site enzyme immunoassay. *Endocrinol*. 40:717-723.
- Grootenhuys AJ, Tinnerman ALA, Hordijk PL, and DeJong FH (1990):** Inhibin in immature rat Sertoli cell conditioned medium: a 32K-dalton α 3-B dimer. *Mol. Cell Endocrinol*; 70:109-116.
- Gudson, J.P. and Anderson, S. (1977):** A clinical evaluation of the "role over" test for pregnancy induced hypertension. *Amer J Obstet, Gynecol*; 127:1-3.
- Guthrie JR, Dennerstein L, Hopper JL, and Burger HG (1996).** Hot flushes, menstrual status and hormone levels in a population based sample of mid-life women. *Obstet. Gynecol*; 88(3):437-442.
- Hall, J.E.; Welt, C.K. and Cramer, D (1999):** W. Inhibin-A and inhibin-B reflect ovarian function in assisted reproduction but are less useful at predicting outcome. *Hum. Reprod*, 14(2):409-415.
- Hall JE, Schtoenfeld DA, Martin KA, and Growley Jr WF(1992):** Hypothalamic GnRH secretion and FSH dynamics during the luteal-follicular transition. *J. Clin. Endocrinol. Metab*; 74:600-607.
- Halligan, A., Bonnar, J., Sheppard, B., Darling, M. and Walshe, J. (1994):** Haemostatic, fibrinolytic and endothelial variables in normal pregnancies and pre-eclampsia. *Br J Obstet Gynaecol*, 101:488-492.
- Halverson LM, and Declerney AH (1996):** Inhibin, activin and follistatin in reproductive medicine. *Fertil. Steril*; 65(3):459-469.
- Hauth IC, Ewell MG, Levine RI, Esterhitz JR, Sibai B, Curet LB, Catalano PM, Morris CD (2000):** Pregnancy outcomes in healthy nulliparas who developed hypertension. *Obstet Gynecol* 95:24.
- Hauth I, Sibai B, Caritis S, VanDorsten P, Lindheimer M (1998):** The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units Maternal serum thromboxane B₂ concentrations do not predict

- improved outcomes in high-risk pregnancies in a low-dose aspirin trial. *Am J Obstet Gynecol* 179:1 193.
- Hauth IC. Goldenberg RL. Parker CR Jr. Philips B . Copper RL. DuBard MB. Cutter CR (1993):** Low-dose aspirin therapy to prevent preeclampsia. *Am J Obstet Gynecol* 168:1083.
- Hayward C. Livingstone I. Holloway S. Liston WA. Brock Dill (1992):** An exclusion map for preeclampsia: Assuming autosomal recessive inheritance. *Am J Hum Genet* 5(1):749.
- He W, Chang J and Zliang J.(1990):** A research on serum, urine and tumor tissue hyaluroic acid assays for detecting malignant ovarian tumors. (*Chinese Journal of Obstetrics and Gynecology* 30: 16 1-3
- Healy, D. and Wallace, E.M (1996):** Inhibins and activins: roles in clinical practice. *Br. J. Obst. Gynecol*; 103:945-956.
- Healy, D.; Poison, D., Yohkacitiya, T. and DeKrester, D (1990):** Inhibin and related peptides in pregnancy. *Bailliere 's Clinical Endocrinology-and Metabolism*; 4(2):233-247.
- Hee JP, MacNaughton J, and Banagh M (1993a):** FSH induces dose-dependent stimulation of immunoreactive inhibin secretion during the follicular phase of the human menstrual cycle. *J. Clin. Endocrinol. Metab*; 76:1340-1343.
- Hee JP, McNaughton J, Baigali Al, and Burger HG (1993b):** Perimenopausal pattern of gonadotrophins, immunoreactive inhibin, estradiol and progesterone. *Maturitas*; 18:9-20.
- Herbert V, Shaw S. Jayatileke E, Stopler-Kasdan T (1994):** Most free-radical injury is iron-related: It is promoted by iron, hemin, holoferitin and vitamin C, and inhibited by desferrioxamine and apoferitin. *Stern Cells* 12:289.
- Hess, L.W., O'Brien, W.F., Holmberg, J.A. and Winkel, C.A. (1986):** Plasma and amniotic fluid concentrations of fibronectin during normal pregnancy. *Obstet Gynecol* 68:25
- Hess, L.W., O'Brien, W.F., Holmberg, J.A., Winkel, C.A., Monaghan, W.P. and Hemming, V.G. (1986):** Plasma and amniotic fluid concentrations of fibronectin during normal pregnancy. *Obstet Gynecol*, 68:25-8.
- Hibbard, B.M. (1988):** Hypertensive disorders in pregnancy. *Principles of Obstetrics* (1st ed). Butterworth, London Edinburgh, p. 194-217.
- Hibbard, L.T. (1983):** Maternal mortality due to acute toxemia. *Obstet Gynecol*, 42:263.

- Hidaka, A., -Tomoda, S., Kitanada, T., Nakamoto, O. and Sugawa, T.(1990):** Prediction of pregnancy induced hypertension. VIIth World Congress of hypertension in pregnancy, Perugia, Italy, Scientific Program.
- Hinselmann H: Die Eklampsie. Bonn, F Cohen, (1924):** quoted from Cunningham, F.G., MacDonald, P.C. and Gnat, N.F. (2001): Hypertensive disorders in pregnancy. Williams Obstetrics. 21st ed. Norwalk, Connecticut: Appleton and Lang, 658-72.
- Hoff C. Peevy K, Giattina K. Spinnato IA, Peterson RDA (1992):**Maternal—fetal HUA-DR relationships and pregnancy-induced hypertension. Obstet Gynecol 80:1007.
- Hubel CA, McLaughlin MK. Evans RW, Hauth IA. Sims CI, Roberts JM (1996):** Fasting serum triglycerides, free fatty acids, and malondialdehyde are increased in preeclampsia, are positively correlated. and decrease within 48 hours postpartum. Am I Obstet Gynecol 174:975.
- Hubel CA, Roberts JM. Taylor RN, Musci Ti, Rodgers GM, McLaughlin MK (1989):** Uipid peroxidation in pregnancy: New perspectives on preeclampsia. Am I Obstet Gynecol, 161:1025.
- Hughes EG, Robertson DM, Handelsrnan DJ, Hayward 5, Healy DL, and DeKrester DM (1990):** Inhibin and estradiol responses to ovarian hyperstimulation: effects of age and predictive value for in-vitro fertilization outcome. J. Gun. Endocrinol. Metab; 70(2): 358-363.
- Hughes, E.C. (1972):** (ed). Obstet Gynecol terminology company. Philadelphia: Davis, F.A. 422.
- Hutt, R., Ogunniui, L.S.O., sullivan, M.H.F. and Elder, M.G. (1994):** Increased platelet volume and aggregation precede the onset of pre-eclampsia. Obstet Gynecol, 83:146-9.
- Illingworth PJ, Groome NP, and Duncan-Coliui W (1996).** Measurement of circulating inhibin forms during the establishment of pregnancy. J.Gun. Endocrinol. Metab; 8 1:1471-1475.
- Illingworth PJ, Reddi K Smith KB, and Baird DT (1991):** The source of inhibin secretion during the human menstrual cycle. J. Gun. Endocrinol. Metab; 73:667-673.
- Irgens-Moller, L., Hemmingsen, L. and Holm, J. (1986):** Diagnostic value of microalbuminuria in preeclampsia. Chin Chim Acta 157:295.
- Irion O, Masse I, Forest JC, Moutquin JM (1998):** Prediction of preeclampsia; low birthweight for gestation and prematurity by uterine artery blood flow velocity waveform analysis in low risk nulliparous women. Br I Obstet Gynaecol 105:422, 1998

- Irvine S, C'awood E, Richardson D, MacDonald E, and Aitken J. (1996):** Evidence of deteriorating semen quality in the United Kingdom: birth cohort study in 577 men in Scotland over 11 years. *Br. Med. J*; 3 12:467-471.
- Ishida, H., Tashiro, H. and Watanabe, M (1990):** Measurement of inhibin concentration in men: study of changes after castration and comparison with androgen levels in testicular tissues, spermatic venous blood and peripheral venous blood. *J. Clin. Endocrinol. Metab*; 70: 1019-1022.
- Jobling M, Mamers P, Healy DL, McLachlan V, Burger HG, Quinn M, Rome R, and Day AJ. A (1994):** prospective study of inhibin in granulosa cell tumors of the ovary. *Gynecol. Oncol*; 55:285- 289.
- Jackson, S., Lmhof, R. and Manning, N. (1990):** Prediction of Preeclampsia or (I.U.G.R) with Doppler analysis *Am J Obstet Gynecol* 162:110.
- Jaffe, E.A. and Wekster, B. (1979):** Recovery of endothelial cell prostacyclin production after inhibition by low doses of aspirin. *J Clin Invest*, 63:532-5.
- Janbu, T. (1989):** Uterine Doppler in high-risk pregnancies. *Acta Obstet Gynecol Scand* 68:387.
- Jann, R. and Spatgestosenbei (1954):** Hydrops fetus et placentae infolge Rhesus-Inkompatibilitat. *Arch Gynaecol*, 184:731-48. Quoted from: National High Blood Pressure Education Program Working Group Report on High Blood Pressure in Pregnancy. *Am S Obstet Gynecol*, 163:1689-712.
- Kallberg, B.E., Ryden, G. and Wichman, K. (1984):** Changes in the renin-angiotensin-aldosterone and kallikrein-kinin systems during normal and hypertensive pregnancy. *Acta Obstet Gynecol Scand*, 118:17-24.
- Kassar, N.S., Addridge, J. and Quirk, B. (1980):** Role-over test. *Obstet Gynecol*, 55:411.
- Kaupila, A., Bangait, M.; Burger, H. and Martikauten, H. (1992):** GnRH agonist analogue therapy in advanced recurrent granulosa cell tumors: further evidence of a role of inhibin in monitoring response to treatment. *Gynecol. Endocrinol* 1992; 6(4)271-274.
- Kazerooni T, Hamze-Nejadi S. (2003):** Calcium to creatinine ratio in a spot sample of urine for early prediction of pre-eclampsia. *Int J Gynaecol Obstet*. Mar;80(3):279-83.
- Kettel LM, Roseff SJ, Bangah ML, Burger et, HG, and Yenr SSC (1991):** Circulating levels of inhibin in pregnant women at term: simultaneous disappearance with estradiol and progesterone after delivery. *Clin. Endocrinol*; 34:19-23.

- Khong, T.Y., DeWolf, F., Robertson, W.B. and Brosens, I. (1986):** In adequate maternal vascular response to placentation in pregnancies complicated by preclampsia and small for gestational age infants. *Br S Obstet Gynaecol*, 93:1049.
- Kilby, M.D., Broughton-Pipkin, H. and Symonds, E.M. (1994):** Pharmacological prophylaxis in pregnancy induced hypertension. In Studd J (ed.) *Progress in Obstet and Gynaecol*, Vol 11:53-74.
- Kilpatrick DC, Uiston WA, Gibson F, Uivingstone I (1989):** Association between susceptibility to preeclampsia within families and HLA-DR4. *Lancet* 2:1063, 1989
- Kingsley DM (1994):** The TGF-B superfamily: new members, new receptors and new genetic tests of function in different organisms. *Genes Dev*; 8:133-146.
- Klien NA, Battaglia DE, Miller PB, Branigan FE, Giudice LC, and Soules MR (1996):** Ovarian follicular development and follicular fluid hormones and growth factors in normal women of advanced reproductive age. *J. Clin. Endocrinol. Metab*; 81(5):1946-1951.
- Knuist M, Bonsel 61, Zondervan HA, Treffers PE (1998):** Low sodium diet and pregnancy-induced hypertension: A multicentre randomized controlled trial. *Br J Obstet Gynaecol* 105:430.
- Koullapis, E.N., Nicolaides, K.H., Collins, W.P., Rodeck, C.H. and Campbell, S. (1982):** Plasma prostanooids in pregnancy induced hypertension. *Br J Obstet Gynecol*, 89:617-21.
- Kyle PM, Campbell 5, Buckley D, Kissane I, de Swiet M, Albano I, Millar 16, Redman CW (1996):** A comparison of the inactive urinary kallikrein: creatinine ratio and the angiotension sensitivity test for prediction of pre-eclampsia. *Br J Obstet Gynaecol* 103:981.
- Kyle, P.M., Clark, S.J., Buckly, D., Kissane, J. et al. (1993):** Second trimester ambulatory blood pressure in nulliparous pregnancy: a useful screening test for pre-eclampsia? *Br J Obstet Gynaecol*, 100:914-919.
- Knight PG (1996):** Roles of inhibins activins and follistatin in the female reproductive system. *Front. Neuroendocrinol*; 17(4): 47-509.
- Knight PG, Groome N, and Beard AJ. (1991):** Development of a two-site immunoradiometric assay for dimeric inhibin using antibodies against chemically synthesized fragments of the A- and B subunits. 1. *Endocrinol*; 129:R9-R12.
- Krummen LA, Woodruff 1K, DeGuzman G, Cox, El, Baly DL, Mann E, Gary 5,**

- Wong WL, Cossum P, and Mather JR (1993):** Identification and characterization of binding proteins for inhibin and active in human serum and follicular fluids. *Endocrinology*; 132: 43 1-443.
- Krummen LA, Toppari J, and Kim WH (1989):** Regulation of testicular inhibin subunit mRNA levels in-vivo: effects of hypophysectomy and selective FSH replacement. *Endocrinology*; 125:163-167.
- Labarrere C (1988):** Acute atherosclerosis. A histopathological hallmark of immune aggression? *Placenta* 9:108.
- Lahlou N, Cit abbert-Buffet N, Christin-Maitre 5, LeNestour E, Roger M, and Bouchard P (1999):** Main inhibitor of follicle stimulating hormone in the luteal follicular transition: inhibin-A, estradiol or inhibin-B *Hum. Reprod*; 14(5):1 190-1193.
- Laivuori H. Kaaja R. Ylikorkala 0, Hiltunen T, Kontula K (200):** 677 C—FT polymorphism of the methylenetetrahydrofolate reductase gene and preeclampsia. *Obstet Gynecol* 96:277.
- Lambert-Messerlian GM, Sailer DN, lumber, MB French CA, Peterson C J, and Canick LA (1998):** Second trimester maternal serum inhibin-A levels in fetal trisomy 18 and Turner syndrome with and without hydrops. *Prenat. Diagn*; 18(10): 1061-1067.
- Landesman R, Douglas RG. Holze E (1954):** The bulbar conjunctival vascular bed in the toxemias of pregnancy. *Am I Obstet Gynecol* 68:170.
- Lao, T.T. and Chin, R.R.H. (1988):** Blood pressure during 1st and 2nd trimesters as predicted preeclampsia. *Am J Obstet Gynecol* 159:1017.
- Lappohn RE, Burger HG, Bouma F, Ban gab M, and Krans,, M (1992):** Inhibin as a marker for granulosa cell tumor. *Acta Obstet. Gynecol*; 55:61-65.
- Lappohn RE, -Burger HG, Bouma L, Bangah M, Krans M. and DeBruijnH. WA (1989):** Inhibin as a marker of granulosa cell tumors. *N. Engl. J. Med*; 321:790-793.
- Lazarchick, J., Stubbs, T.M., Romein, L., Van Dorsten, J.P. and Loadholt, C.B. (1986):** Predictive value of fibronectin levels in normotensive gravid women destined to become pre-eclamptic. *Am S Obstet Gynecol*, 154:1050-2.
- Leaf, A. and Weber, P.C. (1988):** Cardiovascular effects of n-3 fatty acids. *N. Eng S Med*, 3 18:549-557.
- Lee W, Gonik B. Cotton DB (1987):** Urinary diagnostic indices in preeclampsia-associated oliguria: Correlation with invasive hemodynamic monitoring. *Am I Obstet Gynecol* 156:100.

- Lim KH, Friedman SA, Ecker IL, Kao U, Kilpatrick SI (1998):** The clinical utility of serum uric acid measurements in hypertensive disease of pregnancy. *Am J Obstet Gynecol* 178:1067.
- Lenton EA, DeKrester DM, Woodward AJ, and Robertson DM (1991):** Inhibin concentrations throughout the menstrual cycles of normal, infertile and older women compared with those during spontaneous conception cycles. *J. Clin. Endocrinol. Metab*; 73 (6):1 180-1190.
- Levine RI, Hauth IC, Curet LB, Sibai BM, Catalano PM, Morris CD, Der Simonian R, Esterlitz JR, Raymond EG, Bild DE, Clemens ID, Cutler IA (1997):** Trial of calcium to prevent preeclampsia. *N Engl J Med* 337:69.
- Lindheimer M, Woodruff TK (1997):** Activin A, inhibin A, and preeclampsia. *Lancet* 349:1266.
- Lin, S.Y and Goodfriend, T.L. (1970):** Angiotensin receptors. *Am J Physiol*, 218:1319-28.
- Li W, Olofsson JI, Jeung EB, Krisinger J, Yuen BH, and Leung P. (1994):** GnRH and cAMP positively regulate inhibin subunit mRNA level in human placental cells. *Life Sci*; 55: 1717-1724.
- Lockwood GM, Ledger WL, Barlow DII, Groorne NY, and Muttukrishna S (1998a):** identification of the source of inhibins at the time of conception provides a diagnostic role for them in very early pregnancy. *Am. J. Reprod. Immunol*; 40(5):303-308.
- LockWood G M, Muttukrishna S. and Ledger WL. (1998b):** Inhibins and activins in human ovulation, conception and pregnancy. *Human Reprod. Update*; 4(3): 284-295.
- Lokeshwar BL, Flurkadli KS, Sheth AR, and Block NL. (1993):** Human prostatic inhibin suppresses tumor growth and inhibits clonogenic cell survival of a model prostatic adenocarcinoma the Dunning rat tumor. *Cancer Research*; 53:4855.
- Long, P.A., Oats, J.N. and Gbeischer, N.A. (1981):** Labor onset preeclampsia. *J Obstet Gynecol* 21 :16
- Lubarsky SL, Barto JR, Friedman SA, Nasreddine S, Ramadan MK, Sibai BM (1994):** Late postpartum eclampsia revisited. *Obstet Gynecol* 83:502.
- MacGillivray, I. (1983):** Pre-eclampsia the hypertensive disease of pregnancy. W.B. Saunders, London, p. 64.
- Maddocks S, and Sharp RM. Assessment of the contribution of Leydig cells to the secretion of inhibin by the rat testis. *Mol Cell Endocrinol* 1989; 67:113-118.**

- Magann, E.F., William, C.R., Rerry, K.G., Chauban, S.P. (1994):** Factors relevant to mode of preterm delivery with syndrome of HELLP (hemolysis, elevated liver enzymes, and low platelets). *Am J Obstet Gynecol*, 170:1828-34.
- Massague J.** The transforming growth factor beta family. *Ann. Rev. Cell Biol* 1990; 6: 597-641.
- Masse Jacques F, Jean-Claude, M, Marcoux S (1993):** A proxpective study of several potential biological markers for early detection of the developing preeclampsia. *Am. J. Obstet. Gynecol.*, 169: 501-508.
- Mastrogiannis DS. O'Brien WF. Krammer I. Benoit R (1991):** Potential role of endothelin-1 in normal and hypertensive pregnancies. *Am I Ohstet Gvnecol* 165:1711.
- Mattar F. Sibai BM (2000):** Eclampsia: VIII. Risk factors for maternal morbidity. *Am I Ohstet Gvnecol* 182:307.
- Mayo KE, Cerelli CM, and Spiess I (1986):** Inhibin-A subunit eDNA from porcine ovary and human placenta. *Proceedings of the NationalAcademy of Science (USA)*; 83: 5849-5853.
- McCartney CP. Schumacher GFB. Spargo BH (1971):** Serum proteins in patients with toxemic glomerular lesion. *Am I Ohstet Gynecol* 111:580.
- McCartney CP (1969):** Pathological anatomy 01 acute hypertension ,of pregnancy. *Circulation (Suppl 2)* 30:37.
- Mc Cluggage WG, Ashe P, McBride H, Maxwell P, and Sloan JM. (1998):** Localization of the cellular expression of inhibin in trophoblastic tissue. *Histopathology* 32(3):252-256.
- McKay DO (1965):** Disseminated Intravascular Coagulation. New York, Harper & Row.
- McLachlan, R.I.; Matstimoto, A.M. and Burger, H G (1988).** Relative roles of follicle stimulating hormone and luteinizing hormone in the control of inhibin secretion in normal men. *J. Gun, invest*; 82:1-5.
- McLachlan, R.I.; Healy, D.L. and Robertson, D.M. (1987a)** Circulating inlmuoreactive inhibin in the luteal phase and early gestation of women undergoing ovulation induction. *Fertility and Sterility*; 48:1001-1005.
- McLacblan RI, Robertson DM, Heaiy DL, Burger HG, and DeKrester DM (1987b):** Circulating immunoreactive inhibin levels during the normal human menstrual cycle. *J. Clin. Endocrinol. Metab*; 65: 954-96 1.
- McLachlan, R.I.; Robertson, D.M.; Burger, H G. and DeKrester, D.M (1986a):** The

radioimmunoassay of bovine and human follicular fluid and serum inhibin. *Mol. Cell Endocrinol*; 46:175.

McLachlan RI, Robertson DM, DeKrester DM, and Buigert, HG (1986b): Plasma inhibin levels during gonadotropin induced ovarian hyperstimulation for IVF; a new index of follicular function. *Lancet*; 1:1233-1234.

McNatty KP, Hillier SG, Van den Boogaard AMJ, Trimbach-Ketner TGM, Reichart JrLE, and VanHall EV (1983): Follicular development during the luteal phase of the human menstrual cycle. *J. Clin. Endocrinol. Metab*; 56:1022.

McNaughton J. Banal M, McCloud P, Nee J, and Burger HG (1992): Age-related changes in FSH, LH, estradiol and immunoreactive inhibin in women of reproductive age. *Gen. Endocrinol*; 36:336-345.

McParland, P. (1993): Low dose aspirin in pregnancy. *Contemp Rev Obstet Gynaecol*, 5:30-8.

Medhamurthy R, Guller MID, Gay VL, Negro-Vilar A, and Plant TM. Evidence that inhibin plays a major role in the regulation of FSH secretion in the fully adult male Rhesus monkey (*Macaca Mulatta*). *Endocrinology* 1991; 129:389-395.

Melton DA, and Weeks DL (1987): A maternal mRNA localized to the vegetal hemisphere in *Xenopus* egg codes for a growth factor related to TGF- β . *Cell*; 51:861-867.

Meunier H, Cajander SB, and Roberts V.J. (1988): Rapid changes in the expression of inhibin and BB subunits in ovarian cell types during the rat oestrus cycle. *Mol. Endocrinol*; 2:1352-1363.

Mersol-Barg MS, Miller KF, Choi GM, Lee AC, and Kim MH (1990). Inhibin suppresses hCG secretion in term, but not first trimester, placenta. *J. Clin. Endocrinol. Metab*; 71:1294-1298.

Meuttier H, cajander SB, and Roberts VJ (1988): Rapid changes in the expression of inhibin and BB subunits in ovarian cell types during the rat oestrus cycle. *Mol. Endocrinol*; 2:1352-1363.

Mikhail MS, Anvaeghunam A. Garfinkel D. Palan PR. Iltasu I. Romm SL (1994): Preeclampsia and antioxidant nutrients: Decreased plasma levels of reduced ascorbic acid, α -tocopherol, and beta carotene in women with preeclampsia. *Am J Obstet Gynecol* 171:150.

Millar 1GB. Campbell 5K. Alhano 1DM. Higgins BR. Clark AD (1996): Early prediction of pre-eclampsia by measurement of kallikrein and creatinine on a random urine sample. *Br J Obstet Gynaecol* 103:421.

- Minami 5, Yarnoto Al, and Nakano R (1992):** Immunohistochemical localization of inhibin activin subunits in human placenta. *Obstet. Gynecol*; 80:410-414.
- Montag TW.** Tumor markers in gynecologic oncology. *Obstet. Gynecol. Surv* 1990; 45:94-105.
- Moore A, Krurninen LA, and Mather .JP. (1994):** Inhibins, activins, their binding proteins and receptors: interactions underlying paracrine activity in the testis. *Mol. Cell. Endocrinol*; 100:81-86.
- Moore, T.J., Taylor, T. and Williams, G.H. (1984):** Human platelet angiotensin II receptors: regulation by the circulating angiotensin level. *J Clin Endocrinol Metab*, 58:778-82.
- Moore, T.J. and Willimas, G.H. (1981):** Angiotensin II receptors on human platelets. *Circ Res*, 51:314-20.
- Morgan T. Craven C. Ualouel JM. Ward K (1999):** Angiotensinogen variant is associated with abnormal physiologic change of the uterine spiral arteries in first-trimester decidua. *Am J Obstet Gynecol* 180:95.
- Morgan U. Baker P. Broughton Pipkin F. Kalsheker N (1995):** Preeclampsia and the angiotensinogen gene. *Br J Obstet Gynaecol* 102:489.
- Morris NH. Eaton BM. Dekker S (1996):** Nitric oxide, the endothelium, pregnancy and pre-eclampsia. *Br J Obstet Gynaecol* 103:4.
- Moutquin, J.M., Rainville, C. and Giroux, L. (1985):** A prospective study of blood pressure in pregnancy: prediction of preeclampsia. *Am J Obstet, Gynecol* 151:19.
- Muttukrishna 5, North RA, Morris J, (2000):** Serum inhibin A and activin A are elevated prior to the onset of pre-eclampsia. *Hum Reprod* 2000;15:1640-1645.
- Muttukrishna S, Giwnberlain P, Evans LW, Asselin J, Groome NP, and Ledger WL (1999):** Amniotic fluid concentration of dimeric inhibins, activin-A and follistatin in pregnancy. *Eur. J. Endocrinol*; 140(5):420-424.
- Muttukrishna 5, Knight PG, Groome NP, Redman CW, Ledger WL (1997):** Activin A and inhibin A as possible markers for pre-eclampsia. *Lancet*;349:1285-1288.
- Muttukrishna S, Fowler PA, Groome NP, Mitchell GG, Robertson WR., and Knight PG (1994):** Serum concentrations of dimeric inhibin during the spontaneous menstrual cycle and after treatment with exogenous gonadotrophin. *Hum Reprod*; 9:1634-1642.
- Myatt U, Brewer AS. Uangdon O. Brockman DE (1992):** Attenuation of the vasoconstrictor effects of thromboxane and endothelin by nitric oxide in the human

- fetal—placental circulation. *Am J Obstet Gynecol* 166:224.
- Myatt, L. (1986):** Eicosanoids and blood pressure regulation. In: Hypertension in pregnancy proceedings of the 16th study group for the RCOG, vol.3. USA: Perinatology Press, pp. 167-87.
- Nakamura, T., Ito, M. and Matsui, K. (1986):** Significance of angiotensin sensitivity test for prediction PIH. *Obstet Gynecol* 67:388
- Naeye RU, Friedman EA: (1979):** Causes of perinatal death associated with gestational hypertension and proteinuria. *Am J Obstet Gynecol* 133:8.
- National Center for Health Statistics (2000):** Birth, marriages, divorces, and deaths: Provisional data for June 1999. National Vital Statistics Report, Vol 48, No. 8, 2000
- National High Blood Pressure Education Program: Working Group Report on High Blood Pressure in Pregnancy (2000).** *Am J Obstet Gynecol* 183:51.
- Neilson, J.B. (1987):** Dopplar ultrasound: commentary. *Br J Obstet Gynecol*, 94:929.
- Ness RB, Roberts JM (1996):** Heterogeneous causes constituting the single syndrome of preeclampsia: A hypothesis and its implications. *Am J Obstet Gynecol* 175:1365. 1996
- Neuman, M.M., Roan-El, R., Langer, R., Bukovsky, I. and Caspi, E. (1990):** Maternal death caused by HELLP syndrome (with hypoglycemia) complicating mild pregnancy induced hypertension in twin gestation. *Am J Obstet Gynecol*, 162:3727.
- Nishida M, Jirni S, Haji L, Hayashi I, Kai I, and Tasaka H (1991):** Juvenile granulosa cell tumor in association with a high serum inhibin level. *Gynecol. Oncol*; 40:90-94.
- Nishi K, Takayanagi R, Yanase T, Haji M, Hasegawa Y, and Nawata H (2000).** Inhibin-like immunoreactivity produced by the adrenal gland is circulating in-vivo. *Fukuok-Igaka-Zasshi*; 91(1): 8-20 (English Abstract).
- North RA, Taylor RS, Schellenberg I-C (1999):** Evaluation of a definition of preeclampsia. *Br J Obstet Gynaecol* 106:767.
- Nova A, Sibai BM, Barton JR, Mercer BM, Mitchell MD (1991):** Maternal plasma level of endothehin is increased in preeclampsia. *Am J Obstet Gynecol* 165:724.
- O'Brien, W.F. (1990):** Predicting preeclampsia. *Obstet Gynecol*, 75:445-52.
- Oian, P., Maetau, J.M., Nadleland, H. and Fadnes, H.O. (1986a):** Transcapillary fluid balance in preeclampsia. *Br J Obstet Gynecol*, 93:235.
- Oian, P., Monard-Honsen, H.P. and Maltau, J.M. (1986b):** Serum uric acid correlates with B 2-microglobulin in preeclampsia. *Acta. Obstet Gynecol Scand* 65:103.

- Ohashi M, Hasegawa Y, Haji M, Igarashi M, and Nawata H (1990).** Production of immunoreactive inhibin by virilizing ovarian tumor (Sertoli-Leydig tumor). *Clin. Endocrinol. Oxf*; 33(5):613-618.
- Okamoto S, Heav DL, Howlett DI, Rogers PAW, Leeto JF, Trounson AO, and Wood EC (1986):** An analysis of plasma estradiol concentration during clomiphene citrate-human menopausal gonadotropin stimulation in an IVF and embryo transfer program. *J. Clin. Endocrinol. Metab*; 63(3):736-740.
- Okonofua, F.E., Onwaidiegwu, U., Odunsi, A.O., Hussain, S. and O'Brien, S. (1990):** Diastolic blood pressure in patients with negative and positive stand up pressor test. *Obstet Gynecol*, 76:332.
- Olsen SF, Secher NI, Tabor A, Weber T, Walker 11, Gluud C (2000):** Randomized clinical trials of fish oil supplementation in high risk pregnancies. *Br J Obstet Gynaecol* 107:382.
- Oney, I. and Kaulhausen, H. (1982):** The value of the angiotensin sensitivity test in the early diagnosis of hypertensive disorders in pregnancy. *Am J Obstet Gynecol*, 142:17-20.
- Otani S, Usuki S, Saitoh T, Yanagisawa M, Iwasaki H, Tanaka I, Suzuki N, Fujino M, Goto K, Masaki T (1991):** Comparison of endothelin-1 concentrations in normal and complicated pregnancies. *J Cardiovasc Pharmacol* 17:S308.
- Paarlberg KM, DeJong CLD, Van Geijn HP, van Kamp 01, Heinen AG, Dekker GA (1998):** Total plasma fibronectin as a marker of pregnancy-induced hypertensive disorders: A longitudinal study. *Obstet Gynecol* 91:383.
- Page, E.W. and Christianson, R. (1976):** Influence of blood pressure changes with and without proteinuria upon outcome of pregnancy. *Am J Obstet Gynecol* 127:821-33.
- Palmer SK, Moore UG, Young DA, Cregger B, Berman JC, Zamudio S (1999):** Altered blood pressure and increased preeclampsia at high altitude (3100 meters) in Colorado. *Am J Obstet Gynecol* 180:1161.
- Palmer RMJ, Ashton DS, Moncada S (1988):** Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* 333:664.
- Pearlstone AC, Fournet N, Gainbone JC, Pang SC, and Buyalos RP. (1992):** Ovulation induction in women age 40 years and older: the importance of basal FSH level and chronological age. *Fertil. Steril*; 58:674-679.
- Pedersen, E.B. (1988):** Autonomic nervous system and vascular reactivity in normal and hypertensive pregnancies. In Rubin P.C. ed. *Handbook of hypertension*, Vol.10.

Amsterdam Elsevier, p. 155-157.

- Pedersen, A.K., Fitzgerald, G.A. (1984):** Dose-related kinetics of aspirin presystemic acetylation of platelet cyclooxygenase. *N Eng J Med*, 311:1206-11.
- Pedersen, E.G., Johannesen, P., Kristensen, S., et al., (1984):** Calcium, parathyroid hormone and calcitonin in normal pregnancy and preeclampsia. *Gynecol Obstet, Invest*, 18:156-64.
- Pekonen, F., Rasi, V., Ammala, M., Viinikka, L. and Ylikorkala, O. (1986):** Platelet function and coagulation in normal and pre-eclamptic pregnancy. *Thromb, Res*; 43:553-560.
- Pelkey TJ Frierson HF, Mills SF, and Stoler MH (1999):** Detection of the α -subunit of inhibin in trophoblastic neoplasia. *Hum. Pathol*; 30(1):26-31.
- Perry, I.J. and Beevers, D.G. (1994):** The definition of preeclampsia. *Br J Obstet Gynecol*, 101:587-591.
- Perry, I. J., Wilkinson, L.S., Shinton, R.A. and Beevers, D.G. (1991):** Conflicting views on the measurement of blood pressure in pregnancy. *Br J Obstet Gynaecol*, 98:241-243.
- Petraglia F, Luisi S, Benedetto C, Zonca M, Florio P, Casarosa F, Volpe A, Bernasconi S, and Genazzani (1997).** Changes of dimeric inhibin-B levels in maternal serum throughout healthy gestation and in women with gestational diseases. *J. Clin. Endocrinol. Metab*; 82(9):2991 -2995.
- Petraglia F, Florio P, Benedetto C, Gallo C, Woods RJ, Genazzani AR, Lowry P (1996):** High levels of corticotropin-releasing factor (CRF) are inversely correlated with low levels of maternal CRF-binding protein in pregnant women with pregnancy-induced hypertension. *J Clin Endocrinol Metab* 81:852.
- Petraglia, F.; Anesclzi, M.M. and Calza, L (1993):** Inhibin and activin in human fetal membranes: evidence for a local effect on prostaglandin release. *J. Clin. Endocrinol. Metab*; 77:542-548.
- Petraglia F, Calza L, and Garuti GC (1990).** Presence and synthesis of 7 inhibin subunits in human decidua. *J. Clin. Endocrinol. Metab*; 71 :487-492.
- Petraglia F, Fawcenko P, Lim ATW, Rivier U, and Vale W (1987):** Localization, secretion and action of inhibin in human placenta. *Science*; 237:187-189.
- Phalen, J.P. (1977):** Enhanced prediction of pregnancy-induced hypertension by combining supine pressor test with mean arterial blood pressure of middle trimester. *Am J Obstet Gynecol*, 129:397-400.

- Phipps MG, Hogan JW, Pdpert JF, LambertMesserlian G.M, Canick LA, and Seifer DB (2000):** Progesterone, inhibin and hCG multiple marker strategy to differentiate viable from non-viable pregnancies. *Obstet. Gynecol*; 95(2):227-231.
- Powers RW. Evans RW. Ness RB. Cromhlehholme WR. Roberts JM (2000):** Homocysteine is increased in preeclampsia but not in gestational hypertension. *Int J Gynecol Investig (Ahstr 375)*. 7(1) (Suppl).
- Puls LE, Hamous J, Morrow MS, Schmneyer A, MacLaughlin DI, and C'astracane D.** Recurrent ovarian sex cord tumor with annular tubules: tumor marker and chemotherapy experience. *Gynecol. Oncol* 1994; 54:396-401.
- Qu, J. and Thomas, K. (1993):** Regulation of inhibin secretion in human placental cell culture by epidermal growth factor, TGFs and activin. *J.Gun. Endocrinol. Metab*; 77(4): 925-930.
- Qu J, and Thomas K (1992):** Changes in bioactive and immunoactive inhibin levels around human labor. *1. Clin. Endocrinol Metab*; 74:1290-1295.
- Raab W, Schroeder O, Wagner R, Gigg W (1956):** Vascular reactivity and electrolytes in normal and toxemic pregnancy. *Int Chin Endocrinol* 16:1196.
- Ramsay JE, Jamieson N, Greer IA, Sattar N. (2003):** Paradoxical elevation in adiponectin concentrations in women with preeclampsia. *Hypertension*. Nov;42(5):891-4. Epub Sep 29. (Abstract)
- Rappaport VI, Hirata O, Kim Yap H, Jordan SC (1990):** Anti-vascular endothelial cell antibodies in severe preeclampsia. *Am J Obstet Gynecol* 162::138.
- Raty R, Koskinen P, Alanen A, Irjala K, Matinlauri I, Ekblad U (1999):** Prediction of preeclampsia with maternal mid-trimester total renin, inhibin A, AFP and free b-hCG levels. *Prenat Diagn*;19:122— 127.
- Reddi K, Wickings EJ, McNeilly AS, Baird DI, and Hillier SG (1990).** Circulating bioactive follicle stimulating hormone and immunoreactive inhibin during the normal human menstrual cycle. *Hum. Endocrinol*; 33:547-557.
- Redman CWG. Sacks OP. Sargent IL (1999):** Preeclampsia: An excessive maternal inflammatory response to pregnancy. *Am J Obstet Gynecol* 180:499.
- Redman, C.W.G., Bonnar, J. and Beilin, L.J. (1978):** Early platelet consumption in preeclampsia. *Br Med J*, 1:467-469.
- Redman, C.W.G., Allington Bolton, F.G. and Stirrat, G.M. (1977a):** Plasma and Beta. Thromboglobulin in preeclampsia, *Lancet* ii:248.
- Redman, C.W.G., Denson, K.W.E., Beilin, L.J., Bolton, F.G. and Stirrat, G.M.**

- (1977b): Factor VIII consumption in preeclampsia. *Lancet*, ii:1249-1252.
- Reith, A., Booth, N.A., Moore, N.R., Cruickshank, D.J. and Bennett, B. (1993):** Plasminogen activator inhibitors. *Obstet Gynecol*, 100:370-74.
- Regan CU, Fitzgerald DI (1997):** Altered isoprostane to prostacyclin ratio in preeclampsia. *Am J Obstet Gynecol* 176:510.
- Reith, A., Booth, N.A., Moore, N.R., Cruickshank, D.J. and Bennett, B. (1993):** Plasminogen activator inhibitors. *Obstet Gynecol*, 100:370-74.
- Renier MA, Vereecken A, and Buytaert P (1998):** Inhibins, activins and follistatins: a review of complex regulators of the reproductive system. *Eur. J Contracept. Reprod. Health Care*; 3(3): 129-135.
- Risbridger GP, Robertson DM, and DeKrester DM.** Current perspectives of inhibin biology. *Acta Endocrinol. (Copenh.)* 1990; 122:673-682.
- Roberts VJ, Barth S, El-Roeiy A, and Yen SSC (1993):** Expression of inhibin and activin subunits and follistatin mRNA and proteins in ovarian follicles and the corpus luteum during the human menstrual cycle. *J. Clin. Endocrinol. Metab*; 77:1402-1410.
- Roberts V, Meunier H, Sawchenko PE, and Vale W (1989a):** Differential production and regulation of inhibin subunits in rat testicular cell types. *Endocrinology*; 125:235-239.
- Roberts JM, Taylor RN, Musci TI, Rodgers GM, Hubel CA, McLaughlin MK (1989b):** Preeclampsia: An endothelial cell disorder. *Am J Obstet Gynecol* 161:1200.
- Robertson DM, Cahir N, Findlay JK, (1997):** The biological and immunological characteristics of inhibin A and B forms in human follicular fluid and plasma. *J Clin Endocrinol Metab*; 82:889-896.
- Robertson DM, Buiger HG, Sullivan J, Gahir N, Groome N, Poncelet F, Franchi, P, Woodruff I, and Matlier J (1996).** Biological and immunological characterization of inhibin forms in human plasma. *J. Clin. Endocrinol. Metab* 1996; 81(2):669-676.
- Robertson DM, Sullivan J, Watson AI, and Cahir N (1995):** Inhibin forms in human plasma. *J. Endocrinol*; 144:261-269.
- Robertson DM, Devos FL, Foulds LM, McLachlan RL, Burger HG, Morgan FJ, Hearn MTW, and DeKrester DM (1986):** Isolation of a 31K-dalton form of inhibin from bovine follicular fluid. *Mol Cell Endocrinol*; 44:271-277.
- Robertson DM, Hayward S, and Iren DC (1988):** Radioimmunoassay of rat serum inhibin: changes after gonadectomy. *Molecular and Cellular Endocrinology*; 58:1-8.
- Robertson DM, Foulds LM, and Leversha L. (1985):** Isolation of inhibin from bovine

follicular fluid. *Biochem. Biophys. Res Commun*; 126:220- 226.

Robson SC: Magnesium sulphate (1996): The timing of reckoning. *Br J Obstet Gynaecol* 103:99. 1996

Rochat, R., Koonin, L., Atrash, H. and Jewett, J. (1988): Mortality in the United States: Report from the Maternal Mortality Collaborative. *Obstet Gynecol*, 72:91-7.

Rodriguez, M.H., Maskaki, D.I., Mestman, J., Kumar, D. and Rude, R. (1988) Calcium/creatinine ratio and microalbuminuria in the prediction of pre-eclampsia. *Am, J, Obstet Gynecol*; 159:1452-5.

Rogers BB, Bloom SU, Ueveno KJ (1999): Atherosclerosis revisited: Current concepts on the pathophysiology of implantation site disorders. *Obstet Gynecol Surv* 54:189.

Rombauts L, Verltieven G, Meuleman C, Koninx PR, Poncelet E, and Francitmont P (1996): Dimeric inhibin-A and α -subunit immunoreactive material in maternal serum during spontaneous and in vitro fertilization pregnancies. *J. Clin. Endocrinol. Metab*; 81(3): 985-989.

Romero, R., Lockwood, C., Oyarzun, F. and Hobins, J.C. (1988): Toxemia New concepts in old disease. *Semin Perinatol*, 12:302-323.

Roth, G.J., and Majerus, P.W. (1975): The mechanism of aspirin on human platelets I acetylation of a particulate fraction protein. *J Chin Invest*, 56:624-32.

Roseff SJ, Bangah ML, Kettel LM, Vale W Rivier J, Burger HG, and Yen SSC (1989). Dynamic changes in circulating inhibin levels during the luteal-follicular transition of the human menstrual cycle. *J. Clin. Endocrinol. Metab*; 69:1033-1039.

Saftlas, A.F., Olson, D.R., Franks, A.L., Atrash, ILK. and Pokras, R. (1990): Epidemiology of pre-eclampsia and eclampsia in the United States, 1979-1986. *Am J Obstet Gynecol*, 163:460-5.

Sanchez-Ramas, L.; Stephan Sandroni, Frank J. Andres and Kaunitz, A.M. (1991a): Calcium excretion in preeclampsia. I. *Obstet. Gynecol.*, 77:510-5 13.

Sanchez-Ramas, L.; David, C. Jones and Mark T. Cullen (1991b): urinary calcium as an early marker for preeclampsia.. *Obstet. Gynecol.*, 77:685-688.

Santoro N, Adel T, and Skurnick JH (1999): Decreased inhibin tone and increased activin-A secretion characterize reproductive aging in women. *Fertil. Steril*; 71(4): 658-662.

Schiff E, Friedman SA, Stampfer M, Kao U, Barrett PH, Sibai BM (1996): Dietary consumption and plasma concentrations of vitamin F in pregnancies complicated by

- preeclampsia. *Am J Obstet Gynecol* 175:1024.
- Schiff E, Ben-Baruch O, Peleg E, Rosenthal T, Alcalay M, Devir M, Mashiach S (1992):** Immunoreactive circulating endo thein-1 in normal and hypertensive pregnancies. *Am J Obstet Gynecol* 166:624.
- Schneyer AL, Sluss PM, Whitconib RW, Martin KA, Sprengel R, and Crowley WF (1991):** Precursors of c~inhibin modulate FSH receptor binding and biological activity. *Endocrinology*; 129: 1987-1999.
- Schneyer AL, Mason AJ, Burton LE Zeinger IR, and crowley WE. (1990):** Immunoreactive inhibin subunit in human serum.: implications for radioimmunoassay. *J. Clin. Endocrinol. Metab* 70: 1208-1212.
- Scott RS, Burger HG, and Quigg H. (1980):** A simple and rapid in-vitro bioassay for inhibin. *Endocrinology*; 107:1536-1542.
- Scully RE (1977):** Sex cord stromal tumors. In: *Pathology of the Female Genital Tract*. A. Blaustein, Ed., Springer-Verlag, New York; pp:505- 526.
- Sebire NJ, Roberts L, Noble P, Wallace E, Nicolaides KH (2000):** Raised maternal serum inhibin A concentration at 10 to 14 weeks of gestation is associated with pre-eclampsia. *Br J Obstet Gynaecol*;107: 795 -797.
- Seifer DB, Lambert MG, Hogan 3M. Gardiner AC, Blazar AS, and Berk CA (1997):** Day-3 serum inhibin-B is predictive of assisted reproductive technologies outcome. *Fertil. Steril*; 67(1): 110- 114.
- Seifer DB, Gardiner AC, Mess erlian GL, and Schneyer AL(1996).** Differential secretion of dimeric inhibin in cultured luteinized granulosa cells as a function of ovarian reserve. *J. Clin. Endocrinol. Metab*; 81(2)236-739.
- Sibai BM, Caritis S, Hauth J; National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. (2003):** What we have learned about preeclampsia. *Semin Perinatol*. Jun;27(3):239-46.
- Sibai BM. Hauth I. Caritis S. Uindheimer MD. MacPherson C, Klebanoff M. VanDorsten JP. Landon M. Miodovnik M. Paul R. Meis P. Thurnau O. Domhrowski NI. Roberts I, McNelhis D (2000).** The National Institute of Child Health and Human Development Network of Maternal—Fetal Medicine Units: Hypertensive disorders in twin versus singleton gestations. *Am J Obstet Oynecol* 182:938.
- Silver HM, Seebeck M, Carlson R (1999):** Comparison of total blood volume in normal, preeclamptic, and non-proteinuria gestationah hypertensive pregnancy by simultaneous measurement of red blood cell and plasma volumes. *Am J Obstet Gynecol* 179:87.
- Sibai BM, Uindheimer M~ t-lauth I. Caritis 5, VanDorston P. Klebanoff M,**

- MacPherson C, Landon M. Miodovnik M. Paul R. Meis P. Domhrowski M (1998):** Risk factors for preeclampsia, abruptio placentae, and adverse neonatal outcome among women with chronic hypertension. *N Engl J Med* 339:667.
- Sibai BM, Ewell M, Levine RI, Kleharioff MA, Estechitz I, Catalano PM, Goldenberg RU, Ioffe O (1997):** Risk factors associated with preeclampsia in healthy nulliparous women. *Am J Obstet Gynecol* 177:1003.
- Sibai BM, Barton JR, Akl S, Sarinoglu C, Mercer BM (1992):** A randomized prospective comparison of nifedipine and bed rest alone in the management of preeclampsia remote from term. *Am J Obstet Gynecol* 167:879.
- Sibai, B.M.; Mabie, W.C.; Shamsa, F. and Villar, M.A. (1990a):** A comparison of no medication versus methyldopa or labetalol in chronic hypertension during pregnancy. *Am J Obstet Gynecol*; 162: 960.
- Sibai, B.M., Villar, M.A. and Mabie, B.C. (1990b):** Acute renal failure in hypertensive disorders of pregnancy: Pregnancy outcome and remote prognosis in thirty-one consecutive cases. *Am J Obstet Gynecol*; 162:777-783.
- Sibai, B.M. (1988):** Pitfalls in diagnosis and management of preeclampsia. *Am J Obstet Gynecol*, 159:1-5.
- Sibai BM, Mahie BC, Harvey Ci, Gonzalez AR (1987):** Pulmonary edema in severe preeclampsia—eclampsia: Analysis of thirty-seven consecutive cases. *Am J Obstet Gynecol* 156:1174.
- Sibai BM, El-Nazer A, Gonzalez-Ruiz A (1986):** Severe preeclampsia—eclampsia in young primigravid women: Subsequent pregnancy outcome and remote prognosis. *Am J Obstet Gynecol* 155:1011.
- Silverman LA, and Gitelman, SF (1996).** Immunoreactive inhibin, Mullerian inhibitory substance and activin as biochemical markers for juvenile granulosa cell tumors. *J. Pediatr*; 129 (6): 918-921.
- Simmons LA, Hennessy A, Gillin AG, Jeremy RW (2000):** Uteroplacental blood flow and placental vascular endothelial growth factor in normotensive and preeclamptic pregnancy. *Br J Obstet Gynaecol* 107:678.
- Spitz B, Magness RR, Cox SM, Brown CEL, Rosenfeld CR, Gant NF (1988):** Low-dose aspirin. I. Effect on angiotensin II pressor responses and blood prostaglandin concentrations in pregnant women sensitive to angiotensin II. *Am J Obstet Gynecol* 159:1035.
- Staff AC, Ranheim T, Khoury I, Henriksen T (1999):** Increased contents of phospholipids, cholesterol, and lipid peroxides in decidua basalis in women with preeclampsia. *Am J Obstet Gynecol* 180:587.

- Stamilio DM, Sehder VIM, Morgan MA, Propert K, Macones GA (2000):** An antenatal clinical and biochemical markers predict the development of severe preeclampsia? *Am J Obstet Gynecol* 182:589.
- Steinberger A, and Steinberger E (1976):** Secretion of an FSH-inhibiting factor by cultured Sertoli cells. *Endocrinology*; 99:918-921.
- Strevens H, Wide-Svensson D, Grubb A, Hansen A, Horn T, Ingemarsson I, Larsen S, Nyengaard JR, Torffvit O, Willner J, Olsen S. (2003):** Serum cystatin C reflects glomerular endotheliosis in normal, hypertensive and pre-eclamptic pregnancies. *BJOG*. Sep; 110(9):825-30.
- Strickland DM, Guzik DS, Cox K, Gant NF, Rosenfeld CR (1986):** The relationship between abortion in the first pregnancy and the development of pregnancy-induced hypertension in the subsequent pregnancy. *Am J Obstet Gynecol*, 154:146.
- Stone, J.L., Lockwood, C.J., Berkowitz, G.S., Alvarez, M., Laapinski, R. and Berkowitz, R.L. (1994):** Risk factors for severe preeclampsia. *Obstet Gynecol*, 83(3): 357-61.
- Stroud, C.S., Campbell, S.K., Albano, J.D.M. and Millar, J.G.B. (1990):** A simple automated method of urinary kallikrein estimation. VII th World congress of Hypertension in Pregnancy, Perugia, Italy. Scientific Program, p.306.
- Sugino K, Kurosaiva N, and Nakamura T (1993):** Molecular heterogeneity of follistatin, an activin binding protein. *J.Biol. Chem*; 268: 15579-15587.
- Talledo OE, Chesley LC, Zuspan FP (1968):** Renin-angiotensin system in normal and toxemic pregnancies, 3. Differential sensitivity to angiotensin II and norepinephrine in toxemia of pregnancy. *Am J Obstet Gynecol* 100:218.
- Taufield, P., Ales, K, Resnick, L., Druzin, M., Gertner, J. (1987):** Hypocalciuria in preeclampsia. *N, Eng J~Med*.316:715- 718.
- Thurnau, G.R., Dyer, A., Depp, 1.1.1. OR, and Martin, A.O. (1983):** The development of a profile scoring system for early identification and severity assessment of pregnancy induced hypertension.
- Tjoa ML, Mulders MA, van Vugt JM, Blankenstein MA, Oudejans CB, van Wijk IJ. (2003):** Plasma hepatocyte growth factor as a marker for small-for-gestational age fetuses. *Eur J Obstet Gynecol Reprod Biol*. Sep 10;110(1):20-5
- Toner JP, Philput CB, Jones GS, and Muasher SJ. (1991):** Basal FSH level is better predictor in IVF performance than age. *Fertil. Steril*; 63:1-11.
- Tovanabutra 5, Illingworth PJ, Ledger WL, Glasier AF, and Baird DT (1993).** the

relation between peripheral immunoreactive inhibin hCG, estradiol and progesterone during human pregnancy. *Gun. Endocrinol*; 38: 101-107.

Trudigner, B.J. and Giles, W.B. Cook, C.M. (1985): Uteroplacental blood flow velocity time waveforms in normal and complicated pregnancy. *Br J Obstet Gynecol* 92:39.

Trupin LS, Simon LP, Eskenazi B (1996): Change in paternity: A risk factor for preeclampsia in multiparas. *Epidemiology* 7:240.

Tunbridge, R.D.G. and Donnai, P. (1983): Pregnancy-associated hypertension, a comparison of its prediction by roll-over test and plasma noradrenaline measurement in 100 primigravidae. *Br J Obstet, Gynecol* 90:1027-1032.

Turn T, Framaa M, Hildan K, and Ritvos O. The tissue distribution of activin beta-A and beta-B subunits and follistatin mRNA suggests multiple sites of action for the activin-follistatin system during human development. *J. Gun. Endocrinol. Metab* 1994; 78:1521-1524.

Underwood, P., Kesler, K., O'Lane, J. and Callagan, D.A. (1967): Parental smoking empirically related to pregnancy outcome. *Obstet Gynecol*, 29:1-8.

Vale, W; River, G. and Vaughan, J (1994). The role of inhibin and activin In: *The Physiology of Reproduction*. E. Knobil and J.D. Neill, Eds., New York: Raven Press; pp: 1861- 1878.

Vale W, and Vaughan JM (1993): macroglobulin is a binding protein of ~ inhibin and activin. *Endocrinology* 1993; 132: 2038-2050.

Vekemans, M.; Delvoye, P.; L 'Hermite, Al. and Rob, C (1977): Serum prolactin levels during the menstrual cycle. *J. Gun. Endocrinol. Metab* 1977; 44:989.

Ventura SJ, Martin IA, Curtin SC, Mathews TJ, Park MM: Births (1998): Final data for 1998. *National Vital Statistics Reports*; Vol 48, No. 3. Hyattsville, MD, National Center Health Statistics.

Volhard F (1918): Die doppelseitigen haematogenen Nierenerkrankungen. Berlin. Springer, English abstract.

Wakatsuki M, Shintani Y, Abe M, Liu ZH, Shitsukawa K, and Saito S (1996). Immunoradiometric assay for follistatin serum immunoreactive follistatin levels in normal adult and pregnant women. *J. Clin. Endocrinol. Metab*; 81(2):630-634.

Walker II (2000): Pre-eclampsia. *Lancet* 356:1260.

Wallace EM, Gow SM, and Wu EGW (1993): Comparison between testosterone enanthate-induced zoospermia and ligozoospermia in a male contraceptive study I: plasma luteinizing hormone, follicle stimulating hormone, testosterone, estradiol and

inhibin concentrations. *J. Clin. Endocrinol. Metab*; 77:290-293.

Wallenburg, H.C., Dekker, G.A., Makovitz, J.W. and Rotmans, P. (1986): Low dose aspirin prevents pregnancy-induced hypertension and pre-eclampsia in angiotensin sensitive primigravidae. *Lancet*, 1:1-3.

Wallace EM, Grant VE, Sivanston IA, and Grooine NP (1995): Evaluation of maternal serum dimeric inhibin-A as a first trimester marker of Down's syndrome. *Prenat. Diagnosis*; 15:359-362

Wallace EM, Riley SC, Crossley JA, (1997): Dimeric inhibins in amniotic fluid, maternal serum and fetal serum in human pregnancy. *J Clin Endocrinol Metab*;82:218-222.

Walsh SC (1994): Lipid peroxidation in pregnancy. *Hypertension* 13:1.

Walsh, S.W. (1985): Preeclampsia: An imbalance in placental prostacyclin and thromboxane production. *Am J Obstet Gynecol*, 152:335.

Wang, Q. F.; Farnwomth, P. G.; Findlay, J.K. and Burger, H.G (1988): The effect of pure 31 K-daltons bovine inhibin on the specific binding of GnRH to rat anterior pituitary cells in culture. *Endocrinology*; 123:216.

Ward K, Hata A, Jeunemaitre X, Helm C L, Nainikawa C, Farrington PF. Suzumori K, Tomoda S. Berrebi S, Sasaki M, Corvol P, Lifton RP, Lalouel JM (1993): A molecular variant of angiotensinogen associated with preeclampsia (comment). *Nat Genet* 59: 1-7.

Watt HC, Wald NJ, and Huttlv, WJ (1998): The pattern of maternal serum inhibin-A concentrations in the second trimester of pregnancy. *Prenat. Diag*; 18(8):846-848.

Weerasekera DS, Peiris H. (2003): The significance of serum uric acid, creatinine and urinary microprotein levels in predicting pre-eclampsia. *J Obstet Gynaecol. Jan*;23(1):17-9.

Weiner, C.P. and Brandt, J. (1982): Plasma antithrombin III activity in normal pregnancy. *obstet Gynecol*, 56:601- 603.

Weiner CP, Thompson LP, Liu KZ, Herrig IE (1992): Endotheliumderived relaxing factor and indomethacin-sensitive contracting factor alter arterial contractile responses to thromboxane during pregnancy. *Am I Obstet Gynecol* 166:1171.

Weiner C, Martinez E, Zhu LK, Ghodsi A, Chestnut D (1989): In vitro release of endothelium-derived relaxing factor by acetylcholine is increased during the guinea pig pregnancy. *Am I Obstet Gynecol* 161:1599.

- Welt CK, McNichol DJ, Taylor AE, and Hall IE (1999):** Female reproductive aging is marked by decreased secretion of dimeric inhibin. *J. Clin. Endocrinol. Metab*; 84(1):105-111.
- Welt CL, Martin LA, Taylor AE, Lambert GM, Growley WE, Smith JA, Schoenfeld DA, and Hall JE. (1997):** Frequency modulation of follicle-stimulating hormone during the luteal follicular transition:evidence for FSH control of inhibin-B in normal women. *J. Clin. Endocrinol. Metab*; 82(8):2645-2652.
- Wenstrom KD, Chu DC, O'Brien J, and Boots L (1998):** Maternal serum alpha-fetoprotein and dimeric inhibin-A detect aneuploidies other than Down's syndrome. *Am. J. Obstet. Gynecol*; 179(4):966-970.
- Woodruff TK, and Mather JP (1995):** Inhibin, activin and the female reproductive axis. *Annu. Rev. Physiol*; 57:219-244.
- Woodruff TK, Lyon RJ, Hansen SE, Rice GC, and Mather JP (1990).** Inhibin and activin locally regulate rat ovarian folliculogenesis. *Endocrinology*; 127: 3196-3205.
- Ying SY (1988):** Inhibins, activins and follistatins: Gonadal proteins modulating the secretion of follicle-stimulating hormone. *Endocrine Rev*; 9:267-293.
- Ylikorkala, O. Viinikka, L. and Lehtovirta, P. (1987):** Effect of nicotine on fetal prostacyclin and thromboxane in humans. *Obstet Gynecol*, 66:102-5.
- Ylikorkala, O. and Makela, U.M. (1985):** Prostacyclin and thromboxane in gynecology and obstetrics. *Obstet Gynecol*, 152:318-29.
- Zeeman GG, Alexander JM, McIntire DD, Devaraj S, Leveno KJ. (2003):** Homocysteine plasma concentration levels for the prediction of preeclampsia in women with chronic hypertension. *Am J Obstet Gynecol*. Aug;189(2):574-6.
- Zelinski Wooten MB, Hutchinson JS, Chandrasekhar YA, Wolf DP, and Stouffer RL (1992):** administration of human luteinizing hormone after follicular development: further titration of LH surge requirements for ovulatory changes in primate follicles. *J. Clin. Endocrinol. Metab*; 75:502.
- Zhang I, Klebanoff MA, Levine RI, Pun M, Moyer P (1999):** The puzzling association between smoking and hypertension during pregnancy. *Am J Obstet Gynecol* 181:1407.