

REFERENCES

REFERENCES

Abraham L., Oosting J. and Hart M.(1987):

Bell's palsy, factors affecting the prognosis in 200 patients with reference to hypertension and diabetes mellitus
Clin. Otolaryngol., 1987 12; 349- 355.

Adkins W.Y. and Osguthorpe J.D.(1991):

Management of trauma of the facial nerve. Otolaryngol.Clin. North America, 24(3):357-612.

Adour K.K.(1980):

Corticosteroid therapy for acute facial paralysis. Quted from controversies in Otolaryngology. Edited By J. Snow-Philadelphia; W.B. Saunders pp.395-400

Adour K.K.(1991):

Medical management of idiopathic bell's palsy otolaryngol. Clin. North America 24:(3):663-673.

Adour K.K and Wingered J.(1974):

Idiopathic facial paralysis; factors affecting severity and outcome
Neurol. 24, 1112-1116.

Adour K.K., Byl F.M., Hilsinger R.L. and Khan Z.M.(1978)

The true nature of Bell's palsy. Analysis of 1000 consecutive patients. Laryngoscope, 88; 787-801.

Adour K.K., Hilsinger R.L. and Callan E.J.(1985):

Facial paralysis and Bell's palsy; a protocol for differential diagnosis.
Am. J. Otol. Supp 1.68-73

Aoagagi M., Koike Y. and Ichige A.(1988):

Results of facial nerve decompression. Acta Otolaryngol. Stockh. (supp 446); 101-105

Apostalides C. and Carr, M(2000):

Assesment and management of facial palsy; comparison of Nova scotia family physicians and otolaryngologists (midline) J. Otolaryngol. 2000 feb. 29(1): 17-22.

Ardic F.N., Ardic F., Topulglu J., Oncels., Uguz M.Z. and Tpologu D(1997):

Electroneurgraphy in the late period of Bell's palsy. Acta Otolaryngol. May; 117(3) :325-8.

Arriaga M.A., Luxford W.M., Atkins J.S. and Kwarter J.A. (1993):

Predicting facial nerve outcome after acoustic neuroma surgery .Otolaryngol. Head and Neck surgery, 108: 220-224

Barker A., Jalinous R. and Freestone I.(1985):

Non- invasive magnetic stimulation of the human motor cortex Lancet, 1:1106-1107.

Barry S. and May M. (1997):

Disorders of facial nerve, Quoted from scott-Brown's Otolaryngology 6th edition otology 3-24-1138. Butterworth-Heinemann Oxford-Ox2 8DP

Baxter A.(1971):

Dehiscence of the Fallopian canal. J. of Laryngol. Otol., 85 587-594

Benedet E., TalmiY- and Kronenberg J.(1998):

Preoperative Electroneurography in parotid surgery. Head and Neck.20(2): 124-31.

Blumenthal F. and May M.(1986):

Electrodiagnosis in facial nerve lesions Quoted from "The Facial Nerve"- MayM.4:28-33. New Yourk, Theme-Stratton.

Boongird E. (1978):d 18

Electrophysiologic findings and prognosis in Bell's palsy
Muscle-Nerve, 1461-466.

Brandenburg N.A. and Annegers J.F.(1993):

Incidence and risk factors for Bell's palsy.
Neuroepidemiology, 12(6): 313- 325.

Cajal S.R.Y.(1894)

La fine structures des centres nerveux. Proceedings of the
Royal Society, Series 55: 444- 468.

Quoted from Scott Brown's Otorhinolaryngology the edition 3-
24:1137.

Butterworth- Heinemann- Oxford- OX2 8dP

Campbell M.J. and Machin D.(1993):

Medical statistics. A common sense approach. 2nd Ed. John
Wiley & sons New- York; Chichester; Brisbane ; Toronto &
Singapore.

Cawthorne I. And Hyenes D.R. (1956):

Facial palsy. British medical Journal 197-200

**Clark J.R., Carison R.D., Sasaki C.I., Pachner A.R. and Steere
A.C.(1985):**

Facial Paralysis in Lyme Disease. Laryngoscope, 95: 1341-
1344.

Cracco R.(1987):

Evaluation of conduction in central motor pathways.
Neurosurgery 20: 199- 203.

Cramer H.B. and Kartush J.M. (1991):

Testing facial nerve function.
Otorhinolaryngology Clinics North America: 34(555- 570).

Crosby E.C., Humphrey. I. And Lauer E.W. (1962):

Correlative Anatomy of the nervous system.
New Yourk. Macmillan.

Daniel W.W. (1987):

Biostatistics, A foundation for analysis in the health science. 4th
Ed.. John Wiley & sons. New-York; Chichester; Brisbane;
Toronto & Singapore.

Danielides V., Skevas A., Kastantodakis I., Anastasopolos D.(1996)

Role of MST in facial nerve injury as prognostic tool
Eur. Arch Otorhnolaryngol. 210: 34 –37.

Delgado.T.E., Buchiet W.A. and Rosen holta H.R. (1979)

Intraoperative monitoring of facial muscle evoked responses
obtained by intracranial stimulation of the facial nerve. A more
accurate technique for facial nerve dissection.
Neurosurgery, 4:418 –421.

De Med iros J., Nobreg A J., de- Andrade and Novo N.(1996):

Facial Electroneurography in Bell's palsy variability in the
early stage and comparison between interpretation methods
(midline) Arq-Neuropiquitar, 54(3) :397-401.

Desphande A.D.(1990):

Recurrent Bell's palsy in pregnancy
J. Laryngol. Otol. 104: 713-714.

Devrise P.P.(1977):

Prednisone in idiopathic facial paralysis (Bell's palsy).
ORL Journal of Otorhinolaryngol. and related specialities, 39:
257-271.

Devriese P.P, Schumacher I., Schede A. and Houtkoopey J.(1990):

Incidence, prognosis and recovery of Bell's palsy. Clin
Otolaryngol, 15:15-27.

Dickins J.R., Smith J.I. and Graham S.S.(1988)

Herpes Zoster Oticus. Treatment with intravenous acyclovir
Laryngoscope . 98:776- 779.

Dobie R.A.(1993):

Tests of facial nerve function. In Otolaryngology Head and
Neck surgery. Cumming's C.W. 7-149 :2718-2725

Dorsey D.L. and Cammann W.R.(1993):

Obestitric anaesthesia in patients with Bell's palsy
Anaesthesia –Analg, 77:81-3.

Durcan D.J., Shea J.J. and Sluckx J:P(1967):

Bifurcation of the facial nerve. Arch Otolaryngol 86:619-631.

Eicher S.A., Coker N.J., Alford B.R.(1990):

A comparative study of the fallopian canal at the meatal
foramen and labyrinthine segment in young children and adults.
Arch Otolaryngol. Head an Neck surgery ,116:1030-1035.

Engstrom M., Thumas K.A., Naeser P. and Jonson L.(1993):

Facial nerve inhancement in Bell's palsy. Arch Otolaryngol.
Head and Neck surgery, 119:221-5.

Esslen E.(1977):

Investigations on the localization and pathogenesis of
meatolabyrinthine palsies. In The Acute facial Paralysis B
New Yourk-springer verlage

Esslen E.(1983):

Electrodiagnosis of facial palsy. In "surgery of the facial Nerve.
Urban and Schwartzenburg, Munich P.P 49-51.

Estreme S.A., Haghighis., levy W.J.(1988):

Motor Evoked potentials of facial musculature in dogs.
Laryngoscope, 98 :1012-1015.

Finestone A.J. and Byers K.(1994):

Acute facial paralysis ; is it-a stroke or something else.
Geriatrics 49(4):50-2.

Fisch U.(1969):

Disorders of skull base region. Ed. by Hamberg. C. and Wersol
J. Stockholm Alm quist and wike sell

Fisch U.(1981):

Surgery of Bell's palsy.
Arch. Otolaryngol. 107:1-11

Fisch U(1984):

Prognostic value of electrical tests in acute facial paralysis
Am J. Otol, 5:494-8.

Fisch U (1986):

Extracranial surgery for facial hypokinesis in the facial nerve .
May M.(ed.), Thieme strotton, New yourk Ch. 27:509-23.

Fisch U and Esslen .E(1972):

Total intratemporal exposure of the facial nerve.
Arch. Otolaryngol., 95 :335.

Fowler A.P.(1963):

The pathological finding in a case of facial paralysis.
Transactions of the American Academy of ophthalmology and
Otolaryngology 67, 187-107

Fragon H and Loh G.M(1989):

Facial nerve function, Assess ement and evaluation.
Arch Otolaryngol .3 :113-113

Ganang W.F.(1999):

Physiology of nerve and muscle Review of medical physiology.
19th ed PP. 35-45
Long medical publication-Los Altos-California

Gates G.A.(1987):

Facial paralysis
Otolaryngol. Clinic. North America.

20(1):113-131.

Gates G.A.(1993)

Nerve excitability testing
Laryngoscope 103: 379 – 385.

Gilliat R.W. and Taylor J.C.(1959):

Electrical changes following section of the facial nerve.
Proc Royal Soc Med, 52:1080.

Glasscock M.E., Wet R.J. , Jackson C.G. and Dickens J.R. (1979):

Rehabilitation of the face following traumatic injury of the facial nerve.
Laryngoscope, 89:1389 –1404.

Glocker F.X., Magistris M.R., Rosler KM. And Hess C.W.(1994):

Magnetic stimulation and electrical stimulation of the facial nerve in Bell's palsy.
Electroencephalog. Clin- Neurophys., 93 (2):113-20

Guerrier Y(1977):

Surgical anatomy particularly vascular supply of the facial nerve.
Arch otolaryngol. 4:117-120.

Gutnick H. N., Kelleher M.J., and Prass R.L.(1990):

A model of waveform reliability in facial nerve EnoG.
Otolaryngol- Head and Neck Surg, 103:344.

Halvorson D.J., Coker N.J. and wan y-Bennett L.T.(1993):

Histologic correlation of degenerating facial nerve with EnoG.
Laryngoscope, 103:178-184.

Hammerschlage P.E., Konie N., Osten W.F.(1987):

Intraoperative real time monitoring of brain stem facial evoked response.
Laryngoscope, 97:57-62.

Harner S.G., Daube J.R., Abersold M.J. and Beatty C.W.(1987):

Improved preservation of facial nerve function with use of electrical monitoring during removal of acoustic neuroma.
Mayo Clinic Proceedings, 62:92- 102

Quoted from scott-Brown's otolaryngol 6th ed, 3-24 Buter
warth -Heinmann-Oxford-Ox2 8DP

Hauser W.A., Karene W.E. and Annis J.(1971):

Incidence and prognosis of Bell's palsy in the population of
Rochester, Minesota. Mayo Clinic proceedings 46, 258-264.

Quoted from scott- Brown's otolaryngol. 6th ed.

3-24 Buter worth Heinemann Oxford -Ox2 8DP

Hialsinger R.L., Adour K.K. and Doty H.E.(1975):

Idiopathic facial paralysis, pregnancy and the menstrual cycle
Annals of Otorhinolaryngol, 84:433-442.

Hoffman W.Y.(1992):

Reanimation of the Paralyzed face.

Otolaryngol. Clinics North America 25(3: 649-667).

Hosomi Y.(1994):

Histopathological study of the recovery process in exprimental
facial nerve palsy induced by freezing

Nippin -Jibbin KoKa- Gakka-Kawtho, 97(9): 1683-
91(midline).

House.J.W(1983):

Facial nerve grading system.

Laryngoscope. 93: 1056-1069.

House W.A., Karene W. E. and Annis J(1971):

Facial nerve grading system.

Laryngoscope 93: 1056- 1069.

House. J.W. and Brackmann.D.E.(1985):

Facial nerve grading system

Otolaryngol. Head Neck Surg. 93(2:146-147).

Hunt J.R.(1907):

On herpotic inflammations of the geniculate ganglion. A new
syndrome and its complications. Journal of Nervous and
Mental Diseases 34:73-96.

Quoted from Scott Brown's Otolaryngol. 6th ed . 3-24
Buterwoa rth - Heinemann- Oxford- Ox2 8DP

Joachim H.Z., Bialik V. and Eliachar I. (1980):

Early diagnosis in Bell's palsy, Laryngoscope 90: 1705 –1708.

Kandel E.R.(1977):

Neuronal plasticity and the modification of behaviour. In Hand book of physiology .Edited by American Physiology Society. Section 1 Vol.1 part 2. Quoted from scott Broun's Otolaryngol- 6th ed. 3-24 Buter worth Heinmnn- Oxford- Ox2 –8DP.

Kandler R.H. and Jarratt J.A.(1991):

Magnetic stimulation in Bell's palsy
J. Neurol. Neurosurg. Psyc, 54(11):1022.

Kartush J.M., Garcia P. and Telian S.A.(1987):

The source of far field antidromic facial nerve potentials. Am. J. Otolaryngol. 8: (99-204).

Kashima K. (1990):

Antidromically evoked facial nerve response in Guinea pigs, a basis for clinical application in patents with facial plealsy
Eur –Arch. Otorh nolaryngol., 247 :151-155.

Kohsyu H. (1995):

Clinical investigation of transcranial magnetic stimulation of the facial nerve.
Nippon –Jibiin Koka-Gakkai-Kaiho(medline)
98(9):1416 –25.

Langwor thy E.P. and Tavener D. (1963):

The prognosis in facial palsy. Brain, 86 :465.

Lee S.H., Ito J. and Yamamoto E.(1991):

A Horseradish peroxidase study of the fiber orientation in the facial nerve .
Euro Arch. Otolaryngol, 248;(6): 366-9

Litton W. Spae Krause C. and Jhon W.(1969):

Relationship of the facial canal to the annular sulcus:
Laryngoscope, 79 :1584- 1604.

Luetge C. and Whittaker C.K.(1991):

The Benefits of VII-VII neuroanastomosis in acoustic tumour surgery.

Laryngoscope, 101: 127-1275.

Mattox D.E.(1993):

Management of the facial nerve disorders.

Otolaryngol Clin North America, 24:3.

May M.(1986a):

Anatomy of the facial nerve. In "The Facial Nerve" May M.(ed) Theme Stratton. New york Ch.2 PP 15-62.

May M.(1986b):

Microanatomy and pathophysiology of the facial nerve, in "The Facial Nerve" May M.(ed) Thiem Stratton. New York Ch.3 PP.63-74.

May M.(1987):

Disorders of the facial nerve.

Scott Brown's Otolaryngol. 6th ed. Ch. 24 vol.3 PP 560.

Buterworth – Heinmann. Oxford-OX2 8DP.

May M. (1990):

Science is for the moment and truth is for ever

Otolaryngol. Head Neck 102:1-2.

May M. and Hardin (1977):

Interpretation of neurologic finding in facial palsy transaction of the American Academy of Ophthalmology and Otorhinolaryngol. 84 :710-722.

Outoted from Scott Brown's Otolaryngol. 6th ed.

3-24 Buterworth Hienmunn oxford OX2 8DP.

May M. and Klein S.R.(1991):

Differential diagnosis of facial nerve palsy Otolaryngol. Clin. North America

24 (3):613-645.

May M. and Shambough G.E.(1991):

Facial nerve paralysis. In Otolaryngology.

Paparella M.M. 4th ed. Vol. II Ch.14.

May M., Harvey J.E. and Marcovitz W.E.(1971):

The prognostic accuracy of maximum stimulation test compared to that of the nerve excitability test in Bell's palsy
Laryngoscope, 81:931-938.

May M., Blumenthal F. and Taylor F.H.(1981):

Bell's Palsy ; surgery based up on prognostic indicator and results. Laryngoscope 91:2092 :2103.

May M., Blumenthal F. and Klein S.R.(1983):

Acute Bell's palsy; Prognostic value of EMG, MST and other electrical tests. AMJ Otol. 5:1-7.

May M., Klein S.R. and Taylor F.H.(1985):

Idiopathic facial palsy ; Natural history defies steroids or surgical treatment.
Laryngoscope, 95: 406- 409.

March M. A. and Coker N.J.(1991):

Surgical decompression of idiopathic facial palsy
Otolaryngol. Clin North America
24(3:675 -689)

Martin H. and Helsper J.I.(1957):

Spontaneous return of function following surgical section or excision of the seventh cranial nerve in parotid tumour surgery
Annals of surgery, 146: 715 -727.
Quoted from Scott Brown's Otolaryngology 3-24 Buterwarth – Heinmann-Oxford –OX2 8DP.

MC Ouarrie H. and Grafestien D.(1973):

Different criteria of facial nerve injury. Acta Otolaryngol, 2:
416-420.

Metson R.(1988):

Facial nerve evoked potentials in cat. Otolaryngol. Head and Neck surgery, 98:130-137.

Meitson R., Rebez E. West. C and thornaton A.(1991):

Magnetic stimulation of facial nerve.
Laryngosocpe, 101:25-30.

Meyer B. U., Brutton T.C. and Beneck R.(1989):

Investigations of unilateral facial weakness.
Magnetic stimulation of the proximal facial nerve and of the
face associated motor cortex.
J. Neurol, 236 :102.

Meyerhoff J.W.(1993):

Indications and techniques of facial nerve monitoring during
surgery.
Head and Neck surgery, 108; 201- 204.

Morgan M. and Nathwani D.(1992):

Facial palsy and infection.
Clinical infectious diseases, 14 : 203-271.

Mori H., Horicuchi M., Kithura K, Tanaka T (1974)

Bell's palsy and glucose metabolism.
Practical Otolaryngol 67, 955-960.

**Nadol J.B., Chiony G.M., Ojemon R.G., Mckenna M.J., Martuza R.L.
and Montgomery W.W.(1992):**

Preservation of hearing and facial nerve function in resection of
acoustic neuroma. Laryngoscope, 102:1153- 1158.

Nager G.T. and Proctor B.(1991):

Anatomic variations and anomalies involving the facial canal.
Otolaryngol. Clin. North America, 24(3):531 –553.

Patten J.P.(1987):

Neuroanatomy and applied neurophysiology. Scott –Brown's
Otolaryngol. Volume 3 Chapter 24-6th ed.
Butterworth –Heinmann Oxford- OX2 8DP.

Peitersen E.(1977):

Spontaneous course of Bell's palsy. Acta Otolaryngol, 5:97-99.

Peitersen E.(1982):

The natural history of Bell's palsy.
American Journal of Otology, 4:107-111.

Quoted from Scott Brown's Otolaryngol. 6th ed 3-24 Butterworth-Hiemann Oxford- OX2- 8DP.

Perez C., Santos S., Soto A., Frade C., Castro C. and Laballa I.(1999):
Prognostic value of stapedius reflex and Electroneurography in facial paralysis. Acta Otolaryngol. Esp.; 50(3): 187- 90 (midline).

Proctor B.(1991):
The Anatomy of the facial nerve.
Otolaryngol. Clin North America, 24(3):449-504.

Qui. W., Yins. S., Stucker F.G. and Hoasjoe D.K.(1997):
Neurophysiological evaluation of acute facial paralysis in children.
Int. J. Pediatric Otolaryngol. 39 (3): 223-36(midline)

Rimplianen I., Karna P., Lariene J. and Hakkenen V.(1992):
On Bell's palsy Acta otolaryngol. "Stockh.", 112 :311-316.

Romanes G.J. (1979):
Cunningham's Manual Practical Anatomy. The English Language Book Society. Fourteenth edition 3:187.

Rosenthal U.(1983):
Auditory brainstem response abnormalities in Bell's palsy
Otolaryngol. Head Neck surgery, 91:412-414.

Rosler K.M., Hess C.W. and Schmid U.D.(1989):
Investigation of facial motor pathways by electrical and magnetic stimulation.
J. Neurol. Neurosurg. Psyc, 52:1149-1156.

Saito H., Takida I. And kishimoto S.(1992 a):
Facial nerve to facial canal cross sectional area ratio in children.
Laryngoscope 102:(10)1172-1176.

Saito H., Taked T., Hay; I. and Nakteni H.(1992b):
Facial nerve stimulation through Stensen's duct.
Acta Otolaryngol.(Stockh), 112:102-1016.

Schaitkin B. and May M.(1993):

Evaluation and management of facial nerve disorders.
Current opinion in Otolaryngology, Head and Neck surgery
1:79-83.

Schriefer I.N.(1988):

Evaluation of facial nerve condition by transcranial magnetic stimulation.
J. of Neur., Neurosurg., Psyc, 51:60-66.

Schuknecht h.F. and Shinozaki-Hori N.(1985):

Patterns of deg. of the facial nerve.
Am J. Otol.(suppl.), 47: 54.

Schwarting S., Schrocler M., Stener t.E. and Goebel H.tt.(1984)

Morphology of denervated human facial muscles.
Otolaryngol. 46:248-256.

Seddon H.J.(1943):

Three types of nerve injury . Brain 66:237-288.
Quoted from Scott Brown's Otolaryngol 6th ed.
Buter worth- Hienmann Oxford -Ox2 8DP.

Seki Y., Krainl., Yamada I. And Kimura J.(1990):

Transcranial magnetic stimulation of the facial nerve.
Neurosurgery 26: 286- 290.

Sillmann J.S., Niparko J.K., Lee S.S. and Kileny P.R.(1992):

Prognostic value of evoked and standard Electromyography in acute facial paralysis Otolaryngol., Head and Neck surgery
107:3.

Skevas A.J., Danielides V.G. and Assima koupolos D.A.(1990):

Role of facial nerve latency test in the prognosis of Bell's palsy. Laryngoscope 100: 1083-1083.

Stephanian E., Sekhar I.N., Janeka I.P. and Hissch B.(1992):

Facial nerve repair by interposition nerve graft results in 22 patients. Neurosurgery 31:73:77.

Sunderland, S. (1986):

Nerve and nerve injuries.

Forth ed. London Churchill/ Livingston pp. 88-89, 96- 97, 135.

Quoted from Scott Brown's Otolaryngol. 6th ed. 3-24 Butterworth
-Hienmanm- Oxford -OX2 8DP

Sunderland, S.(1991):

Nerve and nerve injuries.

2nd ed. London Churchill/ Livingston pp. 88-89, 96- 97, 135.

Quoted from Scott Brown's Otolaryngol. 6th ed. 3-24 Butterworth
-Hienmanm- Oxford -OX2 8DP

Thurner K.H. and Schrott- Ficher A.(1993):

A quantitative study of nerve fibres in the human facial nerve.

Eur-Arch-Otolaryngol- 250 :(3) 1616-171.

Van Buskirk C.(1995):

The seventh nerve complex- Journal of comparative neurology
82:303-333.

Quoted from Scott Brown's Otolaryngol. 3-24(38) 6th ed.

Butterworth- Hienmann Oxford- OX2 8DP

Vassillios G., Dani lides V., Antonios M., and Skevas I(1991):

Value of facial nerve latency test in prognosis of childhood
Nerv. Syst. 8:126-128.

Watanabe Y., Ishikawa M., Shojakka H., and Mizukoshik.(1993):

Facial nerve palsy; a complication of parotid surgery.

Act Otolaryngol.(suppl), 504; 137-139.

Waylonis G.W. and Johonson E.W.(1964):

Facial nerve conduction delay. Arch. Phys. Med. Rehab.
45:539.

Welling D.B., Glasscock M.E. and Gantz B.J.(1992):

Avulsion of the anomalous facial nerve at stapedectomy.

Laryngoscope, 102:729-733.

Wetmore S.G.(1991):

Surgical landmarks for the facial nerve Otolaryngol. Clin.
North America, 24(3): 505- 530.

Windmill I.M.; Martinez S.H; Shields C.B.(1989):

Magnetically evoked facial nerve potential Otolaryngol. Head and Neck surg. 100:345-347.

Wright A.(1987):

Anatomy and ultrastructure of the human ear.

Scott Brow's otolaryngology 6th ed volume 3- Chapter (1)
PP.530-535. Oxford-OX2 8DP.

Zhigang C., Guangyan Y., Daquan M., Jing T., Zhaohui Y., and Xiaoming Zhang (1998):

Experimental studies on traumatic facial nerve injury.

The Journal of Laryngology and Otology March 1998,
Vol 112, 243-247.