

References

/ Adams D H., Hubscher J., Shaw R., et al., (1989) :

Intercellular adhesion molecule-1 on liver allografts during rejection. *lancet*, 2: 1122.

Adams D H., Hunschen SG., Shaw J., et al., (1991) :

Increased expression of intercellular adhesion molecule-1 on bile ducts in primary biliary cirrhosis and primary sclerosing cholangitis. *Hepatology*, 14(3) : 426.

Adams D H., Mainolfi E., Burra P., et al., (1992) :

Detection of cICAM-1 in chronic liver diseases. *Hepatology*, 16: 810.

Adams D H., Mainolfi E., Burra P., et al., (1993) :

detection of circulating intercellular adhesion molecule-1 after liver transplantation, evidence of local release within the liver during graft rejection. *Transplantation*, 55:83.

Alexander G., and Williams R., (1983) :

The management of hepatitis. *Postgraduate doctor. Middle East.*, sept. 4 : 442.

Alexander F., Rossi RL., O'Bryan M., et al., (1984):

Biliary carcinoma, a review of 109 cases. Am . J. surg, 147.
503.

Almeida JD .,Rubenstein D., and Scott EJ., (1971) :

New antigen-antibody system in Australia-antigen positive hepatitis. Lancet, 11 : 1225.

Alter M., Coleman P., and Alexander W.J., (1989) :

Importance of heterosexual activity in the transmission of hepatitis B and non-A, non-B hepatitis.
JAMA, 262 : 1201 .

Altomonte M., Collizi F.,Esposito G., et al., (1992) :

Circulating ICAM-1 as a marker of disease progression in cutaneous melanoma. N. Engl. J. Med, 327 : 959.

Aoyagi Y., Saitoh A., Suzuki Y., et al., (1991) :

The usefulness of simultaneous determinations of glucosylation and fucosylation indices of alpha fetoprotein in differential diagnosis of neoplastic diseases of the liver . Cancer, 67 : 2390.

Arnaout MA., (1990) :

Leukocyte adhesion molecule deficiency :its structural basis, pathophysiology and implication for modulating the inflammatory response. Immunol Rev, 114:145.

Becker J C., Dummer R., Hartmann AA., et al., (1991)

Shedding of ICAM-1 from human melanoma cell lines induced by IFN- γ and TNF- α : functional consequences on cell mediated cytotoxicity . J. Immunol, 147 : 4398.

Becker J C., Dummer R., Schwinn A., et al., (1992) :

Circulating ICAM-1 in melanoma patients : Induction by interleukin-2 therapy. J. Immunol, 12 :147.

Benjamine E., (1982) :

Relife of obstructive jaundice leads to decrease of alkaline phosphatase enzyme level. The obstruction of biliary tree, in clinical results. J. Interv. Radiol, 5:287.

Blumgart L H., (1990) :

Tumours of the gall bladder and extrahepatic bile ducts. In Maringot's abdominal operations. 19th edition Vol. II. Schwarts and H.Ellis (eds). Prentice- Hall international Inc. London, Toronto, Tokyo and New jersy.pp. 1130-1143.

Boyd AW., Wawryk SO., Burns GF., et al., (1988) :

Intercellular adhesion molecule - 1 has a central role in cell-cell mediated immune mechanisms. Proc. Natl. Acad. Sci. U.S.A., 85 : 3094 .

Brustein M., Kraal G., Mebius RE., et al., (1992) :

Identification of a soluble form of the ligand for the lymphocyte homing receptor. J. Ex. Med, 176 : 1415.

Burra P., Hubscher sc, Shaw J., et al., (1990) :

Is the ICAM/LFA-1 pathway of leukocyte adhesion involved in the hepatocyte damage of alcoholic hepatitis ? Hepatology , 12:394 (abstr).

Cainzos M., Potel ., Puente JL., (1988) :

Allergy in jaundiced patients. Br. J. Surg, 75:142.

Chen C J., Chang Y C., Tsai S F., et al., (1988) :

Epidemiological studies of multiple risk factors in hepatocellular carcinoma in Taiwan. In Vyas GN., Dienstag J.L., and Hoofnagle JH (eds): Viral hepatitis and liver disease, New York.pp 723.

Christie A B., (1987) :

Viral hepatitis, enterovirus infection (Coxsackie viruses and echoviruses) in Manning L. (ed) : Infectious diseases, epidemiology and clinical practice, Churchill livingstone, Edinburgh, 655.

Cobb R R., Dubins J S., Warner J., et al., (1992) :

Functional expression of soluble ICAM-1 by baculovirus infected cells. Biochem. Biophys. Res. Commun. 185:1022.

Conn HO.,(1982) :

Cirrhosis, in : Conn HO.,(ed.) diseases of liver., 5th ed. lippincott company- philadelphia. 23 : 847

De Fougères AR., and Springer T A., (1992) :

Intercellular adhesion molecule-3 (ICAM-3) a third adhesion counter receptor for LFA-1 on resting lymphocytes. J. Exp. Med, 175 : 185.

De Fougères AR., Stacker S A., Schwarting., et al., (1991) :

Characterization of ICAM-2, and evidence for third counter receptor for LFA-1. J. Exp. Med, 174 : 253.

Deinhardt F., (1980) :

Predictive value of markers of hepatitis B virus infection. J. Infect. Dis, 4 : 141.

Deinhardt F., and Gust L D., (1982) :

Viral hepatitis. bull. WHO, 60:661.

Deusch J., and Ahnen DJ., (1993) :

Primary hepatocellular carcinoma, in : Gitnieck G.,(ed):
Current hepatology. Mosby St. louis, Baltimore, Boston,
Chicago, London, Philadelphia, Sydney, Toronto. Vol: 13.
pp 197.

Dougherty G J., Murdoch S., Hogg N., (1987) :

The function of human intercellular adhesion molecule-1
(ICAM-1) in the generation of an immune response. Eur. J.
Immunol, 18 : 35.

Dustin ML., Garcia-Aguilar J., Hibbs ML., et al., (1989) :

Structure and regulation of the leukocyte adhesion receptor
LFA-1 and its counter receptors, ICAM-1 and ICAM-2.
Cold Spring Harbor symp quant. Biol, 54 :753.

Dustin ML., Rothlein R., Bhan A K., et al., (1986) :

Induction by IL-1 and interferon, Tissue distribution, Biochemistry, and Function of a natural adherence molecule (ICAM-1). J. Immunol, 137:245.

Dustin ML, Springer TA., (1989) :

T cell receptors cross linking transiently stimulates adhesiveness through LFA-1. Nature, 341 : 619.

Dustin ML., Staunton DE., Springer T A., (1988) :

Supergene Families meet in the immune system. Immunol Today, 9 : 213.

Edelman G.M., (1985) :

Cell adhesion and the molecular process of morphogenesis. Annu. Rev. Biochem, 54 : 135.

Edmondson HA., and Craig J R., (1987) :

Neoplasms of the liver. In : Schiff, L and Schiff, E.R. (eds): Diseases of the liver. 6th ed., ch. 33-1109. Philadelphia.

Esteban J., Gonzalez A., and Hernandez J., (1990) :

Evaluation of antibodies to hepatitis C virus in a study of transfusion associated hepatitis. N. Engl. J. Med, 323 : 1107.

Faull R J., and Russ G R., (1989) :

Tubular expression of intercellular adhesion molecule-1 during renal allograft rejection. Transplantation, 48 :226.

Fawcett J., Holness CLL., Needham LA., et al.,(1992) :

Molecular cloning of ICAM-3, a third ligand for LFA-1 constitutively expressed on resting leukocytes. Nature, 360 : 481.

Fried M., Shindo M., and Fong T., (1992) :

Absence of hepatitis C viral RNA from saliva and semen of patients with chronic hepatitis C. Gastroentrology, 102 : 1306.

Gearing A J H., and Newman W., (1993) :

Circulating adhesion molecules in disease. Immunol. Today, 14:506 .

Gholson E., and Bconr., (1993) :

Acute viral hepatitis, in: Manming S., (ed) :Essentials of clinical hepatology. Mosby year book Inc. Chicago, London, Philadelphia, pp. 55.

Gianotti F., (1973) :

Papular acrodermatitis of childhood : An Australian antigen disease. Arch. Dis. Child, 49 : 794.

Giavazzi, R., Chirivi R G., Arofalo A., et al., (1991) :

Soluble ICAM-1 is released by melanoma cells and is associated with tumour growth in nud mice. Cancer Res, 51: 5003.

Govindarrajan S., Chin K P., Redeker A G., et al., (1984a) :

Fulminant B viral hepatitis, role of delta agent. Gastroenterol, 86:1417 .

Govindarrajan S., Lim B., and Peters R L., (1984 b) :

Immunohistochemical localization of delta antigen associated with HBV in liver biopsy sections. Histopathology, 8 : 63.

Greve J M., Davis G., Meyer A M., et al., (1989) :

The major human rhinovirus receptor is ICAM-1. *Cell*, 56:839.

Griffiths C E M., Voorhees J J., and Nickoloff B J., (1989) :

Characterization of ICAM-1 and HLA-DR expression in normal and inflamed skin : modulation by recombinant gamma interferon and tumour necrosis factor. *J. Am. Acad. Dermatol*, 20 : 617.

Hamazaki K., Gochi A., Shimamura H., et al., (1996) :

Serum levels of circulating intercellular adhesion molecule-1 in hepatocellular carcinoma. *Hepatogastroenterology*, 43(7) : 229.

Harris CC., and Sun T., (1984) :

Multifactorial etiology of human liver cancer. *Carcinogenesis*, 6 : 697.

Hart I R., and Saini A., (1992) :

Biology of tumor metastasis. *Lancet*, 339 : 1453.

Hemler ME., (1988) :

Adhesive protein receptors on hematopoietic cells. *Immunol. Today*, 9 : 109.

Hollinger F B., Kan NC., and Oefinger P E., (1983) :

Post transfusion hepatitis type A. J. AM. Med. Assoc, 250:
2313.

***Holzmann B., Lechmann J M., Ziegler-Heitbrock H W., et al.,
(1988) :***

Glycoprotein P 3.53, associated with tumor progression in
malignant melanoma, is a novel leukocyte activation
antigen. Int.J. Cancer, 41 : 542.

Hoofnagle J H ., (1981) :

Serologic markers of HBV infection. Ann. Rev. Med, 142 :
400.

Hoofnagle JH., and Schafer D F., (1986) :

Serologic markers of HBV infection. Semin. Liver. Dis, 6.

Horiike N, Onji M, Kumon I., (1993) :

Intercellular adhesion molecule-1 expression on hepatocyte
membrane of patients with chronic hepatitis B and C.liver,
13 : 10.

Hutchinson WL., Johnson PJ., DU MQ., et al., (1991) :

Serum and tissue α -L-Fucosidase activity in the pre-clinical and clinical stages of hepatocellular carcinoma. *clin. Sci*, 81(2) : 177 ,

Hynes RO., (1987) :

Integrins : a family of cell surface receptors. *Cell*, 48: 249.

Hynes RO, (1992) :

Integrins : Versatility, modulation, and signaling in cell adhesion. *Cell*, 69: 11.

Hyodo I., Jinno K., Tanimizu M., et al., (1993) :

Detection of circulating intercellular adhesion molecule-1 in hepatocellular carcinoma. *Int. J. Cancer*, 55(5) : 775.

Ideo G., Bellati G., Pedraglio E., (1990) :

Intrafamilial transmission of hepatitis C., *Virus. Lancet*, 553 : 353

Ingoldby CJH., and Bumgart LH., (1984):

The Significance of preoperative endotoxaemia in obstructive jaundice. *Br.J. Surg*, 10 : 300.

Jaffe S H., Froedlander D R., Tastuzaki F., et al., (1990) :

Differential effects of the cytoplasmic domains of cell adhesion molecules on cell aggregation and sorting out. Proc. Natl. Acad. Sci. USA, 87 : 3587.

Jandrassik L., et al., (1938) :

Measurment of serum bilirubin. Biochem . Z. 297-81.

Jurgen C., Reinhold E., Brocker E., et al.,(1993) :

Soluble intercellular adhesion molecule-1 inhibits MHC-restricted specific T-cell-tumour interaction. Journal of Immunology, 151 : 7224.

Jutila MA., Rott L., Berg E L., et al., (1989) :

Function and regulation of the neutrophil MEL-14 antigen in vivo: comparison with LFA-1 and Mac-1. J. Immunol, 143:3318.

Kamby C., Dirksen H., and Vejborg M., (1987) :

Incidence and aetiologic aspects of the occurrence of liver metastasis in recurrent breast cancer. cancer, 59 : 1524.

Kelsey P B., and Dienstag J L., (1984) :

Viral hepatitis,in : Kass EH and platt R., Current therapy in infectious diseases. B.C. Decker Inc, pp. 117.

Kind PR N., King EG., (1954) :

Measurment of serum Alkaline phosphatase .J. Clin. Path,
7:322.

Kishimoto T K., Jutila M A., Berg EL., et al., (1989):

Neutrophil Mac-1 and Mel-14 adhesion proteins regulated
by chemotactic factors. Science, 245:1238.

Kishimoto T K., Larson R S., Corbi A L., et al., (1989) :

The leukocyte integrins : LFA-1, Mac-1 and P 150, 95.
Adv. Immunol, 46 :149.

Kithier K., (1980) :

Tumour associated antigns, in : Sonnenwirth AC., Jarett
L.,(ed) : Gradwohl's clinical laboratory methods and
diagnosis, 8 th edition. Mosby. St. Louis, Toronto, London,
PP. 1307.

Kohl S., Springer TA., schmalstieg FC., et al., (1984) :

Defective natural killer cytotoxicity and polymorphonuclear
leukocyte antibody dependent cellular cytotoxicity in
patients with LFA-1/OKM-1 deficiency. J. Immunol, 133 :
2972.

Krensky AM., Robbins E., Springer TA., et al., (1984) :

LFA-3 antigens are involved in CTL-target conjugation. J. Immunol, 132:218.

Kryger P., Aldershvile J., Mathresen L R., et al., (1982) :

Copenhagen hepatitis acute programme, acute type B. hepatitis among HBsAg negative patients detected by anti HBc IgM. Hepatology, 2 : 50.

Kurzinger K., Reynolds T., Germain RN., et al., (1981) :

A novel lymphocyte function associated antigen (LFA-1) : cellular distribution, quantitative expression, and structure. J. immunol, 127 : 596.

Larson R S., Springer T A., (1990) :

Structure and function of leukocyte integrins. Immunol. Rev, 114:181.

Laurent F., Vitvitski L J., Berby F., et al., (1992) :

Importance of PCR in the diagnosis of hepatitis C. Rev. Fr. Transfus. Hemobiol, 25 : 216.

Lawrence W., and Bond ., (1992) :

Tumour markers, in : Richard CT, Albert B., David CH., et al., clinical laboratory medicine. St. Louis, Baltimore, Boston, Chicago, London, Philadelphia, Sydney, Toronto. Mosby, pp : 322.

Leaper Al (1990) :

Tumours of small intestine and doudenum, in : Schwartz S.I., and Ellis H., (eds): Maringots abdominal operations, 9th edition vol. I, Prentic Hall inter national Inc, pp. 640.

Lemon S M., Gates N L., Simms T E., et al., (1981) :

IgM antibody to HBc Ag : as a diagnostic parameter of acute infection with HBV. J. Infect. Dis, 143 : 803.

Liebman HA., Furie BC., Tong MJ., et al., (1984) :

Des - carboxy (abnormal) prothrombin as a serum marker of hepatocellular carcinoma. N. Eng. J. Med. 310:1258.

Locarnini S A., Coulepis A G., Gust I D., et al., (1982) :

Biophysical and biochemical characterization of HAV. Intervirology, 18 : 107.

Macswheen RNM., (1980) :

Pathology of viral hepatitis and its sequelae. Clinical gastroenterology, 9 : 23.

Magnius L O., and Espmark J A., (1972) :

New specificities in Australia antigen positive sera distinct from Le Bouvier subdeterminants. J. Immunol, 109:1017.

Makgoba K., Sander ME., and Shaw S. (1989) :

The CD2-LFA-3 and LFA-1-ICAM-1 pathways-relevance to T cell recognition. Immunol Today, 10:417.

Mandi Y., Nagy I., (1995) :

Circulating ICAM-1 in alcoholic liver cirrhosis Int. Arch. Allergy. Immunol. 106 (3) : 302.

Marlin S D., and Springer T A., (1987) :

Purified intercellular adhesion molecule-1 is a ligand for lymphocyte function associated antigen-1. Cell, 51: 813.

Marlin SD., Staunton D E., Springer T A., et al., (1990) :

A soluble form of intercellular adhesion molecule-1 inhibits rhinovirus infection. Nature, 344 : 70.

Martz E., (1987) :

LFA-1 and other accessory molecules functioning in adhesion of T and B lymphocytes. Hum immunol, 18 : 3.

Matsumoto Y., Suzuki T., Asada I., et al., (1982) :

Clinical classification of hepatoma in Japan according to serial changes in serum α - fetoprotein level. Cancer, 49: 354.

McEver R P.,(1995) :

Cell adhesion, in : Hoffman R., Benzjr EJ., Shattil SJ., et al (editors) : Hematology : basic principles and practice. churchill livingstons. New York, Edinburgh, London. 1995 : pp: 53.

Mcpherson GAD., Benjamine IS., Hodgson HJF., et al., (1982) :

Preoperative percutaneous transhepatic biliary drainage : The results of a controlled trial. Br. J. Surg, 71 : 371.

Melnick J L., (1982) :

Classification of HBV as enterovirus type 72 and of HBV as hepadnavirus type 1. Intervirology, 18 : 105.

Miller LJ, Schwarting R, and Springer TA, (1986) :

Regulated expression of the Mac-1, LFA-1, p 150-95 glycoprotein family during leukocyte differentiation. *J. Immunol*, 137 : 2891.

Miller R. and Purcell R. (1990) :

Hepatitis C virus shares amino acid sequence similarity with pestiviruses and flaviviruses as well as members of two plant virus super-groups. *Proc. Natl . Acad.Sci. (U.S.A)*, 87 : 2057.

Mosley J W., (1975) :

The epidemiology of viral hepatitis, an overview. *Am. J. Med. Soci*, 270: 253.

Neimark S., and Rogers AI., (1982) :

Viral hepatitis : a review . *Am. J. Gastroenterology*, 77 : 431.

Nickoloff BJ., Griffiths CEM., Barker JNWN., (1990):

The role of adhesion molecules, chemotactic factors and cytokines in inflammatory and neoplastic skin disease. *J. invest. Dermatol*, (94) : 151.

Nouri Aria KT., Koskinas J., Tibbs CT., et al., (1995) :

Serum intercellular adhesion molecule-1 levels in chronic hepatitis C: association with disease activity and response to interferon .Gut, 36:599.

Okner R K., (1985) :

Hepatitis in : Wyngaarden J.B., and Smith L.H., Cecil textbook of medicine, 17th edition W.B. saunders company, p:778.

Okner RK., (1988) :

Approaches to the diagnosis of jaundice, in : Wyngaarden J.B., Smith L.H (eds), Cecile textbook of medicine 18th edition.W.B. saunder company international edition philadelphia. pp : 817.

Okuda K., (1976) :

Clinical aspects of hepatocellular carcinoma, in : Okuda K., and Peters RL.,(eds),analysis of 154 cases in hepatocellular carcinoma. New York, John Wiley & Sons. pp 1210.

Okuda K., and Okuda H., (1991) :

Primary liver carcinoma, in : Neil M., Jean-Pierre B., Johannes B., Mario R., and Jaun R., (eds). Oxford textbook of clinical hepatology . Oxford , NewYork, Tokyo, Oxoford university, pp 1019 .

Oppenheimer Maks N., Davis LS., Bogue DT., et al., (1991) :

Differential utilization of ICAM-1 and VCAM-1 during the adhesion and transendothelial migration of human T lymphocytes. J. immunol, 147 : 2913.

Pellegrini CA., (1987) :

Pathophysiology of obstructive jaundice, in : Way L., and Pellegrinic CA., (eds). surgery of the gall bladder and bile ducts. 2nd edition, W. S Saunders Company. pp 103.

Pellegrini CA., and Thomas MJ., (1984) :

Recurrent biliary strictures patterns of recurrence and outcome of surgical therapy. Am.J. surg, 147:175.

Pirisi M., Fallet E., Fabris C et al., (1994) :

Circulating intercellular adhesion molecule-1 concentration in liver disease. Am. J. clin. pathol, 102 : 600.

Pober JS., Gimbrone MA., and La pierre LA., (1986) :

Overlapping patterns of activation of human endothelial cells by interleukin-1, tumour necrosis factor, and immune interferone. J. immunol, 137 : 1893.

Popper H., (1977) :

Pathologic aspects of cirrhosis . Am. J. Path, 87 : 228 .

Reesink H., and Vander poel C., (1989) :

Blood transfusion and hepatitis still at heart. Bull. J. 58 : 16.

Rizzetto M. (1983) :

The Delta agent. hepatology 3 : 729.

Rizzetto M., Hoyer B H., Purcell R H., et al., (1984) :

HDV infection, in : Vyas G.N. Deintage J.L. and Hoofanagle J.H. (eds). viral hepatitis and liver disease. Grune stration Inc, Orlando, Florida. pp. 371.

Rizzetto M., Shih J., and Gerin J L., (1980) :

Epidemiology of hepatitis B virus-associated delta agent. lancet, 146: 1215.

Robbins S L., Cotran R S., and Kumar V., (1984) :

Liver and biliary tract, in Robbins SL., Cotran RS., and Kumar V., (eds) : Pathologic basis of disease. 3 rd ed : Philadelphia : W.B. Saunders Co, pp : 884.

Robertson BH., Khanna B., Nainan OV. et al., (1991) :

Epidemiologic patterns of wild - type hepatitis A virus determined by genetic variation. J. infect. disease, 163 : 286.

Robinson W S., (1980) :

Genetic variation among hepatitis B and related viruses. Ann. N.Y. Acad. Sci, 354:371.

Rodkey FL., (1964) :

Measurment of serum albumin. Clin chem, 10 : 606.

Roitt IM., Brostoff J., and Male DK., (1993) :

Cell migration and inflammation : in : Roit IM., Brostoff J., Male DK (editors) : Immunology, Mosby. Hong Kong. 1993 Chap 13. pp: 13.3 .

Rooney MT., and Henry J B., (1991) :

Molecular markers of malignant neoplasms in :Henry JB.,
(ed.) :Clinical diagnosis & management by laboratory
methods.18th ed. Philadelphia, London,Toronto, Monterial,
Sydney, Tokyo, W.B. saunders company, pp : 285.

Rothelin R., Czaikowski M., O'Neil MM., et al., (1988) :

Induction of intercellular adhesion molecule 1-on primary
and continuous cell lines by pro-inflammatory cytokines. J.
Immunol, 141 : 1665.

Rothelin R., Dustin ML., Marlin SD., et al., (1986) :

A human intercellular adhesion molecule-1 (ICAM-1)
distinct from LFA-1. J. Immunol , 137 : 1270.

Rothlein R., Mainolfi E A., Czajkowski M., et al., (1991) :

A form of circulating ICAM-1 in human serum. J Immunol,
147 : 3788.

Rouslahti E., and pierschbacher M D., (1986) :

Arg - Gly- Asp : a versatile cell recognition signal . Cell,
44: 517.

Runyon BA., (1993) :

Bacterial infections in patients with cirrhosis. J. Hepatol, 8:271.

Sabbour M S., (1978) :

Viral hepatitis, in : Sabbour MS., (ed) : Infectious diseases, Dar El-Maaref. Egypt. pp. 172.

Saiki P., Scharf S., Falonna F., et al., (1985) :

Enzymatic amplification of beta-globin genomic sequences & restriction site analysis for diagnosis of sickle cell anaemia. Science, 230 pp :1350.

Sanchez- Madrid F J., Nagy E., Robbins P., et al., (1983) :

A human leukocyte differentiation antigen family with distinct alpha subunits and a common beta subunit : The lymphocyte function associated antigen (LFA-1), the C3bi complement receptor (OKM-1/Mac-1) and the p150, 95 molecule. J exp. Med, 158 : 1785 .

Sato H., Niessler A K., Liu Z Z., et al., (1994) :

Proinflammatory cytokines and endotoxin stimulate ICAM-1 gene expression and secretion by normal human hepatocytes. Immunology, 82 :571.

Sawabu N., and Haltosi N., (1987) :

Serological markers in hepatocellular carcinoma, in: Okuda-Ishak(ed) neoplasms of the liver. New York, Springer Verlag, pp 227.

Scharschmidt BF, and Way LW., (1987) :

The jaundiced patient differential diagnosis and clinical approach, in : Way L W., and Pellegrini CA.,(eds) surgery of the gall bladder and bile ducts. W.B. Saunders Company pp. 215.

Schochetman G., Ou C., Jones W., (1989) :

Polymerase chain reaction. J. Infect Dis, 158 : 1154.

Schwartz SI., (1989) :

Diseases of the extrahepatic biliary tract, in : Schwartz S.I., Sires G.I. and Spences I.C (eds). principles of surgery. 5th edition. McGraw Hill Book Company Vol. II, pp. 1381.

Seth R., Raymond F D., and MakGoba M W., (1991) :

Circulating ICAM-1 isoforms : diagnostic prospects for inflammatory and immune disorders. Lancet, 338: 83.

Shattock A G., and Fielding J F., (1983) :

Exacerbation of chronic liver disease due to HBsAg after delta infection. Br. Med. J, 286-1279.

Sherlock S (1985) :

Jaindice, in : Sherlock S.,(ed.) Diseases of the liver and biliary system, 7th edition, Black well scientific publications. pp 199.

Sherlock S., (1989 a) :

Virus hepatitis, in: Sherlock S.,(ed) : Diseases of the liver and Biliary system. Blackwell scientific publication Oxford-London-Edinburgh, Melbourne. pp 712.

Sherlock S., (1989 b) :

Hepatic tumours in: Sherlock S.(ed) diseases of the liver and Biliary system, 8 the edition. Oxford, London, Edinburgh, Boston, Melbourne, Black well scientific publications, pp 584.

Sherlock S., (1993) :

Diseases of the liver and biliary system. blackwell scientific publication Oxford-London Edenbargh, Melbourne.

Sherlock S., and Dooley J., (1993) :

Chronic hepatitis in Diseases of the liver and biliary system.
9th ed. Blackwell scientific publication. Oxford, London,
Edinburgh, Boston, Melbourne, pp. 293.

Sheu JC., Sung JL., and Chen DS., (1985) :

growth rate of asymptomatic hepatocellular carcinoma and
its clinical implications. Gastroenterology, 89 : 259.

Shevach EM., (1993) :

Accessory Molecules : in Paul W E., (ed.) : Fundamental
immunology. Raven Press. New York. 1993 : Chap 15. pp
: 531.

Shijubo N., Imai K., Aoki S., et al., (1992) :

Circulating ICAM-1 in sera of patients with idiopathic
pulmonary fibrosis. Clin. Exp. Immunol, 89 : 58.

Shimizu Y., Minemura M., Tsukishiro T., et al., (1995) :

Serum concentration of ICAM-1 in patients with
hepatocellular carcinoma is a marker of the disease
progression and prognosis. Hepatology, 22(2) : 525.

Shimizu Y., Van Seventer G A., Horgan K J., et al., (1990) :

Regulated expression and binding of three VLA (beta 1) integrin receptors on T cells. *Nature*, 345:250.

Simmons D., Makgoba M W., and Seed B., (1988) :

ICAM, an adhesion ligand of LFA-1, is homologous to the neural cell adhesion molecule NCAM. *Nature*, 331 : 624

Simpson- KJ., Jones Al., forbes Howie A., et al., (1995) :

Soluble adhesion molecules and interleukin-2 receptor concentration in patients with autoimmune chronic hepatitis. *Eur. J. Gastroenterol-Hepatol*, 7 (5) : 455.

Smedile A., Raimondo G., Fassone M., et al., (1982) :

Radioimmunoassay detection of IgM antibodies to HBV associated delta antigen . *J. of medical virology*, 9 : 131.

Smith C W., Marlin S D., Rothlein R., et al., (1989) :

Cooperative interactions of LFA-1 and Mac -1 with intercellular adhesion molecule-1 in facilitating adherence and transendothelial migration of human neutrophils in vitro. *J. Clin. Invest*, 83 : 2008.

Sobel RA ., Mitchell ME., and Fondren G., (1990) :

Intercellular adhesion molecule-1 (ICAM-1) in cellular immune reactions in human central nervous systems. Am. J. pathol, 136 : 1309.

Soceide O., Gzerniak A., Bradpiece J., et al., (1986) :

Characteristics of fibrolamellar carcinoma : A study of nine cases and view of the literature, Am. J. surg, 151 : 518.

Soulier JP., Gozin D., and Lefrere JJ., (1986) :

A new method to assay des-gamma carboxy prothrombin : results obtained from 85 cases of hepatocellular carcinoma. gastroenterology, 91 : 1258

Springer T.A., (1990) :

Adhesion receptors of the immune system. Nature, 346:425.

Springer TA., Dustin ML., Kishimoto TK ., et al., (1987) :

The lymphocyte function associated antigen LFA-1, CD2, and LFA-3 molecules : cell adhesion receptors of the immune system. Annu. Rev. immunol, 5 : 223.

Staunton D E., Dustin M L., and Springer T A., (1989) :

Functional cloning of ICAM-2, a cell adhesion ligand for LFA-1 homologous to ICAM-1. *Nature*, 339 : 61.

Staunton D E., Marlin S D., Stratowa C., et al., (1988) :

Primary structure of (ICAM-1) demonstrates interaction between members of the immunoglobulin and integrin supergen families. *Cell*, 52 : 925

Suehiro T., Sugimachi K., Matsumata T., et al., (1994) :

protein induced by vitamin K absence or antagonist II as a prognostic marker in hepatocellular carcinoma; comparison with Alpha-fetoprotein. *Cancer* , 72 : 2464.

Szasz G., Persijn JP., (1974) :

Measurment of Gamma-Glutamyl Transferase. *Clin. Chem. Clin Biochem*, 12 : 228

Taketa K., (1990) :

Lectin-reactive profiles of alpha fetoprotein characterizing hepatocellular carcinoma and related conditions. *Gastroenterology*, 99: 508.

Tedder R., Gilson R., and Briggs M., (1991) :

Hepatitis C, virus evidence for sexual transmission. BMJ, 302 : 1299.

Than Than MC., Gee Jo D., and Blumgart LH., (1977) :

Prolyhydroxylase in the skin of patients with obstructive jaundice. Journal of clinical pathology, 30: 1044.

Thomas H C., Potter B J., Elias E., et al., (1979) :

Metabolism of complement in acute type B hepatitis, HBsAg positive glomerulonephritis, polyarthritis nodosum and HBsAg positive and negative chronic active liver disease. Gastroenterology, 76 : 673.

Thomson A W., Sato H S., Nussler AK., et al., (1994) :

Circulating intercellular adhesion molecule-1 (cICAM-1) in autoimmune liver disease and evidence for production of ICAM-1 by cytokine stimulated human hepatocytes. Clin. Exp. Immunol, 95:83.

Tomassini J E., Graham D., Dewitt CM., et al., (1989) :

cDNA cloning reveals that the major group rhinovirus receptor on Hela cells is intercellular adhesion molecule-1. Proc. Nat. Acad. Sci, USA. 86 : 4907

Torii A., Harada A., Nakao A., et al., (1993) :

Expression of ICAM-1 in hepatocellular carcinoma. J. Surg. Oncol, 53(4) : 239.

Tsujisaki M., Imai K., Hirata H., et al., (1991) :

Detection of circulating intercellular adhesion molecule-1 antigen in malignant diseases. Clin. Exp. Immunol, 85(1):3.

Van Nessel C., and Miedema F., (1988) :

Regularity properties of LFA-1 α and B chains in human T lymphocyte activation. Nature, 333 : 850.

Volpes R., Joost J., Valee R J., et al., (1990) :

Immunohistochemical study of adhesion molecules in liver inflammation. Hepatology, 12 : 59.

Wacholtz MC., Patel SS., lipsky P E., (1989) :

Leukocyte function associated antigen-1 is an activation molecule for human cells . J. Exp. Med. 170 : 431.

Walter JB., and Israel Ms., (1979) :

Cell response to injury. general pathology. churchill livingstone 5 th ed. pp 223.

Wang D., Liebowitz D., Wang F., et al., (1988) :

Epstein Barr virus latent infection membrane protein alters the human B lymphocyte phenotype :

Deletion of the amino terminus abolish activity, J Virology, 62 : 4173.

Waterson A P., (1982) :

Viral hepatitis, in : McConnell J (ed) Clinical aspects of immunology 4th ed. Blackwell scientific public, Oxford, London, Edinburgh, Boston. Chap. 56, pp. 1515.

Weetman A P., Cotten S., Magko BA., et al., (1989) :

Expression of an intercellular adhesion molecule (ICAM-1) by human thyroid cell . J. Endocrinol, 122 : 185.

Weishselbaum TE., (1946) :

Measurment of total protein in serum. Amer. J. Clin. Path, 16 : 40.

Weisiger R., (1990) :

Non-A., non-B hepatitis. Promise and implications of hepatitis C assay. Modern Medicine. volume 7:615.

Whitby LG., Smith A F., Beckett GJ., et al., (1993) :

Plasma enzyme testes, in : Whitby LG, Smith AF, Beckett G J, and Walker SW (eds).Diagnosis in lecture notes on clinical biochemistry, oxford, Blackwell scientific publication the fifth edition, pp 78.

WHO (1977) :

Advances in viral hepatitis .Tech . Rep. series, 602 : 7.

Wilkinson JH., (1976):

Measurment of AST in: Principles and practice for diagnosis
enzymology. Year Book Medical publishers.

Williams A F., Barclay A N., (1988) :

The immunoglobulin superfamily. Domains for cell surface
recognition. Ann. Rev. Immunol, 6 : 381.

Wright T., Hsu H., Donegan E., (1991) :

Hepatitis C virus not found in fulminant non-A Non-B
hepatitis. Ann. Intern. Med, 115 : 111.

Wroblewski F., and LA Due JS., (1956):

Measurment of serum ALT. Proc. Soc. Exper. Biol. and
Med, 91:569

Zohrens G., Armbrust T., Pirzer U., et al., (1993) :

Intercellular adhesion molecule-1 concentration in sera of
patients with acute and chronic liver disease : relationship
to disease activity and cirrhosis. Hepatology, 18 : 798.