
REFERENCES

Abman, S.H.; Burchell, M.F.; Schaffer, M.S. and Rosenberg, A.A. (1989):

Late sudden unexpected death in the hospitalized infants with BPD. *Am. J. Dis. Child*, 143: 815.

Abman, S.H.; Kinesell, J.P.; Schaffer, M.S. et al. (1993):

Inhaled Nitric oxide in the management of a premature newborn with severe respiratory distress and pulmonary hypertension. *Pediatr.* 92: 606-609.

Abman, S.H.; Wolfe, R.R.; Accurso, F.J. et al. (1985):

Pulmonary vascular response to oxygen in infants with severe bronchopulmonary dysplasia. *Pediatrics*, 75-80.

Adeniyi-Jones, S.; McCormick, M.; Ramdial, H. et al. (1996):

RT-PCR for cytokine mRNA expression by airway cells from term infants with acute respiratory failure. *Pediatr. Res.*, 39: 323A.

Ahumada, C.A. (1996):

Continuous distending pressure in Goldsmith, J.P. and Karotkin E.H. (eds.) *Assisted ventilation of the neonate*, 3rd, Philadelphia, W.B. Saunders Co. 6: 728.

Aloan, C.A. and Hill, T.V. (1997):

Respiratory care of the newborn and child 2nd ed., J.B. Lippincott Company.

American Association for Respiratory Care (1991):

Clinical practice guidelines for pulse oximetry. *Respir. Care*, 26: 1406.

American Association of Respiratory Care, Clinical Practice guideline (1992):

Humidification during mechanical ventilation. *Resp. Care*, 37 (8): 887-890.

American Association for Respiratory Care Clinical Practice Guideline, (1991):

Oxygen therapy in the acute care hospital.
Respir. Care, 36 (12): 1410-1413.

Anderson, H.L.; Snedecor, S.M. Otsu, T. et al. (1993):

Multicenter comparison of conventional venoarterial access versus venovenous double-lumen catheter access in newborn infants undergoing extra corporeal membrane oxygenation. J. Pediatr. Surg., 28: 530.

Antunes, M.J. (1996):

Inhalation nitric oxide therapy: In Spitzer, A.R. Intensive care of the fetus and neonate. St. Louis Mosby-Year book. Inc., 51: 594.

Arensman, R.M. and Adolph, V.R. (1993):

Extra corporeal life support in children. In Arensman R.M. and Cornish, J.D. (eds): Extracorporeal life support. Boston, Blackwell Scientific, P. 275.

Arief, A.L.; Leach, W.; Park, R. and Lazarowitz, V.C. (1995):

Systemic effects of NaHCO₃ in experimental lactic acidosis in dogs. Am. J. Physiol., 242: F-586.

Arnada, J.V.; Stern, C. and Dunbar, J.S. (1972):

Pneumothorax with pneumoperitoneum in a newborn infant. Am. J. Dis. Child, 123: 163-166.

Arnold, J.H.; Troog, R.D.; Orav, E.J. et al. (1994):

Tolerance and dependence in neonates sedated with fentanyl during ECMO. Anesthesiology, 73: 1136.

Avery, M.E.; Gatewood, O.B. and Brumleg, G. (1966):

Transient tachypnea of the newborn. Am. J. Dis. Child, 111: 380.

Avery, M.E.; Hurd, S.S. and Tooley, W.H. (1986):

Is chronic lung disease in prematurely born infants preventable? Pediatr. Resp., 20: 314A.

Avery, M.E. and Taeusch, H.W. (1984):

Air leak and air trapping. In Avery, M.E. and Taeusch, H.W. (eds): Schaffer's diseases of the newborn, ed. 5 W.B. Saunders Co., 25: 217.

Bakken, E. and Desautels, D.A.C. (1993):

Oxygen therapy: In Koff, P.B.; Eitzman, D.V. and Neu, J. (eds): Neonatal and pediatric respiratory care, 2nd ed. St. Louis, Mosby, Year book, 13: 199.

Bancalari, E. and Gerhardt, T. (1986):

Bronchopulmonary dysplasia. Pediatric Clinic North Am., 33:1.

Bandy, K.P.; Nick, J.J. and Donn, S.M. (1992):

Volume controlled ventilation for severe neonatal respiratory failure, Neonatal intensive care, 5: 70.

Bark, H.; LeRoith, D.; Nyska, M. et al. (1980):

Elevation in plasma ADH level during PEEP ventilation in the dog: Mechanisms involved. Am. J. Physiol., 239: E474.

Barrington, K.J.; Feiner, N.H.; Peters, K.L. and Barton, J. (1986):

Physiological effects of doxapram in idiopathic apnea of prematurity. J. Pediatr., 108: 125.

Bartlett, R.H. (1990):

Extracorporeal life support for cardiopulmonary failure. Curr. Probl. Surg., 27 (10): 261-705.

Bartlett, R.H.; Andrews, A.F.; Toomasian, T.M. et al. (1982):

ECMO for newborn respiratory failure: Forth-five cases. Surgery, 92: 425.

Bartlett, R.H.; Gazzaniga, A.; Toomasian, J. et al. (1986):

ECMO in neonatal respiratory failure 100 cases. Ann. Surg., 204: 236-245.

Bateman, L.; Hooper, R.; Losh, T. and Miller, L. (1995):

Continuous positive airway pressure. In: Barnhart, S.L. and Cervinske, M.P. (eds.): Perinatal and Pediatric Respiratory care, Philadelphia, W.B. Saunders.

Baumgart, S. (1996):

Extra corporeal membrane oxygenation: In Spitzer, A.R.,: Intensive care of the fetus and neonate. St. Louis, Mosby Year book. Inc. 52: 602.

Baumgart, S.; Hirschl, R.B.; Butler, S.Z. et al. (1992):

Diagnosis-related criteria in the consideration of ECMO in neonates previously treated with HFJV. Pediatrics, 89: 491.

Beasley, J.M. and Jones, S.E. (1981):

CPAP in bronchiolitis, Br. Med. J., 283: 1506.

Beck, R.; Anderson, K.D.; Pearson, G.D. et al. (1986):

Criteria for ECMO in a population of infants with PPHN. J. Pediatr. Sur., 21: 297.

Behnke, M.L. (1993):

Patient assessment. In Koff, P.B., Eitzman, D.V. and Neu, J. (eds.): Neonatal and Pediatric Respiratory care, 2nd ed. St. Louis Mosby-Year book.

Behrman, R.E. (1996):

Pathophysiology of body fluids. In Nelson, W.B. and Vaughan, V.C. (eds.): Nelson Textbook of Pediatrics, ed. 15, W.B. Saunders, 12: 852, PP. 176-194.

Berkus, M.D.; Langer, O.; Samueloff, A. et al. (1994):

Meconium-stained amniotic fluid: Increased risk for adverse neonatal outcome. Obstet. Gynecol., 84: 115.

Bernstein, G. (1993):

Synchronous and patient-triggered ventilation in newborns. Neo. Intens. Care, Sep./ Oct.: 54.

Bernstein, G.; Cleary, J.P.; Heldt, G.P. et al. (1993):

Response time and reliability of three neonatal patient-triggered ventilators. Am. Rev. Respir. Dis., 148: 385.

Bernstein, G.; Mannino, F.L.; Heldt, O.P. et al. (1996):

Prospective randomized multicenter trials comparing synchronized and conventional intermittent mandatory ventilation (SIMV Vs. IMV) in neonates. Part 2. J. Pediatr., 128 (4): 453.

Bertrand, J.M.; Riley, S.P.; Popkin, J. et al. (1985):

The long term pulmonary sequelae of prematurity. The Role of familial hyperactivity and the RDS. N. Engl. J. Med., 312: 742-745.

Betit, P. and Thompson, J.E. (1993):

Mechanical ventilation. In Koff, P.B.; Eitzman, D.V. and Neu, J. (eds.): Neonatal and pediatric respiratory care, 2nd ed. St. Louis, Mosby Year book, 18: 282.

Bhatia, J. and Ziegler, E.E. (1983):

Retinol-binding protein and prealbumin in cord blood of term and preterm infants. Early Hum. Dev., 8: 219-233.

Bhutani, V.K. and Abbasi, S. (1992):

Long term pulmonary consequences in survivors with BPD. Clinics in Perinatology Vol. 19 (No. 3), PP. 649-671.

Bland, R.D.; Kim, M.H.; Light, M.J. et al. (1980):

High frequency mechanical ventilation in severe hyaline membrane disease an alternative treatment? Crit. Care Med., 8: 275.

Bohan, D.J. et al. (1980):

Ventilation by high frequency oscillation. J. Appl. Physiol., 48: 710.

Bomsel, F. (1973):

Pulmonary sequelae of HMD. Iatrogenic Factors? Ann. Radiol., 16: 70.

Boon, A.W. (1987):

Respiratory emergencies. In Black J.A. (eds.), Pediatric Emergencies, ed. 2 Butterworths, 23: 231-258.

Borakat, G. and Liebman, J. (1985):

Pulmonary aspects of cardiovascular disease. In Lough, M.D., Doershuk, C.F. and Stern, R.C. (eds.): Pediatric Respiratory therapy, ed. 3, year book Medical publisher, 4: 107-118.

Boros, S.J. (1986):

Principle of ventilator care. In Thibeault, D.W. and Gregory, G.A. (eds.): Neonatal Pulmonary care, ed. 2, Norwalk, Conn. Appleton. Century-Crofts, 15: 367.

-
- Boros, S.J.; Mammel, M.C.; Coleman, J.M. et al. (1985):**
Neonatal HFJV: Four years experience. *Pediatrics*, 75: 657.
- Bower, L.K. and Wilson, J.M. (1995):**
Extracorporeal life support: In Barnhart, S.L. and Czervinske, M.P. (eds.): *Perinatal and Pediatric respiratory care*. W.B. Saunders Company, 21: 376.
- Boyden, E.A. (1976):**
The programming of canalization in fetal lungs of man and Monkey. *Am. J. Anat.*, 145: 125.
- Boyle, K.M.; Baker, V.L.; Cassaday (1995):**
Neonatal pulmonary disorders. In Barnhart SL, Czervinske MP (eds) *Perinatal and Pediatric Respiratory care*, Philadelphia, W.B. Saunders.
- Bregman, J. and Farrell, E.E. (1992):**
Neurodevelopmental outcome in infants with BPD. *Clinics in Perinatology*, Vol. 79, (No. 3), PP. 673-694.
- Brown, E.R.; Stark, A.; Lawson, E. and Avery, M.E. (1978):**
BPD: Possible relationship to pulmonary edema. *J. Pediatr.*, 92: 982.
- Bruce, M.E.; Martin, R.J. and Boat, T.F. (1984):**
Concentration of alpha-protease inhibitors and alpha-2-macroglobulin in serum and lung secretion of intubated infants. *Pediatr. Res.*, 18: 35.
- Burchfield, D. and Neu, J. (1993):**
Neonatal parenchymal disease. In Koff PB, Eitzman D, Neu J (eds): *Neonatal and pediatric respiratory care*, 2nd ed. St. Louis, Mosby, year book.
- Burgerss, W.R. and Chernick, V. (1996):**
Respiratory therapy in newborn infants and children, New York, Thieme.
- Burton, G.G. (1984):**
Respiratory care, a guide to clinical practice, Philadelphia, J.B. Lippincott Co.
-

Butler, W.J. et al. (1980):

Ventilation by high frequency oscillation in humans. *Anesth. Anal.*, 59: 577.

Burton, G.W. and Ingolod, K.V. (1984):

Beta caroten. An Usual type of antioxidant. *Science*, 224: 569.

Cameron, D.; Arnst, R.; Clay, M. and Silverman, M. (1991):

Acrosol delivery in neonatal ventilator circuits. *Pediatric Pulmonology*, 10: 208-213.

Carlo, W.A.; Beoglos, A.; Chatburn, R.L. et al. (1989):

High frequency jet ventilation in neonatal pulmonary hypertension. *Am. J. Dis. Chil.*, 143: 233.

Carlo, W.A. and Martin, R.J. (1996):

Principles of neonatal assisted ventilation. *Pediatr. Clin. Nor. Am.*, 33: 221-237.

Carlo, W.A.; Siner, B.; Chatburn, R.L.L. et al. (1991):

Early randomized intervention with HFIV in RDS. *J. Pediatr.*, 117: 765.

Caron, E.; Maguie, D.P. et al. (1990):

Current management of pain, sedation and narcotic physical dependency of the infant in ECMO. *J. Perinatal. Neonatal. Nurse*, 4: 63-74.

Carter, J.M.; Gerstmann, D.R.; Clark, R.H. et al. (1990):

HFOV and ECMO for the treatment of acute neonatal respiratory failure. *Pediatrics*, 85: 159.

Cassady, G. (1993):

Transcutaneous monitoring in the newborn infants. *J. Pediatr.*, 103: 837.

Cassidy, S.S.; Goaffney, F.A. and Johnson, R.L. (1981):

A prespective on PEEP. *N. Engl. J. Med.*, 304: 421.

Castillo, L.; DeRojas-Walker, T.; Yu, Y.M. et al. (1995):

Whole body arginine metabolism and nitric oxide synthesis in newborns with persistant pulmonary hypertension. *Pediatr. Res.*, 38: 17.

-
- Chai, P.J.; Skaryack, L.A.; Ungerleider, R.M. et al. (1995):**
Jugular ligation does not increase intracranial pressure but does increase bihemispheric cerebral blood flow and metabolism. Crit. Care Med., 23: 1864.
- Chan, K.N.; Wong, Y.C. and Silverman, M. (1990):**
Relationship between infant lung mechanics and childhood lung function in children of very low birth weight. Pediatr. Pulmon., 8: 74-81.
- Chamedes, L. (1996) (ed.):**
Textbook of advanced life support. American heart association, American Academy of Pediatrics.
- Chang, H.K. (1984):**
Mechanism of gas transport during ventilation by High frequency oscillation, J. Appl. Physiol., 56: 553.
- Chatburn, R.L. (1988):**
Similarities and differences in the management of acute lung injury in neonates (IRDS) and in adults (ARDS). Respi. Care, 33 (7): 539-553.
- Chatburn, R.L. (1992):**
Classification of mechanical ventilators. Respir. Care, 37: 1009.
- Cheifetz, I.M.; Craig, D.M.; Kern, F.H. et al. (1996):**
Nitric oxide improves transpulmonary vascular mechanics but does not change intrinsic right ventricular contractility in an acute respiratory distress syndrome model with permissive hypercapnia. Crit. Care Med., 24: 1554-1561.
- Chevalier, J.Y.; Durandy, Y.; Batisse, A. et al. (1990):**
Preliminary report: Extracorporeal lung support in neonatal acute respiratory failure. Lancet, 335: 1364-1366.
- Chishty, A.L.; Alvi, Y.; Iftikhar, M. et al. (1996):**
Meconium aspiration in neonate: combined obstetric and paediatric intervention improves outcome. J. Pakistan Med. Assoc., 46: 104.
-

Clarck, L.C.; Hoffmann, R.E.; Davis, S.L. (1994):

Response of rabbit lung as a criterion of safety for fluorocarbon breathing and blood substitutes. In Chang, M.S.; Riess, R.M., Winslow, R.M., (eds.): Biomaterial, artificial cells and immobilization biotechnology, Marcel Dekker, New York, 292.

Clark, L.C. and Gollan, F. (1966):

Survival of mammals breathing organic liquids equilibrated with oxygen at atmosphere pressure. Science, 152: 1755-1756.

Clark, R.H.; Yoder, B.A. and Sell, U.S. (1994):

Prospective randomized comparison of high frequency oscillation and conventional ventilation in respiratory distress syndrome. Pediatrics, 89:5.

Cleary, J.P.; Bernstein, G.; Mannino, F.L. et al. (1995):

Improved oxygenation during SIMV in neonates with RDS: A randomized, cross over study. J. Peds., 126 (3): 407.

Cleary, G.M. and Wiswell, T.E. (1998):

Meconium-stained amniotic fluid and meconium aspiration syndrome. Pediatric Clinic of North America, Volume 45, No. 3, June, P. 511.

Cleveland, D.V. (1987):

Continuous distending pressure. In Aloia, C.A. and Hill, T.V. (ed.): Respiratory care of the newborn, J.B. Lippincott Co., 12: 219.

Cleveland, D.V. (1987):

Oxygen therapy. In Aloia, C.A. and Hill, T.V. (ed.): Respiratory Care of the newborn. J.B. Lippincott C., 11: 207.

Cloherly, J.P. and Jose, J.H. (1986):

Meconium aspiration. In Cloherly, J.P. and Stark, A.R. (ed.): Manual of neonatal care, ed. 2, Little, Brown and Company Inc., 14: 203.

Coghill, C.J. et al. (1991):

Neonatal and pediatric HFV: Principles and Practice. Respir. Care, 36: 596.

-
- Corbet, A.; Bucciarelli, R.; Goldman, S. et al. (1991):**
Decreased mortality rate among small premature infants treated at birth with a single dose of synthetic surfactant a multicenter controlled trial. *J. Pediatr.*, 118: 277.
- Cornish, J.D. and Arensman, R.M. (eds.) (1992):**
Extracorporeal life support. Blackwell, Boston.
- Cornish, J.D.; Heiss, K.F.; Clark, R.H. et al. (1993):**
Efficacy of venovenous ECMO for neonates with respiratory and circulatory compromise. *J. Pediatr.*, 122: 105.
- Crone, R.X. (1981):**
Assisted ventilation in children. In Gregory, G.A. (ed.): *Respiratory failure in the child*. New York, Churchill Livingstone, 3: 17.
- Cunningham, M.D. and Wood, B.R. (1996):**
Monitoring of pulmonary function. In: Goldsmith, J.P.; Karotkin, E.H. (eds.): *Assisted ventilation of the neonate*, 3rd Ed. Philadelphia, W.B. Saunders, P. 273.
- Curits, S.E.; Fuhrman, B.P. and Howland, B.F. (1990):**
Airway and alveolar pressures during perfluorocarbon breathing in infant lamb. *J. App. Physiol.*, 68: 2322-2328.
- Czervinske, M.P. (1993):**
Arterial blood gas analysis and other cardiopulmonary monitoring. In: Koff, P.B.; Eitzman, D.V. and Neu, J. (eds.): *Neonatal and Pediatric Respiratory Care*, 2nd ed. St. Louis-Mosby, Year book.
- Czervinske, M.P. (1993):**
Continuous positive airway pressure. In: Koff, P.B., Eitzman, D.V. and Neu, J. (eds.): *Neonatal and pediatric respiratory care*, 2nd ed. St. Louis-Mosby Year book.
- Cummings, J.J.; D'Eugenio, D.B. and Gross, S.J. (1989):**
A controlled trial of dexamethazone in preterm infants at high risk for BPD. *N. Engl. J. Med.*, 320: 1505-1510.
-

- Davila, D. and Davila, T. (1981):**
Thiazide diuretics inhibit contraction of isolated smooth muscle, *Pharmacology*, 22: 108-112.
- Davis, J.M.; Richter, S.E.; Kending, T.W. et al. (1992):**
High frequency jet ventilation and surfactant treatment of newborns with severe respiratory failure. *J. Pediatr. Pulmonol.*, 413: 108.
- Day, R.W.; Guarin, M.; Lynch, J.M. et al. (1996):**
Inhaled nitric oxide in children with severe lung disease: Results of acute and prolonged therapy with two concentrations. *Crit. Care Med.*, 24: 215-221.
- DeJour, P. (1981):**
Principles of comparative respiratory physiology. Amsterdam, Elsevier, North Holland, P. 147.
- de Lemos, R.H.; Coulson, J.J.; de Lemos, J.A. et al. (1992):**
Rescue ventilation with high frequency oscillation in premature baboons with hyaline membrane disease. *Pediatr. Pulmonol.*, 12: 11.
- de Lemos, R.A.; Coulson, J.J.; Meredith, K.S. et al. (1989):**
A comparison of ventilation strategies for the use of HFOS in treatment of HMD. *Acta. Anaesthesiol. Scand. Suppl.*, 90: 102.
- DeMaio, J.G.; Harris, M.C.; Deuber, C. and Spitzer, A.R. (1989):**
Effect of blood transfusion on apnea frequency, in growing premature infants. *J. Pediatr.*, 114: 1039.
- Deming, D.D. (1995):**
Respiratory assessment of neonatal and pediatric patients. In Wilkins, R.L., Krider, S.J. and Sheldon, R.L., (eds.): *Clinical assessment in Respiratory care*, 3rd ed. St. Louis: Mosby Year-book.
- Demirakca, S.; Dotsch, J.; Knothe, C. et al. (1996):**
Inhaled Nitric oxide in neonatal and pediatric acute respiratory distress syndrome: Dose response, Prolonged inhalation, and weaning. *Crit. Care Med.*, 24: 1913-1919.

-
- Demling, R.H. and Will, J.A. (1988):**
The effect of furosemide on the pulmonary transvascular fluid filtration rate. *Crit. Care Med.*, 6: 317-319.
- Deoras, K.S.; Coppola, D.; Wolfson, M.R. et al. (1990):**
Liquid ventilation in neonates: Tissue histology and morphometry. *Pediatr. Res.*, 27: 29A.
- Donn, S.M. and Nicks, J.J. (1996):**
Special ventilatory techniques and modalities: patient triggered ventilation. In Goldsmith, J.P.; Karotkin, E.H. (eds.): *Assisted ventilation of the Neonate*, 3rd Philadelphia: W.B. Saunders, P. 215.
- Drazen, J.M.; Kamm, R.D.; Slutsky, A.S. (1984):**
High frequency ventilation, *Physiol. Rev.*, 64: 505.
- Dubin, S.B. (1992):**
Assessment of fetal lung maturity by laboratory methods. *Clin. Lab. Med.*, 12 (3): 603.
- Durand, D.J. and Phillips, B.L. (1996):**
Blood gases: Technical aspects and interpretation in Goldsmith, J.P.; Kartokin, E.H. (eds.): *Assisted Ventilation of the neonate*, 3rd Philadelphia, W.B. Saunders, P. 257.
- Durand, M. and Rigatto, H. (1981):**
Tidal volume and respiratory frequency in infants with BPD, *Early Hum. Dev.*, 5: 66.
- Edwards, D.K. (1979):**
Radiological aspects of BPD. *J. Pediatr.*, 95:823.
- Edward, D.K. (1983):**
Bronchopulmonary dysplasia today. In Milunsky, A.; Friedman, E.A., and Gluck, L. (eds.): *Advances in perinatal medicine*, Vol. 3, New York, Plenum Medical Book Company, PP. 117-162.
- Edwards, D.K.; Dyer, W.M. and Narthway, W.H.Jr. (1977):**
Twelve years experience with BPD pediatrics, 59: 839-846.
-

Gerard, P.; Fox, W.W.; Outerbridge, E.W. (1975):

Early versus late introduction of continuous negative pressure in the management of idiopathic. Respiratory distress syndrome. J. Pediatr., 87: 591.

Gerlach, H.; Pappert, D.; Lewundowski, R. et al. (1993):

Long-term inhalation with evaluated low doses of nitric oxide for selective improvement of oxygenation in patient with adult respiratory distress syndrome. Inten. Care Med., 19: 443-449.

Gerstmann, D.R.; Minton, S.D.; Stoddart, R.A. et al. (1996):

The provo multicenter early HFOV trials: improved pulmonary and clinical outcome in RDS. Pediatrics, 98: 1044.

Gerstman, D.R.; Koichiro, N.; Kinsella, J.P. et al. (1989):

Left carotid artery and coronary arterial flow partitioning during neonatal ECMO. Pediatr. Res., 25: 37A.

Giaid, A. and Saleh, D. (1995):

Reduced expression of endothelial nitric oxide synthase in the lungs of patients with pulmonary hypertension. N. Engl. J. Med., 333: 214.

Goldman, S.L.; Gerhardt, T.; Sonni, R. et al. (1983):

Early prediction of chronic lung disease by pulmonary function testing. J. Pediatr., 102: 613.

Goldman, A.P.; Tasker, R.C.; Haworth, S.G. et al. (1996):

Four patterns of response to inhaled nitric oxide for persistent pulmonary hypertension of the newborn. Pediatrics, 98: 706-713.

Goldsmith, J.P. and Karotkin, E.H. (1996):

Introduction to assisted ventilation. In Goldsmith, J.P and Karotkin, E.H. (eds.); Assisted ventilation of the neonate, 3rd, Philadelphia, W.B. Saunders Co., 1:4.

Gomella, T.L. (1988):

Neonatology, basic management, on-call problems, diseases, drugs. Appleton and lange.

-
- Goodin, T.H.; Kaufman, R.J. and Richard, T.J. (1993):**
Comparative pulmonary toxicity of three perfluorochemicals (PFCs) in the rat and baboon. Proceeding of the fifth international symposium on blood substitutes. San Diego.
- Goodstein, M.H.; Greenspan, T.S.; Friss, H.E. et al. (1992):**
Extrauterine pulmonary development in the healthy preterm infant: hidden morbidity at discharge. *Pediatr. Resp.*, 31: 248A.
- Gottschark, S.K.; King, B. and Schuth, C.R. (1980):**
Basic concepts in PPV of newborn. *Perinatology, Neonatology*, 4:15.
- Gramlich, T. (1995):**
Oxygen therapy. In Barnhart, S.L., and Czervinske, M.P., (eds): *Perinatal and pediatric respiratory care*. Philadelphia, W.B. Saunders.
- Graziani, L.J.; Spitzer, A.R.; Mitchell, D.G. et al. (1992):**
Mechanical ventilation in preterm infants: neurosonographic and developmental studies. *Pediatrics*, 90: 515.
- Greenough, A. and Robertson, N.R.C. (1999):**
Respiratory distress syndrome: In Greenough, A.; Milner, A.d. and Robertson, N.R.L. (eds.): *Neonatal respiratory disorders*, 4th Arnold, 18: 238.
- Greenspan, J.S. (1993):**
Liquid ventilation: A developing technology, *Neonatal Network*, 12:23.
- Greenspan, J.S. (1996):**
Physiology and clinical role of liquid ventilation therapy. *J. Perinatol.*, 16 (Suppl): 47.
- Greenspan, J.S. and Antunes, M.J. (1996):**
Liquid ventilation: In Spitzer, A.R., *Intensive care of the fetus and neonate*. St. Louis, Mosby, Year book Inc., 53: 619.
- Greenspan, J.S.; Wolfson, M.R.; Rubenstein, S.D. et al. (1990):**
Liquid ventilation in human preterm neonate. *J. Pediatr.*, 117: 106-111.
-

Gregory, G.A. (1986):

Devices for applying CDP. In Thibeault, D.W. and Gregory, G.A. (eds.): Neonatal pulmonary care, ed. 2, Norwalk, Conn. Appleton-Century-Crofts, 12: 307.

Gregory, G.A.; Kitterman, J.A. and Phibbs, R.H. (1971):

Treatment of idiopathic respiratory distress syndrome with CPAP, N. Eng. J. Med., 284: 1333-1338.

Griffith, T.M.; Edwards, D.H.; Lewis, M.J. et al. (1984):

The nature of endothelium-derived vascular resistance. Pediatrics, 60: 444-449.

Gross, N.J. and Skorodin, M.S. (1984):

Anticholinergic antimuscarinic bronchodilators. Am. Rev. Respir. Dis., 129: 856-870.

Hagedorn, M.I.; Gardner, S.L. and Abman, S.H. (1989):

Respiratory disease. In Merenstein and Gardner, S.L. (eds.): Hand book of neonatal intensive care, C.V. Mosby Co. PP., 365-426.

Hageman, J.R. and Caplan, M.S. (1995):

An introduction to the structure and function of inflammatory mediators for clinicians. Clin. Perinatal., 22: 257.

Hall, R.T. and Rhodes, P.G. (1975):

Pneumothorax and pneumomediastinum in infant with idiopathic RDS receiving continuous positive airway pressure. Pediatrics, 55: 493.

Halliday, H.L.; Pumpit, F.M. and Brady, J.P. (1980):

Effect of inspired oxygen on echographic effect of inspired oxygen vascular resistance and myocardial contractility in BPD. Pediat., 65: 538.

Halliday, H.L.; McClure, G. and Reid, M. (1989):

Respiratory problems. In Halliday, H.L.; McClure, G. and Reid, M. (eds): hand book of neonatal intensive care, ed. 3, Balliere, Tindall, 10: 123.

- Hansell, D.R. (1997):**
Extracorporeal membrane oxygenation: In Aloia, C.A.; Hill, T.V. (eds.): Respiratory care of the newborn and child. Ed. 2, Lippincott Co., 25: 505-538.
- Hardesty, R.L.; Griffith, B.P.; Bebski, R.F. et al. (1981):**
ECMO: Successful treatment of persistent fetal circulation following repair of CDH. J. Thorac. Cardiovasc. Surg., 81: 556-563.
- Harris, D.G.; Coggin, R.R.; Roby, J. et al. (1983):**
Liquid ventilation in dogs: an apparatus for normobaric and hyperbaric studies. J. Appl. Physiol., 54: 1141-1148.
- Harris, T.R. and Wood, B.R. (1996):**
Physiologic principles. In: Goldsmith, J.P. and Karotkin, E.H., (eds.): Assisted ventilation of neonates 3rd ed., Philadelphia. W.B. Saunders, P. 48.
- Harrison, M.R. and Delorimier, A.A. (1981):**
Congenital diaphragmatic hernia. Sur. Clin. North Am., 61: 1023.
- Haworth, S.G. and Ried, L. (1977):**
Structural study of pulmonary circulation and of heart in total anomalous venous return in early infancy. Br. Heart J., 39: 80.
- Hazinski, T.A. (1990):**
BPD. In Chernick, V. (ed.) Disorders of respiratory tract in children. Philadelphia, W.B. Saunders, PP. 300-320.
- Heffner, J.E. and Replne, J.E. (1989):**
Pulmonary strategies of anti-oxidant defence. Am. Rev. Respir. Dis., 140:531.
- Heicher, D.A.; Kastind, D.S. and Harrod, J.R. (1981):**
Prospective clinical comparison of two methods for mechanical ventilation of neonates rapid rate and short inspiratory time versus slow rate and long inspiratory time. J. Pediatr., 98: 957.

Heldt, G.P. and Bernstein, G. (1993):

Patient-initiated mechanical ventilation. In Boynton, B.R., Carlo, W.A.; Jobe, A.H. (eds): New Therapies for neonatal respiratory failure. A physiological Approach Amsterdam: Elsevier, P. 152.

Heldt, G.P.; Merrit, T.A.; Golembeski, D. et al. (1994):

Distribution of surfactant, lung compliance and areation of preterm rabbit lungs after surfactant therapy and conventional and HFOV. *Pediatr. Resp.*, 31: 270.

Hickey, P.R.; Buckley, M.J. and Philbin, D.M. (1983):

Pulsatile and non-pulsatile cardiopulmonary bypass. Review of a counter productive controversy. *Ann. Thorac. Surg.*, 36: 720-737.

The HIFI Study Group (1989):

HFOV compared with conventional mechanical ventilation in the treatment of respiratory failure in preterm infants. *N. Engl. J. Med.*, 320: 88.

HIFO Study Group (1993):

Randomized study of HFOV in infants with severe RDS. *J. Pediatr.*, 122: 609.

Higgins, R.D.; Richter, S.E. and Davis, J.M. (1991):

Nasal continous positive airway pressure facilitates extubation of VLBW neonates. *Pediatrics*, 88: 999-1003.

Higgins, S.T.; Wu, A.M.; Sen, N. et al. (1996):

Meconium increases surfactant secretion in isolated rat alveolar type II cells. *Pediatr. Res.*, 39: 443.

Hill, K.V. (1997):

Oxygen therapy: In Aloian, C.A.; Hill, T.V., (eds): Respiratory care of newborn and child, ed. 2. J.P. Lippincott Company, 14: 277-290.

Hinds, C.J. (1987):

Artificial ventilation. In Hinds, C.J. (ed.): Intensive care: A concise textbook, Bailliere Tindall, Co., 13: 227.

Hird, M.; Greenough, A. and Gasmu, H. (1990):

Gas trapping during high frequency positive pressure ventilation using conventional ventilators.
Early Hum. Dev., 22: 51.

Hirschl, R.B.; Coburn, C.; Butter, S.M. et al. (1994):

Early onset sepsis neonatorum with listeria-monocytogenes emerging as treatable disease when provided prolonged extracorporeal life support (ECLS). Arch. Pediatr. Adolesc. Med., 148: 513-517.

Ho, T.F. (1989):

Instrumentation in acute pediatrics. In Yip, W.C. and Tay, J.S. (eds.): A practical manual on acute pediatrics, Singapore, P.G., Publishing, Co. 29: 395.

Hocker, J.R.; Pippa, M.S.; Rabalais, G.P. et al. (1992):

ECMO and early onset group B-Streptococcal sepsis. Pediatr., 89:1.

Hodgman, J.E. (1981):

Chronic lung disorders. In Avery GB (ed): Neonatology: Pathophysiology and management of the newborn, 2nd ed. P. 407, Philadelphia, JB Lippincott.

Hoffman, G.M.; Ross, R.A.; Day, S.E. et al. (1997):

Inhaled nitric oxide reduces the utilization of ECMO in PPHN. Crit. Care Med., 25: 352.

Hogman, M.; Frostell, C.; Arnberg, H. et al. (1993):

Bleeding time prolongation and NO inhalation. Lancet, 341: 1664.

Holtzman, R.B.; Hageman, J.R. and Yoger, R. (1989):

Role of ureaplasma urealyticum in BPD. J. Pediatr., 114: 1061.

Huber, G.L. and Drath, D.B. (1981):

Pulmonary oxygen toxicity. In Gilbert, D.L. (ed) Oxygen and Living Processes. An Interdisciplinary Approach. New York, Springer-Verlag, P. 273.

Hudak, B.; Arlen, M.; Hudack, M. et al. (1989):

Home oxygen therapy for chronic lung disease in extremely low birth weight infants. *Am. J. Dis. Child*, 143: 357-360.

Hummer, M.; Viguerat, C.E.; Suter, P.M. et al. (1980):

Urinary antidiuretics hormone excretion during mechanical ventilation and weaning in man.

Anesthesiology, 52: 395.

Hummler, H.; Gerhardt, T.; Gonzales, A. et al. (1996):

Influence of different methods of synchronized mechanical ventilation on ventilation, gas exchange, patient effort and blood pressure fluctuation in premature infants. *Pediatr. Pulmonol.*, 22: 305.

Hustead, V.A.; Gutcher, G.r.; Anderson, S.A. and Zachman, R.D. (1984):

Relationship of Vit. A (Retinol) status to lung disease in the preterm infant. *J. Pediatr.*, 105: 610.

Jackson, J.C.; Trug, W.E.; Standaert, T.A. et al. (1994):

Reduction in lung injury after combined surfactant and HFV. *Am. J. Respir. Crit. Care Med.*, 150: 543.

Johns, R.A.; Linden, J.M. and Peach, M.J. (1989):

Endothelium-dependent relaxation and GMP accumulation in rabbit pulmonary artery are selectively impaired by moderate hypoxia. *Circ. Res.*, 65: 1508-1515.

Jonzon, A. (1991):

Indications for CPAP and respiratory therapy. *Int. J. Technol. Assess. Health Care*, 7 (Suppl. 1), 26-30.

Kamper, J. and Ringsted, C. (1990):

Early treatment of idiopathic respiratory distress syndrome using binasal continous positive airway pressure. *Acta. Pediatr. Scand.*, 79: 581-586.

Kao, L.C.; Warburton, D.; Sargent, C.W. et al. (1983):

Furosemide acutely decrease airway resistance in chronic BPD. *J. Pediatr.*, 103: 624-629.

Kattener, E.; Metez, B.; Waiss, E. and Obladen, M. (1992):

Accelerated lung maturation following maternal steroid treatment in infants born before 30 weeks gestation. *J. Perinatal Med.*, 20 (6): 449.

Kattwinkel, J.; Nearman, H.S.; Faaroff, A.A. et al. (1976):

Apnea of prematurity: comparative therapeutic effects of cutaneous stimulation and nasal CPAP, *J. Pediatr.*, 86: 588.

Keszler, M.; Donn, S.M. and Bucciarelli, R.L. (1991):

Multicenter controlled trial comparing high frequency jet ventilation and conventional mechanical ventilation in new born infants with pulmonary interstitial emphysema. *Journal of pediatrics*, 119: 85-93.

Keszler, M.; Ryckman, F.C.; McDonald, J.V. et al. (1992):

A prospective multicenter randomized study of high to low positive-end expiratory pressure during ECMO. *J. Pediatr.*, 120: 107-113.

Kinsella, J.P.; Ivy, D.D. and Abman, S.H. (1994):

Inhaled nitric oxide improves gas exchange and lowers pulmonary vascular resistance in severe experimental hyaline membrane disease. *Pediatr. Res.*, 36: 402.

Kinnsella, J.P.; McQueston, J.A.; Rosenberg, A.A. et al. (1992):

Hemodynamic effects of exogenous nitric oxide in ovine transitional circulation. *Am. J. Physiol.*, 32: H875-880.

Kinsella, J.P.; Neish, S.R.; Ivy, D.D. et al. (1993):

Clinical responses to prolonged treatment of PPHN with low doses of inhaled nitric oxide. *J. Pediatr.*, 123: 103.

Klein, M.D.; Andrews, A.F.; Wesley, J.R. et al. (1985):

Veno-venous perfusion in ECMO for newborn respiratory insufficiency: A clinical comparison with veno-arterial perfusion. *Ann. Surg.*, 201: 520-526.

Koff, B.P.; Eitzman, D.V. and Neu, J. (eds.) (1993):

Neonatal and pediatric respiratory care. 2nd ed. St. Louis, Mosby Yew book.

Kojima, T.; Hattori, K.; Fujiwara, T. et al. (1994):

Meconium induced lung injury mediated by activation of alveolar macrophages. *Life Sciences*, 54: 1559.

Kolbow, T.; Borelli, M.; Spatula, R. et al. (1988):

Single catheter venovenous membrane lung bypass in the treatment of experimental ARDS. *Trans. Am. Soc. Artif. Internal Organ*, 34: 35-38.

Kolbow, T. and Bowman, R.L. (1963):

Construction and elimination of an alveolar membrane artificial heart-lung. *Trans. Am. Soc. Artif. Internal Organ.*, 9: 238.

Korons, S.B. (1996):

Complications. In Goldsmith, J.P. and Karotkin, E.H. (eds): *Assisted ventilation of the neonate, 3rd*, Philadelphia, W.B. Saunders Co., 13: 245.

Krauss, A.N. (1979):

Ventilation-perfusion relationships in neonates. In Thibeault, D.W.; Gregory, G.A. (eds); *Neonatal pulmonary care*. Reading M.A., Addison-Wesley Publishing Co., P. 54.

Kravath, R.E. (1992):

Blood gases, hydrogen ions and pH. In Finberg, L.; Kravath, R.E. and Fleischman, A.R. (eds.) *Water and electrolytes in pediatrics*, Philadelphia, W.B. Saunders, Co. 6: 35.

Kraybill, E.N.; Runyan, D.K.; Bose, C.L. and Khan, J.H. (1989):

Risk factor for chronic lung disease in infants with birth weights of 751 to 1000 gms. *J. Pediatr.*, 115: 115.

Krummel, T.M.; Greenfield, L.J.; Kirkpatrick, B.V. et al. (1984):

Alveolar arterial oxygen gradients versus the neonatal pulmonary insufficiency index for prediction of mortality in ECMO candidates, *J. Pediatr. Surg.*, 14: 380-384.

Kurlinski, J.P. and Gluck, L. (1985):

Respiratory problems. In Paxon, C.L., (ed.) *Van Leeuwen's newborn medicine*, ed. 2, Chicago, Year book Medical Publishers, Inc.

Kylstra, J.A.; Pajanelli, C.V. and Lanphier, E.H. (1962):

Pulmonary gas exchange in dog ventilated with hyperbarically oxygenated liquid. *J. Appl. Physiol.*, 1: 177-184.

Landers, S.; Hansen, T.N.; Corbet, A.J.S.; Stevener, M.J. and Rudolph, A.D. (1986):

Optimal constant positive airway pressure assessed by arterial alveolar or difference for CO₂ in HMD. *Pediatr. Res.*, 20 (9): 884-889.

Lawson, E.E.; Birdwell, R.L. and Huang, P.S. (1979):

Augmentation of pulmonary surfactant secretion by lung expansion at birth. *Pediatr. Res.*, 13: 611.

Leach, C.; Greenspan, J.S.; Rubenstein, S.D. et al. (1996):

Partial liquid ventilation with perflurocarbon in premature infants with severe RDS. *N. Engl. J. Med.*, 33 5: 761.

Levene, M.I.; Tudehope, D. and Thearic, J. (eds.) (1987):

Essentials of neonatal medicine. Blackwell Scientific Publication.

Liley, H.G. and Stark, A.R. (1991):

Respiratory distress syndrome/ hyaline membrane disease. In Cloherty, J.P. and Stark, A.R. (eds) *Manual of neonatal care*, ed. 3, Little, Brown and Company, PP. 189-194.

Lillehei, C.W. (1993):

History of development of extracorporeal circulation. In Arensman, R.M. and Cornish, J.D. (eds.): *Extracorporeal Life support*. Boston, Blackwell Scientific, P. 26.

Lindberg, L.; Rydgién, G.; Larsson, A. et al. (1997):

A delivery system for inhalation of nitric oxide evaluated with chemiluminescence, electrochemical fuel cells, and capnography, *Crit. Care Med.*, 25: 190-196.

Lucas, G.N.; Dissanayake, P. (1995):

A study of meconium St aimed and meconium aspiration syndrome in a maternity hospital. *Ceylon Med. J.*, 40: 62.

Lucey, J.F. and Dangman, B. (1984):

A reexamination of the role of oxygen in RLF.

Pediatrics, 72: 82

-
- Macklin, C.C. (1939):**
Transport of air along sheath of blood vessels from alveoli to mediastinum: clinical implications.
Arch. Intern. Med., 63: 913.
- Macklin, M.T. and Macklin, C.C. (1944):**
Malignant II: of the lung and mediastinum as an important complication in many respiratory diseases and other conditions: An interpretation of the clinical literature in the light of laboratory experiment. Medicine, 23: 281.
- Malik, A.S. and Hillman, D. (1994):**
Meconium aspiration syndrome and neonatal outcome in a developing country. Ann. Trop. Pediatr., 14: 47.
- Mammel, M.C. and Boros, S.J. (1996):**
High frequency ventilation. In Goldsmith, J.P.; and Karotkin, E.H.; (eds): Assisted Ventilation of neonate ed. 3. Philadelphia: W.B. Saunders, P. 199.
- Mannino, F.L. and Merritt, T.A. (1986):**
The management of RDS in Thibeault, D.W. and Gregory G.A. (eds): Neonatal pulmonary care second edition, Appleton-Century-Crofts, 18: 427.
- Margraf, L.R.; Tomashefski, J.F.Jr.; Bruce, M.C. et al. (1991):**
Morphometric analysis of the lung in bronchopulmonary dysplasia. Am. Rev. Respir. Dis., 143: 391.
- Martin, R.J. and Carlo, W.A. (1986):**
Role of upper airway in pathogenesis of apnea in infants. Respir. Care, 31: 615.
- Mathew, O.P. and Thach, B.T. (1980):**
Selective bronchial obstruction for treatment of bullous interstitial emphysema. J. Pediatr., 96: 475.
- Matsuoka, J.; Kawano, T. and Miyasaka, K. (1994):**
Role of HFV in surfactant depleted lung injury as measured by granulocytes. J. Appl. Physiol., 76: 539.
- Mayes, L.; Perkett, E. and Stahlman, M.T. (1983):**
Severe BPD: A retrospective view. Acta, pediatr. Scand., 72: 225.

McCord, J.M. and Fridovich, I. (1969):

Superoxide dismutase an enzymic function for erythrocyte hemocuprein (hemocuprein). *J. Biol. Chem.*, 144: 6049.

McGettigan, M.C.; Adolph, V.R.; Ginsberg, H.G. and Goldsmith, J.P. (1998):

New way to ventilate newborns in acute respiratory failure. *Pediatric clinic of North America*. Vol. 45, No. 3, June, P. 475.

Mehta, A.; Wright, B.M.; Callen, K. et al. (1986):

Patient triggered ventilation in newborn. *Lancet*, ii: 11.

Melnick, G.; Pickoff, A.S.; Ferrer, P.L. et al. (1980):

Normal pulmonary vascular resistance and left ventricular hypertrophy in young infants with BPD: An echocardiographic and pathologic study. *Pediatrics*, 66: 589.

Melions, J.N.; Bove, E.L.; Dekeon, M.R. et al. (1991):

HFJV improves cardiac function after the fontan procedure. *Circulation*, 84 (Suppl. III); 364.

Meliones, J.N.; Chieftetz, I.M. and Wilson, B.G. (1995):

The use of airway graphic analysis to optimize mechanical ventilation strategies. Palm. Spring, CA, Products Corporation.

Meliones, J.N. and Hansell, D.R. (1994):

Extracorporeal membrane oxygenation: The role of blood components. In Chambers, L.A., Issett, L.A. (eds.): Supporting the pediatric transfusion Recipient. Bethesda. M.D., American Association of blood banks.

Melions, J.N.; Martin, L.D.; Barnes, S.D.; Wilson, B.G. and Wetzel, R.C. (1995):

Respiratory support. In Nicholas, D.G. et al. (eds) Critical heart disease in infants and children, St. Louis, CV, Mosby.

Meliones, J.N.; Mohler, F.W. et al. (1993):

Hemodynamic instability after the initiation of ECMO: Role of ionized calcium. *Crit. Care, Med.*, 10: 1247.

Melions, J.N.; Snyder, R.; Dekeon, M. et al. (1992):

Effect of ventilation on diastolic filling after cardiac surgery. *J.A.C.C.*, 19: 52a.

-
- Meliones, J.N.; Wilson, B.G.; Leonard, R.L. and Spitzer, A.R. (1995):**
Pathophysiology based approach to neonatal respiratory care.
In Dantzker, D.R.; MacIntyre, N.R.; Bakow, E.D. (eds):
Comprehensive Respiratory Care. Philadelphia, W.B.; Saunders.
- Merdith, K.S. (1995):**
High frequency ventilation: In Barnhart, S.L. and Czervinske,
M.P., (eds): Perinatal and Pediatric respiratory care. W.B.
Saunders Company, 20: 354.
- Merenstein, G.B. and Gardner, S.L. (eds.) (1989):**
Hand book of neonatal intensive care 2nd ed., C.V. Mosby Co.,
PP. 365-426.
- Miller, M.J.; Difiore, J.M.; Strohl, K.P. and Martin, R.J. (1990):**
Effect of nasal CPAP in supraglottic and total pulmonary
resistance in preterm infants.
J. Appl. Physiol., 68: 141-146.
- Miller, R.W.; Pollack, M.M.; Murphy, T.M. and Fink, R.J. (1986):**
Effectiveness of CPAP in the treatment of bronchomalacia in
infants a bronchoscopic documentation. Crit. Care, Med. 14 (2):
125-127.
- Milligan, D.W.A.; Issler, H.; Massam, M. and Reynolds, E.O.R. (1984):**
Treatment of neonatal pulmonary interstitial emphysema by
lung puncture, Lancet I, 1010-1011.
- Minton, S.; Gerstmann, D. and Steddard, R. (1995):**
Early intervention in RDS. Cardiopulmonary Review: current
Applications and Economics, Yorba Linda, CA, Sensormedics,
Critical Care.
- Mishoe, S.C.; Brooks, C.W.; Dennison, F.H.; Hill, K.V. and Frye, T.
(1995):**
Octave waveband analysis to determine sound frequencies and
intensities produced by nebulizers and humidifiers used with
hoods. Respiratory Care, 40: 1120.
- Mizuno, K.; Takeuchi, T.; Itabashi, K. et al. (1994):**
Efficacy of SIMV on weaning neonates from the ventilator.
Acta. Paediatr. Jpn., 36: 162.
-

-
- Moa, G. and Nilsson, K. (1993):**
Nasal CPAP: experiences with a new technical approach. *Acta. Pediatr.*, 82 (2): 210-211.
- Mochalova, L.D.; Khodav, D.A. and Zhukova, T.P. (1993):**
Cerebral circulation control in healthy full-term neonates. *Acta. Pediatr. Scand. Supplementum*, 311: 20.
- Moncada, S.; Palmer, R.M.J. and Higgs, E.A. (1991):**
Nitric oxide: physiology, pathophysiology and pharmacology, *Pharmacol. Rev.*, 43: 109-142.
- Montogia, J.P.; Merz, S.I. and Bartlett, R.H. (1991):**
A standardized system for describing flow/pressure relationships in vascular access devices. *Trans. Am. Soc. Artif. Internal Organ*, 34: 4-8.
- Moore, G.C. (1984):**
CPAP in Levin, D.L., Morriss, F.C. and Moore, G.C. (eds.): *A practical guide to pediatric intensive care*, ed. 2, C.V. Mosby Co. 94: 542.
- Moore, G.C. and Perkin, R.M. (1984):**
Assisted ventilation. In Levin, D.L.; Morriss, F.C. and Moore, G.C. (eds.): *A practical guide to pediatric intensive care*, ed. 2, C.V. Mosby Co. 97: 545.
- Morrery, J.P. et al. (1982):**
Improvement in lung mechanics as a function of age in the infant with severe BPD. *Pediatr. Res.*, 16: 290.
- Moskowitz, G.D.; Shaffer, T.H. and Dubin, S.E. (1973):**
Liquid breathing trials and animal studies with ademonad-regulated liquid breathing system. *Med. Instrum.*, 9: 28-33.
- Motoyama, E.K.; Fort, M.D.; Klesh, K.W. et al. (1987):**
Early onset of airway reactivity in premature infants with BPD. *Am. Rev. Res. Dis.*, 136: 50-57.
- Moulton, S.L.; Lynch, F.B.; Cornish, J.D. et al. (1991):**
Carotid artery reconstruction following neonatal extracorporeal membrane oxygenation. *J. Pediatr. Surg.*, 26: 794-799.

Narula, A. and Raju, T.N.K. (1985):

BPD pathophysiology and management. In: Vidyasager, D. and Sarnik, A.P. (eds): Neonatology and pediatric intensive care. PSG Publishing Company, Inc. Littleton, Massachusetts, 24: 314.

Neijens, H.J.; Kerrebijn, K.F. and Smalhout, B. (1978):

Successful treatment with CPAP of two infants with bronchomalacia. Acta Pediatr. Scand., 67: 293-296.

Nelin, L.D. and Hoffman, G.M. (1998):

The use of inhaled nitric oxide in a wide variety of clinical problems. Pediatric clinics of North America, Vol. 45, No. 3, P. 531.

Nelin, L.D.; Moshin, J.; Thomas, C.J. et al. (1994):

The effect of inhaled nitric oxide on the pulmonary circulation of neonatal pig: Pediatri. Res., 35: 20-24.

Nellcor, (1992):

Miniaturized mainstream capnography: Nellcor's ultra cap pulse oximeter and capnograph. Capnography Reference Note Number 4, Pleasanton, CA.

Neonatal Inhaled Nitric Oxide study group (1997):

Inhaled nitric oxide in full-term and nearly full-term infants with hypoxic respiratory failure. N. Engl. J. Med., 336: 597.

Newman, J.H.; Loyd, J.E.; English, D.K. et al. (1983):

Effects of 100% oxygen on lung vascular function in awake sheep. J. Appl. Physiol., 54: 1379.

Newth, C.J. (1979):

Recognition and management of respiratory failure pediatric. Clin North Am., 26: 617-643.

Nichols, P.E. and Rogers, M.C. (1987):

Developmental physiology of the respiratory system. In Rogers, M.C. (eds.): Textbook of pediatric intensive care, Williams and Wilkins, 4: 88-112.

-
- Normand, I.C.S.; Olver, R.E.; Reynolds, E.O.R. et al. (1971):*
Permeability of lung capillaries and alveoli to men-electrolytes in the fetal lamb. *J. Physiol. (London)*, 219: 330.
- North, A.J.; Moya, F.R.; Mysore, M.R. et al. (1995):*
Pulmonary endothelial nitric oxide synthase gene expression is decreased in a rat model of congenital diaphragmatic hernia. *Am. J. Respir. Cell Mol. Biol.*, 13: 676-682.
- Northway, W.H.Jr. (1992):*
An introduction to bronchopulmonary dysplasia. *Clinics in perinatology*, Vol. 19 (No 3): PP. 489-496.
- Northway, W.H.Jr.; Rosan, R.C. and Porter, D.Y. (1967):*
Pulmonary disease following respiratory therapy of hyaline membrane disease, *N. Engl. J. Med.*, 276: 357.
- Nussbaum, E. (1997):*
Respiratory failure. In Nussbaum, E. (ed.): *Pediatric intensive care 3rd*, Futura Publishing Company, Inc. 19: 349.
- Ogata, E.S.; Gregory, G.A.; Kitterman, T.A.; Phibbs, R.H.; Tooley, W.. (1976):*
Pneumothorax in respiratory distress syndrome. Incidence and effect on vital signs, blood gases and pH. *Pediatrics*, 58: 177-183.
- Ogawa, Y.; Miyasaka, K.; Kawano, T. et al. (1993):*
A multicenter randomized trial of high frequency oscillatory ventilation as compared with conventional mechanical ventilation in preterm infants with respiratory failure. *Early Hum. Dev.*, 32:1.
- Ormazabale, M.; Kirkpatrick, B. and Muller, D. (1980):*
Alteration of alveolar-arterial oxygen gradient (A-a) DO₂ in response to tolazoline as a predictor of outcome in neonates with PPH. *Pediatr. Res.*, 14: 607.
- Ortega, M.; Ramos, A.; Atkinson, J. et al. (1987):*
Oxygenation index can predict outcome in neonates who are candidates for ECMO. *Pediatr. Res.*, 34: 39.
-

-
- Ortiz, R.M.; Cilley, R.E. and Bartlett, R.H. (1987):**
ECMO in pediatric respiratory failure. *Pediatr. Clin. North Am.*, 34: 39.
- Otsu, T.; Merz, S.I.; Holtquist, K.A. et al. (1989):**
Laboratory evaluation of double lumen catheter for venovenous neonatal ECMO. *Trans. Am. Soc. Artif. Internal Organ*, 35: 647-650.
- Pagtakhan, R.D. and Chernick, V. (1983):**
Intensive care for respiratory disorder. In Kendig, E.L. and Chernick, V. (eds.) *disorders of respiratory tract in children*, ed. 4, W.B. Saunders, 8: 145-165.
- Palmer, R.M.J.; Ferrige, A.G. and Moncada, S. (1987):**
Nitric oxide release accounts for the biological activity of endothelium derived relaxing factor. *Natur.*, 327: 524.
- Paranka, M.S.; Clark, R.H.; Yoder, B.A. et al. (1995):**
Predictors of failure of HFOV in term infants with severe respiratory failure. *Pediatrics*, 95: 400.
- Patel, C.A. and Klein, T.M. (1995):**
Outcome of infants with birth weights less than 1000g with respiratory distress syndrome treated with high frequency ventilation and surfactant replacement therapy. *Arch. Pediatr. Adolesc. Med.*, 149: 317.
- Pascoe, D.J. (1984):**
Evaluation of the child in the emergency room. In: Pascoe, D.J. and Grossman, M. (ed.): *Quick Reference to Pediatric emergencies*, ed. 3, J.B. Lippincott, Co. 2:2.
- Perkin, R.M. and Levin, D.L. (1991):**
Adverse effects of PPV in children. In Gregory, G.A. (ed.) *Respiratory failure in the child*. Churchill Livingstone, 10: 163.
- Perlman, J.M.; Goodman, S.; Kreusser, K.L. et al. (1985):**
Reduction in intraventricular hemorrhage by elimination of fluctuating cerebral blood-flow velocity in preterm infants with RDS. *N. Engl. J. Med.*, 312: 1353.
-

-
- Perlman, J.M. and Volpe, J.J. (1989):**
Movement disorder of premature infants with severe BPD: A new syndrome. *Pediatrics*, 84: 215.
- Pernoll, M.L. et al. (1986):**
Diagnosis and management of the fetus and neonates at risk, a guide for team care, St. Louis, The C.V. Mosby, Co.
- Phillips, B.L.; Mcquitty, J. and Drand, D.J. (1996):**
Blood gases: technical aspects and interpretation in Goldsmith, J.P. and Karotkin, E.H. (eds): Assisted ventilation of the neonate, 3rd, Philadelphia, W.B. Saunders Co. 11: 213.
- Pilchak, T.M. (1992):**
Anesthesia and monitoring equipment for pediatrics, CRNA: Clin. Forum. Nurse Anesthesia, 3: 64.
- Pitt, B.R.; Lister, G. and Gillis, N.C. (1987):**
Hemodynamic effects on lung metabolic function. In Ryan, U.S. (ed.) pulmonary endothelium in health and disease (lung biology in health and disease, Vol. 32) New York Marcel Dekker, P. 65.
- Pokora, T.; Bing, D.; Mammel, M. et al. (1983):**
Neonatal high frequency jet ventilation. *Pediatrics*, 72:27.
- Primaiano, F.P.; Chatburn, R.L. and Lough, M.D. (1982):**
MAP: Theoretical considerations. *Crit. Care Med.*, 19: 378-382.
- Primhak, R.A. (1983):**
Factor associated with pulmonary air leaks in premature infant receiving mechanical ventilation. *J. Pediatr.*, 102: 764.
- Puybasset, L.; Stewart, T.; Rouby, J. et al. (1994):**
Inhaled nitric oxide reverses the increase in pulmonary vascular resistance induced by permissive hypercapnia in patients with acute respiratory distress syndrome. *Anesthesiology*, 80: 1254.
- Quattromani, F.L.; Foley, L.C.; Bowen, A.D.; Weisman, L. and Hernandez, J. (1981):**
Facial relationship of thymus: Radiologic pathologic correlation in neonatal pneumomediastinum. *AJR*, 37: 1209.
-

Raffin, T.A.; Simon, L.M. and Braun, D. (1980):

Impairment of phagocytosis by moderate hyperoxia (40-60% oxygen) in lung macrophages. *Lab. Invest.*, 42: 622.

Ramin, K.D.; Leveno, K.J.; Kelly, M.A. et al. (1996):

Amniotic fluid meconium: A fetal environmental hazard. *Obstet. Gynecol.*, 87: 181.

Ranieri, V.M.; Grasso, S.; Mascia, L. et al. (1997):

Effects of proportional assist ventilation on inspiratory muscle effort in patients with chronic obstructive pulmonary disease and acute respiratory failure. *Anesthesiology*, 86: 79.

Rao, J.P.; Manno, B.L. and Robertson, A.F. (1978):

Usefulness of CPAP in differential diagnosis of cardiac from pulmonary cyanosis in newborn infants. *Arch. Dis. Child*, 53: 456.

Redmond, C.R.; Graves, E.D.; Falterman, K.W. et al. (1987):

Extracorporeal membrane oxygenation for respiratory and cardiac failure in infants and children. *J. Thorac. Cardiovasc. Surg.*, 93: 199.

Redmond, C.R.; Goldsmith, J.P.; Sharp, M.J. et al. (1986):

Extracorporeal membrane oxygenation for newnates. *J. La. State Med. Soc.*, 138: 40.

Rennie, J.M.; South, M. and Morley, C.J. (1987):

Cerebral blood flow velocity variability in infants receiving assisted ventilation. *Arch. Dis. Child*, 62: 1247.

Reynold, E.O.R. and Taghizadeh, A. (1974):

Improved prognosis of infants mechanically ventilated for HMD. *Arch. Dis. Child*, 49: 505.

Rhodes, P.G.; Graves, G.R.; Patel, D.M.; Campbell, S.B. and Blumenthal, B.I. (1983):

Minimizing pneumothorax and bronchopulmonary dysplasia in ventilated infant with hyaline membrane disease. *J. Pediatrics*, 103: 634.

Robert, J.D.; Polaner, D.M.; Lang, P. et al. (1992):

Inhaled nitric oxide in PPHN. *Lancet*, 340: 818-819.

Roberts, R.J.; Weesner, K.M. and Bucher, T.R. (1983):

Oxygen induced alterations in lung vascular development in the newborn lung. *Pediatr. Res.*, 17: 368.

Roberton, N.R.C. (1993):

Disorders of the respiratory tract. In Roberton, N.R.C. (ed.) *Neonatal intensive care*, ed. 3, Edward Arnold 10: 77-162.

Rossant, R.; Flake, K.J.; Lopez, F. et al. (1993):

Inhaled nitric oxide for adult respiratory distress syndrome. *N. Engl. J. Med.*, 328: 399-405.

Rosenfeld, W.; Evans, H.; Concepcion, L. et al. (1984):

Prevention of BPD by administration of bovine superoxide dismutase in preterm infants with RDS. *J. Pediatr.* 105: 781-785.

Rossaint, R.; Gerlach, H.; Schmidt-Ruhnke, H. et al. (1995):

Efficacy of inhaled nitric oxide in patients with severe ARDS. *Chest*, 107: 1107-1115.

Rossi, C.; Nascimento, S.D.; Femanda, M. et al. (1997):

Should obstetricians clear the airways of the newborn infants with meconium stained amniotic fluid (MSAF)? *Pediatr. Res.*, 41: 173 A.

Roussos, C. and Macklem, P.T. (1992):

The respiratory muscles. *N. Engl. J. Med.*, 307: 786-797.

Rush, M.G. and Hazinski, T.A. (1992):

Current therapy of BPD. *Clinics in perinatology*, Volume 19 (No. 3): PP. 563-590.

Rush, M.g.; Engelhardt, B.; Parker, R.A. et al. (1990):

Double blind, placebo-controlled trial of alternative day furosenide therapy in infant with chronic BPD. *J. Pediatr.*, 117: 112-118.

- Ryan, C.A.; Finer, N.N. and Peters, K.L. (1989):**
Nasal intermittent positive pressure ventilation offers no advantages over nasal CPAP in Apnea of prematurity. *AJDC*, 143 (10): 1196-1198.
- Saga, S.; Modell, J.H.; Calderwood, H.W. et al. (1973):**
Pulmonary function after ventilation with fluorocarbon liquid P12-F (caroxin-F). *J. App. Physiol.*, 34: 160-4.
- Saldanha, R.L.; Cepeda, E.E. and Poland, R.L. (1982):**
The effect of vitamin E prophylaxis on the incidence and severity of BPD. *J. Pediatr.*, 117: 112-118.
- Salman, N.H.; Fuhrman, B.P.; Papo, M.L. et al. (1993):**
Oxygenation and lung mechanics during 24 hours trials of perfluorocarbon associated gas exchange (PAGE) in pig lets. *Pediatr. Res.*, 33: 40A.
- Sargent, J.W. and Seffl, R.J. (1970):**
Properties of perfluorinated liquid. *Fed. Proc.*, 29: 1699-1703.
- Sarnaik, A.P. (1985):**
Respiratory emergencies in children. In Vidyasagar, D. and Sarnaik, A.P. (eds.) *Neonatal and Pediatric intensive care*, PGS, Publishing Co. 4: 58.
- Scanalan, C.L. and Thalken, R. (1995):**
Medical gas therapy. In Scanalan, C.L. Spearman, C. and Sheldon, R.L. (eds.): *Egan's fundamentals of respiratory care*, 6th ed. St. Louis-Mosby, Year book.
- Scanlon, J.W. (1994):**
The very low birth weight infant. In Avery GB, Fletcher MA, MacDonald MG (eds): *Neonatology: Pathophysiology and management of the newborn*, 4th ed. Philadelphia, JB Lippincott.
- Schmidt-Nowara, W. (198):**
Continous positive airway pressure for long term treatment of sleep apnea. *Am. J. Dis. Child*, 138: 82.
- Schreiner, R.L. and Kisling, J.A. (1997):**
Practical neonatal respiratory care, New York, Ravin Press.

Schumacher, R.E.; Barks, J.D.E.; Johnston, M.J. et al. (1988):

Right sided brain lesions in infants following ECMO.
Pediatrics, 82: 755-760.

Sedin, G. (1991):

CPAP and mechanical ventilation. Int. J. Technol. Assess Health Care, 7 (Suppl. 1), 31-40.

Seely, J.F. and Zagelbaum, G.L. (1996):

Arterial blood gas analysis. In Zagelbaum, G.L. and Pare, J.A.P. (eds.): Manual of acute respiratory care, Boston, Little, Brown, 2: 19.

Shabarek, F.M.; Cocanour, C.S.; Xue, H. et al. (1996):

Meconium stimulates alveolar macrophage prostaglandin E2 production but not procoagulant activity. Pediatr. Res., 39: 49A.

Shaffer, T.H.; Greenspan, J.S. and Wolfson, M.R. (1994):

Liquid ventilation. In Boynton, B.R. (ed): New therapies for neonatal respiratory failure, a physiologic approach. Cambridge, New York, 279-301.

Shaffer, T.H. and Wolfson, M.R. (1996):

Liquid ventilation: An alternative ventilation strategy for management of neonatal respiratory distress. Eur. J. Pediatr., 155 (Suppl.): 30.

Shaffer, T.H.; Wolfson, M.R.; Greenspan, J.S. et al. (1992):

Animal models and clinical studies of liquid ventilation in neonatal respiratory distress syndrome. In Robertson, B. (ed): Surfactant in clinical practice. Harwood Academic, Switzerland, 187-198.

Shannon, D.C. and Epstein, M. (1986):

Bronchopulmonary dysplasia. In Thibeault, D.W. and Gregory, G.A. (eds.) Neonatal Pulmonary care, ed. 2, Norwalk, Conn. Appleton-Century-Crofts, 28: 697.

Shapiro, B.A.; Harrison, R.A. and Walton, J.R. (1982):

Clinical application of blood gases ed. 3, Chicago, Year book Medical Publishers Inc.

- Shenai, J.P.; Chyti, L. and Stahlman, M.T. (1985):*
Vitamin A status of neonates with BPD. *Pediatr. Res.*, 19: 185-188.
- Shenai, J.P.; Kennedy, K.A.; Chytill, F. et al. (1987):*
Clinical trial of vitamin A supplementation in infants susceptible to BPD. *J. Pediatr.*, 111: 269-277.
- Shenai, J.P.; Rush, H.G.; Stahlman, M.T. et al. (1990):*
Plasma-retinol-binding protein response to vitamin A administration in infants, susceptible to BPD. *J. Pediatr.*, 116: 607-614.
- Simbrauner, G. and Gregor, G.A. (1981):*
Performance of the neonatal ventilators: the effect of changes in resistance and compliance. *Crit Care Med.*, 9: 509.
- Siplovich, L.; Ban-Ziv, J.; Karplus, M.; Mares, A.J. (1979):*
The pericardial "window" A rare etiological factor in neonatal pneumopericardium. *J. Pediatr.*, 94: 75.
- Smith, R.A.; Venus, B.; Wier, D.D.; Mathru, M.; Masood, S.; Goldstein, J.D. et al. (1987):*
Morphometric changes in a dog model of ARDS after early therapy with CPAP. *Resp. Care*, 32 (7): 525-533.
- Sobonya, R.E.; Logvinoff, M.M.; Taussig, L.M. and Theriault, A. (1982):*
Morphometric analysis of the lung in prolonged BPD, *Pediatr. Res.*, 16: 969.
- Sosenko, I.R.S. and Frank, Z. (1991):*
Oxidants and antioxidants. In Cherniak, N. and Mellins, R.B. (eds.): *Basic mechanisms of pediatric respiratory disease: Cellular and integrative*. Philadelphia, BC Decker, P. 315.
- South, M. and Morley, C.J. (1986):*
Synchronous mechanical ventilation of the neonate. *Arch. Dis. Child*, 61: 1190.
- Speidel, B.D. and Dunn, P.M. (1975):*
Effect of CPAP on breathing pattern of infants with RDS. *Lancet*, 1 (7902): 302-304.

Spitzer, A.R. (1996):

High frequency jet ventilation: In Spitzer, A.R., Intensive care of the fetus and neonate. St. Louis Mosby-Yearbook Inc., 49: 571.

Spitzer, A.R. and Fox, W.W. (1996):

Positive pressure ventilation: Pressure limited to time cycled ventilators: In Goldsmith, J.P. Karotkin, E.H. (eds.): Assisted ventilation of the neonate, 3rd Philadelphia, W.B. Saunders Co., 8: 167.

Spitzer, A.R.; Greenspan, J.S.; Antunes, M.J. et al. (1996):

Special ventilatory techniques II: Liquid ventilation, Nitric oxide therapy and negative pressure ventilation. In Goldsmith, J.P. and Karotkin, E.H. (eds.): Assisted ventilation of neonates ed. 3. Philadelphia: W.B. Saunders, P. 229.

Stark, A.R. (1991):

Apnea, In Cloherty JP, Stark AR (eds): Manual of neonatal care 3rd ed., Little. Brown and Company, PP. 457-467.

Stern, L.; Ramos, A.; Outerbridge, E.W. and Beaudry, P.H. (1970):

Negative pressure artificial respiration: Use in treatment of respiratory failure of the newborn. Can Med. Assn. J., 102: 595.

Stewart, A.R.; Finer, N.N. and Peters, R.L. (1981):

Effects of alterations of inspiratory and expiratory pressures and I:E ratios on MAP, blood gases and intracranial pressure. Pediatrics, 67: 474-481.

Stocker, J.T. (1988):

Pathology of acute BPD. In Bancalari, E. and Stocker, J.T. (eds.): BPA Washington, D.C., Hemisphere, PP. 237-278.

Stoddard, R.; Meredith, K.; Minton, S. et al. (1993):

Treatment of congenital diaphragmatic hernia with high frequency oscillatory ventilation. Pediatr. Pulmonol., 15: 367.

Sumner, E. (1990):

Artificial ventilation of children. In Dinwiddie, R. (ed.): The diagnosis and management of pediatric respiratory disease. Churchill Livingstone, 14: 266.

- Suter, P.M.; Fairley, H.B. and Isenberg, M.D. (1975):**
Optimum end-expiratory airway pressure in patients with acute pulmonary failure. *N. Engl. J. Med.*, 292: 284.
- Thibeault, D.W. (1986):**
Pulmonary barotrauma, Intrstitial emphysema, pneumomediastinum and pneumothorax. In Thibeault, D.W. and Gregory, G.A. (eds.): *Neonatal pulmonary care*, ed. 2, Norwalk, Conn. Appleton-Century-Crofts, 21: 499.
- Thompson, J.E.; Farrell, M.F. and McManus, M. (1992):**
Neonatal and pediatric airway emergencies. *Respir. Care*, 37 (6): 582-599.
- Tidona, R.J.; Nizami, A.A. and Cemansky, N.P. (1988):**
Reducing interference effects in the chemiluminescent measurement of nitric oxides from combustion systems. *Air Pollut Control Assoc.*, 38: 806.
- Toomasian, J.M.; Hsu, L.C.; Hirschl, R.B. et al. (1988):**
Evaluation of Duraflo II heparin coating in prolonged extracorporeal membrane oxygenation. *Trans. Am. Soc. Artif. Internal Organ*, 34: 410-414.
- Toro-Figueroa, L.O. and Brink, L.W. (1992):**
Respiratory disease. In Toro-Figueroa, Levin, D.L. and Morriss, F.C. (eds.): *Essentials of pediatric intensive care manual* uality medical Publishing, Inc., 8: 227.
- Torosian, M.B.; Statter, M.B. and Arensman, R.M. (1996):**
Extracorporeal membrane oxygenation: In Goldsmith J.P. and Karotkin, E.H., (eds.): *Assisted ventilation of the neonate*, 3rd, Philadelphia, W.B. Saunders, Co. 13: 241.
- Traverse, J.H.; Korvenranta, H.; Adams, E.M. et al. (1989):**
Cardiovascular effects of high frequency oscillatory and jet ventilation. *Chest*, 96: 1400.
- Trug, W.E. (1996):**
High frequency oscillatory ventilation: In Spitzer, A.R.; *Intensive care of the fetus and neonate*, St. Louis-Mosby Year book Inc., 50: 584.

Trug, W.E. and Jackson, J.C. (1992):

Alternative modes of ventilation in the prevention and treatment of BPD. In Holtzman, R.B., Frank (eds.): BPD, Clin. Perinatol., 19: 621-643.

Tsuno, K.; Prato, P. and Koloblow, T. (1990):

Acute lung injury from MV, at moderately high airway pressures. J. Appl. Physiol., 69: 956.

Valls-Soler, A.; Wolfson, M.R.; Kenchner, N. et al. (1993):

Comparison of natural surfactant and brief liquid ventilation rescue treatment in very immature lambs. Pediatr. Resp., 33: 347A.

Van Marter, L.J.; Leuinton, A.; Allred, E.N.; Pagano, M. and Kuban, K.C.K. (1990):

Hydration during the first day of life and the risk of BPD in LBW infants. J. Pediatr., 176: 942.

Van Meurs, K.P.; Mikesell, G.T.; Scale, S.R. et al. (1990):

Maximum blood flow rates for arterial cannulae used in neonatal ECMO. Trans. Am. Soc. Artif. Internal Organ, 36: 679-681.

Vidyasagar, D. (1975):

Assisted ventilation in infants with MAS.
Pediatrics, 56: 208.

Visveshwara, N.; Freeman, B.; Peck, M. et al. (1991):

Patient-triggered synchronized assisted ventilation of newborns: Report of preliminary study and three year experience, J. Perinatology, 11 (4): 347.

Vogt, J.F.; Gangitano, E.F.; Liberman, G.R. et al. (1989):

Cannulation of cephalad segment of internal jugular vein during ECMO in the neonate. Snowmass, Co., Children's National Medical Center.

Vohr, B.R.; Bell, E.F. and Oh, W. (1982):

Infant with BPD. Am. J. Dis. Chld, 736: 443.

Wilson, B.G. (1997):

Mechanical ventilation of infant and child. In: Aloia, C.A., Hill, T.V. (eds.): Respiratory care of newborn and child, ed. 2, J.B. Lippincott Company, 16: 311-354.

Wilson, B.G. (1995):

Direct measurement via an inline pneumotach is necessary to determine effective tidal volume in children. Respir. Care, 40 (11): 1172.

Wilson, B.G. (1997):

High frequency ventilation: in Aloia, C.A., Hill, T.V., (eds.): Respiratory care of newborn and child, ed. 2. J.B. Lippincott Company, 24: 488-502.

Wilson, J.M.; Bower, L.K.; Farkler, J.C. et al. (1993):

Aminocaproic acid decrease the incidence of intracranial hemorrhage and other hemorrhagic complications of ECMO. J. Pediatr. Surg., 28: 536.

Wilson, B.G.; Cheifetz, I.M. and Meliones, J.N. (1995):

Optimizing mechanical ventilation in infants and children. Palm. Springs, C.A. Bird. Products Corporation.

Wilson, B.G.; Chieftetz, I.M.; Meliones, J.N. (1996):

The use of airway graphic analysis to optimize mechanical ventilation in small infants and children. Palm. Springs, C.A., Bird Products, Corporation.

Wilson, B.G. and Meliones, J.N. (1996):

Continuous AGA reduces intrinsic PEEP during mechanical ventilation. Crit. Care Med., 24 (1): 128A.

Winger, I.M. and Mudge, G.H. (1985):

Diuretics and other agents employed in the mobilization of edema fluid. In Gilman, G.A., Goodman, L.S., Rall, T.W. (eds.): The pharmacological basis of Therapeutics. New York, Macmillan, PP. 887-907.

Winternitz, M.C. and Smith, G.H. (1920):

Preliminary studies in intratracheal therapy. In: Pathology of War gas poisoning Yale university Press, New Haven.

Zach, T.L.; Steinhorn, R.H.; Georgieff, M.K. et al. (1990):

Leukopenia associated with ECMO in the newborns.
J. Pediatr., 116: 440-443.

Zagelbaum, G.L. and Pare, J.A.P. (1996):

Acute ventilatory failure, In Zagelbaum, G.L. and Pare, J.A.P. (eds.): Manual of acute respiratory care, Boston, Little, Brown, 3: 57.

Zangelbaum, G.L. and Pare, J.A.P. (1996):

Initial assessment and emergency management of acute respiratory disorders. In Zagelbaum, G.L. and Pare, J.A.P. (eds.): Manual of acute respiratory care, Boston, Little, Brown, 1:1.

Zayek, M.; Cleveland, D. and Morin, F.C. (1993):

Treatment of PPHN lamb by inhaled nitric oxide.
J. Pediatr., 122: 743-750.

Zelinka, M.A.; Wolfson, M.R.; Calligar, S. et al. (1991):

Direct administration of gentamicin during liquid ventilation of the lamb: Comparison of lung and serum levels to IV administration. Pediatr. Res., 29: 290A.

Zwischenberger, J.B. and Cox, C.S. (1993):

Emergencies during ECMO and their management. In Arensman, R.B., Cornish, J.D. (eds.): Extracorporeal life support. Cambridge, M.A., Blackwell Scientific Publishing, PP. 207-225.