

- Abrams , E. J; Ginsberg , H. ; Grandy ,S.M. (1982):**
Metabolism of cholesterol and TG in non ketotic diabetes mellitus.
Diabetes; (31): 903-910 .
- Ali, K.A.(1989):**
Ketone bodies metabolism in Biochemistry for post graduate
1 st edition.
University Book Center. P86
- Alkinson,M.A.; Maclaren, N.K.; Sharp,D.W.and Riley,W. J.(1990):**
64,000 Mr autoantibodies as predictors of insulin dependent diabetes. Lancet 353:1357 –60
- Andersen,A.R.;Christiansen ,J.S.; Andersen,J.K.;Kreiner,S.& Deckert,T.(1983):**
Diabetic nephropathy in type I(insulin dependent)diabetes:an epidemiological study.
Diabetologia ; (25):496-501
- Allawi,J.and Jarrett,R.J.(1990):**
Microalbuminuria and cardiovascular risk factors in type II diabetes.
Diabet.Med.;6:115-8
- Allen DB,MacDonald MJ(1986):**
Artifacts in glycosylated hemoglobin values in Pediatric Patients.
J.Pediatr;109:695.

Bach,J.F.(1994):

Insulin-diabetes mellitus as an auto immune disease. Endocr.
Rev.15:516.

Bach,J.F.(1988):

Mechanism of auto immunity in Insulin Dependent Diabetes
Mellitus Clin.Exper immuno,72:1-8.

**Baerwinkle,E.,Turner,S.T.;Weinshilboum,R,Johnson,M.;Richelson,
E.and Sing,C.F.(1986):**

Analysis of the distribution of sodium lithium counter transport:
in a sample representative of the general population.
Gen Epidemiol;(3):365-78.

Blaw,E.B.&Mishaal,A.E.(1972):

Rat glomerular glycoprotein composition and metabolism in
amino nucleoside nephrosis.
Proc Soc Exp Biol (NY);(141):164-72

Bedi,H.K.;Vyas,B.R.;Bomb, B.S.;Agarwal M.L. and Bedi ,T.(1977):

Fibrinogen content and fibrinolytic activity of blood in diabetes
before and after anti diabetic drugs.
J.Ass.Phys. India.

Bertolatus, J.A.;Abuyousef,M.and Hansickar,L.G.(1987):

Glomerular sieving of high molecular weight proteinuric rats.
Kidney Int;(31):1257-66.

Bennett H.(1994):

Definition, Diagnosis, and Classification of Diabetes Mellitus and Impaired Glucose.

Tolerance Pediatric Clin North Am.;37:1467.

Betterle,C.;Presotto,F.and Margin,L.(1994):

The natural history of Pre -type I diabetes mellitus in patients with autoimmune endocrine diseases.

Diabetologia,37,95-103.

Bernard,B.A;Yamada,K.M.and Olden,K. (1982):

Carbohydrates selectivity protect a specific domain fibronectin against proteases.

J.Biol.Chem.; (257):8594-7.

Bodansky,H.J.,Cudworth,A.G.;Drury,P.L.and Kohner,E.M.(1982) :

Risk factors associated with severe proliferative retinopathy in insulin dependent diabetes mellitus.

Diabetes Care; (5):97-100.

Boniafacio,E.;Bingley,P.J.and Shattoc,K.M.(1990):

Quantification of islet cell antibodies and prediction of IDD.Lancet335:147-49.

Brenner,B.M.; Hostetter,T.H.and Humes H.D.(1978):

Molecular basis of proteinuria of glomerular origin.

New. Engl. J. Med.(298):826-33.

Brenner,B.M. (1985):

Nephron adaptation to renal injury or ablation.

Am.J. Physiol , 249: F 234-337.

Brown, D.M.;Michael,A.F.& Oemega ,T.R.(1981):

Glycosaminoglycan synthesis by glomeruli in vivo and vitro.

Biochem Biophys Acta; (674):96-104.

**Brancati,F.L.;Whittle,J.C.;Whelton ,P.K.; Seidler, A.J. &
Klage,M.J.(1992):**

The excess incidence of diabetic end stage renal disease among
Blacks.

JAMA.268:3079.

Burg,J.; Esser,R.J.E.;Paerels,G.B.,(1959):

Biochem et Biophys. Acta;(33):241

In Methods of Biochem Analysis. Vol 2,New York interscience ,pp
220-298-1961.

Camerini –Davlos & Cole. (1970) and Belfoire (1979):

In complication of diabetes text book of medicine by Harry, K. and Jarrett, J. 1982, P.225.

Castano ,L.; Ziegler ,A.G.; Shoelson,S. and Eisenbarth, G.S. (1993):

Characterization of insulin autoantibodies in relatives of patients with type I diabetes.

Diabetes, Vol.42, August, 1993, 1202 –1209.

Castano ,L. and Eisenbarth ,G.S. (1990):

Type I diabetes: a chronic auto immune disease of human, mouse and rat. Ann . Rev Immunol., 8:647-79.

Cair,F.A. (1968):

Diabetes and eye.

Oxford: Blackwell; P 84.

Chandramouli, V.& Carter , J.R. (1975):

Cell membrane changes in chronically diabetes rats.

Diabetes ; (24):257 –62.

Carter , A. & Martim , N.H. (1962):

Serum s.a. levels in health & disease.

J. Clin. Pathol ; (15):69 – 72

Chachati , A.;Frenckell , R.;Fordart-Willems, J.;Godon,J.P.and

Lefebvre , P.J. (1987):

Variability of albumin excretion in insulin dependent diabetics.

Diabet. Med ; (4): 441-5.

Chiggeri ,G.M.; Gandiino, G.;Delfino,G.& Queirolo,C.(1985):

Electrical charge of serum & urinary albumin in normal and diabetic humans.

Kidney Int. ; (28): 168-77.

Christiansen , J. (1988):

Early renal hyperfunction and hypertrophy in insulin DDM.

Boston, Martinus Nijhoff publishing ,P 157.

Chavers,B.M.; Bilous , R.W. ; Ellis , E.N. ; Steffes , M.W. and Mauer , S.M. (1989):

Glomerular lesions and urinary albumin excretion in type I diabetes

New. Engl. J. Med.; (320):966-70

Chiari , S.N. & Nath , N. (1984):

S.a. content and sialidase activity of polymorphonuclear leukocytes in D.M

Am Med Sci : 288:1-5

Cohen, A.H. ;Mampaso , F. & Amboni ,L. (1977):

Glomerular podocyte degeneration in human renal disease. An ultrastructural study.

Lab. Invest ;(37): 40-2.

Constam ,G.R. (1965):

Prognosis of diabetes mellitus.

Helv. Med . Acta ; (32): 287

Cohen , M. P. & Ku ,L. (1984) :

Inhibition of fibronectin binding to matrix components by non-enzymatic glycation.

Diabetes ; (33) :970: 4.

Coleman , D.L. ,Kuzana , J.E. and Leiter , E.H. (1990):

Effect of diet on incidence of diabetes in non obese diabetic mice .

Diabetes , 39: 432 –436.

Crook , M.A., Tutt , P.& Pickup, J.C. (1993):

Elevated serum s.a. concentration in NIDDM& its relation to. blood Pressure & retinopathy .

Diabetes Care ; (16) :1,57-60. (J an).

Crook , M. A. ; Tutt , P. ;Simpson , H. & Pickup – J.C. (1993):

Serum sialic acid and acute phase proteins in type I and type II diabetes mellitus.

Clin. Chem . Acta. 19; 219: (1-2):131-8 (June)

Dadone , M. ; Hasstedt , S.J.; Hunt ,S.C. ;Smith ,J. B;Ash,D.O. and

Williams, S.R. (1984) :

Genetic analysis of Sodium – Lithium counter transport in ten hypertension prone Kindreds.

Am. J. Med ; (17): 565 – 77.

Dahlquist G., Kallen B. (1992):

Maternal- child blood group in compatibility and other perinatal events increase the risk for early - onset type I (insulin-dependent) Diabetes mellitus. Diabetologia; 35:671-657.

Damsgaard, E.M.& Mogensen, C.E. (1986):

Microalbuminuria in elderly hyperglycaemia patients & controls. Diabet Med ; (3) :430 -5.

Davis, M.D., Myers , F.L. and Engerman, R.L. (1969):

Clinical observation concerning the pathogenesis of diabetic retinopathy. In Goldberg M. F. ,Fines S.L. (eds) symposium on the treatment of diabetic retinopathy public health service publ. no.1890.

Washington , D.C., US. Govt printing office & pp 7-22.

Deckert, M.; Feldt -Rasmussen , B.;Djurup ,R.and Deckert, M. (1988):

Glomerular size and charge selectivity in insulin dependent diabetes mellitus.

Kidney Int; (33) : 100-6.

Deckert , T.; Feldt - Rasmussen , B.;Broch- Johnsen ,K.;Jensen,T.&Kofoed- Enevoldsen A.(1989):

Albuminuria reflects widespread vascular damage. The stenohypothesis.

Diabetologia ; (32): 219-26.

Dean , B. M. ;Becker ,E. ;McNally , J.M.;Tarn, A.C.;Gale ,E.A.M.;
Bottazo, G.F. (1986):

Insulin auto antibodies in the pre diabetic period: Correlation with
 islet cell antibodies and development of diabetes. Diabetologia 29
 :339-42.

Derby, L.; Laffel , L.B.M.& krolewski ,A.S.(1988):

Risk of diabetic nephropathy declines with age in type I(insulin-
 dependent) diabetes.
 Diabetologia ; (31):485 A.

Dixon ,H.B. (1983):

Terminology for s.a.
 Biochem.J.; (215):711.

Eisnbarth, G.S.;Ziegler, A.G.;Colman , P.A. (1994):

Pathogenesis of IDDM in Joslin 's Diabetes Mellitus. Kohn CR
 and Weir G.C. (eds) , 13 th edition Philadelphia, Hong Kong
 ,London , Sydney Tokyo, P. 216.

Eisenbarth , G.S. (1986):

Type I diabetes mellitus , A chronic auto immune disease .N.
 Engl.J. Med.314: 1360-68 .

Elisabeth ,R. (1990):

Diabetic nephropathy.
 Diabetographia ; (5):4.

Emmelot , P. (1973):

Biochemical properties of normal and neoplastic cell surfaces ; a review.

Eur. J. cancer ; (9): 319-333.

Eylar , E.H. (1985):

On the biological role of glycoproteins.

Journal of theoretical Biology.10,89.

Falk, R.J. , Sateinman ,J.L.& Minier ,S.M. (1983):

Polyantigenic mesangium in b.m. constituents in diabetic nephropathy .

Diabetes; (32), (Supl. 2):34-9.

Fajans ,S.S. (1990):

Classification and diagnosis of diabetes in :Diabetes Mellitus, Theory and practice, 4th edition . Rifkin H. and porte D.J. (eds)
New york ,El sevier ,P. 396-51.

Fajans ,S.S. (1985) :

Advance in Experimental medicine and biology veronic , M (ed)
New York Plenum publishing Comp , P. 65-87.

Fajans ,S.S. (1982):

In the Genetics of Diabetes Mellitus , Serones symposium No.47 –
Koblierling ,J.and Tottersall ,T. (eds). New York , Academic Press

,pp .251-650.

Farquhar ,M.G. (1981):

The glomerular b.m. : a selective macromolecular filter. In Hay
ED (ed.) Cell biology of extracellular matrix.
New York: Plenum pp 335-78.

Feldt-Rasmussen , B. and Mathiesern ,E.R. (1984):

Reliability of urinary albumin excretion in diabetic nephropathy.
Diabet. Nephropathy ; (3):101-3.

Foster , D. W. (1989):

Insulin resistance – a secret killer?
New. Engl. Med ; (320):733-4.

Freednan ,P.; pelers ,J.H. and Kark ,P.N. (1960):

Localization of gamma globulin in the diseased kidney.
Archs Intern.Med.; (105):525-535.

Gallschalk , A. (1955):

N-acetyl neuraminic acid biochemistry.
Nature , 176 , 881

Gamboro ,G. ; Baggio , B. &Crepaldi , G. (1988):

Abnormal erythrocyte charge in diabetes mellitus . Link with
microalbuminuria:
Diabetes ; 37 (6):745-8.

Ganrot ,P.O.; Gydelli ,K.& Ekelund,H.(1967):

Serum concentration of alpha-2 macroglobulin , haptoglobulin&
alpha-1 antitrypsin in diabetes mellitus.
Acta Endocrinol ;(55):537 – 544.

Gasic, G. & Gasic , T.(1962):

S.a. as a tumour marker in sera of some tumor –bearing
individuals
Nature , Lond. ,196-207.

Gavela , M. ; Lipavac – V and Crenek , K.S. (1990):

Total & ganglioside – bound sialic acid content of lymphocytes
from non- insulin dependent diabetic patients.
Acta – diabetol –Lat .; 27 (4):357-64.

Grant , M. ; Russel , B. ; Fitzgerald ,C. & Merimee ,T.J. (1980):

Insulin- like growth factors in vitreous. Studies in control and
diabetic subjects with neovascularization.
Diabetes ; (35):416-20.

Green , A. , Gale , E. A.M. , and Patterson , C.C. (1992):

Incidence of childhood – onset insulin- dependent diabetes
mellitus. Lancet 319: 905.

Habel M A (1993) :

Classification of diabetes mellitus Lancet 1:57.

Harrison ,R. ; Connolly, R,C. and Abdalla, A. (1969):

Intercellular contents : Cytology in Gray 's anatomy 36th ed. :,7.

Longman Groub Limited. Churchill Lvingstone Britain. Nature,
Lond., 224.

Hassab A. , Abdel-Kader E. Sherif A., and Hassan M (1988):

Diabetes Mellitus among school children in Alexandria. Bullet
High Institute Bullet high institute Public health , 18:193

Hayes , G.R.& Lockwood,D.H.(1986)

The role of cell surface sialic acid in insulin receptor function &
action.

J.Biol Chem;261: 2791-98.

Heinrich ,P.C.; Catell,J.V.& Andus,T.(1990):

Interleukin 6 & acute phase response.

Biochem J.:(265): 2791-98.

Hogemann, B.:Voss , B. &Altenwerth , f.J. (1986):

Concentration of collagen and laminin in sera of patients with
diabetes mellitus.

Klin Wochenschr; (64) :382-5.

**Holthofer ,H. ; Vitanen ,I., Pettersson , E.;Alfthan, O;linder, E.and
Miettinen ,A. (1981) :**

Lectins as fluorescence microscopic markers for saccharidase in
the human kidney.

Lab Invest.; (45): 391-399.

Hostetter ,T.H. (1992) :

Diabeti nephropathy (Metabolic vs haemodynamic consideration)
Diabetic Care; (9):1205.

Hostetter,T.H.;Rennke,H.G. and Brenner,B.M. (1982):

The case for intrarenal hypertension in the initiation and
progression of diabetic and other glomerulopathies.
An J Med ; (72): 375-80.

Inda-Ritsumin, S. ;Tomura, S.and Tomitak , T. (1990):

Endothelin in glomerulonephritis.
XIth international congress of nephrology, Tokyo, Japan,
Abstracts:5A.

**Inommata , S.;Nakamoto ,Y.,Inove,M.;Itoth. M.;Ohsawa,Y.and
Masamune,O. (1989):**

Relationship between urinary albumin excretion and renal
histology in NIDDM: with reference to the clinical significance of
microalbuminuria.
J.Diabet.Comp, (3) : 178-88.

Inoue , W.M.&Yagame,M.(1985):

Multi organ analysis of acute phase reactant proteins in patients
with diabetic nephropathy.
Diabetic nephropathy;(4):176-79.

Irvin , W.J.(1986):

Immunologic factors in diseases in : Macleode, J. (ed.) Davidson's
and practice of Medicine 14th ed. Churchrll Livingstone ,

Edinburgh , P. 24.

Jackson , R.; Vardi ,P. and Eisenborth , G.S. (1988)

Dual parameter linear model for prediction of onset of type I diabetes in ICA positive relatives.

Clin.Res. 36:585.

Jacobs , K.;Shownaker ,C.;Rudersdorf ,R. (1985):

Role of the kidney in erythropoiesis.

Nature; (313):806.

Janka,H.U.;Warran, J.H.& R and ,L.I. (1989):

Risk factors for progression of background retinopathy in long-standing insulin- dependent diabetes mellitus.

Diabetes ; (38):460-4

Jarrett ,R.J. (1988):

Is insulin an atherogenic ? (eitorial)

Diabetologia; (31):71-5.

Jarrett,R.J. (1984):

The epidemiology of coronary heart disease and related factors in the context of diabetes mellitus and impaired glucose tolerance in:

Jarrett R.J.,ed. Diabetes and Heart Disease.

Amsterdam: elsevier Scince Publishers B.V, : 1:23.

Jerneld, B. and Algvere ,P.:(1987):

Proteinuria & blood glucose levels in a population with diabetic

retinopathy.

Am. J. ophthalmol.;(104):283-389.

**Jensen ,T.;Borch-Jonsen, K.;Kofaed-Enevoldsen, A; and Deckert ,
T. (1987):**

Coronary heart disease in young type I (insulin-dependent)
diabetic patients with and without nephropathy: incidence and risk
factors.

Diabetologia ;(30):144-8.

**Jones,S.L.; Faria ,J; Tariq ,T.; Mattock , M.B. and Viberti ,G.C.
(1989):**

Sodium-Lithium counter transport activity and serum lipoproteins
in IDDM patients.

Diabet. Med. 6 (suppl.2):28 A.

Johansson ,C.;Sammelsson,U. and Ludvigsson,J. (1994):

A high weight gain early in life is associated with an increased
risk of type I. Diabetes mellitus. Diabetologia, 37:9194

Kanwar, Y.S. & Farquhar, M.G. (1980):

Detachment of endothelium and epithelium from the glomerular
b.m. produced by kidney perfusion with neuraminidase.

Lab. Invest; (42):375-85.

Kanwar,Y.S.; Liu, Z.Z.; Kashiara, N. and Wallner , E.I. (1991):

Current status of structural and functional basis of glomerular
filtration & proteinuria.

Seminar in Nephrol., 11/4: 390-413.

Kanwar ,Y.S. ; Rosenzweig , L.J. Linker , A. & Jakubowski , M. B. (1983):

Decreased de novo synthesis of glomerular proteoglycans in diabetes: biochemical and auto radiographic evidence.

Proc. Natl . Acad. Sci. USA. 80:2272-5.

Kanwar ,Y. (1984):

Biophysiology of glomerular filtration and proteinuria.

Lab. Invest.; (51):7-12

Kannel W.B. and McGee ,D.L. (1979):

Diabetes and glucose tolerance as risk factors for cardiovascular disease: the framingham study.

Diabetes care ; (2): 120-6

Karjalainen , J .; Martin , J. M. and Knip ,M. (1992):

A bovine albumin peptides as a possible trigger of IDDM.

N. Engl. J. Med. ,327:302-307.

Kasem N.S. , Sauka H.M. (1989):

Inheritance of diabetes mellitus. Bullet High Institute Public Health;19: 1.

Katopoids N. ;Hirshout , Y. and Geller , P . (1982):

Lipid associated sialic acid test for the detection of human cancer.

Cancer Res; (42):52-70

Kawasaki, Moriuchi ,R. and Takino,H. (1992):

Autoantibodies to 64 ,000 Mr islet cell protein in long term type I diabetic patients. *Diabetologia* 35:748 –752.

Kemal , M. ; Salbri , A . ; Horstzincke and James, D. (1986):

Significance of serum protein & LSA as a marker for genito urinary malignancies.

Cancer ; (57):1389-1394.

Kerjaschki ,D. ;Vernillo , A.T. & Farquhar, M.G. (1985):

Reduced Sialylation of podocalyxin –the major sialoprotein of the rat kidney glomerulus –in aminonucleoside nephrosis

Am. J. Pathol; (118) :343-9.

Kerjaschki , D.; Sharkey, D.J. &Farquhar ,M.G. (1984):

Identification and characterization of podocalyxin-the major sialoprotein of the renal glomerular epithelial cell.

J. Cel Biol ; (98): 1591-6.

Kessel ,D.and Bosmann, H.B. (1974):

Studies on a cell derived from the L 1210 murine leukaemia with altered surface properties and decreased capacity for tumour production.

Cancer Res. 34:603-608.

Kesson,C.M. and Slotter, S.D. (1979):

Smoking in diabetics.

Lancet; (1): 504 –5.

Khanderia ,U.; Keller ;J and Grossman ,H.B. (1983):

Serum sialic acid as a biological marker for malignant diseases.
J.Sur. Oncol; (23):163

Kida , H. ; Yoshimura , M., Ikeda;Saitou,Y. & Noto,Y.(1991):
Pathogenesis of diabetic nephropathy.
J. Diabetic complications .;(3):84.

Kjellen ,L.;Bielefeld ,D. & Hook , M. (1983):
Reduced sulfation of liver heparan sulfate in experimentally
diabetic rats.
Diabetes ; (32):337-42

Klein ,R.;Klein ,B.E.K.& Moss,S.E.(1985):
Rekinopathy in younger onset diabetic patients.
Diabetes Care;(8):311.

Klenk K,E. and Lavenstein , K (1952):
Z. Physiol. Chem ., 291,147.

Klenk, E. and Walter , H. (1952):
Z. Physiol . Chem. ,291, 259.
In Methods of Biochem. Analysis.Vol.2 New York interscience,
pp 183.

Klein ,R.; Klein , B .E. K. & Moss , S.E. (1985):
Retinopathy in younger onset diabetic patients.
Diabetes Care ; (8) : 311.

Serum sialic acid as a biological marker for malignant diseases.
J.Sur. Oncol; (23):163

Kida , H. ; Yoshimura , M., Ikeda;Saitou,Y. & Noto,Y.(1991):
Pathogenesis of diabetic nephropathy.
J. Diabetic complications .;(3):84.

Kjellen ,L.;Bielefeld ,D. & Hook , M. (1983):
Reduced sulfation of liver heparan sulfate in experimentally
diabetic rats.
Diabetes ; (32):337-42

Klein ,R.;Klein ,B.E.K.& Moss,S.E.(1985):
Rekinopathy in younger onset diabetic patients.
Diabetes Care;(8):311.

Klenk K,E. and Lavenstein , K (1952):
Z. Physiol. Chem ., 291,147.

Klenk, E. and Walter , H. (1952):
Z. Physiol . Chem. ,291, 259.
In Methods of Biochem. Analysis.Vol.2 New York interscience,
pp 183.

Klein ,R.; Klein , B .E. K. & Moss , S.E. (1985):
Retinopathy in younger onset diabetic patients.
Diabetes Care ; (8) : 311.

Klein , B.E.; Moss , S.A. & Klein , R. (1990):

Effect of pregnancy on progression of diabetic retinopathy.
Diabetes Care; (13):34-40.

Klein,R.;Klein,B.E.K.;Moss, S.E.; Davis M.B. & De-Mets,D.L.(1986)

The wisconsin Epidemiologic Study of Diabetic Retinopathy V.
Proteinuria and retinopathy in a population of diabetic persons
diagnosed prior to 30 years of age.
Diabetic Renal –Retinal syndrome ; (3):245-64.

Klein , R.; Klein, B.E.K.&Moss ,S.E. (1989):

The Wisconsin Epidemiologic Study of Diabetic Retinopathy IX.
four year incidence and progression of diabetic retinopathy when
age at diagnosis is less than 30 years.
Arch. Ophthalmol; (107):237-43.

Klein ,R.; Klein , B.E.K. & Moss,S.E. (1985):

Retinopathy in younger onset diabetic patients.
Diabetes Care;(8):311.

Klein ,R.;Klein ,B.E.K.&Moss,S.E.;Davis,M.D.&De-Mets,D.L.(1984)

The Wisconsin Epidemiologic Study of Diabetic Retinopathy II.
Prevalence and risk of diabetic retinopathy when age at diagnosis
is less than 30 years.
Arch. Ophthalmol ;(102):520-6.

Knapik- Kordecko.M.;Rola , K.H. and Warwas,M. (1990):

Glycated Proteins and various acute phase factors in patients with

type II diabetes.

Wiad-Lek 1-15 ; 43 (5-6):193-8.

Kosraba , J.N.;Cruickshanks , K.J. and Lawler- Heavner(1993):

Pathogenesis of D.M. Early exposure to cow 's milk and solid foods in infancy , genetic predisposition and risk of IDDM. Diabetes ,42:288-295 .

Kofaed- Enevoldsem ,A.;Borch-Johnsen , K.,Kreiner , S.; Krup, J.&Deckert,T.(1987):

Insulin dependent diabetic patients in Dinmark. Diabetes;(36):205-9.

Koivisto , V.A.;Huttenen, N.P. and Vierikko ,P. (1981):

Continuous Subcutaneous insulin infusion corrects exercise-induced albuminuria in Juvenile diabetes. Br. Med. J;(282):778-9.

Kremmer ,T.; (Boldizsar,M.;Gesztsek ,A. and Szajani ,B. (1991):

Are evaluation of the lipid-bound s.a. deternination. Clinica Chemica Acta; (203):259-268.

Krolewski, A.S.;Kosinski , E.J. ;Warran, J. M. (1987):

Magnitude and determinants of coronary artery disease in Juvenile-onset, insulin dependent diabetes mellitus. Am. J. Cardial; (59): 750-5.

Krolewski, A.S.;R and,L.I; warran , J. H & Alello,L.M. (1985):

Proliferative diabetic retinopathy (P DR) risk is closely related to haemoglobin (A1 C) level.
Diabetes . 34 (suppl.1):71A.

Krolewski, A.S. ; Canessa, M.; Warran, J.H.; Laffel, M.B:

Christlieb, A.R.; Knowler, W.C. & R and, L.I. (1988)

Predisposition to hypertension and susceptibility to renal disease in insulin dependent diabetes mellitus.
N. Engl. J. Med; (318):140-5.

Kuhn, R. and Prossmer, R. (1957):

In methods in Biochemical Analysis Vol 2.
New York Interscience pp;261:279.
Angew Chem; (69) :534.

Kuglin; Gries, F.A. and Klab (1988):

Immunological markers of diabetes mellitus. Diabetes 37 :130-132

Kushner, I. (1988):

The acute phase response: an overview.
Methods Enzymol (163): 373-83.

Laffel, L.M.B.; Krolewski, A.S.; R and, L.I., Warran, J.H.

Christlieb, A.R. and Elia, J.A. (1987):

The impact of blood pressure on renal function in insulin dependent diabetes.
Kidney Int; (31):207

Larsen , H.W. (1960):

Diabetic retinopathy. An ophthalmoscopic changes and the pathogenic factors in this disease.

Acta Ophthalmol. (Copenh), 60: 1-89.

Lee,L.S.;Hwang,J.Y.;Change,J.J.;Hsu,C.H.;Liao,S.T.&Lo,I.L.(1989)

Apolipoproteins A-I&B in NIDDM.

Taiwan I Hsuch Hui Chih.,88(11-12):1139-1142.

Lindberg , G. ; Eklund , G.A. ; Gullberg , B. & Rastman ,L.(1991):

Serum s.a. concentration & cardio vascular mortality .

Br Med J ; (302):143-46.

Lindberg,G.;Rastam,L.;Gullberg,D.&Eklund,G.A.(1992):

Serua s.a. concentration predicts both CHD& stroke mortality:

Multivariate analysis including 54385 men & women during 20.5

Years follow up.

Int. J.Epidemiol.:(21): 253-57.

Mackiewicz, A.; Ganapath,M.K.;Sultz Scultz, Brabenec,A.&

Kushner ,I.(1990):

TGF-beta regulates production of acute phase proteins.

Proc Natl Acad Sci USA;(87):1491-95.

Maclaren , N. and Athininson , M.(1988):

IDDM environmentally induced England, J.Med.327 :348

Maclaran, Arthkinson M (1992):

Insulin-dependent diabetes mellitus environmentally induced N
Engl J. Med. 327:348:

Maier , V.;Hinz , M.;Schaltz , H.;Nierle, C. & pheiffer , E.F.(1973):
Evidence for a specific glucose receptor by removal of s.a. from
isolated pancreatic islets mice.
Diabetologia ; (9):80

Maloney , J.B.M. & Durg , M.I. (1982):
The effect of pregnancy on the natural course of diabetic
retinopathy.
Am. J. Ophthalmol ; (93):745

**Manduteanu , I. ;Calb , M. ; Lupu , C.;Simionesev , N. and
Simionesv M.(1992):**
Increased adhesion of human diabetic platelets to cultured valvular
endothelial cells.
J. Submicrosc. Cytol. Pathol. , 24 (4):539-47.

Mangili,R.;Bending, J.J.;Scollg-Lilk-Gupta & Vibdrti ,G.C. (1988):
Increased Sodium-Lithium counter-transport activity in red cells of
patients with IDDM and nephropathy.
N engl. J. Med; (318):146-50.

Marshall , S.M. & Alberti , KGMM. (1989):
Comparison of prevalence & associated features of abnormal
albumin excretion in IDDM & NIDDM.
Quart. J. Med. ; (70):61-71

Mathiesen , E.R. ; Saurbrey ,N. ; Hommel., E. &Parving, H.H.
(1986):

Prevalence of microalbuminuria in children with IDDM.

Diabetologia ;(29):640-3

Mathiesen , E.R, Borch –Johnsen , K.; Jensen , D.V. and Deckert, T.
(1989):

Increased survival in patients with diabetic nephropathy.

Diabetologia ; (32):884-6.

Matsumoto ,E.;Matsumoto ,G. Coshima-Aetol (1990):

Serum type IV collagen concentration in diabetic patients with microangiopathy as determined by enzyme immuno assay with monoclonal antibodies.

Diabets;(39):885-90.

Mathiesen , E.R., Qxenball , B.; Johansen , K.;Sendsenp , A. and Deckert ,T. (1984):

Incipient nephropathy in type I (insulin dependent) diabetes.

Diabetologia ; (26):406-10

Mauer , S.M.;Stefles , M.W.;Ellis , E.N.;Sutherland , D.E.R.;Brown D. Mand-Gaetz, F.C. (1984):

Structural-functional relationship in diabetic nephropathy.

J. Clin , Invs; (74):1143-55

McMillan , D. E. (1982):

Further observations on serum viscosity changes in diabetes mellitus.

Metabolism , Vol , 31, No,3.

McMillan , D.E. (1974):

Disturbance of serum viscosity in diabetes mellitus
Journal of clinical investigation; (53):1071-1079.

Melvin ,T., Kin ,Y. &Michael , A.F. (1984):

Selective binding of Ig G and other negatively charged plasma
proteins in normal and diabetic human kidney.
Am. J. Pathol ;(115):443-6

Messina , A.;Davies , D.J. and Dillane , P.C. (1987):

Glomerular epithelial abnormalities associated with the onset of
proteinuria in amino nucleoside nephrosis.
Am J. Pathol , (126) :220-9.

Mogensen , C.E. & Schmitz , O.(1987):

The diabetic kidney. From hyperfiltration and microalbuminuria to
end stage renal failure.
Medical Clinical of North America ; (6):1465.

**Mogensen , C.E; Hansen , K.W.;Osterby ,R.&Damsgarad , E.M.
(1992)**

Abnormal albuminuria in the genesis & prediction of renal disease
in diabetes .
Diabetes. Care ; 15/9:1192.

Mogensen ,C.E.;Mauer , S.M.& Kjellstrand , C.M. (1988):

Diabetic nephropathy. Schrier ,R.W. and Gottschalk ,
C.W.:(eds):Diseases of the kidney.4th ed Little , Brown and Co.

Mogensen , C.E. (1971):

Urinary albumin excretion in early and long term juvenile
diabetes.

Scand J. Clin. Lab. Invest.:(28):183-93.

**Morell , A.G.;Gregoriadis , G.;Scheinbery , I.H.; Hickmon , J.and
Ashwell , G. (1971):**

The role of s.a. in determining the survival glycoproteins in the
circulation.

Journal of biochemical chemistry 246,1461.

Mogensen,C.E. and Christensen , C.K. (1984):

Detection of type VI Collagen in extracellular matrix of renal
glomeruli and its increase in diabetes.

Diabetes;(39):31-7

Mogensen,C.E. and Christensen , C.K. & Vittingus , E. (1983):

The stages in diabetic renal disease with emphasis on the stage of
incipient diabetic nephropathy.

Diabetes;32 (suppl.2) :64-78

Most ,R.S.and Sinnock , P. (1983):

The epidemiology of lower extremity amputations in diabetic
individuals.

.Diabetes Care ; (6):87-91.

Myers , D.;Winetz ,J. A; Chiuf and Michael ,A.S. (1982):

Mechanisms of proteinuria in diabetic nephropathy , a study of glomerular barrier function.

Kidney Int.; (21):633-41.

Nakamura ,Y. and Myers,B.P. (1988):

Charge Selectivity of proteinuria in diabetic glomerulopathy.

Diabetes ; (37):1202-11.

Nelson , R.G.;Kunzelman , C.L. and Saad , M.F. (1989):

Albuminuria in NIDDM & impaired glucose tolerance.

Diabetologia ;(32): 870-6.

Nishida , K.;Nagamura , Y.& Gitsu ,N. (1990):

Change in levels of specific terminal sugar residues of glycoproteins in sera of patients with complication of diabetes & liver injury.

Rinsho , Byori , Jul ;38 (7):835-7

Ozben,T.(1991):

Elevated serum s.a. & urine s.a. levels in renal disease.

Ann. Clin Biochem;(28): 44-48.

Palmer , J. P.;Asplin , C.M. and Clemons , P. (1983):

Insulin auto antibodies in insulin –dependent diabetes before insulin treatment. Science 22 :1337-39.

Palmer , J.P. ;Helgirst , S.and Giatgen , A. (1989):

Interaction of beta cell activity an IL-IB concentration and exposure time in isolated rat islet of Langerhans.

Diabetes , 38:1211-16.

Pappenheimer , J.R;Renkin , E.M. & Barrero , L.M. (1951):

Filtration diffusion and molecular sieving through peripheral capillary membranes . A contribution to the pore theory of capillary permeability.

Am. J.Physiol ; (167): 13-46.

Parthasarathy , N.& Spiro , R.G. (1982):

Effect of diabetes on the glycosaminoglycan component of the human glomerular b.m.

Diabetes , (31):738-41

Parving ,H.H& Hommel E. (1989):

Prognosis in diabetic nephropathy.

Br. Med. J. ; (299): 230-3

Parving , H.H.; Hommel, E. & Mathiesen ,E. (1988):

Prevalence of microalbuminuria , arterial hypertension , retinopathy and nephropathy in patients with insulin –dependent diabetes.

Br. Med. J; (296):156-60.

Paul ,E.&Jenning ,S. (1992):

The mechanism of diabetic microangiopathy.

Diabetographia ;(11):4.

Phelps , R.L.;Sakol ,P.&Metzker , B.E. (1986):

Changes in diabetic retinopathy during pregnancy , corrections
with regulation of hyperglycaemia.

Arch. Ophthalmol ;(104):1806-10

Pyorala , K.;Leakso , M.& Uusitupa ,M. (1987):

Diabetes and atherosclerosis: an epidemiologic view.

Diabetes / Metab. Rev ; (3): 464-524.

Rabinowe S.L.,George ,N.L.& Loughlin ,R.(1986):

D.M. & its complication Am. J. med 81:779-782

Rahman ,M.A;Zafar –Gand & Shera ,A.S. (1990):

Changes in glycated proteins in long term complications of
diabetes mellitus.

Biomed Pharmacother;44(4):229-34

Rand,L.I.;Homme,G.J.;Ederer,F. ant Canner,P.L.(1985):

Diabetic retinopathy Study Research Group, Factors influencing
the development of visual loss in advanced diabetic retinopathy.

Inrest.Ophthalmol.

Vis Sci;(26):893-1991.

Ratzan H.B. (1992):

Aetiology and pathogenesis of diabetes mellitus.

Ped. Clin. N.A.M, 35 (2):620.

Rehman , J. and Rahman ,M.A. (1991):

Studies on glycated plasma proteins in diabetic patients.

JAMA. J- Pak. Med. Assoc;41 (11):16-8.

Riley LJ,Cooper M.; Norins RG (1989):

Alkali Therapy of DKA: Biochemical, Physiology ,and Clinical
Perspective Diabetes.

Metab. Rev. 5:627.

**Rogers, M.E; Williams , D.T ; Niththy-Ananthan , R.;Rampling
,M.;Lil-Heslop,K.E. and Johnston , D.G. (1992):**

Decrease in erythrocyte glycophorin sialic acid is associated with
increased erythrocyte aggregation in human diabetes.

Clin. Sci. Colch , 82 (3)-309-13.

Rosenzweig ,L.J &Kanwar ,Y.S (1982):

Removal of Sulphated (heparan sulphated) or non sulphated
(chyluronic acid) glycosaminoglycans results in increased
permeability of the glomerular basement membrane to 125I-
Bovine serum albumin.

Lab Invest; (47):177-84.

Roseman ,S.;Jourdian, G.W.; Watson ,D.& Rood,R.(1961):

Metabolism of N-acetyl neuraminic acid.

Proc. Natl. Acad. Sci.U.S.; (47):958.

Saifer ,A and Gerstenfeld ,S. (1952):

Measurement of s.a. by diphenylamine reaction .

J. Lab. Clin. Med.; (50):17.

Salhanisk , A.T and Amatruda , J.M. (1988):

Role of sialic acid in insulin action and the insulin resistance of diabetes mellitus.

Am.J. Physiol :255 (2 pt 1):E 173-9.

Salhanide , A.I. and Amatruda , J.M. (1988):

Role of s.a. in insulin activity and the insulin resistance of diabetes mellitus.

Am. J. physiol. ; (255) : E 173 -9 .

Schauer, R. (1985):

S.a , and their role as biological masks.

Trends in Biomedical Science 10 , 357 – 360

Schnitzer , J. A. and Pinnay , E. (1992):

Quantitation of specific binding of orosomucoid to cultured microvascular endothelium.

Am. J. physiol. ;(263) :H 48-55.

Sampson , M. J.; Chambers , J .;Springings , D. and Drurg, P.L (1990):

Intra ventricular septal hypertrophy in type II diabetic patients with microalbuminuria or early proteinuria.

Diabet. Med; (7):126-37

Schober , E.;Pollack., A.;Coradelb H.and lubec,G. (1982):

Glycation of glomerular basement membrane in type I insulin dependent diabetic children.

Diabetologia ;(23):485-7

Schreiber ,A.B.;kenny ,J.;Kowalsk. G.(1985):

A unique family of endothelial cell polypeptide mitogens. The antigenic and receptor cross reactivity of bovine endothelial cell growth factor , brain- derived acidic fibroblast growth factor and eye derived growth factor II.

J. Cell Biol ; (101):1623-6.

Scott ,D.;Berkourtz ,C.and Augusto ,B. (1988):

S.a. prevent loss of large von Willebrand factor multimers by protecting against amino terminal proteolytic cleavage.

Blood , Vol. 72 , No. 5:pp, 1790-1796.

Shimomura,H. and Sipro , R.G. (1987):

Studies on the macromolecular components of human glomerular b.m. and alteration in diabetes , decreased levels of heparan sulphate proteoglycan &Laminin.

Diabetes ;(36):374-81.

Siebert , F.D;Pfaff. M.L. siebert ,M.V. (1948):

Detection of sialic acid by tryptophane-perchloric acid reaction method.

Arch. Biochen. Biophys.:(18):279.

Siliver , H.K.B;Karim ,K.A.Salinas ,A.&Swenertonk, D. (1981):

Significance of s.a. and carcinoembrgonic antigens as monitors of tumour burden among patients with carcinoma of the ovary.

Surg. Gynecol obstet. 153:205-213.

Sirek , O.N. and Sirek ,A. (1971):

Plasma glycoproteins and diabetes mellitus in man in handbuch
des Diabetes mellitus Ed. E.F. feiffer .J.F
Lehmans , Munich, Vol. 2, P. 317.

Skovborg , F.;Aa ,V.;Nielsen ,J. &Ditzel,J. (1966):

Blood Viscosity in diabetic patients.
Lancet.; (1):129:2:805.

Sobenin ,I.A.; Tertou ,U.&Koschinoky T.(1993):

Modified LDL from diabetic patients causes cholesterol
accumulation in human intimal aortic cells.
Atherosclerosis April ,100;(1)41-54.

**Sosenko , J. M. ; Breslow , J. L; Mietinen , O.S. ;Gabbay, K. Y. H.
(1980):**

Hyerglycaemia & Lipid levels.
N. Engl – J . Med. ; 302 (12) : 650 – 654.

Sperling M.A. (1996):

Diabetes Mellitus in:Behrman RE, Kliegman RM and Arvin AM
(editors) .Nelson (Text book of Pediatric),15th ed. Saunders
Company Philadelphia, London Toronto,1646-1664.

Spiro ,R.G. (1969):

Glycoproteins: their biochemistry , biology and role in human

disease.

New England Journal of Medicine ; 281&1043.

Spiro ,R.G. and Parthasarathy ,N.(1982):

Studies on proteoglycan of b.m.: In Kuhn K, Schcene H, Timpl R (eds).

New trends in BM research.

New York Raven Press ;PP :87-98.

Spuda J., Neumann M ;Kvapil M; Dart a Skov a D(1996):

Serum levels of s.a. in patients with type I D.M. in relation to the incidence of late complications.

Cas Lek Cesk ,135:723-5

Srikanta ,S.:Canad ,O.P. and Dakson , R.S. (1983):

D.M. and dyslipidamia.

American , Intern. Med.,99:320-329.

Srinivarsan ,P.R. (1959):

Biochemistry of s.a.

J. Biol. Chem ., 234,716.

Starup , K.; Larsen , H. & Enk , B. (1980):

Flourescein angiography in diabetic children .

Acta – ophthalmol: (58): 347.

Stersk ,G.and Wall,S. (1986):

Determinants of microangiopathy in growth-onset diabetes: with

special reference to retinopathy and glycaemic control.

Acta Pediatr. Scand ; Suppl. (327): 6-45.

Sternberg ,M.;Cohen- Forterre,L. & Pegrou , J.(1985):

Connective tissue in diabetes mellitus: biochemical alterations of the intercellular matrix with special reference to proteoglycans ,collagen, and basement membranes.

Diabetes Metab ;(11):27-50.

Steven ,A.S;Marcus , A.K.;Lawrence ,M.T. and Stephen , G.M. (1991):

Diabetes mellitus in current medical diagnosis and treatment ;30th edition , Churchill, Livingstone ,P.859.

Stuart , J. ; George , A. G ; Davis , A. J. & Hurlow , R.A. (1981):

Haematological stress syndrome in atherosclerosis.

J. Clin. Pathol. ; (34): 464- 467

Sugahara , K. ; Sugimoto ,K.; Nomura ,O.& Usiu ,T.(1980):

Enzymatic assay of serum s.a.

Clin. Chem. Acta;(108):493-8

Svennerholm , L (1958):

Methods for estimation of NANA.

Acta. Chen. Scand. ,12,547.

Svennerholm,L. (1957):

Methods for determination of serum s.a. orcinol method.

Ark. Kemi ,10,577

Taniuchi , K.; Chifu, K. and Hayashi, N. (1981):

Enzymatic method for the determination of sialic acid in serum & its application as a marker of acute phase reactants.

Kobe . J. Med. Sci.; (27) : 91- 102.

Taniuchi , K.; Miyamoto, Y.; Doi, K.; Tanka , T. and Baba , S. (1984):

Plasma acute phase proteins & increased platelet s. a. in diabetic patients.

Kode. J. Med. Sci. 30 : L 7-16.

Tarsio , J.F.; Wigness , B. and Khode ,T.D. (1985):

Non enzymatic glycation of fibronectin and alterations in the molecular association of all matrix and b.m. components in diabetes mellitus.

Diabetes; (34):477-84.

Tarsio ,J.F.;Reger ,L.A.&Furcht , L.T. (1988):

Molecular mechanisms in b.m. complication of diabetes mellitus association.

Diabets ; (37):5320-9.

Tchobroutsky , G. (1978):

Relation of diabetic control to development of microvascular complications.

Diabetologia; (15): 143.

Thomas, M.D. (1992):

Lipid metabolism 11 :Pathways of metabolism of special lipids in text book of Biochemistry With Clinical Correlation , Third edition 3rd ed. W.A. John Willey & Sons Inc. , publication. Philadelphia , Pennsylvania , P 371-373 & 454-456.

Timple ,R.(1984):

Structural models and molecular interaction in basement membranes & cell movement.
Ciba foundation symposium 108. London.UK.
Pitman press:25-43.

Tomlanovich ,S.;Denn ,W.M.,Jones,H.W.;Schwartz ,H.C. and Myers ,B.D. (1987):

Functional nature of glomerular injury in progressive diabetic glomerulopathy.
Diabetes; (36):556 -65

Tomino ,Y.; Inoue, W.; Yagame, M.;Nometa ,Y.;Sakai ,H.;Ito,K.;Nagoaka, K. and Indeda, N. (1985):

Measurement of s.a. and acute phase reactant proteins in sera of patients with diabetic nephropathy.
J.Diabet- Complications.2 (4):175-8.

Tomino , Y. ;Satai,H.;Miura,M.;Suga,T.,Yagame ,M.& Endoh ,M. (1986):

Effect of immumoglobulin deposition of glomerular s.a. in patients with I_g A nephropathy.

Am. J. Nephrol.;(6): 187-192.

Tomino , Y.; Yagame ,M.; Inoue W.; Watanabe,S.;Kaneshigh.

Nomato ,Y.; Sakai , H. and Takedo ,M. (1987):

Double immunofluorescence studies of glomerular sialic acids in patients with diabetic nephropathy.

J. Diabet. Complicons.1(2):61-4.

Tomino, Y.; Inoue, W.; Watanabe,S.; Yagame , M.; Eguchi, K.;

Nomatynd Sakai, H. (1988):

Detection of glomerular sialic acids in patients with diabetic nephropathy. Am. J. Nephrol; (8):21-26.

Trucco R. E; and caputtol , R. (1954):

In Methods of Biochem. Analysis Vol. 2 New York interscience ,pp 356-460

J.Biol. Chem, 206/901.

Tsan, M.F. & McIntyre ,P. A. (1976):

The requirement for membrane s.a. in the stimulation of superoxide production during phagocytosis by human polymorphonuclear leukocytes.

J.Exp. Med; (143):1306-1316.

Viberti ,G.C.; Messent ,J. & Morocutti ,A. (1992):

Diabetic nephropathy.

Diabetes care 15/9:1216.

Viberti , G. C.; Keen, H., Dadds, R. & Bending, J.J (1987):

Metabolic control and progression of diabetic nephropathy.
Diabetologia ; (30):481-2.

Viberti , G.C. and Keen , H. (1984):

The patterns of proteinuria in diabetes mellitus.
Diabetes; (33): 686-92

Viberti , G.C. ; Mactintosh , D. and Keen, H. (1988):

Determinants of the penetration of proteins through the glomerular
barrier in insulin-dependent diabetes mellitus.
Diabetes;32(Suppl. 2):92-5

**Viberti , G.C.; Bilous , R.W.; Mocsinlosh , D.;Bending, J.J. and
Keen, H. (1983):**

Long term correction of hyperglycaemia and progression of renal
failure in IDDM.
Br.Med.J.286-598-602.

Virtanen , S.M.;Rasarent, A.R.O.,A. (1994):

Diet cow's milk protein antibodies and the risk of IDDM in finish
children.
Diabetologia , 37:381-387.

Vittinghus , E. and Mogensen ,C.E. (1982):

Graded exercise and protein excretion in diabetic man and the
effect of insulin treatment.
Kidney Int.; (21):725-9.

Wahle , P.; Deppermann , D.& hasslacher C. (1982):

Biochemistry of glomerular b.m. of the normal&diabetic human.
Kidney Int.; (21):744-9.

Wagner, A.; Genouese , S. and Bosi ,S.(1994):

Low metabolic deterioration toward diabetes in islet cell antibody
+ve patients autoimmune polyendocrine disorder.
J. Diabetes , P.201-205.

Walkar , J.D; Tarig ,T.&Vibertig.(1990):

Sodium-Lithium counter transport activity in red cells of patients
with IDDM and nephropathy and their parents.

Werner ,L.and Odin ,L. (1952):

Quantitation of sialic acid.
Acta. Soc. Med. Upsol .; 57,230.

West ,K.M.;Ahuja ,H.M.S.&Bennett,P.H.(1983)

The role of circulating glucose and triglycerides concentration and
their interactions with the risk factors as determinants of arterial
disease in nine diabetic population samples from the WHO
Multinational Study.
Diabetes Care; (6):361-9.

West , K, M; Erdreich ,L. J. and Slober. , J. A. (1980):

Risk factors for retinopathy and nephropathy diabetes: a detailed
study.
Diabetes ; (29) : 501-508

Wesberg-NG and Micheal AF. (1973):

Human glomerular b.m. chemical composition in diabetes mellitus.

Acta. Med. Scan;(194):39-47

Williamson ,J.R.; Titton , R. G. and Charg., K.(1988):

Basement membrane: abnormalities in diabetes mellitus: relationship to clinical microangiopathy.

Diabetes Metab. Rev;(4):339-370.

Williamson , J.R.; & Rilo , C.(1977):

Current status of capillary b.m. disease in diabetes mellitus.

Diabetes; (26) :65-75.

Williamson ,J.R.;Titton, R.G. and Charg ,K.(1988):

Basement membrane: abnormalities in diabetes mellitus: relationship to clinical microangiopathy.

Diabetes Metab. Rev.:(4):339-370.

Wisemann,M.J.,Visberti ,G.C.,Mackintosh, D.;Jarrett ,R.J.& Keen,H.(1984):

Glycaemia , arterial pressure and microalbuminuria in type I(insulin dependent diabetes mellitus).

Diadetologia ; (26): 401-5.

Winetz ,J.A.;Golbetz ,H.V.;Spencer , R.J.; Lee,J.A. and Myers ,B.D.(1982):

Glomerular function in advanced diabetic nephropathy.

Kidney Int. (21): 744-9

Wu ,V-Y., Wilson , B.& Cohen, M.P. (1987):

Disturbances in glomerular b.m. glycosaminoglycans in experimental diabetes.

Diabetes; (36): 697-83.

Wyngaarden , Smith & Bennett , P.(1992):

Endocrinal reproductive diseases.

In Cecil Text book of medicine, 19th edition, Churchill Livingstone,P 121

Yamamoto , T. ; Nakamura, T. ; Noble , N.A. & Border,W.A. (1993):

Expretion of TGF – beta is elevated in human & experimental diabetic nephropathy.

Proc. Natl . Acad. Sci. USA ; (90) : 1814-18.

Yamashiro , Y.; Taniguchi, H.;Marimoto , Y., Rotama ,S.& Baba ,S.(1989):

Target antigen of islet cell antibody in Japanese diabetes mellitus.

Diabetes Res. Clin. Pract. 1; 6(3): 161-5.

Yokoyama,H.; Jenson ,J.S; and Deckert ,T.(1995):

Serum s .a. concentration is elevated in IDDM. especially in diabetic

nephropathy.

J. Intern . Med; 237(5): 519-23.

Yon ,J. W.; Paok ,C. and Mearther , R. G. (1988):

Pathogenesis of diabetes mellitus.

Lancet, 1:1-4.

Zavaroni , I;Bonora ,E. and Pagliara ,M. (1989):

Risk factors for coronary artery disease in healthy persons with hyperinsulinaemia and normal glucose tolerance.

New Engl. J. Med.;320-702-6

Zeller , K.; J.acobson ,H. and Raskin , P. (1991):

The effect of dietary protein and phosphorus restriction on renal function in diabetic nephropathy.

Kidney Int.; (37): 246.

Zolbina ,E.N.; Dubinkini, U. and Merkushov, A.V. (1992):

Effect of glutamic acid decerboxylase on insulin secretion immunology. 31,289-296