

References ...

Akisu, M. & Kultursay, N.:

Value of the urinary uric acid to creatinine ratio in diagnosis of prinalatal asphyxia. *Acta-paediatr* - Jpn 1998 feb; 40 (): 78 - 81

Angerson, W. J.:

Blood flow in the brain. New York, Oxford university press, 1989.

Armstead, W. M.; Miro, R.; and Busija, D. W.:

Post- ischemic generation of superoxide anion by newborn pig brain. *Am. J. physiol.* 225:401, 1988.

Armstrong, D. L.; Sauls, C. D.; and Goddard, F. J.:

Neuropathologic findings in short-term survivors of intraventricular hemorrhage. 141: 617, 1987.

Azzopardi, J. D.:

Prognosis of newborn infant with hypoxic ischemic brain injury assessed by phosphorus magnetic resonance specterometry. *Pediatr. Res.* 25: 445, 1989

Barratt, T. M.:

Urolithiasis and nephrocalcinosis. In Holliday NA, Barratt, T. M.; Vernier, R. L. (eds): *Pediatric Nephrology*. Baltimore, Williams & Wilkins, P. 706; 1987.

Bechman, J. S.:

The double edge role of nitric oxide in brain function and superoxide mediated injury. *J. Dev. Physiol.* 15: 53, 1991.

Bejar, R.; Worzinak, M.; Allard, M.; et al:

Antenatal origins of neurologic damage in newborn infants. *Am. J. obstet. Gynecol.* 159: 357, 1988.

Berard, E.:

Nephrocalcinosis and prematurity: importance of urate and oxalate excretion. *Nephron*, 69:237-241., 1995

References ...

Breart, G.; Rumeau – Rouquette, C.:

Cerebral palsy and perinatal asphyxia in full term newborn infants. Arch pediatr 3: 70-4, 1996

Burt, R. D.; Vaughan, T. C.; and Daling, J. R.:

Evaluating the risks of cesarean section: Low Apgar score in repeated C-section and vaginal deliveries. Am. J. public Health 78: 1312, 1989.

Burtis, C. A. & Ashwood, E. R.:

Nitrogen metabolism and Renal function: In Tietz Fundamentals of Clinical Chemistry P. 577, 1996.

Carpenter, R. H. (ed.):

Blood supply of the brain, studying of the brain development. 3rd edition, Churchill living Stone Inc. P. 305, 1985.

Carter, B. S.; Haverkanp, A. D.; Merenstein, G. B.; et al.:

The definition of acute perinatal asphyxia. Clinics in perinatol. 20: 287, 1993.

Catlin, E. A.; Carpenter, M. W.; Brann, B. S.; et al.:

The Apgar score revisited, influence of gestational age. J. pediatr. 109: 865, 1986.

Chesley, L. C.; Williams, L. O.:

Renal glomerular and tubular function in relation to hyperuricemia of pre-eclampsia and eclampsia. Am J Obstet Gynecol 50: 367; 1945.

Choi, D. W.:

Cerebral hypoxia: some new approaches and answered questions. J. Neurosci. 10: 3494, 1990.

De-Boer, J.; Klein, H.; Postema, F.; et al.:

Rat strial cation shifts reflecting hypoxic-ischemic damage can be predicted by One – line impedance measurements. Stroke 20: 1377, 1989

References ...

Denes, E. D. and Borthold, J. S.:

Early prognostic value of serum creatinine level in children with posterior urethral valves. *Journal of Urology* .157 (4), 1441 – 3, 1997.

De – Reuck, J. L.:

Cerebral angioarchitecture and perinatal brain lesions in premature and full-term infants. *Acta Neurol. Scand.* 60: 381, 1984.

Dragunow, M.; Sirimanne, E.; fawlor, P.A.; et al.:

A role for calcitonin gene-related peptide during recovery from hypoxic-ischemic brain injury in the infant rat. *Brain Res.* 14: 267, 1992.

Epstein, M. F.:

Resuscitation in the delivery room. In: Cloherty J.P. and Stark A.R. (eds). *Manual of Neonatal care*, 3rd edition, P. 56, 1991.

Ergenekon, E.:

Biol. neonate: 76 (4): p 200-206. Oct.; 1999

Evon, Y. Snyder and John P. Cloherty J. P.:

Manual of neonatal care. P. 515, 1997

Fellman, V. & Raivio K.O.:

Reperfusion injury as the mechanism of brain damage after perinatal asphyxia. *Pediatr res* 41: 599- 06, 1997

Fenichel, G. M.:

Hypoxic-ischemic encephalopathy in the newborn. *Arch. Neurol.* 40: 261, 1983.

Field, D.R.; Parer, J. T.; Aulender, R. A.; et al.;

Cerebral oxygen consumption during asphyxia in fetal sheep. *J. Dev. Physiol.* 14: 131, 1990.

References ...

Hill, A. & Vilpe, J. J.:

Hypoxic-ischemic encephalopathy. In: Textbook of Neurology
MRC Robertson (ed) London, Churchill Livingstone, P. 546, 1986.

Hill, A. and Volpe, J.:

Perinatal asphyxia: Clinical aspects. Clinics in perinatol. 16:
4356, 1989.

Hill, L. M.:

Metabolism of uric acid in normal and toxemic pregnancy.
Mayo Clin Proc 53: 743; 1978.

Huang, C. C.; Wang, S. T.; Chang, Y. C.:

N. Engl. J. Med. 29; 341 (5): P328 – 35, Jul 1999.

Jelinck, J. and Jensen, A.:

Catecholamine concentrations in plasma and organs of the fetal
Guinea pig during hypoxaemia and asphyxia. J. Dev. Physiol.
15: 145, 1991.

Jensen, A.; Hohnmann, M. and Kunzel, W.:

Dynamic changes in organ blood flow and oxygen consumption
during acute asphyxia in fetal sheep. J. Dev. Physiol. 9: 543,
1987.

Jimenez, R.; Vilanena, J. and Figuero, S.:

Prostaglandin and thromboxan B2 levels in CSF of asphyxiated
newborns. Abstract 289, in 14th European congress: June, 1994.

Johnston, M. V. & Silverstein, F. S.:

New insights into mechanisms of neuronal damage in the
developing brain. Pediatr, Neurosci. 12: 87, 1986.

Karlowicz, M. G. & Adelman, R. D.:

Nonoliguric and oliguric acute renal failure in asphyxiated term
neonates. Pediatric Nephrol 9:718- 22. 1995

References ...

Nelson, K. B. and Ellenberg, J.H.:

Obstetric complications as risk factors for cerebral palsy or seizure disorders. JAMA 251:1843,1984.

Nelson, K. B. and Emery, E. S.:

Birth asphyxia and the neonatal brain: what do we know and when do we know it? Clinics in Perinatal. 20:327, 1993.

Netter, F. H.:

Arterial supply of the brain. In the Ciba collection of medical illustration vol. I, nervous system. Ciba Pharm. Company. P. 46, 1988.

O'connor, M.; Harkness, R.; Simmonds, R. and Hytten, F.:

The measurement of hypoxanthine, xanthine, inosine and uridine in umbilical cord blood & fetal scalp blood samples as measure of fetal hypoxia. Br. J. Obstet. Gynaecol 1.88:381, 1981.

Palme-Klinder, C.:

Methods of resuscitation in low Apgar score newborn infants: A national survey. Acta Paediatr. Scand., 81:739, 1992.

Palmer, G.H. & Vanucci, R.C.:

Potential new therapies for perinatal cerebral hypoxia- ischemia. Clin perinatol 20:411-32, 1997.

Palminha, J. M.; Ramos, P.; Larajeire, T.; et al:

Perinatal asphyxia: lipoperoxidation. And HIE. Abstract page 100, In 2nd world congress of perinatal medicine: September, 1993.

Pamela, C.:

Chap biochemistry 2nd (ed). Lippincott's Company 1994

Patel, J.; Edwards, A. D.:

Prediction of outcome after perinatal asphyxia Curr Open Paediatr 9: 128-32; 1997.

References ...

Peacock, M.; Robertson, W. G.:

The biochemical aetiology of renal lithiasis. In: Urinary Calculous Disease. J. E. A. Wickham, Ed. London Churchill Livingstone, 1979.

Peebles, D. M.:

Near infrared spectroscopy and intrapartum fetal surveillance. Abstract page 5, In 2nd World Congress of Perinatology Medicine: September, 1993.

Penning, D. H.; patrick, J. and Jemiono, S.:

Release of glutamate and GABA in the ovine fetal hippocampus. J.Dev. physiol. 16:301,1991.

Perez, R. E.; Spinoza, M. and Riquelene, R.:

Arginine vasopressin mediates cardiovascular responses to hypoxemia in fetal sheep. Am.J. physiol. 256:1011,1989.

Perlman, J. M.:

Intrapartum hypoxia-ischemic cerebral injury and subsequent cerebral palsy. Pediatrics 99: 851-8; 1997.

Perlman, J. M. and Tack, E. D.:

Renal injury in the asphyxiated newborn infants. Relationship to neurologic outcome. J. Pediatr. 113:875, 1988.

Perone, R. D.; Madias, N. E.; Levey, A. S.:

Serum creatinine as an index of renal function: New insights into old concepts. Clin. Chem., 38:1933-1953; 1992.

Philip D. Mayne:

Clinical chemistry in diagnosis and treatment 6th 1994

Phipps, R. H.:

Delivery room management of the newborn. In: Avery, G. E. (ed), Scheffer and Avery's Diseases of the Newborn. 2nd, ed., WB Saunders Co. Philadelphia p:182, 1981.

References ...

Portman, R. G.; Carter, B. S.; Gaylord, M. S.; Murply, M. G.;

Thieme, R. E.; Merenstein, G. B.:

Predicting neonatal morbidity after perinatal asphyxia: Ascoring system. Am. J. Obstet Gynecol 162: 174 - 182, 1990.

Poland, R.L.and Freeman, R. K.:

Guidelines for perinatal care. Relationship between perinatal factors and neurologic outcome . American Academy of Pediatrics: p. 221. 1992.

Pontsky, B.:

Vascular supply of the brain. In Neuroanatomical Pathology .3rd edition Philadelphia WB savnders: p. 8. 1984

Porter, R.:

Aspects of neuronal development. Neurophysiology. 3rd edition philadelphia WB. Savnders. june III. P.149. 1978.

Poulsen, J. P.; Rognum, T. O.; Oyasaeter, S.; Saugstad, O.D.:

Changes in oxypurine concentration in vitrous humour of pigs during hypoxemia and post mortum. Pediatr. Res. 28: 482-484; 1990.

Raffin, C. N.; Harrison, M. and Sick, J. J.:

ELC troencephalogram suppresion and anoxic depolarizationinfluences on cerebral oxygenation during ischemia. J. Cerebr. Bl. flow metab. 11: 407. 1991.

Reid, D. L.; Parer, J. T. and Williams, K. M.:

Effect of severe reduction in maternal placental blood flow distribution in the sheep fetus. J. Dev. physiol. 15:183. 1991.

Richardson, B. S.:

Fetal adaptive response to asphyxia. Cilics in Perinatol. 16 : 595 . 1989.

References ...

Rieselbach, R. E.; Steele, T. H.:

Influence of the kidney upon urate homeostasis in health and disease. *Am J Med* 56: 665; 1974.

Robertson, N. J. and Edwards, A. D.:

(Curr – Opin – Pediatr: 10 (6): 575-80, 1998, Dec.

Robertson, C. M. and Finer, N. N:

Long – term follow up of term neonates with perinatal asphyxia. *Clinics in perinatol.* 20:483. 1993

Rosenberg, A. A.; Maurdauph, E. and White, C. W.:

The role of oxygen free radicals in the postasphyxial cerebral hypoperfusion in newborn lambs. *Pediatr. Res.* 26: 215, 1995.

Russel, Gap.; Jeffer, S. G. and Cooke. R.W. I.:

Plasma hypoxanthine as marker for hypoxic – ischemic induced periventricular leucomalacia? *Arch Dis. Child* 67: 388 – 392; 1992.

Sarnat, H. B. and Sarnat, M. S.:

Neonatal encephalopathy following fetal distress : a clinical and electroencephalographic study. *Arch. Neurol* .33:696. 1976.

Saugstad, O. D.:

Hypoxanthine as an indicator of hypoxia: Its role in health and disease through free radical production . *pediatr. Res* 23:143. 1988.

Seitchik, J.:

Observations on the renal tubular reabsorption of uric acid: I. Normal pregnancy and abnormal pregnancy with and without preeclampsia. *Am J Obstet Gynecol* 65: 981; 1953.

Shankaran, S.; Woldt, E. and Koepke, T.:

Acute neonatal morbidity and long – term central nervous system sequelae of perinatal asphyxia. *Early Hum. Dev.* 25 : 135. 1991.

References ...

Shankaran, S.:

Syndrome perinatal asphyxia. Clin. Perinatal. 20: 287. 1995.

Shy, K. K.; Luthy, D. A. and Bennett, F. C.:

Effect of electronic fetal heart rate monitoring as compared with periodic auscultation on the neurologic development of premature infants. N. Eng. L. J. Med. 322 : 588. 1990.

Silverstein, F. S.; Buckanan, K.; Hudson, C. et al.:

Flunarizine limits hypoxia-ischemia induced morphologic injury in immature rat brain. Stroke, 17: 477, 1986.

Simeone, F. A.; Frazer, G. and Lawner, P.:

Ischemic brain edema in immature rat brain. Stroke, 17: 477, 1979.

Snyder, E. Y. and Cloherty, J. P.:

Perinatal asphyxia. In: Cloherty, J. P. and Stark, A. R. (eds), Manual of Neonatal care, 3rd edition little, Brown and white Company, p. 393, 24 : 161. 1991.

Snyder, E. Y. & Cloherty, J. P.:

Perinatal asphyxia. In: Cloherty J and Stark AR (eds): Manual of Neonatal Care. 4th ed., Lippincott-Raven, Boston, P: 515, 1998.

Sonntage, J. and Prankal, B.:

Serum creatinine concentration, urinary creatinine excretion and creatinine clearance during first 9 weeks of preterm infant with a birth weight below 1500 gm. European Journal of Pediatrics, 155 (9), 815 – 9, 1996.

Sorenson, L. B. and Levinson, D. J.:

Origin and extrarenal elimination of uric acid in man. Nephron 14: 7; 1975

References ...

Thiringer, K.; Thornberg, E. and Kjellmer, I.:

Neurone-specific enolase (NSE) in asphyxiated newborns, relationship to encephalopathy. Abstract 252, In: 14 the European Congress: June, 1994.

Thorp, J. A.; Sampson, J. E.; Parisi, V. M.; et al.:

Routine umbilical blood gas determinations. Am. J. Obstet. Gynecol. 161, 600 1989.

Tsukahara, H.; Hiraoka, M. and Hori, C.:

Urinary uric acid excretion in term and preterm infant. Journal of Pediatrics and Child Health, 33 (4) 330 – 2, 1996 Aug.

Volpe, J. J.:

Hypoxic – ischemic encephalopathy: Clinical aspects. Neurology of the newborn, 2nd edition, Philadelphia. WB Saunders p.236, 1987.

Vaccarino, F. Guidotti, A. and Coasta, E.:

Ganglioside inhibition of glutamate mediated protein kinase C translocation in primary cultures of cerebellar neurons. Proceeding of the National Academy of Science of the U.S.A., 84: 8707, 1987.

Vannucci, R.:

Current and potentially new management strategies for perinatal hypoxic-ischemic encephalopathy. Pediatrics, 85 (6): 961, 1990.

Vanucci, R. C.:

Experimental biology of cerebral hypoxia-ischemia: relation to perinatal brain damage. Pediatr Res 27: 317-26; 1990

Vanucci, R. C. & Perlman, J. M.:

Interventions for perinatal hypoxia-ischemia encephalopathy. Pediatrics 100: 1004-14; 1997.