




REFERENCES

Anderson, T. G. (1970):

Testing of susceptibility to antimicrobial agents and assay of antimicrobial agents in body fluids In: Manual of Clinical Microbiology Blair, J. E.; Lennette, E.H. Trunt, J. P.
Am. Soci. Microbiol. 299-310.

Archer, G.L. and Mayhall, C. G. (1983):

Comparison of epidemiological markers used in the investigation of an outbreak of methicillin resistant staphylococcus infection.

J. Clin. Microbiol., 18: 395-399.

Archer, G.L. and Pennell, E. (1990):

Detection of methicillin resistance in staphylococci by using a DNA probe.

Antimicrob. Agents Chemother. 34: 1720-1724.

Arnold, P.M.; Allyn P.A.; Nichols, E. M.; Hill, D. L.; Pezzlo, M. and Bartlett, R. H. (1982):

Control of methicillin-resistant *Staphylococcus aureus* in a burn unit: Role of nurse staffing.

J. Trauma 22: 954-959.

Ayliffe, G. A. T. and Mitchell, K. (1994):

Isolation of infected patients.

Post graduate doctor Middle East 2: 64-68.

Barber, M. (1961):

Methicillin-resistant Staphylococci.

J. Clin Path. 14: 385-393.

Baron, E. J. and Finnegold, S. M. (1990):

Bailey and Scotts diagnostic microbiology 8th edition.

C.V. Mosby Company. USA. Chap. 18: 253-262, Chap 24: 323-332.

Barry, A. and Jones, R. (1987):

Reliability of high-content disks and modified dilution tests for detecting staphylococcal resistance to the penicillinase-resistant penicillins.

J. Clin. Microbiol. 25: 1897.

Bayliss, B. G. and Hall, E. R. (1965):

Plasma coagulation by organisms other than *Staphylococcus aureus*.
J. Bacteriol. 89: 101-106.

Bernard, A. B. and Robert, E. G. (1974):

Staphylococci in: Microbiology and Infectious Disease McGraw-Hill
Book Company-AB-Lakiston publication. New York. 306.

Bitar, C. M.; Mayhall, C. G.; Lamb, V. A.; Bradshaw, T. J.; Spadora, A. C. and Dalton, H. P. (1987):

Outbreak due to methicillin and rifampin-resistant *Staphylococcus aureus*: epidemiology and eradication of the resistant strain.

Hospital Infect. Control. 8:15-23.

Blumberg, H. M.; Rimland, D.; Carroll, D. J.; Terry, P. and Wachsmuth, I. K. (1991):

Rapid development of ciprofloxacin resistance in methicillin-susceptible and resistant staphylococcus aureus.

Journal of Infectious Diseases. 163: 1279-1285.

Boyce, J. M. (1981):

Nosocomial Staphylococcal infections.

Ann. Intern. Med. 95: 241-242.

Boyce, J. M. (1989):

Methicillin-resistant *Staphylococcus aureus*.

Infect. Dis. Clin. North Am. 901.

Boyce, J. M. (1991):

Should we vigorously try to contain and control MRSA?

Infect. Control Hosp. Epidemiol. 12: 46-54.

Boyce, J. M.; Marguerite, M. and Murray, D. (1994):

Methicillin-resistant *Staphylococcus aureus* (MRSA) A Briefing for acute care hospital and nursing facilities.

Infect Control Hosp. Epidemiology. 15: 105-115.

Brumfitt, W. and Hamilton, M. (1989):

Methicillin resistant *Staphylococcus aureus*.

N. Engl. J. Med. 320: 1188-1196.

Burrows, W. (1973):

Staphylococci in: Textbook of microbiology 20th edition W.B. Saunders Company Philadelphia. London, Toronto. 405-417.

Calia, F. M. (1992a):

Clindamycin in S. L. Gorbach; Bartlett, J. G. and Blacklow, N. R. (eds) (1992):

Infectious Disease Philadelphia: Sunders, 2154-223.

Calia, F. M. (1992^b):

Erythromycin in S. L. Gorbach; Bartlett, J. G. and Blacklow, N. R. (eds) (1992).

Infectious Diseases, Philadelphia, Sunders, 223-231.

Casewell, M. W. (1986):

Epidemiology and control of modern methicillin resistant *Staphylococcus aureus*.

J. Hosp. Infect. 7: 1-11.

Chambers, H. F. (1991):

Treatment of infection and colonization by methicillin resistant *Staphylococcus aureus*.

Infect Control Hosp. Epidemiol. 12: 29-35.

Chambers, H. F. and Hackbarth, C. J. (1987):

Effect of NaCl and nafcillin on penicillin-binding protein 2a and heterogeneous expression of methicillin-resistance in *Staphylococcus aureus*.

Antimicrob. Agents Chemother. 31: 1982-1988.

Chambers, H. F. and Hackbarth, C. J. (1992):

Methicillin resistant *Staphylococcus aureus* genetics and mechanism of resistance in Cafferkey MT (ed): Methicillin resistant *Staphylococcus aureus*:

Clinical Management and Laboratory Aspects. New York Marcel Dekker, 21.

Chambers, H. F. and Sachdeva, M. (1990):

Binding of B-lactam antibiotics to penicillin-binding proteins in methicillin resistant *Staphylococcus aureus*.

J. Infect. Dis. 16: 1170-1176.

Chambers, H. F.; Sachdeva, M. and Kennedy, S. (1990):

Binding affinity of penicillin binding protein 2 a correlates with in vivo activity of B-lactam antibiotics against methicillin resistant *Staphylococcus aureus*.

J. Infect. Dis. 162: 705-10.

Chesbro, W.R.; Wamola, I. and Bartley, C. H. (1969):

Correlation of virulence with growth rate in *Staphylococcus aureus*.

Can. J. Microbiol. 15: Pp.723-725.

Cheessbraugh, M. (1991a):

Medical laboratory manual for tropicl countries, ELBS Cambridge, vol. II. section V. P. 26: 69.

Cheessbraugh, M. (1991b):

Medical laboratory manual for tropicl countries, ELBS Cambridge, vol. II. section V. P. 100-129.

Christie, R. and Keogh, E. V. (1940):

Physiological and serological characteristic of staphylococci of human origin.

J. Path. Bacter. 51: 189-197.

Clemet, D. S.; Macaluso, R.; Leons and Ralph, H. (1988):

High rate of methicillin resistance of *S. aureus* isolated from hospitalized nursing home patients.

Arch. Intern. Med. 148: 569-570.

Cohen, S. H.; Morita, M. M. and Bradford, M. (1991):

A seven year experience with methicillin-resistant *Staphylococcus aureus*.

Am. J. Med. 91 (Suppl. 3B): 233S-237S. 16.

Collopy, B. T.; Dalton, M. F.; Wright, C. and Mallany, C. (1984):

Comparison of the clinical significance of methicillin resistant and methicillin sensitive *Staphylococcus aureus* isolations.

Med. J. Aust., 140: 211-214.

Cookson, B.; Peters, B.; Webster, M.; Phillips, I.; Rohmon and Noble, W. (1989):

Staff carriage of epidemic methicillin-resistant *Staphylococcus aureus*.

J. Clin. Microbiol. 27: 1471-1476.

Coolee, J. G.; Duguid, J. P.; Fraser, A. G. and Marmion, B. P. (1989):

The staphylococci In: Mackie McCartney Practical Medical Microbiology. 30th ed (Reprint).

Churchill Livingstone. Edinburgh, London and New York. 303-315.

Coovadia, Y. M.; Bhana, R. H.; Johnson, A. P.; Haffejee, I. and Marples, R. P. (1989):

A laboratory confirmed outbreak of rifampin-methicillin resistant *Staphylococcus aureus* (MRSA) in a newborn nursery.

J. Hosp. Infect. 14: 303-312.

Cowen, S. T. and Steel, K. J. (1964):

Comparison of differentiating criteria for staphylococci and micrococci.

J. Bact. 88: 804-805.

Craig, W. A. (1992):

Rifampin and related drugs in S.L. Gorbach; Bartlett, J.G. and N.R. Blocklow (eds)

Infectious Diseases. Philadelphia, Saunders 265-271.

Crossley, K.; Loesch, D.; Landesman, B.; Mead, K.; Chearn, M. and Strate, R. C. (1979):

An outbreak of infections caused by strains of *Staphylococcus aureus* resistant to methicillin and aminoglycosides clinical studies.

J. Infect Dis. 139: 273-279.

Cruikshank, R. (1953):

The epidemiology of some skin infections.

Brit. Med. J. I: 55-59.

Dacre, J.; Emmerson, A. M. and Jenner, E. A. (1986):

Gentamycin-methicillin resistant *Staphylococcus aureus*, epidemiology and containment of an outbreak.

J. Hosp. Infect. 7: 130-136.

Davis, B. D.; Dullbecco, R.; Elsen, H. N.; Ginsberg, H. S.; Wood, W. B. and McCarty, M. (1973):

Staphylococci In: Microbiology including immunology and molecular genetics 2nd edition Harper and Row, Publishers, Hagerstown, Maryland New York, Evanston, Sanfrancisco, London. 728-738

de Lencastre, H. A. M. Sà Figueiredo C. Urban J. and A. Tomasz (1991):

Multiple mechanisms of methicillin resistance and improved methods for detection in clinical isolates of *S. aureus*.

Antimicrob. Agents and chemother. 35: 632-639.

Dickson, J. I. and Marples, R. R. (1986):

Coagulase production by strains of *staphylococcus aureus* of differing resistance characters : a camparison of two traditional methods with a lates agglutination system detecting both clumping factor and protein A.

J. Clin. Path. 39: 371-375.

Doern, G.V. (1982):

Evaluation of a commerical latex agglutination test for identification of *staphylococuus aureus*.

J. Clin. Microbiol. 15: 416-420.

Dornbusch, K. and Hallander, H. O. (1973):

Transduction of pencillinase production and methicillin resistance-enterotxin B production in strains of *S. aureus*.

J.Gen Microbiol, 76: 1-11.

Dornbusch, K.; Hallander, H. O. And Lofquist, F. (1969):

Extrachromosomal control of methicillin -resistance and toxin production in *S. aureus*.

J. Bacteriol. 98:351-358.

Drew, W. L.; Barry, A. L. and O'Toole, R. (1972): Reliability of the Kirby-Bauer disc diffusion method for detecting methicillin-resistant strains of *S. aureus*.

Appl. Microbiol. 24: 240-247.

Duckwarth, G. J. (1988):

Diagnosis and managment of methicillin resistant *Staphylococcus aureus* infection.

B. M. J. 1988:307:1049-1052

Duckworth, G. J.; Lothian, J. L. and Williams, J. D. (1988):

Methicillin resistant *Staphylococcus aureus* report of an outbreak in a London Hospital.

J. Hosp. Infect. 11: 1-15.

Duguid, J. P. (1989):

Staphylococcus cluster-forming gram positive cocci In: Collee. J. G.; Duguid J. P.; Fraser A. G. and Marmion B. P (eds)

Mackie and McCartney, Practical Medical Microbiology, Churchill Livingstone, Edinburgh, London, Melbourne and New York. 303-316.

Easmon, C. S. and Goodfellow, M. (1990):

Staphylococcus and *Micrococcus* in Topley and Wilson's principles of bacteriology, virology and immunity 8th edition.

Edward Arnold. A division of Hodder and Stoughton. London, Melbourne, Auckland. 161-186.

Eickoff, T. C. (1982):

Nosocomial infections New Engl.

J. Med. 306: 1545-1556.

Elek, S. (1956):

Experimental staphylococcal infection in the skin of man.

Ann. N.Y. Acad. Sci. 65: 85-90.

Ellison, R. T.; Judson, F. N.; Peterson, L. C.; Cohn, D. I. and Ehret, J. M. (1984):

Oral rifampin and trimethoprim/sulfamethoxazole therapy in asymptomatic carriers of methicillin resistant *Staphylococcus aureus* infection.

West. J. Med. 140: 735-740.

Ena, J. (1993):

The epidemiology of intravenous vancomycin usage in university hospital. A 10-year study.

J. A. M. A 269:598.

Essers, L. and Radebold, K. (1980):

Rapid and reliable identification of *Staphylococcus aureus* by a latex agglutination test.

J. Clin. Microbiol. 12: 641-643.

Fairbrother (1940):

Coagulase production as criterion for the classification of staphylococci. J. Path. Bact 50:83. Cited from. Kloss W.E (1980): Natural populations of the genus staphylococcus.

Ann. Rev. Microbiol. 34: 559-582.

Faller, A. and Schleiffer, K. H. (1981):

Modified oxidase and Benzidine test for separation of staphylococci from micrococci.

J. Clin. Microb. 13: 1031-1035.

Fekety, R. (1987):

The management of the carrier of methicillin resistant *Staphylococcus aureus*.

Curr. Clin. Top. Infect. Dis. 8: 169-180.

Foagali, J. L.; Thong, M. L. and Grant, D. (1992):

Ten years experience with methicillin resistant *Staphylococcus aureus* in a large Australian hospital.

J. Hosp. Infect. 20: 113-119.

Forsgren, A. (1977):

Immunological aspects of protein A. In Schlessinger, D. (ed):
Microbiology. Washington, D. C.
American Society for Microbiology. 353.

Geraci, J. E. (1977):

Vancamycin Mayo.
Clin. Proc. 52: 631-634.

Gerberding, J. L.; Mick, C.; Liu, H. and Chambers, H. F. (1991):

Comparison of conventional susceptibility tests with direct detection
of penicillin-binding protein 2a in borderline oxacillin-resistant
strains of *Staphylococcus aureus*.
Antimicrob. Agents and Chemother. 35: 2574-2579.

Glew, R. (1992):

Vancomycin. In S.L. Gorbach; Bartlett, J. G. and Blacklow N.R.
(eds).
Infectious Diseases, Philadelphia, Sunders.

**Goher, H.; Ayoub, M.; Gharib, M.; El-Mahad, A. and Mokhtar, S.
(1993):**

Determination of methicillin resistance among *Staphylococcus
aureus* using different antibiotic susceptibility tests.
Medical Journal of Cairo University, Vol. 61, No. 1.

Goldmann, D. A. (1992):

Epidemiology of *Staphylococcus aureus* and group A streptococci
in Bennett, J. V. Brachman, P. S. (eds):
Hospital infections, ed. 3. Boston, Little, Brown and Company.
767.

Gram, C. (1884):

Ueber die isolirte farbung de schizomyceten in Schnitt-und Trocken
preparaten.

Fortschritte der Medicine: 2: 185.

**Griehle, H. G.; Krause, S. L.; Pappas, S. A. and Dicostanzo, M. B.
(1981):**

The prevelence of high-level methicillin resistance in multiply
resistant hospital staphylococci.

Medicine. 605: 62-69.

Griffith, L. J. and Mullins, C. G. (1968):

Drug resistance as influenced by inactivated sensitivity disks.

Appl. Microbiol. 16: 656-658.

Hackbarth, C. J. and Chamber, H. F. (1989):

Methiciltin-resistant staphylococci detection methods and treatment
of infection.

Antimicrob. Agents and Chemother. 33: 995-999.

Hajek, V. and Marsalek, E. (1976):

The differentiation of pathagenic staphylococci and suggestion for
their taxonomic classification.

Zentralblatt für Bakteriologie parasitenkunde infektions krankheiten
and hygiene Abt. I. Orig A. 217, 176-182. Ibid, suppl. 5: 11.

**Haley, R. W.; Hightower, A. W.; Khabbaz, R. F.; Thornsberry, C.;
Martone, W. J.; Allen, J. R. and Hughes, J. M. (1982):**

The emergence of methicillin -resistant *Staphylococcus aureus*
infections in United States Hospitals.

Ann. Intern. Med . 97: 297-308.

Hall, L. M. C.; Jorden, J. Z. and Wang, F. (1989):

Methicillin-resistant *S. aureus* from China characterized by digestion of total DNA with restriction enzymes.

Epidem. Inf. 103: 183-192.

Hansen, S.; Freedy, P. (1984):

Variation in the abilities of automated, commercial and reference methods to detect methicillin-resistant (hetero resistant) *Staphylococcus aureus*.

J. Clin. Microbiol. 20: 494.

Harrstein, A. I. and Word, T. T. (1991):

Methicillin resistant *Staphylococcus aureus*.

Infect Control Hosp. Epidemiol. 12: 9-10.

Hartman, B. J. and Tomasz (1986):

Expression of methicillin-resistance in heterogeneous strains of *Staphylococcus aureus*.

Antimicrob. Agents Chemother. 29: 85-92.

Hershow, R. C.; Khayr, W. F. and Smith, N. L. (1992):

A. comparison of clinical virulence of nosocomially acquired methicillin resistant and methicillin sensitive *Staphylococcus aureus* infection in a university hospital.

Infect. Control Hosp. Epidemiol. 13: 587-593.

Hewitt, J. H.; Coe, A. W. and Parker, M. T. (1969):

The detection of methicillin resistance in *Staphylococcus aureus*.

J. Med. Microbiol. 2: 443-454.

Hill, R. L.; Duckworth, G. J. and Casewell, M. W. (1988):

Elimination of nasal carriage of methicillin resistant *Staphylococcus aureus* with mupirocin during a hospital outbreak.

J. Antimicrob. Chemother. 22: 377-384.

Hindler, J. A. and Inderlied, C. (1985):

Effect of the source of Mueller-Hinton agar and resistance frequency on the detection of methicillin-resistant *S. aureus*.

J. Clin. Microbiol. 21: 205.

Hindler, J. A. and Warner, N. L. (1987): Effect of the source of Mueller-Hinton agar on the detection of oxacillin-resistant in *S. aureus* using screening methodology.

J. Clin. Microbiol. 25: 734-735.

Howard, B. J. and Kloos, W. E. (1987):

Staphylococci In: Clinical and Pathogenic Microbiology Howard, B. J.; Klass, J.; Wessfeld, A. S.; Rubin, S. J. and Tiltan, R. C. The C.V. Mosby Company. St. Louis- Washington, D. C. Toronto. 231-244.

Ingerman, M. J. and Santoro, J. (1989):

Vancomycin: A. new old agent.

Infect. Dis. Clin. North. Am. 3: 641.

Jacobs, S. I.; Willis, A. T. and Goodburn, G. M. (1964):

Pigment production and enzymatic activity of staphylococci the differentiation of pathogens from commensals.

J. Path. Bacter. 87: 151-167.

Jawetz, E.; Brooks, G. F.; Melnick, J. L.; Butel, J. S.; Adelberg, E. A. and Ornston, J. N. (1989):

Staphylococci In: Medical Microbiology (Reprint) 18th ed.
Appleton and Lang.-Norwalk. Connecticut\San mateo California.
187-192.

Jawetz, E.; Melinckj, J. L. and Adelberg, E. A. (1987):

"The staphylococci" In: Review of Medical Microbiology.
Appleton and Lang.- Norwalk. Cannecticut\Los Altos California.

Jevons, M. P. (1961):

Celbenin-resistant Staphylococci.
Brit. Med. J. 1: 124-125.

Jevons, M. P.; Coe, A. W. and Parker, M. T. (1963):

Methicillin resistance in staphylococci.
Lancet. 904-907.

Jorgensen, J. H. (1991):

Mechanism of methicillin resistance in *Staphylococcus aureus* and
method for laboratory detection.
Infect, Control. Hosp. Epidemial. 12: 14-19.

Kaatz, G. W.; Seo, S. M.; Dorman, N. J. and Lerner, S. A. (1990):

Emergence of teicoplanin resistance during therapy of
Staphylococcus aureus endocarditis.
J. Inf. Dis. 162: 103-108.

Kahn, F. A. (1993):

Quinolones and Macrolides. Roles in respiratory infection.
Hosp. Pract. 28:149.

Kaye, D. (1980):

The clinical significance of tolerance of *Staphylococcus aureus*.
Ann. Inter. Med. 93: 925.

Kayser, F. H. (1975):

Methicillin-resistant staphylococci.
Lancet, 2: 650-653.

Kilmek, J. J.; Marsik; F. J.; Bartlett, R. C.; Weir; B.; Shea, P. and Quintiliani, R. (1976):

Clinical epidemiologic and bacteriologic observations of an outbreak of methicillin resistant *Staphylococcus aureus* at large community hospital.
Am. J. Med. 61: 340-345.

Kloss, W. E. (1980):

Natural populations of the genus staphylococcus.
Ann. Rev. Microbiol. 34: 559-591.

Kloss, W. E. and Jorgensen, J. H. (1985):

Staphylococci In: Manual of clinical microbiology 4th ed.
Lennette, E.H. Washington, D.C: As M. 143-153.

Kloss, W. E. and Lambe, D. W. (1991):

Staphylococcus, in Balows A. Hausler, W. J.; Herrmann, K. L. (eds): Manual of clinical microbiology. ed 5. Washington D. C. American Society for Microbiology. 222.

Kloss, W. E. and Schleifer, K. H. (1975):

Simplified scheme for routine identification of human staphylococcus species.
J. Clin. Microbiol. 1: 82-88.

Kuhl, S. A.; Pattee, P. A. and Bladwin, J. N. (1978):

Chromosomal map location of the methicillin-resistance determinant in *S. aureus*.

J. Bacteriol. 135: 460-465.

Locksley, R. M.; Cohen, M. L.; Quinn, T. C.; Tompkins, L. S.; Coyle, M. B.; Kirihaara, J. M. and Counts, G. W. (1982):

Multiply antibiotic-resistant *Staphylococcus aureus*: introduction, transmission and evolution of nosocomial infection.

Annals of Internal Medicine, 97: 317-324.

Margaret, M. (1993):

Methicillin resistant *Staphylococcus aureus* past, present and future.
Nursing clinics of North America vol . 28 No. 3.

Mc Gougal, L. (1987):

Acquired oxacillin resistant *Staphylococcus aureus*.
Clin. Microbiol. Newsl. 9:18.

Mc Gowan, J. E. (1985):

Role of the microbiology laboratory in prevention and control of nosocomial infections, 110-122.

Lennette, E.H. Balow, A.; Hansler, W. J. and Shadomy, H. J. (ed)
Manual of clinical microbiology, 4th ed. American Society for Microbiology, Washington, D.C.

Mc Gowan, J. E.; Terry, P. M.; Haung, T.; Houk, C. L. and Davies, J. (1979):

Nosocomial infections with gentamycin-resistant *S. aureus*: plasmid analysis as an epidemiological tool.

Journal of Infectious Diseases, 140: 864-872.

Mulligan, M. E.; Ruane, P. J.; Johnston, L. and wong, J. P. (1987):

Ciprofloxacin for eradication of methicillin resistant staphylococcus colonization.

Am. J. Med. 82(suppl. 4 A) 215-219.

Murray, P. R.; Drew, W. L.; Kobayashi, G. S. and Thompson, J. H. (1990):

Staphylococcus In: Medical Microbiology. International Student Edition: Published by Wolfe Publications Ltd. 47-63

Neu, H. (1983):

The emergence of bacterial resistance and its influence on empiric therapy.

Reviews of Infectious Diseases, 5: 9-20.

Neu, H. C. (1992a):

Use of fluroquinolone.

Infect. Dis Clin. Pract. (IDCP) 1: 1.

Neu, H. C. (1992b):

The crisis in antibiotic resistance.

Science 257: 1064-1073.

Nicas, T. I.; Cole, C. T.; Preston, D. A.; Schabel, A. A. and Nagarajan, R. (1989):

Activity of glycopeptides against vancomycin resistant gram positive bacteria.

Antimicrobial Agents Chemother. 33:1477-1481.

Nolan, C. N. and Beaty, H. N. (1976):

Staphylococcus aureus bacteraemia. Current Clinical Patterns.

Am. J. Med. 60: 495-500.

Novick, R. P. and Bauanchand, D. (1971):

Extrachromosomal nature of drug resistance in *S. aureus*.
Ann. N. Y. Acad. Sci, 182: 279-294.

Ogston, A. (1881):

Br. Med. J. 1: 369-375. Cited from Kloss, W. E. (1980) Natural populations of the Genus *Staphylococcus*.
Ann. Rev. Microbiol. 34: 559-592.

Onesko, K. M. and Wienke, E. L. (1987):

The analysis of impact of a mild, low-iodine lotion soap on the reduction of nosocomial methicillin resistant *Staphylococcus aureus* a new opportunity for surveillance by objectives.
Infect. Control 8: 284-288.

Panlilio, A. L.; Culver, D. H. and Gaynes, R. P. (1992):

Methicillin-resistant *Staphylococcus aureus* in U.S. Hospitals, 1975-1991.
Infect. Control. Hosp. Epidemiol. 13: 582-586.

Passet, J. (1885):

Fortschr. Med. 3: 33-43. Cited from kloss, W.E. (1980): Natural papulation of the Genus *Staphylococcus*.
Ann. Rev. Microbiol. 34: 559-582.

Pasteur (1879):

Cited from Davis, B. D.; Dulbecco, R.; Elsen, H. N. Ginsbery, H. S.; Wood, W. B. and McCarty, M. (1973): *Staphylococci* In: Microbiology including immunology and molcular genetics 2nd edition.

Harper and Row, Publishers. Hagerstown, Maryland New York, Evanston. Sanfrancisco, London.728-738.

Paul, M. O.; Aderibigbe, D. A. and Sule, C. Z. (1982):

Antimicrobial sensitivity patterns of hospital and non-hospital strains of *Staphylococcus aureus* isolated from nasal carriers.

J. Hyg. Camb. 89: 253-260.

Peacock, J. E.; Moorman, D. R.; Wenzel, R. P. and Mandell, G. L. (1981):

Methicillin-resistant *Staphylococcus aureus*: Microbiologic characteristics antimicrobial susceptibilities, and assessment of virulence of an epidemic strain.

J. Infect. Dis. 144: 575-582.

Peterson, I.; Quick, J. and Jensen, B. (1989):

Emergence of ciprofloxacin-resistance in nosocomial methicillin resistant *Staphylococcus aureus* (MRSA) isolates during ciprofloxacin plus rifampin therapy from MRSA colonization the 29th Antimicrobial Agents and Chemotherapy September 20 Houston TX: Abstract.1255-1260.

Piercy, E. A.; Barbaro, D.; Luby, J. P. and Mackowiak, P. A. (1989):

Ciprofloxacin for Methicillin resistant *Staphylococcus aureus* infections.

Antimicrobial Agents-Chemother.128-130.

Putland, R. A. and Guinness (1985):

Antibiotic susceptibility testing of methicillin-resistant *Staphylococcus aureus* isolated in an Australian Hospital.

J. Clin. Microbiol., 22: 822-827.

Rahal, J. J. and Simberkoff, M. S. (1979):

Bactericidal and bacteriostatic action of chloramphenicol against meningeal pathogens.

Antimicrob. Agents Chemother. 16: 13.

Reboli, A. C.; John, J. F. and Levkoff, A. H. (1989):

Epidemic methicillin-gentamycin-resistant *Staphylococcus aureus* in a neonatal intensive care unit.

Ann. J. Dis. Child. 143: 34-39.

Ribner, B. S.; Landry, M. N. and Gholson, G. L. (1986):

Strict versus modified isolation for prevention of nosocomial transmission of methicillin resistant *Staphylococcus aureus*.

Infect. Control. 7: 317-320.

Richmond, A. S.; Simberkoff, M. S.; Schaeffer, S. and Rahal, J. J. (1977):

Resistance of *Staphylococcus aureus* to semisynthetic penicillins and cephalothin.

J. Infect. Dis. 135: 108-112.

Riley, T. V. and Rouse, I. L. (1995):

Methicillin-resistant *Staphylococcus aureus* in Western Australia.

J. Hosp. Inf. 29: 177-188.

Robert Koch (1878):

Cited from Davis, B. D.; Dulbecco, R.; Elsen H. N.; Ginsberg, H. S.; Wood, W. B. and McCarty, M. (1973): *Staphylococci* In: Microbiology including immunology and molecular genetics 2nd edition.

Harper and Row, Publishers, Hagerstown, Maryland, New York, Evanston, San Francisco, London. 728-738.

Rosenbach, F. J. (1884):

Micro organism bei Den Wund-infections Krank heiten des menschen Wies baden: Bergmann 122 pp. Cited from Kloss, W. E. (1980): Natural population of the Genus staphylococcus. Ann. Rev. Microbiol. 34: 559-592.

Rubbo, S. D.; Stratford, B. C. and Dixson, S. (1962):

Spread of a marker organism in hospital ward.
Brit. Med. J. 2: 282-290.

Sabath, L. D. (1982):

Mechanism of resistance to Beta lactam antibiotics in strains of *Staphylococcus aureus*.
Ann. Inter. Med. 97: 297-308.

Sandra, J.; Wadsworth, Ki-Ho Kim, Vilas Satishchandran, Peter Axelrod, Allan L. Truant and Byungse Suh (1992):

Development of new antibiotic resistance in methicillin-resistant but not methicillin-susceptible *Staphylococcus aureus*.
J. Antimicrob. Chemother. 30: 821-826.

Saraglou, C.; Cromer, M. and Bisno, A. L. (1982):

Methicillin resistant *Staphylococcus aureus*. interstate spread of nosocomial infections with emergence of gentamycin methicillin resistant strains.
Inf. Control: 1: 81-89.

Saravolatz, L. D.; Pohlod, D. J. and Arking L. M. (1982):

Community- acquired MRSA infections: A new source for nosocomial outbreaks.
Ann. Inter. Med. 97: 325-329.

Schaberg, D.; Power, G.; Betwold, J. and Forbes, B. A. (1985):

Conjugative plasmids in *S. aureus*.

J. Infect. Dis. Vol. 1152 (1) 43-49.

Selkon, J. B. (1980):

The role of an isolation unit in the control of hospital infection with methicillin-resistant staphylococci.

J. Hosp. Infect. 1: 41-46.

Semilack, J. D.; Wilson, W. R. and Cockerill, F. R. (1991):

Tetracycline, chloramphenicol, Erythromycin, Clindamycin, and metronidazol. Mayo.

Clin. Proceeding. 66:1270.

Sewell, C. M.; Clarridge, J.; Lacke, C.; Weinman, E. J. and Young, E. J. (1982):

Staphylococcal nasal carriage and subsequent infection in peritoneal dialysis patients.

JAMA 248:1493-1495.

Shanson, D. C. (1992):

Antibiotic resistance in *Staphylococcus aureus*. In: Cafferkey MT(ed): Methicillin-resistant *Staphylococcus aureus*:

Clinical Management and Laboratory Aspects, New York, Marcel Dekker. 11.

Shanson, D. C.; Johnstone, D. and Midgley, J. (1985):

Control of hospital outbreak of methicillin resistant *Staphylococcus aureus* infection: value of an isolation unit.

J. Hosp. Infect. 6: 285-292.

Shales, D. M. and Shales, J. H. (1995):

Teicoplanin selects for *Staphylococcus aureus* that is resistant to vancomycin.

Clinical Infectious Diseases. 20: 1071-1073.

Shulman, J. A. and Nahmias, A. J. (1972):

Staphylococcus infections: Clinical aspects In: Cohen, J. O. (ed): The staphylococci, John Wiley and sons Inc, New York chapter. 20: 437-481.

Sjostrom, J. E.; Lofdahl, S. and Philipson, L. (1975):

Transformation reveals a chromosomal locus of gene (s) for methicillin-resistance in *S. aureus*.

J. Bacteriol. 123: 905-915.

Smith, S. M.; Robert, H. K. and Tumang, F. T. (1989):

Ciprofloxacin therapy for methicillin resistant *Staphylococcus aureus* infection or colonization.

Antimicrobial Agent Chemother. 33: 181-184.

Sorrell, T. C.; Packman, D. R.; Shanke, S.; Foldes, M. and Munro, R. (1982):

Vancomycin therapy for methicillin resistant *Staphylococcus aureus*.

Ann. Inter. Med. 97: 344-350.

Standiford, H. C. (1987):

Methicillin resistant *Staphylococcus aureus* infection, it's time to get Tough (editoril).

Infect. Control. 8: 187.

Tewodros, W. and Gedebeu, M. (1984):

Nasal carrier rates and antibiotic resistance of *Staphylococcus aureus* isolated from hospital and non-hospital populations. Addis Ababa.

Trans. Roy. Soc. Trop. Med. Hyg. 78:314-318.

Thompson, R. L.; Cabezudo, I. and Wenzel, R. P. (1982):

Epidemiology of nosocomial infection caused by methicillin resistant *Staphylococcus aureus*.

Ann. Inter. Med. 97: 309-317.

Thornsberry, C.; Caruthers, J. Q. and Baker, C. N. (1973):

Effect of temperature on the in vitro susceptibility of *S. aureus* to penicillinase-resistant penicillins.

Antimicrob. Agents Chemother. 4: 263-269.

Trucksis, M.; Hooper, D. C. and Wolfson, J. S. (1991):

Emerging resistance to fluoroquinolones in *Staphylococcus aureus*.

Annals of Internal Medicine. 114, 424-426.

Tuazon, C. U. (1984):

Skin and skin structure infections in the patients at risk: Carrier State of *Staphylococcus aureus*.

Am. J. Med. 67: 166-171.

Volk, W. A.; Benjamin, D. C.; Kadener, R. J. and Parsons, J. T. (1991):

Staphylococci. In: Essentials of medical microbiology.

J. B. Lippincott Company Philadelphia. 366-374.

Wagner, G. E. (1990):

Indigenous flora and natural barriers to infection, hospital-acquired infections and Staphylococci. In: Microbiology 2nd edition. Kingsbury, D. T, and Wagner, G.E. A Wiley Medical publication; Wiley, J. and Sons. New York. Chichester. Brisbane. Toronto, Singapore. P. 71-73, 81-85, 373-378.

Waldvogel, F. A. (1990):

Staphylococcus aureus (including Toxic shock syndrome) in Mandeli, G. L.; Douglas, R. G. and Bennett, J. E. (eds): Principles and practice of infectious diseases ed 3 New York, Churchill Livingstone. 1489.

Walsh, T. and Hansen, S. L. (1987):

Prospective microbiologic surveillance in control of nosocomial methicillin resistant *Staphylococcus aureus*. Infect. Control. 8: 7-14.

Walter, J. B. and Israel, M. S. (1989):

Hospital infection. In: General Pathology. 6th edition Churchill Livingstone. 255-260.

Watanakunakorn, C. (1985):

Effect of inoculum size on in vitro susceptibility of methicillin-resistant *Staphylococcus aureus* to eighteen antimicrobial agents. Eur. J. Clin. Microbiol. 4: 68-96.

Wellstood, S. (1989):

Comparative evaluation of the MRS test. A 4 to 6 hour screening test for detecting oxacillin-resistant staphylococci. Diagn-Microbiol. Infect. Dis. Vol. 12, 395-399.

Wenzel, R. P. (1985):

Prevention and treatment of hospital acquired infection In: Cecil text book of medicine. Vol 2. 17th edition.

Wyngarden, J. B.; Smith, L. H. and Saunders, W. B. Company.
1485-1491.

Wenzel, R. P.; Nettleman, M. D.; Jones, R. N. and Pfaller, M. A.

(1991):

Methicillin resistant *Staphylococcus aureus* implications for the 1990s and effective control measures.

Am. J. Med. 91 (Suppl 3 B): 221S-227S.

Williams, J. and Wenzel, R. (1987):

Copying with methicillin-resistant staphylococcus infections.

J. Crit. Illness. Pp. 65-68.

Williams, R. E. O. (1963):

Healthy carriage of *Staphylococcus aureus*: its prevalence and importance.

Bacteriol. Rev. 27:56-71.

Winkler, J.; Block, C.; Leibovici, L.; Faktor, J. and Pitlik, S. D.

(1990):

Nasal, carriage of *S. aureus*. Correlation with hormonal status in women.

J. Inf. Dis. 162: 1400-1402.

Wolfgang, K. J. and Willett, H. P. (1976):

Staphylococci In: Zinsser Microbiology. 16th edition.

Appleton-Century Crafts. 412-429.

Wolfgang, K. J.; Willett, H. P. and Amos, D. B. (1984):
Staphylococci. In: Zinsser microbiology. 18th edition.
Appleton-Century Crafts. P. 443-461.

Wolfgang, K. J.; Willett, H. P. and Amos, D. B. (1988):
Staphylococci In: Zinsser Microbiology 19th edition.
Appleton-Century Crafts. P. 343-355.

Zaman, R. and Dibb, W. (1994):
Methicillin resistant *Staphylococcus aureus* (MRSA) isolated in
Saudi Arabia: epidemiology and antimicrobial resistance patterns.
J. Hosp. Inf. 26: 297-300.