

R E F E R E N C E S

- \* Abd Elnabi, E.A. and Mikhael, M.N. (1988):  
Beta-lactamase-producing Branhamella catarrhalis  
causing acute otitis media in children.  
Med. J. Cairo Univ., Vol. 56, 1: 143-147.
- \* Abd Elrehim, S.I.; Attia, M.W.; Nafie, E.; A'ssal,  
A.U. and Arafa, R.M. (1988):  
Bacteriological study of acute and chronic  
suppurative otitis media.  
Az. J. Microbiol., June, Vol. 3.
- \* Ahmad, F.; McLeod, D.T.; Calder, M.A. and Crougham,  
M.J. (1984):  
Antimicrobial susceptibility of Branhamella  
catarrhalis isolates from bronchopulmonary  
infection.  
Antimicrobial Agents and Chemotherapy, 26: 424-  
425.
- \* Ahmad, F.; McLeod, D.T.; Power, J.T. and Calder, M.A.  
(1985a):  
Branhamella catarrhalis prevalence in a hospital  
population.  
J. Hospital Infection, 6: 71-74.
- \* Ahmad, F.; Calder, M.A.; Marshall, T.G. and Crougham,  
M.J. (1985b):  
Urinary tract infection caused by Branhamella  
catarrhalis.  
J. of Infect., 10: 176.

- \* Aitken, J.M. and Thornley, P.E. (1983):  
Isolation of Branhamella catarrhalis from sputum and tracheal aspirate.  
J. Clin. Microbiology, 18: 1262-1263.
  
- \* Alvarez, S.; Jones, M.; Holtsclaw, S.; Guardera, J. and Berk, S. (1985):  
In vitro susceptibilities and beta-lactamase production of 53 clinical isolates of Branhamella catarrhalis.  
Antimicrobial Agents and Chemoth., 27: 646-647.
  
- \* Anon, (1982):  
Branhamella catarrhalis:  
Pathogen or opportunist ?  
Lancet, 8: 1056.
  
- \* Arko, R.J.; Wong, K.H. and Peacock, W.L. (1979):  
Nuclease enhancement of specific cell agglutination in a serodiagnostic test for Neisseria gonorrhoeae.  
J. Clin. Microbiol., 9: 517-519.
  
- \* Austrian, R.; Howie, V.M. and Ploussard, J.H. (1977):  
The bacteriology of pneumococcal otitis media.  
Johns Hopkins Med. J., 141: 104-111.
  
- \* Axelsson, A. and Brorson, J.E. (1973):  
The correlation between bacteriological findings in the nose and maxillary sinus in acute maxillary sinusitis.  
Laryngoscope, 83: 2003.
  
- \* Baginsky, A. (1907):  
Über Meningitis cerebrospinalis pseudoepidemica.  
Berlin Klin. Wschr., 44: 385.

- \* **Bailey, W.R. and Scott, E.G. (1986):**  
Microorganisms in the respiratory tract infections.  
In "Diagnostic Microbiology".  
(Eds. Finegold, S.M. and Baron, E.J.).  
C.V. Mosby Co., 7th. ed., Ch. 8, p. 74.
- \* **Ballinger, J.J. (1985):**  
Acute inflammatory diseases of the middle ear.  
In "Diseases of the Nose, Throat, Ear, Head and Neck."  
(Eds. Lee and Febiger) 13th. ed.  
Philadelphia, p. 1128.
- \* **Barber, M. and Waterworth, P.M. (1962):**  
Antibacterial activity of the penicillins.  
Br. Med. J., 1: 1159-1163.
- \* **Barker, A.E. (1908):**  
Meningitis complicating otitis media, with lumbar puncture.  
Lancet, 1: 854.
- \* **Bartos, L.C. and Murphy, T.F. (1988):**  
Comparison of the outer membrane proteins of  
50 strains of Branhamella catarrhalis.  
J. Infect. Dis., 158: 761-5.
- \* **Baumann, P.; Doudoroff, M. and Stanier, R.Y. (1968):**  
Study of the Moraxella group I genus Moraxella  
and the Neisseria catarrhalis group.  
J. of Bacteriol., 95: 58-73.

- \* **Bauer, A.W.; Kirby, W.M.; Sherris, J.C. and Turck, M. (1966):**  
Antibiotic susceptibility testing by a standardized single disk method.  
American Journal of Clin. Path., 45, 183-190.
- \* **Bergey, D.H.; Harrison, F.C.; Breed, R.S.; Hammer, B.W. and Hinton, F.M. (1923):**  
Bergey's Manual of Determinative Bacteriology.  
Williams and Wilkins, Baltimore.
- \* **Berger, U. (1963):**  
Die anspruchslosen Neisserien.  
Ergebnisse der Mikrobiologie, Immunitätsforschung  
and Experimentelle Therapie, 36: 97-167.
- \* **Biedel, C.S. (1978):**  
Modification of recurrent otitis media by short-term sulfonamide therapy.  
Am. J. Dis. Child., 132: 681-683.
- \* **Bland, R.D. (1972):**  
Otitis media in the first six years of life:  
Diagnosis, bacteriology and management.  
Pediatr., 49: 187.
- \* **Bluestone, C.D. (1975):**  
Obstructive adenoids in relation to otitis media.  
Ann. Otol. Rhinol. Laryngol., 84: 44-48.
- \* **Bluestone, C.D. and Beery, Q.C. (1976):**  
Adenoidectomy in relation to otitis media.  
Ann. Otol. Rhinol. Laryngol., 85: 280-85.

- \* **Bluestone, C.D. (1984):**  
Augmentin therapy for acute otitis media in children.  
Progress and perspectives on Beta-Lactamase Inhibition: A Review of Augmentin.  
N. Engl. J. Med., 306: 1399-1404.
- \* **Bluestone, C.D. and Doyle, W.J. (1985):**  
Eustachian tube function: physiology and role in otitis media.  
Workshope report.  
Ann. Otol. Rhinol. Laryngol., 94: 1-60.
- \* **Bluestone, C.D. (1985):**  
Update on antimicrobial therapy for otitis media and sinusitis in children.  
Cutis, 36: 7-12.
- \* **Bluestone, C.D. (1986):**  
Otitis media and sinusitis in children.  
Role of Branhamella catarrhalis.  
Drugs 31 (Suppl. 3): 132-141.  
ADIS Press Limited.
- \* **Bluestone, C.D. and Klein, J.O. (1988a):**  
Anatomy.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 4-10.
- \* **Bluestone, C.D. and Klein, J.O. (1988b):**  
Definition, Terminology and Classification.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 2-4.

- \* **Bluestone, C.D. and Klein, J.O. (1988c):**  
Epidemiology.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 35-40.
- \* **Bluestone, C.D. and Klein, J.O. (1988d):**  
Physiology, pathophysiology and pathogenesis.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 20-30.
- \* **Bluestone, C.D. and Klein, J.O. (1988e):**  
Microbiology.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 50-52.
- \* **Bluestone, C.D. and Klein, J.O. (1988f):**  
Diagnosis.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.G. Co., Philadelphia, p. 72-80.
- \* **Bluestone, C.D. and Klein, J.O. (1988g):**  
Complication and sequelae.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 200-20.
- \* **Bluestone, C.D. and Klein, J.O. (1988h):**  
Immunology.  
In "Otitis Media in Infants and Children".  
Darlene Cooke.  
Saunders W.B. Co., Philadelphia, p. 60-62.

- \* **Bluestone, C.D. (1988):**  
Management of otitis media in infants and children:  
Current role of old and new antimicrobial agents.  
Pediatr. Infect. Dis. J., 7: 129-136.
- \* **Bodor, F.F.; Marchant, D.D.; Shurin, P.A. and Barenkamp, S.J. (1985):**  
Bacterial etiology of conjunctivitis-otitis media syndrome.  
Pediatr., 76: 26-28.
- \* **Bovre, K. (1967):**  
Transformation and DNA base. Composition in taxonomy, with special reference to recent studies in Moraxella and Neisseria.  
Acta Pathol. Microbio. Scand., 69: 123-44.
- \* **Bovre, K. (1979):**  
Proposal to divide the genus Moraxella into two subgenera, subgenus Moraxella and subgenus Branhamella.  
Inter. J. Syst. Bacteriol., 29: 403-406.
- \* **Bovre, K. (1980):**  
Progress in classification and identification of Neisseria based on genetic affinity.  
In "Microbiological Classification and Identification".  
(Eds. Goodfellow and Board).  
Academic Press, London, New York, p. 55-72.
- \* **Bovre, K. (1981):**  
The family Neisseriaceae.  
In Starr et al. (Eds).  
Springer-Verlag, Berlin, Ch. 122.

- \* **Bovre, K. (1984):**  
Family VIII Neisseriaceae.  
In Krieg (Ed.), Bergey's Manual of Systematic Bacteriology.  
Williams and Wilkins, Baltimore, 9th. ed., Vol. I, p. 288-310.
- \* **Branefors, H.P. (1973):**  
Serological studies of *Haemophilus influenzae* with special reference to otitis media infections.  
Dissertation, University of Goteborg, Sweden.
- \* **Branefors, H.P.; Dahlberg, T. and Nylén, O. (1975):**  
Acute otitis media. A clinical, bacteriological and serological study of children with frequent episodes of acute otitis media.  
Acta Otolaryngol., 80: 399-406.
- \* **Brook, I.; Anthony, B.F. and Finegold, S.M. (1978):**  
Aerobic and anaerobic bacteriology of acute otitis media in children.  
J. Pediatr., 92: 13-15.
- \* **Brook, I.; Pettit, T.H.; Martin, W.J. and Finegold, S.M. (1979):**  
Aerobic and anaerobic bacteriology of acute conjunctivitis.  
Ann. Ophthalmology, 11: 13-16.
- \* **Brook, I. (1981):**  
Bacteriology of paronychia in children.  
Am. J. Surgery, 141: 703-5.

- \* Brook, I. and Schwartz, R. (1981):  
Anaerobic bacteria in acute otitis media.  
Acta Otolaryngol (Stockh.), 91: 111-114.
- \* Brook, I.; Yocum, P. and Friedman, E.M. (1981):  
Aerobic and anaerobic flora recovered from tonsils  
of children with recurrent tonsillitis.  
Ann. Otolaryngology, Rhinology and Laryngology, 90:  
261-63.
- \* Brook, I. and Yocum, P. (1984):  
Bacteriology of chronic tonsillitis in young  
adults.  
Arch. Otolaryngology, 110: 803-5.
- \* Brook, I. and Hirokawa, R. (1985):  
Treatment of patients with a history of recurrent  
tonsillitis due to group A beta-hemolytic strepto-  
cocci.  
Clinical Pediatrics, 24: 331-336.
- \* Brook, I. (1986):  
Direct and indirect pathogenicity of Branhamella  
catarrhalis.  
Drugs 31 (Suppl. 3): 97-102.  
ADIS Press Limited.
- \* Brorson, J.E.; Axelsson, A. and Holm, S.E. (1976):  
Studies on Branhamella catarrhalis with special  
reference to maxillary sinusitis.  
Scand. J. Infect. Dis., 8: 151.
- \* Brorson, J.E. and Malmvall, B.E. (1981):  
Branhamella catarrhalis and other bacteria in  
the nasopharynx of children with longstanding  
cough.  
Scand. J. Infect. Dis., 13: 111.

- \* Brorson, J.E.; Martinell, J. and Wilske, H. (1981):  
Branhamella catarrhalis: antibiotic susceptibility and beta-lactamase production.  
J. Antimicrobial Chemoth., 7: 208-9.
  
- \* Brownlee, R.C.; DeLoache, W.R.; Cowan, C.C. and Jakson, H.P. (1969):  
Otitis media in children: Incidence, treatment and prognosis in pediatric practice.  
J. Pediat., 75: 636-642.
  
- \* Burnet, A.K.; Easton, D.J. and Gordon, A.M. (1975):  
Neisseria catarrhalis septicemia in acute lymphoblastic leukemia.  
Scott. Med. J., 20: 37-38.
  
- \* Bush, K. (1989):  
Characterization of Beta-lactamase: Minireviews.  
Antimicrob. Agents Chemother., 33: 259-263.
  
- \* Calder, M.A.; Croughan, M.J.; Mcleod, D.T. and Ahmad, F. (1986):  
The incidence and antibiotic susceptibility of Branhamella catarrhalis in respiratory infections.  
Drugs, 31 (Suppl. 3): 11-16.
  
- \* Campagnari, A.A.; Gupta, M.R. and Dudas, K.C. (1987):  
Antigenic diversity of lipooligosaccharides of nontypable Haemophilus influenzae.  
Infect. Immun., 55: 882-887.
  
- \* Cantekin, E.I.; Doyle, W.J. and Bluestone, C.D. (1983):  
Effect of levator veli palatini muscle in Eustachian tube function.  
Arch. Otolaryngol., 109: 281-284.

- \* Catlin, B.W. and Cunningham, L.S. (1961):  
Transforming activities and base contents of  
deoxyribonuclease from various *Neisseriae*.  
J. Gen. Microbiol., 26: 303-305.
- \* Catlin, B.W. and Cunningham, L.S. (1964):  
Genetic transformation of *Neisseria catarrhalis*  
by deoxyribonuclease preparations having different  
average base compositions.  
J. Gen. Microbiol., 37: 341-352.
- \* Catlin, B.W. (1970):  
Transfer of the organism named *Neisseria*  
*catarrhalis* to *Branhamella* gen. nov.  
Int. J. Systematic Bacteriology, 20: 155-59.
- \* Chapman, A.J.; Musher, D.M.; Jonsson, S.; Clarridge,  
J.E. and Wallace, R.J. (1985):  
Development of bactericidal antibody during  
*Branhamella catarrhalis* infection.  
J. Infect. Dis., 151: 878-82.
- \* Cheesbrough, M. (1989a):  
Culturing of microorganisms.  
In "Medical Laboratory Manual for Tropical  
Countries".  
English Language Book Society, Vol. II, Ch. 35,  
p. 40.
- \* Cheesbrough, M. (1989b):  
Biochemical testing of microorganisms.  
In "Medical Laboratory Manual for Tropical  
Countries".  
English Language Book Society, Vol II, Ch. 36,  
p. 65.

- \* **Cheesbrough, M. (1989c):**  
Culture media.  
In "Medical Laboratory Manual for Tropical Countries".  
English Language Book Society, Vol. II, Ch. 48,  
p. 398.
- \* **Clarke, R.M. and Haining, R.B. (1936):**  
Neisseria catarrhalis endocarditis.  
Ann. Intern. Med., 10: 117-121.
- \* **Cocchi, P. and Olivelli, A. (1968):**  
Meningitis caused by Neisseria catarrhalis.  
Acta Pediat. Scand., 57: 451-453.
- \* **Coffey, J.D.; Booth, H.N. and Martin, A.D. (1966):**  
Otitis media in the practice of pediatrics:  
bacteriological and clinical observations.  
Pediatr., 38: 25.
- \* **Coffey, J.D.; Booth, H.N. and Martin, A.D. (1967):**  
Neisseria catarrhalis in exudative otitis media.  
Arch. Otolaryngol., 86: 403-406.
- \* **Cohen, L.F.; McNeil, C.J. and Wells, R.F. (1973):**  
Otitis media in the practice of pediatrics.  
JAMA, 223: 1379-1380.
- \* **Collee, J.G. and Miles, R.S. (1989):**  
Tests for identification of bacteria.  
In "Practical Medical Microbiology".  
(Eds. Collee, J.G.; Duguid, J.P.; Fraser, A.G.  
and Marmion, B.P.).  
Churchill Livingstone Co., 13rd. ed., Vol. II,  
Ch. 8, p. 141-160.

- \* Collee, J.G.; Duguid, J.P.; Fraser, A.G. and Marmion, B.P. (1989):  
Practical Medical Microbiology.  
Churchill Livingstone Co., 13rd. ed., Vol. II.
- \* Davies, B.I. (1985):  
Symposium on Branhamella catarrhalis, held at  
Brockham Park.  
Surrey, England.
- \* Davies, B.I. and Maesen, F.P. (1986):  
Epidemiological and bacteriological finding on  
Branhamella catarrhalis respiratory infections  
in the Netherland.  
Drugs, 31 (Suppl. 3): 28-33.
- \* Dever, G.J.; Stewart, J.L. and Davis, A. (1985):  
Prevalence of otitis media in selected populations  
on Pohnpei. A preliminary study.  
Int. J. Pediatr. Otorhinolaryngol., 10: 143-  
152.
- \* De Maria, T.E.; McGhee, R.B. and Lim, D.J. (1984):  
Rheumatoid factor in otitis media with effusion.  
Arch. Otolaryngol., 110: 279-280.
- \* Diamant, M. and Diamant, B. (1974):  
Abuse and timing of use of antibiotics in acute  
otitis media.  
Arch. Otolaryngol., 100: 226-232.
- \* Diamond, L.A. and Lorber, B. (1984):  
Branhamella catarrhalis pneumonia and immuno-  
globulin abnormalities: A new association.  
Am. Rev. Respir. Dis., 129: 876-78.

- \* Doern, G.V. and Morse, S.A. (1980):  
Branhamella catarrhalis: Criteria for laboratory identification.  
J. Clin. Microb., 11: 193-95.
- \* Doern, G.V.; Siebers, K.G.; Hallick, L.M. and Morse, S.A. (1980):  
Antibiotic susceptibility of  $\beta$ -lactamase-producing strains of Branhamella catarrhalis.  
Antimicrob. Agents Chemother., 17: 24-29.
- \* Doern, G.V.; Miller, M.J. and Winn, R.E. (1981):  
Branhamella catarrhalis systemic disease in humans.  
Case reports and review of the literature.  
Arch. Intern. Med., 141: 1690-92.
- \* Doern, G.V. and Gantz, N.M. (1982):  
Isolation of Branhamella (Neisseria) catarrhalis from men with urethritis.  
Sex. Transm. Dis., 9: 202-4.
- \* Doern, G.V. (1985):  
Branhamella catarrhalis: An emerging human pathogen.  
Clin. Microb. Newsletter, 7: 75-78.
- \* Doern, G.V. (1988):  
Branhamella and Neisseria species.  
In "Laboratory Diagnosis of Infectious Diseases, Principles and Practice".  
(Eds. Balous, A.; Housler, W.; Ohoshi, M. and Turano, A.), Ch. 5, p. 217.  
Springer-Verlag Co.

- \* Douer, D.; Danziger, Y. and Pinkhas, J. (1977):  
Neisseria catarrhalis endocarditis.  
Ann. Intern. Med., 86: 116-19.
  
- \* Doyle, R.J.; Nedjat-Haiem, F.; Keller, K.F. and Frascch, C.E. (1984):  
Diagnostic value of interactions between members of the family Neisseriaceae and lectins.  
J. of Clin. Microbiology, 19: 383-387.
  
- \* Dugdale, A.E.; Lewis, A.N. and Canty, A.A. (1982):  
The natural history of chronic otitis media.  
N. Eng. J. Med., 307: 1459-1460.
  
- \* Eliasson, I. (1980):  
A protein antigen characteristic of Branhamella catarrhalis.  
Acta Pathol. Microbiol. Scand., 88: 281-86.
  
- \* Eliasson, I. and Kamme, C. (1985):  
Characterisation of the plasmid-mediated beta lactamases in Branhamella catarrhalis, with special reference to substrate affinity.  
J. Antimicrobial Chemotherapy, 15: 139-149.
  
- \* Eliasson, I. (1986):  
Serological identification of Branhamella catarrhalis.  
Serological evidence for infection.  
Drugs, 31 (Suppl. 3): 7-10.  
ADIS Press Limited.

- \* Eliasson, I. and Kamme, C. (1986):  
Upper respiratory tract infections.  
Ecological and therapeutic aspects of  $\beta$ -lactamase  
production with special reference to Branhamella  
catarrhalis.  
Drugs 31 (suppl. 3): 116-121.
  
- \* Embden, J.; Klingerer, B. and Dessen-Kroon, M. (1980):  
Penicillinase producing Neisseria gonorrhoeae  
in Netherlands: Epidemiology and genetic and  
molecular characterisation of their plasmids.  
Antimicrobial Agents and Chemotherapy, 18: 789-  
797.
  
- \* Enders, G. (1925):  
Der Mikrococcus catarrhalis als Erreger einer  
Sepsis mit Endocarditis und Nephritis.  
MMW, 72: 723-25.
  
- \* Elston, H.R.; Onigley, H.J. and Fitch, D.M. (1970):  
Meningitis due to Neisseria catarrhalis.  
Neb. Med. J., 55: 369-371.
  
- \* Fallon, R.J. and Young, H. (1989):  
Neisseria: Acinetobacter: Branhamella.  
In "Practical Medical Microbiology".  
(Eds. Collee, J.G.; Duguid, J.P.; Fraser, A.G.  
and Marmion, B.P.).  
Churchill Livingstone Co., 13rd. ed., Vol. II,  
Ch. 21, p. 360-73.
  
- \* Fatthi, A. and El-Ashmawy, S. (1980):  
A recent immunological concept of some ear and  
nose diseases.  
J. Laryngol. and Otol., 94: 291-299.

- \* Farmer, T. and Reading, C. (1982):  
Beta-lactamases of Branhamella catarrhalis and their inhibition by clavulanic acid.  
Antimicrobial Agents and Chemotherapy, 21: 506-508.
- \* Farmer, T. and Reading, C. (1986):  
Inhibition of the Beta-lactamases of Branhamella catarrhalis by clavulanic acid and other inhibitors.  
Drugs, 31 (Suppl. 3): 70-78.
- \* Feigin, R.D.; Sanjoaquin, V. and Middelkamp, J.N. (1969):  
Purpura fulminans associated with Neisseria catarrhalis septicaemia and meningitis.  
Ped., 44: 120-23.
- \* Feigin, R. and Spector, G. (1984):  
Otitis Media.  
In "Text Book of Pediatric Infectious Diseases".  
Saunders W.B. Co., Philadelphia, p. 122-137.
- \* Feingold, M.; Klein, J.O.; Haslam, G.E.; Tilles, J.G.; Finland, M. and Gellis, S.S. (1966):  
Acute otitis media in children.  
Am. J. Dis. Child., 111: 361-365.
- \* Freijd, A.; Oxelius, N.A. and Rynnel, D.B. (1985):  
A prospective study demonstrating an association between plasma IgG concentrations and susceptibility to otitis media in children.  
Scand. J. Infect. Dis., 17: 115-120.

- \* **Frosch, P. and Kolle, W. (1896):**  
Die Mikrokokken. In Flügge (Ed.).  
Die Mikro-organismen, 3rd. ed., Part 2, p. 154-155.  
Verlag von Vogel, Leipzig.
- \* **Fox, R.H. and McClain, D.E. (1974):**  
Evaluation of the taxonomic relationship of *Micrococcus cryophilus*, *Branhamella catarrhalis*, and *Neisseriae* by comparative polyacrylamide gel electrophoresis of soluble proteins.  
Int. J. Syst. Bacteriol., 24: 172-76.
- \* **Fox, R.H. and McClain, D.E. (1975):**  
Enzyme electrophoretograms in the analysis of taxon relatedness of *Micrococcus cryophilus*, *Branhamella catarrhalis* and atypical *Neisseria*.  
J. Gen. Microbiol., 86: 210-16.
- \* **Fulghum, R.S.; Daniel, H.J. and Yarborough, J.G. (1977):**  
Anaerobic bacteria in otitis media.  
Ann. Otol. Rhinol. Laryngol., 86: 196-203.
- \* **Giebink, G.S.; Mills, E.L.; Huff, J.S.; Edelman, C.K.; Weber, M.L.; Juhn, S.K. and Quie, P.G. (1979):**  
The microbiology of serous and mucoid otitis media.  
Pediatr., 63: 915-919.
- \* **Goss, C. (Ed.) (1967):**  
Gray's Anatomy of the Human Body.  
Lea & Febiger, Philadelphia, p. 1087.

- \* Goulon, M.; Raphael, J.C.; Barois, A. and Babinet, P. (1973):  
Septicémie et méningite purulente à Neisseria catarrhalis.  
Ann. Med. Interne. (Paris), 124: 741-46.
- \* Gray, B.M.; Convers, G.M. and Dillon, H.C. (1979):  
Serotypes of Streptococcus pneumoniae causing disease.  
J. Infect. Dis., 140: 979-83.
- \* Grimaldi, P.M. (1976):  
The value of impedance testing in diagnosis of middle ear effusion.  
J. Laryngol. Otol., 90: 141-152.
- \* Gronroost, J.A.; Kortenkgas, A.E.; Ojala, I. and Vuori, M. (1964):  
The aetiology of acute ear infection.  
Acta Otolaryngol. (Stockh.), 58: 149: 158.
- \* Hall, I.S. and Colman, B.H. (1981):  
Acute otitis media.  
In "Diseases of the Nose, Throat and Ear".  
Churchill Livingstone Co., 11th. ed., Ch. 32,  
p. 254-267.
- \* Hanson, L.A. and Holm, S. (1961):  
Precipitating antibodies in human sera from different age groups and colostrum as determined by streptococcal antigens with diffusion-in-gel methods.  
Acta. Pediatr. Scand., 50: 7-10.

- \* **Harding, A.L.; Anderson, P.; Howie, V.M.; Polussard, J.H. and Smith, D.H. (1973):**  
Haemophilus influenzae isolated from children with otitis media.  
In "Haemophilus Influenzae".  
(Eds. Sell, S.H. and Karzon, D.T.).  
Vanderbilt University Press, Nashville, p. 21-28.
- \* **Hart, V.K. (1927):**  
The bacteriology of acute ears.  
Laryngoscope 37: 56.
- \* **Henderson, F.W.; Collier, A.M.; Sanyal, M.A.; Watkins, J.M.; Fairclough, D.L.; Clyde, W. and Denny, F.W. (1982):**  
A longitudinal study of respiratory viruses and bacteria in the etiology of acute otitis media with effusion.  
New Eng. J. Med., 306: 1377-83.
- \* **Henriksen, S.D. and Bovre, K. (1968):**  
The taxonomy of the genera Moraxella and Neisseria.  
J. of General Microbiology, 51: 387-392.
- \* **Herbert, W.J. (1989):**  
Management of experimental animals.  
In "Practical Medical Microbiology".  
(Eds. Collee, J.G.; Duguid, J.P.; Fraser, A.G. and Marmion, B.P.).  
Churchill Livingstone Co., 13rd. ed., Vol. II, Ch. 14, p. 277.

- \* Hoi-Dang Van, A.B.; Bouguenec, C.B.; Barthelemy, M. and Labia, R. (1978):  
Novel  $\beta$ -lactamase from Branhamella catarrhalis.  
Ann. Microbiol. (Ins. Pasteur), 129: 397-406.
  
- \* Holmqvist, J. and Larsson, G. (1976):  
Eustachian tube dysfunction.  
Scand. Audiol., 5: 107.
  
- \* Holland, D.F. (1920):  
Generic index of the commoner forms of bacteria.  
Journal of Bacteriology, 5: 215-229.
  
- \* Holten, E. and Jyssum, K. (1974):  
Activities of some enzymes concerning pyruvate metabolism in Neisseria.  
Acta. Pathol. Microbiol.Scand., 82: 843-48.
  
- \* Howie, V.M. and Ploussard, J.H. (1969):  
The "in vivo sensitivity test".  
Bacteriology of middle ear exudate during anti-microbial therapy in otitis media.  
Pediatrics, 44: 940-944.
  
- \* Howie, V.M.; Ploussard, J.H. and Lester, R.L. (1970):  
Otitis media: a clinical and bacteriological correlation.  
Pediatr., 45: 24.
  
- \* Howie, V.M.; Ploussard, J.H.; Sloyer, J.L. and Johnston, R.B. (1973):  
Immunoglobulins of the middle ear fluid in acute otitis media.  
Infect. Immun., 7: 589-593.

- \* Ingvarsson, L.; Lundgren, K. and Ursing, J. (1982a):  
The bacterial flora in the nasopharynx in healthy children.  
Acta Otolaryngol. (Stockh.), 386: 94-6.
- \* Ingvarsson, L.; Lundgren, K.; Olofsson, B. and Wall, S. (1982b):  
Epidemiology of acute otitis media in children,  
Acta Otolaryngol. (Stockh.), 388: 5.
- \* Jawetz, E.; Brooks, G.F.; Melnick, J.L.; Butel, J.S.;  
Adelberg, E.A. and Ornston, L.N. (1989):  
Neisseria. In "Review of Medical Microbiology".  
Lange Medical Book, California, 18th. ed., Ch.  
21, p. 238-43.
- \* Johnson, K.G.; McDonald, I.J. and Perry, N.B. (1976):  
Studies on the cellular and free lipopolysaccharides from Branhamella catarrhalis.  
Can. J. Microbiol., 22: 460-67.
- \* Johnson, M.A.; Drew, W.L. and Montgomery, K. (1979):  
Beta-lactamase-producing Branhamella catarrhalis:  
A respiratory pathogen, in Abstracts of the Annual  
Meeting of the Am. Soc. for Microb. Washington,  
D.C.  
Am. Soc. for Microb., p. 336.
- \* Johnson, M.A.; Drew, W.L. and Roberts, M. (1981):  
Branhamella (Neisseria) catarrhalis - a lower  
respiratory tract pathogen?  
J. Clin. Microbiology, 13: 1066-69.

- \* **Johnson, A.P. (1983):**  
The pathogenic potential of commensal species of *Neisseria*.  
*J. Clin. Pathol.*, 36: 213-223.
- \* **Johnsson, J. and Brorson, J.E. (1983):**  
Influence of beta-lactamase-producing strains of *Branhamella catarrhalis* and *Haemophilus influenzae* on certain beta-lactam antibiotics.  
*J. Antimicrobial Chemotherapy*, 12: 269-271.
- \* **Jones, M. (1974):**  
Comparison of bacteria from ear and upper respiratory tract in otitis media.  
*Arch. Otolaryngol.*, 72: 329-36.
- \* **Jones, R.N. and Sommers, H.M. (1986):**  
Identification and antimicrobial susceptibility testing of *Branhamella catarrhalis* in United States Laboratories, (1983-1985).  
*Drugs*, 31 (Suppl. 3): 34-37.
- \* **Kaleida, P.H.; Casselbrant, M.L. and Bluestone, C.D. (1987a):**  
Antibiotic or myringotomy or both for acute otitis media?  
Abstracts of the 4th. Inter. Symposium on Recent Advances in Otitis Media.  
Columbus, OH., Ohio State University School of Medicine, p. 69-70.
- \* **Kaleida, P.H.; Bluestone, C.D. and Rockette, H.E. (1987b):**  
Amoxicillin-clavulanate potassium in comparison with cefaclor for acute otitis media in infants and children.  
*Pediatr. Infect. Dis. J.*, 6: 265-271.

- \* Kallings, I; Bengtsson, S.; Christensen, P.; Holm, S.E. and Lind, L (1983):  
Antibiotic sensitivity of Haemophilus influenzae, Streptococcus pneumoniae, Streptococcus pyogenes and Branhamella catarrhalis isolated from upper respiratory tract infections in Sweden.  
Scand. J. Infect. Dis., 39: 100-105.
- \* Kallings, I. (1986):  
Sensitivity of Branhamella catarrhalis to oral antibiotics.  
Drugs, 31 (Suppl. 3): 17-22.
- \* Kamme, C. (1970):  
Evaluation of the in vitro sensitivity of Neisseria catarrhalis to antibiotics with respect to acute otitis media.  
Scand. J. Infect. Dis., 2: 117-20.
- \* Kamme, C.; Lundgren, K. and Mardh, P.A. (1971):  
The aetiology of acute otitis media in children.  
Scand. J. Infect. Dis., 3: 217-221.
- \* Kamme, C.; Vang, M. and Stahl, S. (1983):  
Transfer of beta-lactamase production in Branhamella catarrhalis.  
Scand. J. Infect. Dis., 15: 225-226.
- \* Kamme, C. and Nilsson, N.I. (1984):  
Secretory of otitis media.  
Microbiology of the middle ear and nasopharynx.  
Scand. J. Infect. Dis., 16: 291.

- \* Kamme, C.; Vang, M. and Stahl, S. (1984):  
Intergeneric and intrageneric transfer of Branhamella catarrhalis, beta-lactamase production.  
Scand. J. Infect. Dis., 16: 153-55.
  
- \* Kamme, C.; Eliasson, I.; Knutson, B. and Vang, M. (1986):  
Plasmid-mediated Beta - lactamase in Branhamella catarrhalis.  
Drugs, 31 (Suppl. 3): 55-63.
  
- \* Karma, P.; Virtanen, T.; Pukander, J.; Sipila, M.; Luotonen, J. and Herva, E. (1985):  
Branhamella catarrhalis in acute otitis media.  
Acta Otolaryngol. (Stockh.), 99: 285-90.
  
- \* Kaufman, B.A. and Likavec, M.J. (1985):  
Branhamella catarrhalis cellulitis around cerebro-spinal fluid shunt: Case report.  
J. Hosp. Infect., 6: 393.
  
- \* Kenna, M.A.; Bluestone, C.D. and Fall, P. (1987):  
Cefixime vs. cefaclor in the treatment of acute otitis media in infants and children.  
Pediatr. Infect. Dis. J., 6: 992-996.
  
- \* Kinnman, J.; Lee, C.H. and Park, S.H. (1967):  
Bacterial flora in chronic, purulent maxillary sinusitis.  
Acta. Otolaryngol. (Stockh.), 64: 37.
  
- \* Klein, J.O. and Teele, D.W. (1976):  
Isolation of viruses and mycoplasmas from middle ear effusions: A review.  
Ann. Otol. Rhinol. Laryngol., 85: 140-144.

- \* Klein, J.O. (1980):  
Microbiology of otitis media.  
Ann, Otol. Rhinol. Laryngol., 89: 98-102.
- \* Klein, J.O. and Marcy, S.M. (1983):  
Bacterial sepsis and meningitis.  
In "Infectious Diseases of the Fetus and Newborn Infant".  
(Eds. Remington, J.S. and Klein, J.O.).  
Saunders W.B. Co., Philadelphia, p. 679-735.
- \* Knapp, J.S. and Holmes, K.K. (1983):  
Modified oxidation - fermentation medium for  
detection of acid production from carbohydrate  
by *Neisseria* species and *Branhamella catarrhalis*.  
J. Clin. Microbiol., 18: 56-59.
- \* Kovatch, A.L.; Wald, E.R. and Michaels, R.H. (1983):  
 $\beta$ -lactamase-producing *Branhamella catarrhalis*  
causing otitis media in children.  
J. Pediatr., 102: 261-4.
- \* Labia, R.; Barthelemy, M.; Le Bouguennec, C. and Hoi-Dang Van, A. (1986):  
Classification of beta-lactamase from *Branhamella catarrhalis* in relation to penicillinases produced by other bacterial species.  
Drugs, 31 (Suppl. 3): 40-47.
- \* Laxdal, O.E.; Blake, R.M.; Cartmill, T. and Robertson, H.E. (1966):  
Etiology of acute otitis media in infants and children.  
Can. Med. Assoc. J., 94: 159-163.

- \* Lee, W.S.; Fordham, T. and Alban, J. (1981):  
Otitis media caused by beta-lactamase-producing  
Branhamella catarrhalis.  
J. of Clin. Microbiol., 13: 222.
  
- \* Le Faou, A. and Rio, A. (1983):  
Branhamella catarrhalis en pathologie infectieuses  
des voies aériennes.  
Pathologie et Biologie, 33: 2035-37.
  
- \* Leinonen, M.; Luotonen, J.; Herva, E.; Valkonen, K.  
and Makela, P.H. (1981):  
Preliminary serologic evidence for a pathogenic  
role of Branhamella catarrhalis.  
J. Infect. Dis., 144: 570-74.
  
- \* Levine, L.R. (1985):  
Quantitative comparison of adverse reactions  
to cefaclor vs. amoxicillin in a surveillance  
study.  
Pediatr. Infect. Dis., 4: 358-361.
  
- \* Lewis, D.M.; Birck, H.G.; Schram, J.L. and Lim, D.J.  
(1979):  
Antibody activity in otitis media with effusion.  
Ann. Otol. Rhinol. Laryngol., 88: 393-96.
  
- \* Lewis, V.J.; Weaver, R.E. and Hollis, D.G. (1968):  
Fatty acid composition of Neisseria species as  
determined by gas chromatography.  
J. Bacteriol., 96: 1-3.
  
- \* Liston, T.E.; Foshee, W.S. and Pierson, W.D. (1983):  
Sulfisoxazole chemoprophylaxis for frequent otitis  
media.  
Pediatr., 71: 524-530.

- \* Long, S.; Herrtig, F.; Teter, M. and McGowan, K. (1983):  
Nasopharyngeal flora and acute otitis media.  
Infect. Immun., 41: 987-991.
  
- \* Louis, M.H.; Gabay, E.L.; Finegold, S. and Mathisen, G.E. (1983):  
Branhamella catarrhalis pneumonia.  
West. J. Med., 138: 47-49.
  
- \* Lundgren, K. and Ingvarsson, L. (1986):  
Acute otitis media in Sweden.  
Role of Branhamella catarrhalis and the rationale  
for choice of antimicrobial therapy.  
Drugs, 31 (Suppl. 3): 125-131.
  
- \* Luotonen, J.; Herva, E.; Karma, P.; Timonen, M.;  
Leinonen, M. and Maleka, P.H. (1981):  
Bacteriology of acute otitis media in children  
with special reference to Streptococcus pneumoniae.  
Scand. J. Infect. Dis., 13: 177-183.
  
- \* Maddocks, J.L. and May, J.R. (1969):  
"Indirect pathogenicity" of penicillinase producing  
enterobacteria in chronic bronchial infections.  
Lancet, 1: 793-795.
  
- \* Malkamaki, M.; Honkanen, E.; Leinonen, M. and Makela, H.P. (1983):  
Branhamella catarrhalis as a cause of bacteremic  
pneumonia.  
Scand. J. Infect. Dis., 15: 125-26.

- \* **Malmvall, B.E.; Brorson, J.E. and Johnson, J. (1977):**  
In vitro sensitivity to penicillin V and  $\beta$ -lactamase production of Branhamella catarrhalis (letter).  
J. Antimicrob. Chemother., 3: 374-75.
- \* **Marchant, C.D.; Shurin, P.A.; Turczyk, V.A.; Wasikowski, D.E.; Tutihasa, M.A. and Kinney, S.E. (1984):**  
Course and outcome of otitis media in early infancy: A prospective study.  
J. Pediatr., 104: 826-831.
- \* **Marchant, C.D.; Shurin, P.A. and Johnson, C.E. (1986):**  
A randomized controlled trial of amoxicillin plus clavulanate compared with cefaclor for treatment of acute otitis media.  
J. Pediatr., 109: 891-896.
- \* **Marrs, C.F.; Schoolnik, G. and Koomey, J.M. (1985):**  
Cloning and sequencing of a Moraxella bovis pilin gene.  
J. Bacteriol., 163: 132-9.
- \* **Marrs, C.F.; Ruehl, W.W. and Schoolnik, G. (1988):**  
Pilin gene phase variation of Moraxella bovis is caused by an inversion of the pilin genes.  
J. Bacteriol., 170: 3032-9.
- \* **Mawson, S.R. and Ludman, H. (1979):**  
Otitis Media.  
Diseases of the Ear, A Textbook of Otology.  
Arnold, E., London, 4th. ed., p. 305.

- \* Mawson, S.R. (1979):  
Acute inflammation of the middle ear cleft.  
In "Scott-Brown's Diseases of the Ear, Nose and Throat".  
Butterworths, London, 4th. ed., Vol. II, p. 175.
- \* Maynard, J.E.; Fleshman, J.K. and Tschopp, C.F. (1972):  
Otitis media in Alaskan Eskimo children: Prospective evaluation of chemotherapy.  
JAMA, 219: 597-599.
- \* McCague, J.J.; McCague, N.J. and Altman, C.C. (1976):  
Neisseria catarrhalis urethritis: a case report.  
J. Urol., 115: 471.
- \* McLeod, D.T.; Ahmad, F.; Power, J.T.; Calder, M.A. and Seaton, A. (1983):  
Bronchopulmonary infection due to Branhamella catarrhalis.  
Br. Med. J., 287: 1446-1447.
- \* McLeod, D.T.; Ahmad, F. and Calder, M.A. (1985):  
Infection and Branhamella catarrhalis.  
Biomed. Pharmacother., 39: 170-3.
- \* McLeod, D.T.; Ahmad, F.; Calder, M. and Croughan, M. (1986):  
Bronchopulmonary infection due to Branhamella catarrhalis.  
Clinical features and therapeutic response.  
Drugs, 31 (Suppl. 3): 109-112.

- \* **McNeely, D.J.; Kitchens, C.S. and Kluge, R.M. (1976):**  
Fatal Neisseria (Branhamella) catarrhalis pneumonia  
in an immunodeficient host.  
Am. Rev. Respir. Dis., 114: 399-402.
- \* **Meri, S.; Lehtinen., T. and Palva, T. (1984):**  
Complement in chronic secretory otitis media.  
Arch. Otolaryngol., 110: 774-778.
- \* **Meurman, O.H.; Sarkkinen, H.K.; Puhakka, H.J. and  
Virolainen, E.S. (1980):**  
Local IgA class antibodies against respiratory  
viruses in middle ear and nasopharyngeal secretion  
of children with otitis media.  
Laryngoscope, 90: 304-311.
- \* **Mogi, G.; Yoshida, T.; Honjo, S. and Maeda, S. (1973):**  
Middle ear effusions: Quantitative analysis of  
immunoglobulins.  
Ann. Otol. Rhinol. Laryngol., 82: 196-202.
- \* **Moon, C.N. and Hahn, M. (1978):**  
Pneumatic otoscopy and impedance studies in middle  
ear disease diagnosis.  
Laryngoscope, 88: 1439-1442.
- \* **Mortimer, E.A. and Watterson, R.L. (1956):**  
A bacteriologic investigation of otitis media  
in infancy.  
Pediatr., 17: 359-366.
- \* **Murphy, T.F. (1989):**  
The surface of Branhamella catarrhalis: a  
systematic approach to the surface antigen of  
an emerging pathogen.  
J. Pediatr. Infect. Dis., 8: 75-77.

- \* Mygind, N.; Melstrup, K.I.; Thomson, J.; Thomsen, V.F.; Josefsson, K. and Sorenson, H. (1981):  
Penicillin in acute otitis media: A double blind placebo-controlled trial.  
Clin. Otolaryngol., 6: 5-13.
- \* Nash, D.R.; Wallace, R.J.; Shurin, P.A. and Steingrube, V.A. (1986):  
Isoelectric focusing of Beta-lactamase from sputum and middle ear isolates of Branhamella catarrhalis in the United States.  
Drugs, 31 (Suppl. 3): 48-54.
- \* Newing, W.J. and Christie, R. (1947):  
Meningitis: Isolation of an organism resembling Neisseria catarrhalis from cerebrospinal fluid; report of a case.  
Med. J., 1: 306-307.
- \* Nicotra, B.; Rivera, M.; Luman, J.I. and Wallace, R.J. (1985):  
Branhamella catarrhalis as a lower respiratory tract pathogen in patients with chronic lung disease.  
Arch. Intern. Med., 146: 890.
- \* Nilson, B.W.; Poland, R.L.; Thompson, R.S.; Morehead, D.; Baghdassarian, A. and Carver, D.H. (1969):  
Acute otitis media: treatment results in relation to bacterial etiology.  
Pediatr., 43: 351-55.
- \* Ninane, G.; Joly, J.; Piot, P. and Kraytman, M. (1977):  
Branhamella (Neisseria) catarrhalis as a pathogen.  
Lancet, 2: 149.

- \* Ninan, G.; Joly, J. and Kraytman, M. (1978a):  
Bronchopulmonary infection due to Branhamella catarrhalis: 11 cases assessed by transtracheal puncture.  
Br. Med. J., 1: 276-78.
- \* Ninan, G.; Joly, J.; Kraytman, M. and Piot, P. (1978b):  
Bronchopulmonary infection due to beta-lactamase producing Branhamella catarrhalis treated with amoxycillin/clavulanic acid.  
Lancet, 2: 257.
- \* Noguchi, T.T.; Nachum, R. and Lawrence, C.A. (1963):  
Acute purulent meningitis caused by chromogenic Neisseria - a case report and review of the literature.  
Med. Arts. Sci., 11: 11-18.
- \* O'Callaghan, C.H.; Morris, A.; Kirby, S.M. and Shingler, A.H. (1972):  
Novel method for detection of beta-lactamase by using a chromogenic cephalosporin substrate.  
Antimicrobial Agents and Chemotherapy, 1: 283-288.
- \* Odio, C.M.; Kusmiesz, H.; Shelton, S. and Nelson, J.D. (1985):  
Comparative treatment trial of Augmentin versus cefaclor for acute otitis media with effusion.  
Pediatr., 75: 819-826.
- \* Orsini, A.; Tamalet, J. and Chanas, P. (1954):  
Septicemia mortelle à Neisseria catarrhalis.  
Sem. Hop. Paris, 62: 3498-9.

- \* Paradise, J.L.; Bluestone, C.D. and Felder, H. (1969):  
The universonality of otitis media in fifty infants  
with cleft palate.  
Pediatr., 44: 35-42.
- \* Paradise, J.L. and Bluestone, C.D. (1974):  
Early treatment of universal otitis media in  
infants with cleft palate.  
Pediatrics, 53: 48-54.
- \* Paradise, J.L. (1976):  
Pediatrician's view of middle ear effusions:  
more questions than answers.  
Ann. Otol. Rhinol. Laryngol., 85: 20-23.
- \* Paradise, J.L.; Bluestone, C.D. and Rogers, K.D. (1987):  
Efficacy of adenoidectomy for recurrent otitis  
media: Results from parallel random and nonrandom  
trials.  
Pediatr. Res., 21: 286-290.
- \* Park, C.H.; Lopez, J.S. and Cook, C.B. (1978):  
Methods for detecting beta-lactamase.  
Antimicrobial Agents and Chemotherapy, 13: 318-  
320.
- \* Percival, A.; Corkill, J.E. and Rowland, J. (1977):  
Pathogenicity of, and beta-lactamase production  
by Branhamella catarrhalis.  
Lancet, 2: 1175.
- \* Perrin, J.M.; Charney, E. and MacWhinney, J.B. (1974):  
Sulfisoxazole as chemoprophylaxis for recurrent  
otitis media.  
N. Eng. J. Med., 291: 664-667.

- \* Pfister, L.E.; Gallagher, M.V. and Potterfield, T.G.  
(1965):  
Neisseria catarrhalis bacteremia with meningitis.  
JAMA: 193: 399-401.
  
- \* Philippon, A.; Riou, J.Y.; Sotolongo, F. and  
Guibourdenche, M. (1986):  
Detection, distribution and inhibition of  
Branhamella catarrhalis beta-lactamases.  
Drugs, 31 (Suppl. 3 ): 64-69.
  
- \* Piot, P. and Roberts, M. (1979):  
Beta-lactamase production in commensal Neisser-  
iaceae.  
Lancet, 1: 619.
  
- \* Pollard, J.A.; Richard, J.; Wallace, J.; Donald, R.N.  
and Wilson, R.W. (1986):  
Incidence of Branhamella catarrhalis in the sputa  
of patients with chronic lung disease.  
Drugs, 31 (Suppl. 3): 103-108.
  
- \* Pollock. A.A. and Holzman, R.S. (1976):  
Neisseria catarrhalis endocarditis.  
Ann. Intern. Med., 85: 206-7.
  
- \* Poxton, I.R.; Brown, R. and Wilkinson, J.F. (1989):  
PH measurement and Buffers.  
In "Practical Medical Microbiology".  
(Eds. Collee, J.G.; Duguld, J.P.; Fraser, A.G.  
and Marmion, B.P.).  
Churchill Livingstone Co., 13rd. ed., Vol. II,  
Ch. 5, p. 93.

- \* **Prellner, K.; Christensen, P.; Hovelius, B. and Rosen, C. (1984):**  
Nasopharyngeal carriage of bacteria in otitis prone and non otitis-prone children in day-care centers.  
Acta Otolaryngol., 98: 343.
  
- \* **Proctor, B. (1977):**  
Embryology and anatomy of the Eustachian tube.  
Arch. Otolaryngol., 86: 503-526.
  
- \* **Puczynsk, M.S.; Stankiewicz, J.A. and O'Keefe, J.P. (1987):**  
Single dose amoxicillin treatment of acute otitis media.  
Laryngoscope, 97: 16-18.
  
- \* **Pukander, J.; Luotonene, J.; Sipila, M. and Karma, P. (1982):**  
Incidence of acute otitis media.  
Acta Otolaryngol. (Stockh.), 93: 447-453.
  
- \* **Rey, A. (1974):**  
Branhamella.  
In Krieg (Ed.), Bergey's Manual of Systematic Bacteriology.  
Williams and Wilkins, Baltimore, 8th. ed., p.432.
  
- \* **Rifkind, D.R.; Chanock, R.M.; Kravetz, H.; Johnson, K. and Knight, V. (1962):**  
Ear involvement (myringitis) and primary atypical pneumonia following inoculation of volunteers with Eaton agent.  
Ann. Rev. Respir. Diseases, 85: 479-489.

- \* Riou, J.Y.; Guibourdenche, M. and Popoff, M.Y. (1983):  
A new taxon in the genus *Neisseria*.  
*Annales de Microbiologie (Institut Pasteur)*,  
134: 257-267.
  
- \* Riou, J.Y. and Guibourdenche, M. (1986):  
*Branhamella catarrhalis*: New methods of bacterial  
diagnosis.  
*Drugs*, 31 (Suppl. 3): 1-5.  
ADIS Press Limited.
  
- \* Rogers, J.R.; Adamson, D.M.; Spaulding, L.J. and  
Blevins, W.L. (1982):  
Abstr. Annu. Meet.  
*Am. Soc. Microbiol.*, Ch. 13, p. 273.
  
- \* Rohn, D.D.; Vatman, F. and Cantekin, E.I. (1983):  
Incidence of organisms in otitis media.  
*Ann. Otol. Rhinol. Laryngol.*, 96 (Suppl. 107):  
17.
  
- \* Rudin, R.; Welin, L.; Svardsudd, K. and Tibblin, G.  
(1985)  
Middle ear disease in samples from the general  
population.  
*Acta Otolaryngol. (Stockh.)*, 99: 53-59.
  
- \* Ruokonen, J.; Sandelin, K. and Makinen, J. (1979):  
Adenoid and otitis media.  
*Ann. Otol. Rhinol. Laryngol.*, 88: 166-171.
  
- \* Russell, R.R. and McDonald, I.J. (1976):  
Comparison of the cell envelope proteins of *Micro-*  
*coccus cryophilus* with those of *Neisseria* and  
*Branhamella* species.  
*Can. J. Microbiol.*, 22: 309-12.

- \* Russell, R R.; Johnson, K.G. and McDonald, I.J. (1978):  
Immunological relationships of some non pathogenic  
Neisseria.  
Can. J. Microbiol., 24: 189-91.
  
- \* Saito, A.; Yamaguchi, K.; Shigeno, Y.; Kohno, S.;  
Shigeno, H.; Kusano, N. and Hara, K. (1986):  
Clinical and bacteriological evaluation of  
Branhamella catarrhalis in respiratory infections.  
Drugs, 31 (Suppl. 3): 87-92.
  
- \* Schaefer, O. (1971):  
Otitis media and bottle feeding.  
An epidemiological study of infants in Canadian  
Eskimos.  
Can. J. Public Health, 62: 478-89.
  
- \* Schachter, J.; Grossman, M.; Holt, J.; Sweet, R.;  
Goodner, E. and Mills, J. (1979):  
Prospective study of chlamydial infection in  
neonates.  
Lancet, 2: 377-379.
  
- \* Schalen, L.; Christensen, P.; Kamme, C. and Pettersson,  
K. (1980):  
High isolation rates of Branhamella catarrhalis  
from the nasopharynx in adults with acute  
laryngitis.  
Scand. J. Infect. Dis., 12: 277-80.
  
- \* Schalen, L.; Christensen, P.; Eliasson, I.; Kamme,  
C. and Schalen, C. (1985):  
Inefficacy of penicillin V in acute laryngitis  
in adults; evaluation from results of double-  
blind study.  
Annals of Otology, Rhinology and Laryngology,  
94: 14-17.

- \* Schoutens, E.; DeKoster, J.P.; Tombroff, M. and Yourrassowsky, E. (1973):  
Use of transtracheal aspiration in the bacteriological diagnosis of bronchopulmonary infections.  
Biomedicine, 19: 160.
- \* Schreiner, A.; Digranes, A. and Myking, O. (1972):  
Transtracheal aspiration in the diagnosis of lower respiratory tract infections.  
Scand. J. Infect. Dis., 4: 49.
- \* Schreiner, A.; Bjerkestrand, G.; Digranes, A.; Halvorsen, J.F. and Kommerndal, T.M. (1978):  
Bacteriological findings in the transtracheal aspirate from patients with acute exacerbation of chronic bronchitis.  
Infection, 6: 54-57.
- \* Schreiner, A. (1984):  
Pathogenicity of Branhamella catarrhalis.  
Acta. Otolaryngol. (Stockh.), 407: 40-42.
- \* Schwartz, R.; Rodriguez, W.J.; Khan, W.N. and Ross, S. (1977):  
Acute purulent otitis media in children older than 5 years. Incidence of Haemophilus as a causative organism.  
Am. Med. Assoc., 238: 1032-37.
- \* Schwartz, R.; Rodriguez, W.J.; Mann, R.; Khan, W. and Ross, S. (1979):  
The Nasopharyngeal culture in acute otitis media.  
A reappraisal of its usefulness.  
JAMA, 241: 2170-2173.

- \* **Scott, A. and Brown, W.G. (1972):**  
Diseases of the Ear, Nose and Throat.  
(Eds. Battantyne, J. & Greves, J.).  
Butterworths, London, 3rd. ed., Vol. II, p.265.
- \* **Scott, A.C. (1989):**  
Laboratory control of antimicrobial therapy.  
In "Practical Medical Microbiolog".  
(Eds. Collee, J.G.; Duguid, J.P.; Fraser, A.G.  
and Marmion, B.P.).  
Churchill Livingstone Co., 13rd. ed., Vol. II,  
Ch. 9, p. 17.
- \* **Senturia, B.H.; Bluestone, C.D.; Lim, D.J. and Klein,  
J.O. (1980):**  
Recent advances in otitis media with effusion.  
Ann. Otol. Rhinol. Laryngol., 89: 361-362.
- \* **Shah, K.N. and Desai, M.P. (1969):**  
Ascaris lumbricoides from the right ear.  
Indian Pediatr., 6: 92-93.
- \* **Shambaugh, G.E. (1967):**  
Surgery of the ear.  
Saunders, W.B. Co., Philadelphia, London, Toronto,  
2nd. ed., p. 203.
- \* **Shambaugh, G.E. and Quie, P.G. (1973):**  
Acute otitis media and Mastoiditis.  
In "Otolaryngology".  
Saunders W.B. Co., Philadelphia, London, Toronto,  
Vol. II, p. 113.

- \* Shurin, P.A.; Pelton, S.I. and Klein, J.O. (1976):  
Otitis media in newborn infants.  
Ann. Otol. Rhinol. Laryngol., 85: 216-221.
- \* Shurin, P.A.; Howie, V.M.; Pelton, S.I.; Ploussard, J.H. and Klein, J.O. (1978):  
Bacterial etiology of otitis media during the first six weeks of life.  
J. Pediatr., 92: 893-97.
- \* Shurin, P.A.; Marchant, C.D.; Kim, C.H.; Van Hare, G.F. and Johnson, C.E. (1983):  
Emergence of beta-lactamase-producing strains of Branhamella catarrhalis as important agents of acute otitis media.  
Pediatr. Infect. Dis., 2: 34-38.
- \* Shurin, P.A.; Marchant, C.D.; Van Hare, G.F. and Johnson, C.E. (1985):  
Branhamella catarrhalis - an indirect or causal pathogen?  
(Abstract), 14th. International Congress of Chemotherapy, Kyoto, Japan, June.
- \* Shurin, P.A. (1986):  
Branhamella catarrhalis.  
Drugs, 31 (Suppl. 3): 38-39.
- \* Shurin, P.A. and Van Hare, G.F. (1986):  
Therapy of acute otitis media caused by Branhamella catarrhalis.  
Preliminary report.  
Drugs, 31 (Suppl. 3): 122-124.

- \* Simpson, I.N. and Plested, S.J. (1983):  
The origin and properties of satellite bands  
seen in isoelectric focusing.  
J. Antimicrobial Chemotherapy, 12: 127-131.
- \* Slevin, N.J.; Aitken, J.M. and Thornley, P.E. (1984):  
Clinical and microbiological features of Bran-  
hamella catarrhalis, bronchopulmonary infections.  
Lancet, 1: 782-783.
- \* Slocombe, B. (1986):  
Inhibition of beta-lactamase in Branhamella catarr-  
halis.  
Drugs, 31 (Suppl. 3): 79-81.
- \* Sly, R.M.; Zambie, M.F.; Fraser, M. and Fernandes,  
D.A. (1980):  
Tympanometry in Kindergarten children.  
Ann. Allergy, 44: 1-7.
- \* Sng, E.H.; Yeo, K.L., and Rajan, V.S. (1981):  
Simple method for detecting penicillinase producing  
Neisseria gonorrhoeae.  
Br. J. Venereal Dis., 57: 141-142.
- \* Sorenson, H. (1977):  
Antibiotics in suppurative otitis media.  
Otolaryngol. Clin. North. Am., 10: 45-50.
- \* Spark, R.P.; Dahlberg, P.W. and LaBell, J.W. (1979):  
Ophthalmia neonatorum: Branhamella (Neisseria)  
catarrhalis conjunctivitis.  
Infection, 7: 55-56.

- \* **Sprinkle, P.M. and Veltri, R.W. (1971):**  
Microbiology of the middle ear and nasopharynx.  
Ann. Otol. Rhinol. Laryngol., 80: 854-864.
  
- \* **Srinivasan, G.; Raff, M.J. and Templeton, W.C. (1981):**  
Branhamella catarrhalis pneumonia - report of  
two cases and review of the literature.  
Am. Rev. Resp. Dis., 123: 553-555.
  
- \* **Stickler, G.B.; Rubenstein, M.M.; McBean, J.B.;  
Hedgecock, L.D.; Hugstad, J.A. and Griffing,  
T. (1967):**  
Treatment of acute otitis media in children.  
Am. J. Dis. Child., 114: 123-126.
  
- \* **Stobberingh, E.E.; Davies, B.I. and Boven, C.P. (1984):**  
Branhamella catarrhalis: antibiotic sensitivities  
and beta-lactamases.  
J. Antimicrobial Chemotherapy: 13: 55-64.
  
- \* **Stobberingh, E.E.; Van-Eck, H.J.; Houben, A.W. and  
Van Boven, C.P. (1986):**  
Analysis of the relationship between Ampicillin  
resistance and beta-lactamase production in  
Branhamella catarrhalis.  
Drugs, 31 (Suppl. 3): 23-27.
  
- \* **Sugita, R.; Kawa Mura, S. and Hareda, K. (1983):**  
Chemotherapy of acute otitis media.  
J. Otolaryngology, 83: 1381-87.
  
- \* **Sweeney, K.G.; Verghese, A. and Needham, C.A. (1985):**  
In vitro susceptibilities of isolates from patients  
with Branhamella catarrhalis pneumonia compared  
with those of colonizing strains.  
Antimicrobial Agents and Chemotherapy, 27: 499-  
502.

- \* Sykes, R.B. (1978):  
Methods for detecting beta-lactamases.  
In "Laboratory Methods in Antimicrobial Chemotherapy".  
(Eds. Reeves, D.S.; Phillips, I.; Williams, J.D. and Wise, R.).  
Churchill Livingstone, Edinburgh, Ch. 7, p. 64-69.
- \* Taylor, G.D. (1972):  
The bifid uvula.  
Laryngoscope, 82: 771-778.
- \* Teele, D.W.; Klein, J.O. and Rosner, B.A. (1980):  
Epidemiology of otitis media in children.  
Ann. Otol. Rhinol. Laryngol., 89: 5-6.
- \* The Oxoid Manual (1982):  
Culture Media, Ingredients and Other Laboratory Services.  
5th. ed. Oxoid Limited.
- \* Thornley, E.P.; Aitken, J.M.; Slevin, N.; MacVicar, J. and Drennan, C.J. (1982):  
Branhamella catarrhalis infection of the lower respiratory tract: Reliable diagnosis by sputum examination.  
Br. Med. J., 285: 1537-1538.
- \* Thornley, E.P.; Aitken, J.M.; Nichol, G.M. and Slevin, N.J. (1986):  
Amoxycillin-Clavulanic acid combination in broncho-pulmonary infection due to beta-lactamase-producing Branhamella catarrhalis.  
Drugs, 31 (Suppl. 3): 113-114.

- \* Tilles, J.G.; Klein, J.O.; Jao, R.L.; Haslam, J.E.;  
Feingold, M.; Gellis, S.S. and Finland, M. (1967):  
Acute otitis media in children.  
New Eng. J. Med., 227: 613-18.
  
- \* Timmermans, F.J. and Gerson, S. (1980):  
Otitis media in bottle fed infants and children.  
Can. Med. Assoc. J., 122: 545-47.
  
- \* Van Buchem, F.L.; Dunk, J.H. and Vanthof, M.A. (1981):  
Therapy of acute otitis media: Myringotomy, anti-  
biotics, or neither?  
A double-blind study in children.  
Lancet, 2: 883-87.
  
- \* Van Dishoeck, H.E. and Franssen, M.G. (1957):  
The incidence and correlation of allergy and  
chronic maxillary sinusitis.  
Pract. Otolary., 19: 502.
  
- \* Van Hare, G.F.; Shurin, P.A.; Marchant, C.D.; Johnson,  
C.E. and Kim, C.H. (1987):  
Acute otitis media caused by Branhamella catarrha-  
lis: biology and therapy.  
Rev. Infect. Dis., 9: 16-27.
  
- \* Vedros, N.A. (1984):  
Genus I Neisseria.  
In Krieg (Ed.), Bergey's Manual of Systematic  
Bacteriology.  
Williams and Wilkins, Baltimore, Vol.1: 290-96.
  
- \* Veltri, R.W. and Sprinkle, P.M. (1973):  
Serous otitis media: Immunoglobulin and lysozyme  
levels in middle ear fluids and serum.  
Ann. Otol. Rhinol. Laryngol., 82: 297-301.

- \* Virgil, M.; Howie, J.H.; Ploussard, L. and Richard, L. (1970):  
Otitis Media: a clinical and bacteriological correlation.  
Pediatr., 45: 29-35.
  
- \* Wald, E.R.; Milmlone, G.J.; Bowen, A.; Salamon, N. and Bluestone, C.D. (1981):  
Acute maxillary sinusitis in children.  
N. Eng. J. Med., 304: 749.
  
- \* Wallace, R.J.; Steele, L.C.; Brooks, D.; Lumen, J.I. and Wilson, R.W. (1985):  
Amoxicillin-clavulanic acid in the treatment of lower respiratory tract infection caused by beta-lactamase - positive Haemophilus influenzae and Branhamella catarrhalis.  
Antimicrobial Agents and Chemotherapy, 27: 912-917.
  
- \* Wallace, R.J. and Musher, D.M. (1986):  
Branhamella catarrhalis in chronic lung diseases.  
Chest, 90 (3): 447-450.
  
- \* Wardle, J.K. (1986):  
Branhamella catarrhalis as an indirect pathogen.  
Drugs, 31 (Suppl. 3): 93-96.
  
- \* Warren, W.S. and Stool, S.E. (1971):  
Otitis media in low birth weight infants.  
J. Pediatr., 79: 740-743.
  
- \* West, M.; Berk, S.L. and Smith, J.K. (1982):  
Branhamella catarrhalis pneumonia.  
South. Med. J., 75: 1021-1023.

- \* **Wilson, G.S. and Miles, A. (1975):**  
Neisseria, Veillonella and Moraxella.  
In "Topley and Wilson's Principles of Bacteriology,  
Virology and Immunity".  
Arnold, E. Co., London, 6th. ed., Vol. I, p.684.
- \* **Wilson, G.S. and Wilkinson, A.E. (1983):**  
Neisseria; Branhamella and Moraxella.  
In "Topley and Wilson's Principles of Bacteriology,  
Virology and Immunity".  
Arnold, E. Co., London, 7th. ed., Vol. II, p.156.
- \* **Wistreich, G.A. and Barker, R.E. (1971):**  
The presence of fimbriae (pili) in three species  
of Neisseria.  
J. Gen. Microbiol., 65: 167.
- \* **Wong, D.T. and Ogra, P.L. (1980):**  
Immunology of tonsils and adenoids - an update.  
Int. J. Pediatr. Otorhinolaryngol., 2: 181-191.
- \* **Wright, J. (1970):**  
Bacteriology of ear, nose and throat diseases.  
J. Laryngol. Otol., 84: 383.
- \* **Wright, P.F.; McConnell, K.B.; Thompson, J.M. and  
Sell, S.H. (1985):**  
A longitudinal study of the detection of otitis  
media in the first two years of life.  
Int. J. Pediatr. Otorhinolaryngol., 10: 245-  
252.

- \* Yamaguchi, T.; Urasawa, T. and Kataura, . (1984):  
Secretory immunoglobulin and antibodies to respiratory viruses in middle ear effusion of chronic otitis media with effusion.  
Ann. Otol. Rhinol. Laryngol., 93: 73-75.
- \* Young, H.; Paterson, I.C.; McDonald, D.R. (1976):  
Rapid carbohydrate utilization test for the identification of Neisseria gonorrhoeae.  
British Journal of Venereal Diseases, 52: 3.
- \* Zoeller, C.; Andrieu, G.; Crosnier, R. and Passa, J. (1933):  
Un case de septicemia à Neisseria catarrhalis .  
Rhinopharyngite, nephrite, meningite.  
Bul. Soc. Med. Hop. Paris, 49: 1104-10.