

REFERENCES

- Abita, J., Cast, H., Milsten, S., Kufman, S., and Saimost, G. (1985):**
Urinary neopterin and biopterin levels in patients with AIDS-related complex.
Lancet, II: 51.
- Adcock, E.W., Teasdale, E., and August, C.B., (1973):**
Human chorionic gonadotrophin: its possible role in maternal lymphocyte suppression.
Science, 181: 845.
- Alanen, A., and Lassila, O. (1982):**
Deficient nature killer cell function in pre-eclampsia.
Obstet. Gynecol., 60: 631.
- Alanen, A., Kekomaki, R., Kero, P., et al. (1984):**
Circulating immune complexes in hypertensive disorders of pregnancy.
J. Reprod. Immunol., 6: 133.
- Albert, A. (1957):**
Transformation of purines into pyridines. Biochem. J., 65: 5310.
Quoted from Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G. and Werner, E. (1989): Neopterin as a marker for activation of cellular immunity: Immunological basis and clinical application.
Advances in clinical chemistry, 27: 82.
- Allebes, W., Knop, S. Herold, M., Huber, C. and Capel, P. (1991):**
Immune therapy with monoclonal anti-idiotypic antibodies: Tumour reduction and lymphokine production.
Leuk. Res.; 15(4) : 215.
- Altohek, A. (1961):**
Electron microscopy of renal biopsies in toxemia of pregnancy.
JAMA; 175:791.

American College of Obstetricians and Gynecologists (1986):

Management of pre-eclampsia.

Technical bulletin No. 91, February.

Amino, N., Tanizawa, O., and Mijai, K. (1978):

Changes of serum immunoglobulins, IgG, IgA, IgM and IgE, during pregnancy.

Obstet. Gynecol., 52: 415.

Anderson, J.M., (1972):

Nature's transplant: The transplantation of viviparity. New York: Appleton-Century-Crofts.

Armarego, F., and Randles, D. (1983):

Aerobic oxidation of 5,6,7,8 tetrahydroneopterin. In Biochemical and Clinical Aspects of Petridines. Curtius, H., Pfeleiderer, W., and Wachter H. (eds), pp. 423-427 - de Gruyter, Berlin, and New York.

Armstrong, N.P., Teisner, B., Redman, J.G., et al. (1986):

Complement Cascade in pre-eclampsia.

Br. J. Obstet. Gynecol., 93:811.

Arthur, J., and Ammann, M.D. (1987):

Immunodeficiency diseases. In: Basic and Clinical Immunology. Edited by Stites, D.P., Stobo, J.D., and Vivian Wells, J. 6th ed. By Appleton & Lange, Norwalk, Connecticut/Los Altos, California. P: 329.

Atrash, H.K., Koonin, L.M., Lawson, H.W., Franks, A.L., and Smith, J.C. (1990):

Maternal mortality in the United States, 1979-1986.

Obstet. Gynecol., 76: 1055-1060.

Baboonian, C., and Griffiths P. (1983):

Is pregnancy immunosuppressive? humoral immunity against viruses.

Br. J. Obstet. Gynecol., 90:1168.

Bach, F.H., and Hirschorn, K. (1964):

Mixed lymphocyte culture.

Science; 145:1315.

Bach, F.H., and Voynow, N.K. (1966):

One-way stimulation in mixed leucocyte cultures.

Science, 153:545 - 547.

Bach, F.H., and Amos, D.B. (1967):

Hu-1: major histocompatibility locus in man.

Science, 156: 1506-11508.

Bach, F.H., (1982):

Transplantation immunity and cytotoxicity phenomena. In: Bach, F.H., and Schwartz, R.S., (eds). Immunology, 2nd ed. New York, John Wiley & Sons, 399.

Bailey, K. and Ayling, J. (1975):

High pressure liquid chromatography of substituted petridines. In chemistry and Biology of Petridines. Pfeleider, W. (eds), pp. 633. de Gruyter, Berlin and New York. In Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G., and Werner, E. (1989): Neopterin as a marker for activation of cellular immunity: Immunologic basis and clinical application. Advances in Clin. Chem. 27: 82-139.

Bailey, K., Herrod, H.C., Younger, R., et al. (1985):

Functional aspects of T-lymphocyte subsets in pregnancy.

Obstet. Gynecol., 66:211.

Bain, B., Vas, M.R.; and Lowenstein, L. (1964):

Blood, 23:108. Quoted from Manual of Laboratory Immunology. Edited by Miller, L.E., Ludke, H.R., Peacock, J.E.; and Tomar, R.H. Second edition, pp 410-412, Lea and Febiger. Philadelphia, London, (1991).

Baker, D.A., Hameed, C., Telani, N., et al. (1987):

Lymphocyte subsets in the neonates of pre-eclamptic mothers.

Am. J. Reprod. Immunol. Microbiol., 14: 107.

Balasch, J., Mirapeix, E., Borche, L., et al. (1981):

Further evidence against pre-eclampsia as an immune complex disease.

Obstet. Gynecol., 58: 435.

Balow, J.E., and Rosenthal, A.S. (1973):

Glucocorticoid suppression of macrophage migration inhibitory factor.

J. Exp. Med., 137: 1031.

Barak, M., Merzbach, D., and Gruener, N. (1989 a):

Neopterin measured in serum and tissue culture supernatants by competitive enzyme-linked immunosorbent assay.

Clin. Chem. 35(7):1467.

Barak, M. Merzbach, D., and Gruener, N. (1989 b):

The effect of immunomodulators on PHA or gamma IFN induced release of neopterin from purified macrophages and peripheral blood mononuclear cells.

Immunology Letters, 21:317.

Barton, J.R., Sibai, B.M., Whybrew, W.D.; and Mercer, B.M. (1992):

Urinary endothelin-1: Not a useful marker for pre-eclampsia.

Am. J. Obstet. Gynecol., 168: 599-601.

Barton, J., and Sibai, B. (1991):

Care of the pregnancy complicated by HELLP syndrome.

Obstet. Gynecol. Clin. North. Am.; 18(2):165-179.

Beer, A.E. (1978):

Possible immunologic bases of pre-eclampsia/eclampsia.

Semin. Perinatol; 2:39.

Beer, A.E., Quebbeman, J.F., Ayero, J.V., and Haines, R. (1981):

Major histocompatibility complex antigens, maternal and paternal immune responses and chronic habitual abortions in human.

Am. J. Obstet. Gynecol. 141:981.

Beer, A.E., and Need, J.A. (1985):

Immunological aspects of pre-eclampsia/eclampsia.
Birth defects; 21: 131-154.

Beer, A.E., (1988):

Immunology of reproduction. In: Samter, M., Talmage, D.W.,
Frank, M.M., Austen, K.F., and Claman, H.N., (eds).
Immunological diseases. Boston: Little, Brown. 329.

Belizan, J.M., Villar, J., and Repke, J. (1988):

The relationship between calcium intake and pregnancy induced
hypertension: up-to-date evidence.
Am. J. Obstet. Gynecol.; 158: 898-902.

Bell, J.L., and Baron, D.N. (1968):

Protein precipitation using trichloroacetic acid, followed by Biuret
Reaction.
Proc. Assoc. Clin. Biochem; 5:63.

Benester, B., and Wood, E.J. (1970):

Immunoglobulin levels in normal pregnancy and pregnancy
complicated by hypertension.
J. Obstet. Gynecol. Br. Cwith, 77:518.

Bergholtz, B.O., and Thorsby, E. (1977):

Macrophage-dependent response of immune human T-
lymphocytes to PPD in vitro.
Scand. J. Immunol., 6:779.

Berzofsky, J.A., and Berkower, I.J. (1984):

In: Fundamental Immunology. Edited by Paul, W.E., New York,
Raven Press.

Beverley, P.C., and Gallord, R.E. (1981):

Distenctive functional characteristics of human T-lymphocytes
defined by E-rosetting or a monoclonal antibody.
Eur. J. Immunol.; 11:329.

- Bichler, A., Fuchs, D., Hausen, A., Hetzel, H., Koing, K. and Wachter, H. (1982):**
Urinary neopterin in patients with genital cancer.
Clin. Biochem., 15:38.
- Bichler, A., Fuchs, D., Hetzel, H., Reibnegger, G.; and Wachter, H. (1983):**
Measurement of urinary neopterin in normal pregnant and non pregnant women and in women with benign and malignant genital tract neoplasmas.
Arch. Gynecol., Vol.: 233(2), P: 121-130.
- Biggers, J.D., and Borland, R.M. (1976):**
Physiological aspects of growth and development of the preimplantation on mammalian embryo.
Ann. Rev. Physiol, 38:95.
- Birkeland, S.A., and Kristofferson, K. (1977):**
Cellular immunity in pregnancy; blast transformation and rosette formation of maternal T and B lymphocytes. Across-section analysis.
Clin. Exp. Immunol., 30:408.
- Birkeland, S.A., and Kristofferson, K. (1979):**
Pre-eclampsia a state of mother-fetus immune imbalance.
Lancet; ii:720.
- Birkeland, S.A., and Kristofferson, K. (1980):**
Lymphocyte transformation with mitogens and antigens during normal human pregnancy: a longitudinal study.
J. Immunol., 11: 321.
- Bitterlich, G., Szabo, G., Werner, E., Larcher, C., Fuchs, D., Hausen, A., Reibnegger, G., and et al. (1988):**
Selective induction of mononuclear phagocytes to produce neopterin by interferons (IFNs).
Immunobiology, 176: 228.

- Blau, N., Joller, P., Atares, M., Cardesa-Gracia, J., and Niederweiser, A. (1985):**
Increase of GTP cyclohydrolase 1 activity in mononuclear blood cells by stimulation of heterozygotes of GTP cyclohydrolase deficiency.
Clin. Chem. Acta; 148:47.
- Blau, N., Schoedon, G., and Curits, H. (1989):**
Biosynthesis and significance of neopterin in the immune system.
Eur. J. Cancer Clin. Oncol., 25(4):603.
- Bloom, J., Suffredini, A., Parrillo, J., and Palestine, A. (1990):**
Serum neopterin levels following I.V. endotoxin administration to normal humans.
Immunobiology, 181 (4-5):317.
- Blumer, R.; and Hancock, R., (1977):**
Depletion of circulating T-lymphocytes in pregnancy.
Clin. Exp. Immunol.; 28:302.
- Bocci, V., Paulesu, L., and Ricci, M.G. (1985):**
The physiological interferon response: IV production of interferon by the perfused human placenta at term.
Proc. Soc. Exp. Biol. Med.; 180: 137-143.
- Bolis, P.F., Bianchi, M.M., La Fianza, A., et al. (1987):**
Immunogenetic aspects of pre-eclampsia.
Biol. Res. Preg., 8:42.
- Bolton, A.E., and Hunter, W.M. (1986):**
In: Handbook of Experimental Immunology, Vol. 1 Edited by Weir, D.M. Oxford, Blackwell Scientific Publications.
- Bomford, R., Mason, S., and Swash, M. (1975):**
Hutchison's Clinical Methods. Bailliere Tindall, The English Language Book Society, England, p: 95.

Boyum A. (1968):

Separation of lymphocytes and erythrocytes by centrifugation.
Scand. J. Clin. Lab. Invest., 20, Suppl. 97.

Brabin, B.J. (1985):

Epidemiology of infection in pregnancy.
Rev. Infect. Dis., 7:579.

Bradbury, S., Billington, W.D., Kirby, D.R., and Williams, E.A. (1970):

Histochemical characterization of the surface mucoprotein of normal and abnormal human trophoblast.
Histochem. J., 2:263.

Bradley, B.A., and Festenstein, H. (1978):

Br. Med. Bull.; 34:223. Quoted from Manual of Laboratory Immunology. Edited by Miller, L.E., Ludke, H.R., Peacock, J.E.; and Tomar, R.H. Second edition, pp. 31-37. Lea and Feibger, Philadelphia, London, (1991).

Brinkman II, C.R. (1992):

Hypertensive disorders of pregnancy. In: Hacker, N., F., and Moore, J.G. (eds): Essential of Obstetrics and Gynecology, 2nd ed., p. 163-174. USA by W.B. Saunders Company.

Broadbent, B., Camargo, E., Jaffe, H., and Lederman, H. (1991):

Recombinant human interferon-gamma as adjunct therapy for aspergillus infection in a patient with chronic granulomatous disease.

J. Infect. Dis., 163 (4): 908.

Brumester, G., Hahn, G., Kersten, W., Platzner, E., and Kalden, J. (1988):

Activation of blood monocytes and synovial macrophages by recombinant interferon gamma and tumour necrosis factor - alpha in patients with reumatoid arthritis.
Immunobiology, 178: 90.

Bulmer, J.N., Morrison, L., Johnson, P.M., and Meager, A. (1990):

Immunohistochemical localization of interferons in human placental tissues in normal, ectopic and molar pregnancy.

Am. J. Reprod. Immunol., 22: 109-116.

Campbell, D., Mac Gillivray, I., and Carr-Hill, P., (1985):

Pre-eclampsia in second pregnancy.

Br. J. Obstet. Gynecol., 92:131.

Cappelli, N., Archidiacon, G., Santamouro, S., Gattini, M., et al. (1977):

Behaviour of serum immunoglobulins (IgG, IgA, IgM).

Minerva. Gynecol., 29:631.

Carr, M.C., and Stites, D.P. (1972):

Reactivity of maternal lymphocytes to phytohemagglutinin.

Lancet, i: 1073.

Carr, M.C., Stites, D.P., and Fudnberg, H.H. (1973):

Cellular immune aspects of the human fetal maternal relationship

II. In vitro response of gravida lymphocytes to phytohemagglutinin.

Cell. Immunol., 8:448.

Cauchi, N.M., Gilbert, C.L., and Brown, J.B. (1984):

The clinical pathology of pre-eclampsia. In the clinical pathology of pregnancy and of the newborn infant. pp. 351, ed. Publisher Edward Arnold.

Chakravorty, A.P. (1967):

Fetal and placental weight changes in normal pregnancy and pre-eclampsia.

J. Obstet. Gynecol. Br. Common; 74:247.

Chard, T., Craig, P.H., Menabawey, M., and Lee, C. (1986):

Alpha interferon in human pregnancy.

Br. J. Obstet. Gynecol., 93:1145.

Chen, S.L. (1990):

Cells and tissues of the immune system. In: Clinical Immunology, Principles and Laboratory diagnosis. Edited by Sheehan, C.J.B., Lippincott Company. Philadelphia. P.: 19-29.

Chesley, L.C., Annitto, J.E.; and Cosgrove, R.A. (1968):

The familial factor in toxæmia of pregnancy.
Obstet. Gynecol.; 32: 303.

Chesley, L.C., (1974):

A short history of eclampsia.
Obstet. Gynecol., 43:599.

Chesley, L.C., (1985):

Diagnosis of pre-eclampsia.
Obstet. Gynecol.; 65: 423.

Chesley, L.C., and Cooper, D. (1986):

Genetics of hypertension in pregnancy: Possible single gene control of pre-eclampsia and eclampsia in the descendants of eclamptic women.
Br. J. Obstet. Gynecol.; 93: 898-908.

Chisolm, J.; and Handorf, C. (1987):

Increased absorption of and sensitivity to Cadmium during late pregnancy: is there a relationship between markedly decreased maternal cadmium binding protein (metallothionein) and pregnancy induced hypertension.
Med. Hypotheses; 24: 346-351.

Clark, S. (1989):

Severe pre-eclampsia: the role of invasive hemodynamic monitoring.
Female Patient; 14(12): 52-56.

Clemens, L.E., Stiteri, R.K., and Stites, D.P. (1979):

Mechanism of immunosuppression of progesterone on maternal lymphocyte activation during pregnancy.
J. Immunol., 122: 1978.

- Clements, P.J., Yu, D.T., Levy, J.; and Pearson, C.M., (1976):**
Human lymphocyte subpopulations: the effect of pregnancy.
Proc. Soc. Exp. Biol. Med.; 152: 664.
- Clements, R., Peter, J., Agopian, M., Telian, N.; and Furst, D., (1990):**
Elevated serum levels of soluble IL-2 and neopterin in diffuse and limited scleroderma: Effect of chlorambucil.
J. Rheumatol.; 17(7): 103-112.
- Conrad, F., Fuchs, D., Hausen, A. et al., (1987):**
In Biochemical and Clinical Aspects of Pteridines :Pfleider, W., Wachter, H.; and Blair, J.A., (eds), pp. 233-241, de Gruyter.
- Contractor, S.F., and Davies, H. (1973):**
Effect of human chroionic somatotropin and HCG on phytohemagglutinin-induced lymphocyte transformation.
Nature, 243:284.
- Cooper, D.W., Deane, E.M., Marshall, P., and Gallory, E.D.(1988):**
C₃ allotypes in pregnancy hypertension and eclampsia.
Hum. Hered., 38:52.
- Cornfield, D.B., Jancks, J., Binder, R.A., and Rath, C.E. (1979):**
T and B lymphocytes in pregnant women.
Obstet. Gynecol., 53:203.
- Cotton, D.B., Lee, W., Hunta, J.C., and Dorman, K.E. (1988):**
Hemodynamic profile of sever pregnancy induced hypertension.
Am. J. Obstet. Gynecol.; 158:523-530.
- Coulam, C.B., Silverfield, J.C., Kazmar, R.E. et al. (1983):**
T-lymphocyte subsets during pregnancy and the menstrual cycle.
Am. J. Reprod. Immunol., 4:88.

Cunningham, F.G., MacDonald, P.C., Gant, N.F., and Gilstrap III, L.G. (1993):

Hypertensive disorders in pregnancy. In: Williams Obstetrics. 19th ed. p. 763-764. USA, by Appleton & Lange.

Currie, G.A., Van Doornick, W., and Bagshawe, K.D. (1968):

Effect of neuraminidase on the immunogenicity of early mouse trophoblast.

Nature (London), 219:191.

Dadak, C., Ulrich, W., and Sinzinger, H. (1984):

Morphologic changes in the umbilical arteries of babies born to pre-eclamptic mothers.

Placenta, 5:419.

D'Amelio, R., Bilotto, P., Pachi, A., and Aiuti, F. (1979):

Circulating immune complexes in normal pregnant women and in some conditions complicating pregnancy.

Clin. Exp. Immunol., 37:33.

Davey, D. (1985):

Hypertensive diseases in pregnancy. In Studd, J. (ed), Progress in Obstetrics and Gynecology. Vol.5. Churchill Livingstone. p. 89.

Davey, D.A., and Mac Gillivray, I. (1988):

The classification and definition of the hypertensive disorders of pregnancy.

Am. J. Obstet. Gynecol.; 158:892.

de Jong, C., Dekker, G., and Sibai, B. (1991):

The renin-angiotensin-aldosterone system in pre-eclampsia.

Clin. Perinatol.; 18(4): 683-711.

Dekker, G., and Sibai, B. (1991):

Early detection of pre-eclampsia.

Am. J. Obstet. Gynecol.; 165: 160-172.

Desbuquois, B., and Aurbach, G.D. (1971):

Radioimmunoassay Technique.

J. Clin. Endocrinol. Metab., 33:732.

De Voe, J.S.; and O'Shaughnessy, R., (1984):

Clinical manifestations and diagnosis of pregnancy-induced hypertension.

Clin. Obstet. Gynecol.; 27(4): 836-852.

Dewhurst, J. (1981):

Integrated obstetrics of gynecology for postgraduates. 3rd ed.

London. Blackwell Scientific Publications. p: 259.

Dick, H.M., and Kissmeyer - Nielsen, F.K. (1979):

Histocompatibility techniques. Amesterdam, Elsevier, North Holland. Biomedical Press.

Dionigi, R., Zonta, A., Albertario, F., et al. (1973):

Cyclic variation in the response of lymphocytes to phytohemagglutinin in healthy individuals.

Transplantation, 16: 550.

Dodson, M.G., Kerman, R.H., Lange, C.F., Sefani, S.S.; and O'Leary, J.A. (1977):

T and B cells in pregnancy.

Obstet. Gynecol.; 49:299.

Dubey, D.P., Yunis, I., and Yunis, E.J. (1986):

In Manual of Clinical Laboratory Immunology, 3rd ed. Edited by Rose, N.R., Friedman, H., and Fahey J.I. Washington, D.C., American Society for Microbiology.

Dunlop, W., and Davison, J.M. (1977):

The effect of normal pregnancy upon renal handling of uric acid.

Br. J. Obstet. Gynecol.; 84:13.

Easterling, T., and Benedetti, T. (1989):

Pre-eclampsia: a hyperdynamic disease model.

Am. J. Obstet. Gynecol.; 165:160-172.

El Roeity, A., and Gleicher, N. (1988):

Immunologic concepts of pre-eclampsia. In: Rubin, P.C. (ed.).
Handbook of hypertension. Vol. 10. Amsterdam, El Sevier
Science, p: 257.

**Farrant, I., Clark, J.C., Lee, H., Knight, S.C., and O'Brien, J.
(1980):**

Conditions for measuring DNA synthesis in PHA stimulated
human lymphocytes in 20 µl hanging drops with various cell
concentrations and periods of culture.
J. Immunol. Methods, 33:301-312.

Faulk, W.P., Demalyer, E.M., and Davis, A.J. (1974):

Some effect of malnutrition on the immune response in man.
Am. J. Clin. Nutr.; 27:638.

Feeny, J.G., Tovey, L.A. and Scott, J.S. (1977):

Influence of previous blood transfusion on the incidence of
pre-eclampsia.
Lancet, I: 874.

Feeny, J.G. (1980):

Changed paternity. In: Bonnar, J., MacGillivray, I., Symonds, E.M.
(eds): Pregnancy hypertension. Baltimore: University Park Press,
p:41.

Feinberg, B.B., and Gonik, B. (1991):

General precepts of the immunology of pregnancy.
Clin. Obstet. Gynecol., 34:3-16.

**Felmayer, W., Werner, E., Fuchs, D., Hausen, A., Reibnegger,
G., and Wachter, H. (1989):**

Characteristics of IFN-induced tryptophan metabolism in human
cells in vitro.
Biochemic. Biophysica. Acta, 1012: 140-147.

Finn, R., Hill, C.A., and Goven, A.J. (1972):

Immunological responses in pregnancy and survival of fetal homograph.

Br. Med. J., 3:150.

Foad, B.S.I. et al. (1974):

Clin. Exp. Immunol.; 17:657. Quoted from Manual of Laboratory Immunology. Edited by Miller, L.E., Ludke, H.R., Peacock, J.E.; and Tomar, R.H. Second edition, pp.88-90. Lea and Feiber-Philadelphia - London, (1991).

Freedolph, D.A., Ushyima, R.N., and Larson, C.L. (1974):

Recurrent herpes genitalis, treatment with BCG.

Obstet. Gynecol., 26:521.

Friedman, S.A., and Neff, R.K., (1977):

Pregnancy hypertension, PSG Publishing Co., Inc., London, Mass.

Friedman, S.A. (1988):

Pre-eclampsia: a review of the role of prostaglandins.

Obstet. Gynecol.; 71:122.

Friedman, S.A., Taylor, R.N., and Roberts, J.M. (1991):

Pathophysiology of pre-eclampsia.

Clin. Perinatol.; 18: 661-82.

**Fuchs, D., Hausen, A., Huber, C., Margreiter, R., Reibnegger, G.,
Spielberger, M., and Wachter, H. (1982):**

Pteridin ausscheidung als marker Hoppeseyler's Z. Physiol. Chem. 363:664. Quoted from Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G., and Werner, E. (1989): Neopterin as a marker for activation of cellular immunity. Immunological basis and clinical application. Advances in Clin. Chem. 27: 82-139.

Fuchs, D., Hausen, A., Knosp, O. et al. (1983):

Neopterin evaluation in patients with pulmonary tuberculosis. In Curtius, H.C., Pfeleiderer, W., Wachter, H. (eds.): Biochemical and Clinical Aspects of Pteridines, Vol.2, Berlin, de Gruyter, 1983, p. 281.

Fuchs, D., Hausen, A., Reibnegger, G., Werner, E., and Wachter, H. (1988):

Neopterin in clinical medicine.
Lancet, i:702.

Fuchs, D., Forsman, A., Hagbera, L., Larsson, M., Norkrans, G., Reibnegger, G., Werner, E., and Wachter, H. (1990):

Immune activation and decreased tryptophan in patients with HIV-1 infection.
J. IFN. Res., 10 (6): 599-603.

Fudenberg, H.H., Good, R.A., Goodman, H.C., Hitzig, W. et al. (1971):

Primary immunodeficiencies: Report of a W.H.O. Committee.
Pediatrics, 47:927.

Fukshima, T., and Nixon, J. (1979):

Oxidation and conversion of reduced forms of biopterin. In "Chemistry and Biology of Pteridines". Kislink and Brown (eds.), pp. 31-34, El-Sevier, North Holand, Amesterdam. In Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G., and Werner, E. (1989): Neopterin as a marker for activation of ceullar immunity. Immunological basis and clinical application. Advances in Clinical Chemistry., 27: 82-139.

Gall, S.A. (1983):

Matenral adjustments in the immune system in normal pregnancy.
Obstet. Gynecol. 26:521.

Gang, N.F., Schwartz, E.S., and Majerovics, A. (1974):

Studies on the placenta II nephrotoxicity of antibodies produced against a chorionic glycoprotein.
Am. J. Obstet. Gynecol.; 120: 73.

- Ganter, C., Grassmayer, K., Hausen, A., and Wachter, H. (1977):**
Ausscheidung einer fluoreszierenden-substanz in Harn Ehrlich
Ascites-tumour tragender Mause. *Mikrochim. Acta*, 2: 33-38.
Quoted from Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G.,
and Werner, E. (1989): Neopterin and activation of cellular
immunity. *Advances in Clinical Chemistry*, 27: 82-125.
- Gauges, J.H., Jones, E., and Curzen, P. (1975):**
Spontaneous lymphocyte transformation in pregnancies
complicated by pre-eclampsia.
Am. J. Obstet. Gynecol., 121:542.
- Gehrz, R.C., Christianson, W.R., Linner, K.M., et al. (1981):**
A longitudinal analysis of lymphocyte proliferative responses to
mitogens and antigens during human pregnancy.
Am. J. Obstet. Gynecol., 140:665.
- Gerrad, T., Dyer, D., Enterline, J.; and Zoon, K. (1989):**
Products of stimulated monocytes enhance the activity of
IFN-gamma.
J. IFN-Res.; 9:115-124.
- Gilbert, E.S., and Harmon, J.S. (1993):**
Hypertensive disorders. In: *High risk pregnancy delivery*. in
Gilbert, E.S., and Harmon J.S. (eds). p.: 375-386, USA, by
Mosby-Year book, Inc.
- Gill, T.J., and Repetti, C.F. (1979):**
Immunological and genetic factors influencing reproduction.
Am. J. Pathol. 95: 463.
- Gitlin-David (1975):**
Normal biology of AFP.
Ann. N.Y Acad. Sci. 259:7.

Glassman, A.B., Bennett, C.E., Christopher, J.B. and Self, S. (1985):

Immunity during pregnancy: Lymphocyte sub-populations and mitogen responsiveness.

Ann. Clin. Lab. Sci., 15:357.

Gleichner, N., and Theofilopoulos, A.N. (1979):

Immune complexes in pregnancy.

Lancet, 27:216.

Glistrip, L., and Gant, N. (1990):

Pathophysiology of pre-eclampsia.

Semin. perinatol.; 4 (2): 147-151.

Gonik, B., Loo, L.S., West, S., and Kohl, S. (1987):

Natural killer cytotoxicity to herpes simplex virus - infected cells in human pregnancy.

Am. J. Reprod. Immunol. Microbiol., 13:23.

Goodlin, R.C., Cotton, D.B.; and Haesslein, H.C. (1979):

Sever edema-proteinuria hypertension gestosis.

Am. J. Obstet. Gynecol., 135:76.

Graybill, J.R., and Alford, R.H. (1976):

Variability of sequential studies of lymphocyte blastogenesis in normal adults.

Clin. Exp. Immunol., 25:28.

Greaves, M.F., Owen, J.J., and Raff, M.C. (1973):

T and B lymphocytes, origins, properties and roles in immune response. American Elsevier Publishing Company, Inc., New York.

Gregory, C.D., Shah, L.P., Lee, H., et al. (1985):

Cytotoxic reactivity of human natural killer cells during normal pregnancy a longitudinal study.

J. Clin. Lab. Immunol., 18:175.

Griffin, J.F., and Wilson, E.W. (1978):

Abnormal immune reactivity in pre-eclamptic patients.
Proceedings of the university of Otago Medical School. Vol. 65,
pp.9.

Griffith, P.D., Buie, K.J., and Heath, R.B., (1980):

Persistence of high titre antibodies to the early antigens of
cytomegalovirus in pregnant women.
Arch. Virol. 64: 303.

Gudson, J.P. (1969):

Fetal and maternal immunoglobulin levels during pregnancy.
Am. J. Obstet. Gynecol., 103:895.

Gudson, J.P. (1976):

Maternal immune response in pregnancy. In: Scott, J.S., and
Jones, W.R. (eds): Immunology of human reproduction.
Academic Press, London, 118.

Gudson, J.P., Heise, E.R., and Herbst, G.A. (1977):

Studies of lymphocyte populations in pre-eclampsia.
Am. J. Obstet. Gynecol.; 129:255.

Guo, F., and Qian, S.Z. (1989):

Preliminary study on the immune function of red cells in patients
with pregnancy-induced hypertension.
Chung-Hau-Fu. Chan-Ko-Tsa-Chin, Jul, Vol. 24 (4), P: 215-6,
252, ISSN: 0529-567X (abstract).

Guyton, A. (1991):

Textbook of medical physiology, 8th ed., Philadelphia, W.B.
Saunders.

Haeger, M., Bengtsson, A., Karlsson, K., et al. (1989):

Complement activation and anaphylatoxin (C3a and C5a)
formation in pre-eclampsia and by amniotic fluid.
Obstet. Gynecol., 73:551.

- Haeger, M., Unander, M., and Bengtsson, A. (1990):**
Enhanced anaphylatoxin and terminal C5b-9 complement complex formation in patients with the syndrome of hemolysis, elevated liver enzymes and low platelet count.
Obstet. Gynecol., 76:698.
- Haeger, M., Unander, M., Hansson, B.N., et al. (1992):**
Complement, neutrophils and macrophage activation in women with severe pre-eclampsia and HELLP Syndrome.
Obstet. Gynecol., 79:19.
- Hagihara, M., Nagatsu, T., Ohhashi, M.; and Muira, T. (1990):**
Concentrations of neopterin and biopterin in sera from patients with rheumatoid arthritis or systemic lupus erythematosus and in synovial fluid from patients with RA or OA.
Clin. Chem.; 36 (4): 705-708.
- Hansch, G.M., Seitz, M., Martinotti, G., Betz, M., Rauterberg, E., Gemsa, D. (1984):**
Macrophage release arachidonic acid, prostaglandin E2, and thromboxane in response to late complement components.
J. Immunol.; 133: 2145-2150.
- Hansen, J.A., Douglas, M., and Edmond, J. (1986):**
In Manual of Clinical Laboratory Immunology. 3rd ed. Edited by N.R. Rose, H. Friedman and J.L. Fahey. Washington D.C., American Society for Microbiology.
- Harmon, W., Parkman, T., Gavin, E., Grupe, R., Yunis, E., and Levey, R.H. (1982):**
Comparison of cell mediated lympholysis and mixed lymphocyte culture in the immunologic evaluation for renal transplantation.
J. Immunol. 129: 1573-1577.
- Harrison, M.R. (1972):**
Immunocompetence of maternal lymphocytes.
Lancet, 2: 1319.

- Hartzman, R.J., Segall, M. Bach, M.L., and Bach, F.H. (1971):**
Histocompatibility matching. VI. Miniaturizations of the mixed leucocyte culture test: a preliminary report.
Transplantation, 11: 268-273.
- Hausen, A., Fuchs, D., Grunewald, K., Huber, H., Konig, K., and Wachter, H. (1981):**
Urinary neopterin as a marker for hematological neoplasia.
Clin. Chem. Acta., 117: 297-305.
- Hausen, A, Fuchs, D., Konig, K., and Wachter, H. (1982 a):**
Determination of neopterin in human urine by reversed-phase high performance liquid chromatography.
J. Chromatography Biomedical Application, 227:61-70.
- Hausen, A., Fuchs, D., Reibnegger, G., Ernst, Werner, R. and Wachter, H. (1989):**
Neopterin in clinical use.
Pteridines, Vol.1, pp. 3-10.
- Hausen, G.J., Lidor, A., Zakuth, V. et al. (1987):**
Immunocompetence in pregnancy: production of interleukin-2 by peripheral blood lymphocytes.
Cancer. Detect. Prev. Suppl. 1:39.
- Hawkins, T., Jones, M.P.; and Gallery, E. (1993):**
Secretions of prostanoids by platelets and monocytes in normal and hypertensive pregnancies.
Am. J. Obstet. Gynecol., 168: 661-667.
- Herberman, R.B., and Holden, H. (1978):**
Natural cell mediated immunity.
Adv. Cancer Res.; 27: 305.
- Hibbard, B.M. (1962):**
Hydramnios and pre-eclamptic toxemia.
Clin. Obstet. Gynecol., 5:1044.

Hill, A.B. (1977):

Textbook of Medical statistics: 10th ed. HOlder and Stroughten,
London.

Hill, J.A., Hsia, S., Doran, M., et al. (1986):

Natural killer cell activity and antibody dependent cell-mediated
cytotoxicity in pre-eclampsia.
J. Reprod. Immunol., 205:212.

Hirschberg, H., Kaakinen, H., and Thorsby, E. (1976):

Typing for HLA-D determinants. Comparison of typing using
homozygous stimulating cells and primed cultures.
Tissue Antigens, 16: 335-367.

Hoff, C., Stevens, R.G., Mendenhall, H., and et al. (1990):

Association between risk of pre-eclampsia and HLA-DR.
Lancet. 335:660.

Hoff, C., Peevy, K., Giattina, B.S.N., et al. (1992):

Maternal-fetal HLA-DR relationship and pregnancy-induced
hypertension.
Obstet. Gynecol., 80:1007.

Howvert, M.H., Class, F.H., Gmeling, F.H. et al. (1985):

Humoral immunity in normal and complicated pregnancy.
Eur. J. Obstet. Gynecol. Reprod. Biol., 19 (4), 205.

Hubel, C.A., Roberts, J.M., Taylor, R. (1989):

Lipid peroxidation in pregnancy: new perspective on
pre-eclampsia.
Am. J. Obstet. Gynecol.; 161: 1025-1034.

Huber, C.H., Fuchs, D., Hausen, A. et al. (1983):

In Spitzzy, K., Karrer, K. (eds): Proceedings of the 13th
International Congress of Chemotherapy, Vol. 18. Vienna, pp.
267.

- Huber, C., Fuchs, D., Hausen, A., Margretier, R., Reibnegger, G., Spielberger, M., and Wachter, H. (1983 a):**
Petridine as a new biochemical marker to detect human T-cells activated by allogeneic or modified self major histocompatibility complex (MHC) determinants.
J. Immunol., 130: 1047.
- Huber, C., Niederwieser, D., Schonitzer, D., Fuchus, D, Hausen, A., Kosp, O., and Reibnegger, C. (1983 b):**
Petridines in the assessment of lymphocyte activation.
Biochemical and Clinical Aspects of Petridines, (2) Walter de Gruyter and Co., Berlin. New York pp: 177-183.
- Huber, C., Batchelor, J., Fuchs, D., Hausen, A., Lange, A., Niederwieser, D., Reibnegger, G., Swetly, P., Troppmair, J., and Wachter, H. (1984):**
Immune response associated production of neopterin. Release from macrophages primarily under control of IFN gamma.
J. Exp. Med., 160: 310-316.
- Huber, C., Troppmair, J, Tilg, H, Woloszczuk, W., et al. (1985):**
Assessment of endogenous interferon release by evaluation of neopterin levels: A new concept for monitoring of human allograft recipients.
Transplantation proceedings, Vol. XVII, No. 6: 2513-2515.
- Hughes, E.C. (1972):**
Obstetric-Gynecologic Terminology. In: Hughes, E.C.(ed.). Philadelphia. Davis.
- Hulka, J.F., and Mohr,K. (1967):**
Interference of cortisone - induced homograft survival by progestins.
Am. J. Obstet. Gynecol., 97: 407.
- Hyland, K. (1985):**
Estimation of tetrahydro-dihydro-and fully oxidised pterins by high performance liquid chromatography using electrochemical and fluorometric detection.
J. Chromatography, 343: 35-41.

- Irino, T., Okuda, T., and Grollman, A. (1967):**
Changes induced in the glomeruli of the kidney of rats by placental extracts as observed with the electron microscope.
Am. J. Pathol., 50: 421.
- Jenkins, D.M., Acres, M.G., and Peters, J. (1972):**
Human chronic gonadotrophin and the fetal allograft.
Am. J. Obstet. Gynecol., 114:13.
- Jenkins, D.M., Need, J.A., and Rapah, S.M. (1977):**
Deficiency of specific HLA antibodies in sever pregnancy pre-eclampsia.
Clin. Exp. Immunol., 27: 485.
- Jenkins, D.M., Need, J.A., Scott, J.S., et al. (1978):**
Human leucocyte antigens and mixed lymphocyte reaction in sever pre-eclampsia.
Br. Med. J.; I: 542.
- Jenkins, D.M. (1990):**
Immunology of gestosis. Proceedings on the 22nd international congress of pathophysiology of pregnancy. Budapest. p: 30.
- Johnson, N., Moodley, J., and Hammond, M.G. (1988):**
Human leucocyte antigen status in African women with eclampsia.
Br. J. Obstet. Gynecol., 95: 877.
- Johnson, U., and Gustavii, B. (1987):**
Complement components in normal pregnancy.
Acta. Pathol. Microbiol. Immunol. Scand. [C] 95:97.
- Jones, T., and Brown, G. (1967):**
The biosynthesis of pteridines from guanosine triphosphate.
J. Biochem.; 242: 3989-3997.
- Kalmus, H. (1946):**
Genetical antigenic compatibility as a possible cause of the toxemias occurring late in pregnancy.
Ann. Eugen., 13: 127.

Kandile, D., El Nahas, M., El Shika, Z., et al. (1980):

Plasma protein pattern and immunoglobulins in normal pregnancy and pre-eclamptic toxemia.

J. Egypt. Soc. Obst. Gyn., 45:13.

Kaufman, S.S., Berlow, G.K., Summer, S., Schulman, S., Orloff, S., et al., (1978):

Hyperphenylalaninemia due to a deficiency of biopterin. A variant form of phenylketonuria.

N. Engl. J. Med. 299:673.

Kaufman, S.S. (1981):

Regulatory properties of pterin-dependent hydroxylases; variation on a theme. In Usdin, E., Weiner, N., Youdium, M. (eds). Function and regulation of monoamine enzymes.

New York: Macmillan, pp. 165-173. Quoted from the Lancet,ii Neopterin in clinical medicine (1988).

Kay, M.D., Bonnard, G.D., West, W.H., and Herberman, R.B. (1977):

A functional comparison of human Fc-receptor-bearing lymphocytes active in natural cytotoxicity and antibody dependent cellular cytotoxicity.

J. Immunol. 118: 2058.

Kaye, D.M., and Jones, W.R. (1971):

Effect of HCG on in vitro lymphocyte transformation.

Am. J. Obstet. Gynecol., 109: 1296.

Kern, P., Kerbs, H.; and Dietrich, M. (1984 b):

Serum neopterin: Screening test to exclude transfusion hazards by blood donors with cytomegalovirus infection, AIDS. pp. 10-13, eighteen Congresses Intern. Society of Blood Transfusion.

Kern, P., and Dietrich, M. (1985):

Increase in serum neopterin levels in patients receiving recombinant IL-2 and recombinant IFN-alpha. In Biochemical and Clinical Aspect of Pteridines. Curtius, H.: Pfeleiderer, W. and Wachter, H. (eds). pp. 335. Walter de Gruyter and Co., Berlin, New York.

Kilpatrick, D.C., Liston, W.A., Jazwinska, E.G., and Smart, G.E. (1987):

Histocompatibility studies in pre-eclampsia.
Tissue Antigens, 29:232.

Kilpatrick, D.C., Liston, W.A., Gibson, F., and Livingstone, J. (1989):

Association between susceptibility to pre-eclampsia within families and HLA-DR4.
Lancet, 2: 1063.

Kilpatrick, D.C., Gibson, F., Livingston, J., and Liston, W.A. (1991):

Pre-eclampsia is associated with HLA-DR4 sharing between mother and fetus.
Tissue antigens. 35:178.

Kirby, D.R.S., Billington, W.D., Bradbury, S., and Goldstein, D.J. (1964):

Antigen barrier of the mouse placenta.
Nature, 204: 548.

Kirkwood, E.M. (1975):

Is pre-eclampsia an immune complex disease?
Lancet. 2: 1100.

Kitzmler, J.L., and Benirschke, K. (1973):

Immunofluorescent study of placental bed vessels in pre-eclampsia of pregnancy.
Am. J. Obstet. Gynecol.; 115: 248.

Kitzmler, J.L., Stoneburger, L., Yelonosky, P.F., and Lucas, W.E. (1973):

Serum complement in normal pregnancy and pre-eclampsia.
Am. J. Obstet. Gynecol., 117:312.

Klapper, D.G., and Mendenhall, W.H. (1971):

Immunoglobulin D concentration in pregnant women.
J. Immunol., 107: 912.

Klonoff-Cohen, H., Savitz, D., Cefalo, R., and McCann, M. (1989):

An epidemiology study of contraception and pre-eclampsia.
JAMA; 262: 3143-3147.

Knox, G.E., Stagno, S., Volanakis, J.E., and Huddleston, J.F. (1978):

A search for antigen-antibody complexes in pre-eclampsia.
Further evidence against immunologic pathogenesis.
Am. J. Obstet. Gynecol., 132:87.

Kong, T.Y., DeWolf, F., Robertson, W.B., and et al., (1986):

Inadequate maternal vascular response to placentation in pregnancies complicated by pre-eclampsia and by small-for-gestational age infants.
Br. J. Obstet. Gynecol., 93:1049.

Koslowski, J.P., Guiget, M., Taquoi, G., et al. (1978):

Complement (C3 and CH50) and immunoglobulins in toxemia of pregnancy.
J. Gynecol. Obstet. Biol. Reprod., 7:923.

Kovar, I.Z., and Riches, P.G. (1988):

C3 and C4 complement components and acute phase proteins in late pregnancy and parturition.
J. Clin. Pathol., 41:650.

Kovats, S., Main, E.K., Librach, C., et al. (1990):

A class I antigen, HLA-G expressed in human trophoblasts.
Science, 248: 220.

Kozuharova, M., Bogdanov, N., Mantov, S., et al. (1985):

Immunological studies in pregnant women with EPH gestosis. In: Abstracts of 11 World Congress of Gynecology and Obstetrics. Berlin (West), 237 (Suppl) PP. 42.

Krause, P.J., Ingardia, C.J., Pontius, L.T., et al. (1987):

Host defense during pregnancy: neutrophil chemotaxis and adherence.
Am. J. Obstet. Gynecol., 157:274.

Kronenberg, M., Siu, S., Hood, E., and Shastri, N. (1986):

The molecular genetics of the T-cell antigen receptor and T-cell antigen recognition.

Ann. Rev. Immunol., 4: 529.

Kwietny, H., and Bergmann, F. (1959):

Separation and identification of pteridines by paper chromatography. J. Chromatography, 2: 162-172. Quoted from Wachter, H., Hausen, A., Reibnegger, G., and Werner, E. eds. (1989): Neopterin as a marker for activation of cellular immunity: Immunological basis and clinical application.

Advances in Clinical Chemistry, 27: 82-139.

Larsen, J., and ODell, W.D. (1986):

In: Manual of Clinical Laboratory. Immunology, 3rd ed. Edited by Rose, N.R., Friedman, and Fahey, J.L. Washington, D.C., American Society for Microbiology.

Lazebnik, N., Kuhnert, B., and Kuhnert, P. (1989):

Zinc, cadmium, and hypertension in parturient women.

Am. J. Obstet. Gynecol. ; 161: 437-400.

Lellé, R.J., Henkel, E., Leinemann, D., and Goeschen, K. (1989):

Measurement of CEA, TPA, Neopterin, CA125, CA153, and C199, in sera of pregnant women, umbilical cord blood and amniotic fluid.

Gynecol. Obstet. Invest., 27: 137-142.

Leohirun, L., Thunasethakul, P., Sumethkul, V., Pholcharoen, T.; and Boonpucknavig, V. (1991):

Urinary neopterin in patients with systemic lupus erythematosus. Clin. Chem.; 37(1): 47-50.

Ling, N.R., and Kay, J.E. (1975):

Lymphocyte stimulation. North Holland Publishing Co., Oxford.

Mabie, W., Ratts, T., and Sibai, B. (1989):

The central hemodynamics of severe pre-eclampsia.

Am. J. Obstet. Gynecol.; 161: 1443-1448.

Mac Gillivray, I. (1958):

Some observations on the incidence of pre-eclampsia.
J. Obstet. Gynecol. Br. Empire, 65:536.

Maltzer, M.C., and Silva, J. (1980):

In vitro defects of phagocyte chemotaxis during pregnancy.
J. Clin. Microbiol., 11: 170.

Maluish, E.; and Strong, D.M. (1986):

In: Mannual of Clinical Laboratory Immunology, 3rd ed. Edited by
Rose, N.R., Friedman, H., and Fahey, J.L. Washington, D.C.,
American Society for Microbiology.

Mansour, M. (1990):

Value of neopterin estimation as a marker of rheumatoid disease
activity.
Thesis submitted to faculty of medicine. Ain Shams University.

**Margreiter, R., Fuchs, D., Hausen, A., Reibnegger, G., et al.
(1983):**

Neopterin as a new biochemical marker for diagnosis of allograft
rejection.
Transplantation;36: 650-653.

Maroulis, G.B., Buckley, R.H., and Yunger, J.B., (1971):

Serum immunoglobulin concentrations during normal
pregnancy.
Am. J. Obstet. Gynecol., 109:917.

Marti, J., Harrman, H. (1977):

Immunogenesis: a new etiologic concept of essential EPH
gestosis, with special consideration of the primigravid patients.
Am. J. Obstet. Gynecol.; 128: 489.

**Martin, J. Blake, P.G., Kenneth, G., Blake, M., Martin, W., Perry,
J., and Mac Caul, J. (1991):**

The natural history of HELLP syndrome: Patterns of disease
progression and regression.
Am. J. Obstet. Gynecol.; 164: 1500-1513.

- Massayeff, R., Gilli, F., and Krebs, B. (1975):**
Evolution of AFP serum levels throughout life in humans and rats and during pregnancy in the rat.
Ann. N Y. Acad. Sci. 259:17.
- Massobrio, M., Benedetto, C., Bertini, E., Tetta, V., and Gaunussi, G. (1985):**
Immune complexes in pre-eclampsia and normal pregnancy.
Am. J. Obstet. Gynecol., 152: 578.
- Masson, P.L., Delire, M., and Cambiaro, C.L. (1977):**
Circulating immune complexes in normal human pregnancy.
Nature, 266: 542.
- Mc Cartney, C.P. (1969):**
The acute hypertensive disorders of pregnancy, classified by renal histology.
Gynecologia; 167: 214-220.
- McLaughlin, P.J., Stirrat, J.M., Redman C.W., and Levinsky, R. (1979):**
Immune complexes in normal and pre-eclamptic pregnancy.
Lancet 1:934.
- Medcalf, R.L., Kuhn, R.J., Iwanov, J.D., and Moulds, R.F. (1983):**
Vaso-activity generated from platelets by immune complexes in pre-eclampsia.
Clin. Exp. Pharmacol. Physiol., 10 (3) : 369.
- Miles, C.E.M., and Hales, C.N. (1968):**
Solid phase RIA or immunoradiometric assay (IRMA).
Nature, 219: 186.
- Miller, L.E.; Ludke, H.R., Peacock, J.E., and Tomar, R.H. (1991):**
In: Manual of Laboratory Immunology. Second edition, pp. 31-37.
Lea and Febiger, Philadelphia. London.
- Moore, M.P., Carter, N.P., and Redman, C.W. (1983):**
Lymphocyte subsets in normal and pre-eclamptic pregnancies.
Br. J. Obstet. Gynecol., 90 (4): 326.

Morse, J.H., Stearns, G., and Arden, J. (1976):

The effect of crude and purified human gonadotrophin on in vitro stimulated human lymphocyte cultures.

Cell Immunol., 25: 178.

Mould, R.F. (1976):

Introductory Medical Statistics. 1st ed. Pitman Publishing Ltd., London.

Murgita, R.A., and Tomasi, T.B. (1975a):

Suppression of immune response by alpha fetoprotein I. Primary and secondary response.

J. Exp. Med., 141:L 269.

Murgita, R.A., and Tomasi, T.B. (1975 b):

Suppression of the immune response by alpha fetoprotein II. MLR and mitogen induced lymphocyte transformation.

J. Exp. Med. 141:440.

Nathan, C., Murray, H., Wiebe, M., and Rubin, B. (1983):

Identification of interferon (IFN)-gamma as the lymphokine that activates human macrophage oxidative metabolism and antimicrobial activity.

J. Exp. Med.; 158: 670-689.

Nathan, C. (1986):

Peroxide and petridine: A hypothesis of the regulation of macrophage antimicrobial activity by IFN-gamma. In "Interferon"

Gresser, I., and Vilcek, J. (eds). New York, Academic Press, PP. 125-143.

Need, J.A., Jenkins, D.M., and Scott, J.S., (1976):

The response of lymphocytes to phytohemagglutinin in women with pre-eclampsia.

Br. J. Obstet. Gynecol.; 83:438.

Need, J.A. (1979):

Immunological phenomena in pre-eclamptic toxemia.

Clin. Obstet. Gynecol., 6:443.

Need, J.A., Scott, L.S., and Jenkins, D.M. (1980):

Serum factors in severe pre-eclamptic toxemia. In: Bonnar, J., Mac Gillivray, I., and Symmonds, E.M. (eds). Pregnancy hypertension. Proceedings of the first congress of the International Society for the study of hypertension in pregnancy, MTP press limited. International Medical Publishers PP. 299.

Need, J.A. Bell, B., and Meffin, E. (1983):

Pre-eclampsia in pregnancies from donor insemination.
J. Reprod. Immunol.; 5: 329-338.

Nelson, J.H., and Hall, J.E. (1964):

Studies on the lymphatic system in humans I. Morphological changes in lymph nodes in pregnancy at term.
Am. J. Obstet. Gynecol., 90:482.

Nesslehut, H., Hinney, B., Rath, W., Batwanz, A., Cillien, N.; and Kuhn, W. (1992):

Changes in the basic immunologic parameters interleukin-2 receptor and CD8-antigen in pre-eclampsia.
Z. Geburtshilfe Perinatol; 196 (2); 63-6 (Abstract).

Newby, C.J., Hayakawa, K., and Herzenberg, L.A. (1986):

In: Handbook of Experimental Immunology I. Edited by Weir, D.M. Oxford, Blackwell Scientific.

Newman, V., and Fullerton, J. (1990):

Role of nutrition in the prevention of pre-eclampsia: Review of the literature.
J. Nurse Midwifery, 35 (5): 282-291.

Niederwieser, D., Huber, C., Gratwohl, A., Bannert, P. et al. (1984):

Neopterin as a new biochemical marker in the clinical monitoring of bone marrow transplant recipients.
Transplantation; 38: 497-500.

**Nisell, H., Carlstrom, K., Cizinsky, S., Grunewald, C., Nylund, L.,
Randamaa, I. (1992):**

Atrial natriuretic peptide concentrations and hemodynamic effects of acute plasma volume expansion in normal pregnancy and pre-eclampsia.

Obstet. Gynecol., 79: 902-907.

Nova, A., Sibai, B.M., Barton, J.R., Mercer, B.M.; and Mitchell, M.D. (1991):

Maternal plasma level of endothelin is increased in pre-eclampsia. Am. J. Obstet. Gynecol.; 165: 724-7.

Nowell, P.L. (1960):

Phytohemagglutinin: an indicator of mitosis in cultures of normal human leucocytes.

Cancer Res. 20: 452-466.

O'Brein, W.F. (1992):

The prediction of pre-eclampsia.

Clin. Obstet. Gynecol. Vol. 35, No.2, June.

Okamura, K., Furukawa, K., Nakakuki, M., et al. (1984):

Natural killer cell activity during pregnancy.

Am. J. Obstet. Gynecol., 149:396.

Oppenheim, J.J., Dougherty, S., Chen, C., and Baker, J. (1975):

Utilization of lymphocyte transformation to assess clinical disorders. In Vyas, G. (ed). Laboratory diagnosis of immunological disorders. Green and Stretton, New York. P. 87-109.

Oppenheim, J.J., and Rosenstreich, D.L. (1976):

Lymphocyte transformation: Utilization of automatic harvesters. P. 573-577. In Bloom, B., and David, J. (ed.), In vitro methods in cell mediated and tumour immunity. Academic Press, Inc., New York.

Oppenheim, J.T., and Schecter, B. (1980):

In Manual of Clinical Laboratory Immunology. 2nd ed. Edited by Rose, N.R., and Friedman, H. Washington, D.D., American Society for Microbiology.

Osa, S.R., and Weksler, M.E. (1977):

Demonstration of significant differences in the proliferative capacity of lymphocytes from normal human subjects.
Cell. Immunol., 32: 391.

Otani, S., Usuki, S., Saitoh, T. et al. (1991):

Comparison of endothelin-1 concentrations in normal and complicated pregnancies.
J. Cardiovasc. Pharmacol., 17 (Suppl. 7): S308-312.

Paloczi, K., Ujhelyi, E., Fuchs, D., and Mihalik, R. (1991):

Correlation of the percentage of activated CD3+DR+ lymphocytes to serum neopterin levels in HIV-seropositive haemophiliacs.
Klin-Wachenschr., 26 : 69 (4), P. 143 (abstract).

Pasca, A.S. (1977):

Impairment of immunity during pregnancy and antiviral effect of amniotic fluid.
Lancet, I:330.

Patterson, E., Borquist, H., Albercht, A., Von Saltza, M., and Stokstad, E. (1955):

A new pteridin in urine required for growth of protozoon Crithidia Fasciculata.
J.Am. Chem. Soc., 77:3167.

Pence, M., Petty, W.M., and Rocklin, R.E. (1975):

Suppression of maternal responsiveness to paternal antigens by maternal plasma.
J. Immunol., 114: 525.

Peterson, R.E., and Imerato Mcbinley, J. (1976):

Cortisol metabolism in the perinatal period. In New, M.I., and Fiser, R.H. (eds); Diabetes and other endocrine disorders during pregnancy and the newborn. Liss New York PP. 141.

Petrucco, O.M., Seamark, R.F., Holmes, K., Forbes, I.J., and Ssymons, R.G. (1976):

Changes in lymphocyte function during pregnancy.
Br. J. Obst. Gyn., 83: 245.

Pfleiderer, W. (1987):

Chemistry and properties of dihydropterins. In "Biochemical and Clinical Aspects of Pteridines", Vol.5, PP. 3-21. Wachter de Grueter and Co. Berlin, New York.

Pope, R.M., Yoshinoya, S., and Persellin, R.H. (1982):

The detection of circulating immune complexes and IgG and IgM rheumatoid factors in normal human pregnancy.
Am. J. Reprod. Immunol., 2: 208.

Present, P.A., and Comstock, G.W. (1975):

Tuberculin sensitivity in pregnancy.
Am. Rev., Respir. Dis., 112: 413.

Pritchard, J.A. (1980):

Hypertensive disorders in pregnancy. In Williams Obstetrics, Sixteen edition. Edited by Pritchard, J.A., and Mac Donald, P.C. New York, Appleton-Century-Crofts, P. 665-700.

Pritchard, J.A., MacDonald, P.C.; and Grants, N.F. (1985):

Hypertensive disorders in pregnancy. In Williams Obstetrics, Seventeen edition, Edited by Pritchard, J.A., and MacDonald, P.C. New York, Appleton-Century-Crofts, p. 525.

Pross, H.F., and Baines, M.G. (1977):

VI. A brief review.
Cancer Immunol. Immunother.; 3:75.

Pulm, J., Thiery, M., and Sabbe, L. (1978):

Distribution of mononuclear cells during pregnancy.
Clin. Exp. Immunol., 31:45.

- Purtilo, D.T., Hallgren, H.M., and Yunis, E.J. (1972):**
Depressed maternal lymphocyte response to phytohemagglutinin in human pregnancy.
Lancet, 1:769.
- Ramon, R. (1982):**
Pre-eclampsia-eclampsia and other gestational edema-proteinuria-hypertension disorders (GEPH). In: Benson, R.C. Current obstetric and gynecologic diagnosis and treatment. Los Altos, California: Lange Medical Publication, p: 738.
- Rappaport, V.J., Hirata, G., Yap, H.K.; and Jordan, S.C. (1989):**
Antivascular endothelial cell antibodies in sever pre-eclampsia.
Am. J. Obstet. Gynecol.; 162:138-46.
- Ray, J.G. (Ed) (1976):**
Manual of tissue typing techniques. D. HEW Publication. No. (NIM), 78-545, P: 22.
- Redman, C.W., Beilin, L.; and Bonnar, L. (1976):**
Renal function in pre-eclampsia.
J. Clin. Pathol.; 29 (Suppl. 10): 91.
- Redman, C.W., Bodmer, W.F., Bodmer, J.G., and et al.(1978):**
HLA antigens in sever pre-eclampsia.
Lancet; 2:397.
- Regan, L., Braude, P.R., and Hill, D.P. (1991):**
A prospective study of the incidence, time of appearance and significance of anti-paternal lymphocytotoxic antibodies in human pregnancy.
Hum. Reprod., 6:294.
- Reibnegger, G., Fuchs, D. Hausen, A., et al. (1984):**
Urinary neopterin in malignant diseases of a childhood. A marker for activity of the cell mediated immunity. Tumour Diagn. Ther. 5:234-237. Quoted from Wachter, H., Fuchs, D., Hausen, A. Reibnegger, G., Werner, E. (eds): (1989): Neopterin as a marker for activation of cellular immunity: Immunological basis and clinical application. Adv. Clin. Chem. 27:82-139.

Reibnegger, G., Egg, D., Fuchs, D., Ginther, R., and Wachter, H. (1986):

Urinary neopterin reflects clinical activity in patients with rheumatoid arthritis.

Arth. Rheum., 29(9) : 1063.

Reinherz, E.L., and Schlossman, S. (1981):

The differentiation and function of human T-lymphocytes.

Cell, 19:821.

Rembold, H., and Buschmann, L. (1963):

Struktur und synthese des neopterins. Chem, Ber. 96: 1406-1410. Quoted from Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G., and Werner, E. (1989): Neopterin as a marker for activation of cellular immunity: Immunological basis and clinical application (Review), Adv. Clin. Chem. 27: 81-141.

Repke, J. (1991 a):

Calcium, magnesium, and zinc supplementation and perinatal outcome.

Clin. Obstet. Gynecol., 34(2): 262-267.

Reynolds, J. and Brown, G. (1964):

The biosynthesis of folic acid. J. Biol. Chem. 239: 317-325. Quoted from Wachter, H., Fuchs, D., Hausen, A., Reibnegger, G., and Werner, E. (1989): Neopterin as a marker for activation of cellular immunity. Immunobiological basis and clinical application. Adv. In Clin. Chem. 27: 82-139.

Richard, E., Kronenberg, L., Howard, M., Rosenblatt, M., et al. (1982):

Interferon: immunology and Clinical Significance.

Annals of Internal Medicine, 96: 80-93.

Rick, P.S., Elliot, J.P.; and Freeman, P.K. (1980):

Use of corticosteroids in pregnancy induced hypertension.

Obstet. Gynecol.; 55:206.

Roberts, J., Taylor, R., Musci, T., Rodgers, G. Hubel, C. and McLaughlin, M. (1989):

Pre-eclampsia: An endothelial cell disorder.
Am. J. Obstet. Gynecol.; 161: 1200-1204.

Roberts, J., Taylor, R., and Goldfien, A. (1991):

Endothelial cell activation as a pathogenetic factor in pre-eclampsia.
Semin. Perinatol.; 15 (1) :86-93.

Robertson, W.B., Borsevis, I., and Dixon, H.G. (1975):

Uteroplacental vascular pathology.
Eur. J. Obstet. Gynecol. Reprod. Biol., 5:47.

Rokos, H., and Rokos, K. (1983):

A radioimmunoassay for determination of D-erythro-neopterin. In "Chemistry and Biology of pteridines" (J.A. Blair, ed), PP. 815-819, de Gruyter, Berlin. and New York.

Rokos, H., Bienhaus, G., Gadow, A., and Rokos, K., (1985):

Determination of neopterin and reduced neopterins by radioimmunoassay. Biochemical and Clinical Aspects of pteridines, Vol.4, Wachter, H., Curtius, H., Pfeleiderer, W. (eds). Walter de Gruyter, Co. Berlin, New York.

Rokos, H., Frisium, H., and Krunze, R. (1986):

Neopterin and dihydroneopterin in serum of controls and patients with various diseases. Circadian rhythm of neopterin. Influence of cortisol? In: Cooper, B. and Whitehead, V. (eds). Chemistry and Biology of pteridines, Walter de Gruyter. Berlin. pp. 411-414.

Romero, R., Lookwood, C., and Hobbins, C. (1988):

Toxemia: new concepts in an old disease.
Semin. Perinatol.; 12 (4): 302-323.

Rote, N.S., and Caudle, M.R. (1983):

Circulating immune complexes in pregnancy, pre-eclampsia and autoimmune disease: evaluation of Raji cell enzyme-linked immunosorbent assay and polyethylene glycol precipitation methods.
Am. J. Obstet. Gynecol., 147: 267.

- Saftlas, A.F., Olson, D.R., Frank, A.L., Atrash, H.K., and Pokras, R. (1990):**
Epidemiology of pre-eclampsia and eclampsia in the United States, 1979-1986.
Am. J. Obstet. Gynecol., 163: 460-465.
- Sakurai, A., and Gotto, M. (1967):**
Neopterin; isolation from human urine.
J. Biochemistry, 61: 142.
- Sargent, I.L., Redman, C.W., and Sirrate, G.M. (1982):**
Maternal cell mediated immunity in normal and pre-eclamptic pregnancy.
Clin. Exp. Immunol., 50(3): 601-609.
- Schena, F.P., Manno, C., Selvaggil, L.G., and Pastone, A. (1979):**
Immune complexes in normal pregnancy and pre-eclampsia.
Lancet. i: 333.
- Schiller, J., Storer, B., and Witt, P. (1991):**
Biological and clinical effects of I.V. TNF-alpha administered three times weekly.
Cancer Research, 51 (6): 165.
- Schoedon, G., Troppmair, J., Fontana, A., Huber, C., Curtius, H., and Niederwieser, A. (1987):**
Biosynthesis and metabolism of pterins in peripheral blood mononuclear cells and leukaemic lines of man and mouse.
Eur. J. Biochem., 166: 303.
- Schreiber, A.D. (1977):**
Clinical immunology of the corticosteroid.
Prg. Clin. Immunol., 3:103.
- Schrocksnadel, H., Herold, M., Steckel-Berger, G.; Fuchs, D., et al. (1992):**
Cellular immunity in pregnancy-induced hypertensive disease.
Geburtshilfe-Frauenheilkd, Oct., 52 (10):592-5 (Abstract).

Scott, J.R., Beer; and Stasny, P. (1976):

Immunogenetic factor in pre-eclampsia and eclampsia.
J. Am. Med. Ass., 233: 402.

Scott, J.R., (1977):

Fetal growth retardation associated with maternal administration
of immunosuppressive drugs.
Am. J. Obstet. Gynecol., 128: 668.

Scott, J.R., Jenkins, D.M.; and Need, J.A. (1978):

Immunology of pre-eclampsia.
Lancet, 1: 704.

Sculier, J., and Body, J. (1991):

Intravenous administration of amphotericin B entrapped in
liposomes: Induction of high serum levels of Tumour Necrosis
Factor (TNF)-alpha.
Ann. Oncol., 2(2): 141-4 (abst.).

Seidman, D.S., Ever-Hadani, P., Stevenson, D.K. et al., (1989):

The effect of abortion on the incidence of pre-eclampsia.
Eur. J. Obstet. Gynecol., Reprod. Biol., 33:109.

Serhal, P., and Craft, I. (1987):

Immune basis for pre-eclampsia: evidence from oocyte
recipients.
Lancet, i: 744.

**Shintaku, M., Isshiki, G., Hase, Y., Tsuruhara, T. and Oura, T.
(1982):**

Normal pterin values in urine and serum in neonates and its
age-related change throughout life.
J. Inherited Metab. Dis., 5:241-242.

Sibai, B.M., El-Nazer, A., and Gonzalez-Ruiz, A.R. (1986):

Sever pre-eclampsia-eclampsia in young primigravidas:
subsequent pregnancy out-come and remote prognosis.
Am. J. Obstet. Gynecol., 155:1011.

Sibai, B.M., (1990):

A protocol for managing severe pre-eclampsia in the second trimester.

Am. J. Obstet. Gynecol.; 163:733-738.

Sibai, B.M. (1991):

Immunological aspects of pre-eclampsia.

Clin. Obstet. Gynecol.; 34:27.

Sibai, B.M. and Mabie, W. (1991):

Hemodynamics of pre-eclampsia.

Clin. Perinatol.; 18(4): 727-747.

Sibai, B.M., Mercer, B., and Sarinoglu, C. (1991):

Severe pre-eclampsia in the second trimester: recurrent risk and long term prognosis.

Am. J. Obstet. Gynecol.; 165: 1408-1412.

Siegel, M., and Greenberg, M. (1955):

Incidence of poliomyelitis in pregnancy.

N. Engl. J. Med., 253: 841.

Siegel, M. and Gleicher, N. (1981):

Changes in peripheral mononuclear cells in pregnancy.

Am. J. reprod. Immunol., 1:154.

Siklos, A., Nemeth, A., Barlallits, L., et al. (1987):

Decreased killer cell activity in pre-eclampsia.

Gynecol. Obstet. Invest, 23: 84.

Simmonds, R.L., Price, A.L., and Ozerkis, A.J. (1968):

The immunologic problems of pregnancy: The effect of estrogen and progesterone on allograft survival.

Am. J. Obstet. Gynecol., 100:908.

Simmons, R.L. (1969):

Histocompatibility and the survival of the fetus.

Transpl. Proc. 1:47.

- Smith, J.K., Caspary, E.A., and Field, E.J. (1972 a):**
Lymphocyte reactivity to antigen in pregnancy.
Am. J. Obstet. Gynecol., 113:602.
- Smith, J.K., Caspary, E.A.; and Field, E.J. (1972 b):**
Immune responses in pregnancy.
Lancet i : 96.
- Spargo, B., Mc Cartney, C.P.; and Winemiller, R. (1959):**
Glomerular capillary endotheliosis in toxemia of pregnancy.
Arch. Pathol.; 68: 593.
- Sridama, V., Yangs, S.L., Moaward, A., and Degroot, L.J. (1983):**
T-cell subsets in patients with pre-eclampsia.
Am. J. Obstet. Gynecol., 147:566.
- Stevenson, A.C., Davison, B.C., and Say, B. (1971):**
Contribution of feto-maternal incompatibility to etiology of pre-eclamptic toxemia.
Lancet; 2: 1286.
- Stirrat, G.M., Redman, C.W.G., and Levinsky, R.J. (1978):**
Circulating immune complexes in pre-eclampsia.
Br. Med. J., i: 450.
- Stirrat, G.M. (1987):**
The immunology of hypertension in pregnancy. In: Sharp, F., Symonds, E.M. (eds). Hypertension in pregnancy. Ithaca, N.Y.: Perinatology Press, 249.
- Stites, D.P., and Rodgers, R.P. (1987):**
In: Basic and Clinical immunology, 6th ed. Edited by Stites, D.P., Stobo, J.D., and Wells, J.V. Wells. Norwalk, Appelton and Lange.
- Stites, D., P., and Terr, A., I. (1991):**
Physiologic and environmental influences on the immune system
In Basic and Clinical Immunology, 7th edition. A large medical book. Appleton and Lange, U.S.A.

Strelkauskas, A.J., Davies, I.J., and Dray, S. (1978):

Longitudinal studies showing alternation in the levels and functional response of T- and B-lymphocytes in human pregnancy.

Clin. Exp. Immunol., 32: 531.

Studd, J.W. (1971):

Immunoglobulins in normal pregnancy, pre-eclampsia and pregnancy complicated by nephrotic syndrome.

J. Obstet. Gynecol. Br. Common W., p: 78.

Sulevnic, V., and Poljakovic, L. (1973):

Immunoglobulins (IgG, IgM, and IgA) in normal and high risk pregnancy.

Acta. Eur. Fertil. 4:89.

Sumioki, H., Shimokawa, H., Miyamoto, S., et al. (1989):

Circadian variations of plasma atrial natriuretic peptide in four types of hypertensive disorders during pregnancy.

Br. J. Obstet. Gynecol.; 96: 922-7.

Teisner, B., Hay, J., and Tucker, M., (1982):

Circulating C3c and C3d during normal pregnancy.

Am. J. Reprod. Immunol. 2: 309.

Thomson, J.S., Crawford, M.K., Reilly, R.W., and Severson, C.D. (1967):

The effect of estrogenic hormones on immune responses in normal and irradiated mice.

J. Immunol. 93: 331.

Thomson, N.C., Stevenson, R.D.; Behan, W.H. et al. (1976):

Immunological studies in pre-eclamptic toxemia.

Br. Med. J. I: 1307.

Thong, Y.H., Steele, R.W., and Vincet, M.M. (1973):

Impaired in vitro cell-mediated immunity to rubella virus during pregnancy.

N. Engl. J. Med., 289:604.

Thorsby, E., and Piazza, A. (1975):

Joint Report II. Typing for HLA-D determinants (LD-1 or MLC). In Kissmeyer Nielsen, F. (ed). Histocompatibility Testing. Copenhagen, Munksgaard, PP. 414-458.

Tillikainen, A., Schroder, J., and Dela Chapelle, A. (1974):

Fetal leucocytes in the maternal circulation after delivery. II. Masking of HLA-antigens. Transplant, 17: 355.

Tizard, I.R. (1988):

In: Immunology: An Introduction, 2nd edition. Saunders College Publishing, New York, Philadelphia.

Toder, V., Blank, M., Gleicher, N., et al. (1983):

Activity of natural killer in normal pregnancy and edema-proteinuria-hypertension gestosis. Am. J. Obstet. Gynecol., 145:7.

Toder, V., Nebel, L. and Gleicher, N. (1984):

Studies of natural killer cells in pregnancy. Analysis of the single cell level. J. Clin. Lab. Immunol., 14: 123.

Troppmair, J., Lange, A., Huber, C., Schoedon, G., Niederwieser, A., and Adolf, G. (1986):

Intracellular changes in pterin releasing peripheral blood mononuclear cells after stimulation with interferon-gamma. In: Wachter, H., Curtius, H., Pfeleiderer, W. (eds). Biochemical and Clinical Aspects of Petridines, Vol.4. Walter de Gruyter; 363.

Tsukimori, K., Maeda, H., Shingu, M., Koyanagi, T., Nobunaga, M.; and Nakano, H. (1992):

The possible role of endothelial cell in hypertensive disorders during pregnancy. Obstet. Gynecol.; 80:229-233.

Urbaniak, S.J., White, A.G., Barclay, G.R., Wood, S.M.; and Kay, A.B. (1987):

In: Handbook of Experimental Immunology, Vol.3. Third ed.
Edited by Weir, D.M., Oxford, Blackwell Scientific Publications.

Varga, P., Gunther, W., Bolte, A. and et al. (1991):

Natural lymphocyte cytotoxicity of women with different symptoms
of toxemia.

Eur. J. Obstet. Gynecol. Reprod. Biol.; 39:133.

Varma, T.R. (1982):

Serum uric acid as an index of fetal prgnosis in pregnancies
complicated by pre-existing hypertension in pre-eclampsia of
pregnancy.

Int. J. Gynecol.; 20:401.

**Vazquez-Escobsca, C., Perz-Medina, R., and Gomez-Estrada, H.
(1983):**

Circulating immune comoplexes in hypertensive disease of
pregnancy.

Obstet. Gynecol., 62:45.

Wachter, H., Hausen, A., and Grassmayer, K., (1979):

Erhohte Ausscheidung Von Neopterin in Harn Von patieaten mit
maligen tumoren und mit viruse rkrankungen. Hoppeseyler's
Z-physiol. Chem. 360, 1957-1960. In Wachter, H., Fuchs, D.,
Hausen, A. (eds). (1989): Neopterin as a marker for activation of
cellular immunity. Immunologiocal basis and Clinical application.
Adv. In Clin. Chem. 27: 82-139.

**Wachter, H., Hausen, A., Grassmayer, K. et al. Z. Physiol.
Chem., 360:1957 (1979):** Quoted from Huber, C., Troppmair,
J., Fuchs, D. et al. (1985): Immune response-associated
production of neopterin: Relese from macrophages under control
of lymphokine.

Transplantation Proceddings, Vol. XVII, No. 1, P: 582-5.

Wachter, H., Fuchs, D., Hausen, Reibnegger, G., and Werner, E. (1989):

Neopterin as a marker for activation of cellular immunity. Immunological basis and clinical application. Adv. In Clin. Chem. 27: 82-139.

Wallenburg, H., (1989):

Detecting hypertensive disorders of pregnancy. In Chambers I., Enken, M., and Keirse, M., (editors). Effective care in pregnancy and childbirth, Vol. I, Pregnancy, Oxford England, Oxford University Press.

Wallenburg, H., Dekker, G., and Makovitz, J. (1991):

Effect of low dose aspirin on vascular reactivity in angiotensin sensitive primigravid women. Am. J. Obstet. Gynecol.; 146: 1169-1173.

Wasih, S. (1990):

Physiology of low dose aspirin therapy for the prevention of pre-eclampsia. Semin. perinatol.; 14(2) : 152-170.

Wang, Y., Kay, H., and Killam, A. (1991):

Decreased levels of polyunsaturated fatty acids in pre-eclampsia. Am. J. Obstet. Gynecol.; 164: 812-818.

Weinberg, E.D. (1984):

Pregnancy associated depression of cell mediated immunity. Rev. Infectious Dis., 6:814.

Weinstein, L. (1982):

Syndrome of haemolysis, elevated liver enzymes and low platelet count. A severe consequence of hypertension in pregnancy. Am. J. Obstet. Gynecol.; 142: 159.

Weir, R.J. (1979):

Hypertension and pre-eclampsia in pregnancy. In: Ian Donald (ed). Practical obstetric problems, 5th ed. London, Lippincott, Ltd., 284-320.

- Weir, D.M., and Stewart, J. (1993):**
In: Immunology. edited by Weir, D.M. & Stewart, J. 7th ed.
Churchill Livingstone. London. Medical Division of Longman
Group UK limited. P.: 119-135.
- Werner, E. Bichler, A., Daxenbichler, G., Fuchs, D. Hausen, A.,
Hetzel, H., and et al. (1987 a):**
Determination of neopterin in serum and urine.
Clin. Chem.; 33: 62-66.
- Werner, E., Fuchs, D., Hausen, A., Reibnegger, G. and Wachter,
H. (1987 b):**
Proposed biochemical function of tetrahydroneopterin in
interferon (IFN) - gamma activated macrophages.
Biochemical and Clinical Aspects of Pteridines, 5: 143-152.
- Werner, E., Felmayer, W., Fuchs, D., Hausen, A., Reibnegger,
G., and Wachter, H. (1989):**
Parallel induction of tetrahydrobiopterin biosynthesis and
indoleamine 2,3-dioxygenase activity in human cells and all lines
by interferon (IFN)-gamma.
Biochem. J., 262: 861-866.
- Werthamer, S., Govindaraj, S. and Amaral, L. (1976):**
Placenta transcortin and localised immune response.
J. Clin. Invest., 57: 1000.
- Wide, L. (1969):**
A two-site sandwich variation of IRMA assays.
Acta. Endocrinol. Suppl., 142:207.
- Wohlert, M.S., Hing, J., Pederson, E.B., and et al. (1984):**
Circulating immune complexes and hypertension in pregnancy.
Acta. Obstet. Gynecol. Scand., 63:609.
- Wolfe, S.A., Tracey, D.E., and Henney C.S. (1976):**
Induction of natural killer cells by BCG.
Nature (London). 262:584.

- Woloszczuk, W., Schwarz, M., Phohana, E., Kovarik, J. (1985):**
Relation of neopterin to gamma interferon serum levels in patients with kidney transplants. In: Wachter, H., Curtius, H.C., Pfeleiderer, W. (eds): Biochemical and Clinical Aspects of Pteridines, Vol. 4. New York: Walter de Gruyter, 1985:489.
- Woloszczuk, W., Troppmair, J., Leitner, E., Flener, R., Schwartz, M., Kovarik, J., Pohanka, E., and Huber, C. (1986):**
Relationship of interferon (IFN)-gamma and neopterin levels during stimulation with alloantigens in vivo and in vitro. Transplantation, 4 (6): 716-719.
- Woodroffe, A.J., Foldes, M., McKenzie, P.E., Thompson, A.J., et al. (1979):**
Serum immune complexes and disease. Aust. N.Z.J. Med. 9:129.
- Working Group on High Blood Pressure in Pregnancy (1990):**
National high blood pressure education program working group report on high blood pressure in pregnancy. Am. J. Obstet. Gynecol.; 163: 1689-1712.
- Worthington-Roberts, B., and William, S. (1989):**
Nutrition in pregnancy and lactation, 4th ed. St. Louis, times Mirror/Mosby College Publishing.
- Yalow, R.S.; and Berson, S.A. (1960):**
J. Clin. Invest.; 39: 1157. Quoted from Manual of Laboratory Immunology. Edited by Miller, L.E., Ludke, H.R., Peacock, J.E.; and Tomar, R.H. Second Edition. pp.67-70, Lea and Febiger, Philadelphia. London. (1991).
- Yang, S.L., Kleinman, A.M., and Wei, P.Y. (1975):**
Immunologic aspects of term pregnancy toxemia. A study of immunoglobulins and Complement. Am. J. Obstet. Gynecol., 122:727.

Yunis, E.J., and Amos, D.B. (1971):

Three closely linked genetic systems relevant to transplantation.
Proc. Natl. Acad. Sci. USA, 68: 3031-3035.

Ziegler, I., Hamm, U., and Berndt, J. (1983):

Participation of pterines in the control of lymphocyte stimulation
and lymphoblast proliferation.
Cancer Research, 43: 5356-5359.

Ziegler, I., Schwulera, U., and El Wart, J. (1986):

Pteridines are produced during interleukin (IL)-2 induced T cell
proliferation and modulate transmission of this signal.
Exp. Cell. Res., 167: 531.

Zuspan, F. (1991 b):

New concepts in the understanding of hypertensive diseases
during pregnancy.
Clin. Perinatol.; 18 (4): 653-659.