

REFERENCES

- Alan Murray, Neil, B. McLaughlin, John P. Bourke, J. Colin Daig, Stephen S. Furniss and Ronald, W.F., Campbell (1994): Errors in manual measurement of QT intervals. *Br. Heart J.* 71: 386-390.
- Anderson, M.T., Lee, G.B. and Champion, B.C. (1972): Cardiac dysrhythmia associated with exercise stress testing. *Am. J. Cardiol.*, 30: 763.
- Anne, M. Gillis, Movhieddin Trabouls; John T.Y. Hii, D Georyewyse (1998): Antiarrhythmic drug, effect, on QT interval dispersion in patient under 90 ECG testing for ventricular Cardia. *Am. J. Cardiol.*, 81: 588-593.
- Antman, E.M. and Braunwald, E. (1996): Arrhythmias in acute myocardial infaction. Braunwald E. Heart disease, A Textbook of Cardiovascular Medicine, Fifth ed. W.B. Saunders Company, Philadelphia, p. 1246.
- Antman, E.M. and Rutherford, J.D. (1992): Coronary care medicine: a practical approach. *Journal of American College of Cardiology*, 18: 1.
- Astrand, P.O. and Soltin, B. (1964): Plasma and red cell volume after prolonged sever exercise. *J. Appl. Physiol.*, 19: 819-823.
- Atterhog, j.H.; Ekelund, L.G.; Kaijser, L. (1971): Electrocardiographic abnormalities during exercise 3 weeks to 18 months after anterior myocardial infarction. *Br. Heart J.* 33: 871.

- Barber, M.J., Mueller, T.M. and Davies, B.G. (1985): Interruption of sympathetic and vagal-mediated afferent responses by transmural myocardial infarction. *Circulation*, 72: 623.
- Behar, S., Reicher Ress, H. and Schechter, M. (1993): Frequency and prognostic significance of secondary ventricular fibrillation complicating acute myocardial infarction. *American Journal of Cardiology*, 71: 152.
- Bellet, S.; Eliakim, M.; Deliyannis, S. and Figallo, E.M. (1962): Radioelectrocardiographic changes during strenuous exercise in normal subjects, *Circ.* 25: 686.
- Beregard, S.; Holmgren, A. and Jonsson, B. (1960): The effect of body position on the circulation at rest and during exercise with special reference to the influence on the stroke volume, *Acta Physiol. Scand.*, 49: 279-298.
- Berisso, M.Z., Carratino, L. and Ferroni, A. (1990): Frequency, characteristics and significance of supraventricular tachycardias detected by 24-hours electrocardiographic recording in the late hospital phase of acute myocardial infarction. *American Journal of Cardiology*, 65: 1064.
- Bernard, C. (1992): Exercise stress testing, Braunwald, *Heart Disease*, 161, 179.
- Bigger, J.T., Dresdale, R.J. and Heissenbutiel, R.H. (1977): Ventricular arrhythmias in ischaemic heart disease: mechanism, prevalence, significance and management. *Prog. Cardiovasc. Dis.*, 19: 255.
- Bigger, J.T., Fleiss, J.L. and Kleiger, R. (1984): The multicenter post-infarction research group. The relationships among ventricular

- arrhythmias, left ventricular dysfunction and mortality in the 2 years after myocardial infarction. *Circulation*, 69: 250-258.
- Boinean, J.P. and Cox, J.C. (1973): Slow ventricular activation in acute myocardial infarction: a source of reentrant ventricular contraction. *Circulation*, 48: 702-713.
- Bonoris, P.E; Greenberg, P.S.; Castellant, M.D. and Ellestad, M.H. (1978): Significance of changes in R wave amplitude during treadmill stress angiographic correlation, *Am. J. Cardiol.*, 41: 846-853.
- Boudoulas, H.; Ruff, p.D.; Fulkerson, P.K.; Lewis, R.P.; Dervenogas, S. (1981): Effect of propranolol on postexercise left ventricular ejection time index. *Am. J. Cardiol.* 48: 357-60.
- Bourke, J.P., Richards, D.a.B. and Ross, D.L. (1991): Routine programmed electrical stimulation in survivors of acute myocardial infarction for prediction of spontaneous ventricular tachyarrhythmias during follow-up: results, optional stimulation protocol and cost-effective creening. *Journal of American Colleague of Cardiology*, 18: 780-788.
- Brandes, A.; Bethge, K.P. and Buerschaper, M. (1997): QT dispersion in healthy subjects *J. Am. Coll. Cardiol.* 29(2): Suppl. A: 508A.
- Califf, R.M., McKinnis, R.A. and Burks, J. (1982): Prognostic implications of ventricular arrhythmias during 24-hour ambulatory monitoring in patients undergoing cardiac catheterization for coronary artery disease. *American Journal of Cardiology*, 50: 23-31.

- Campbell, R.W.F. (1994): Arrhythmias in Julian, D. and Braunwald, E. Management of acute myocardial infarction. London, W.B. Saunders, Ltd. p. 223.
- Campbell, R.W.F. (1994): Arrhythmias in Julian D and Braunwald E; Management of aCute Myocardial Infarction. London, W.B. Saunders Ltd. p. 223.
- Campbell, R.W.F., Murray, A. and Julian, D.G. (1981): Ventricular arrhythmias in first 12 hours of acute myocardial infarction: natural history study. British Heart Journal, 46: 351.
- Casico, W.E.; Woefel, A.; Knisley, S.B.; Stock, K.S. et al. (1988): Use dependence of Amiodaron during the sinus tachycardia of exercise in CAD. Am. J. Cardio., 16: 1042.
- Chairman, B.R. (1997): Exercise Stress testing. In Braunwald textbook of Cardiovascular Medicine. [edited by] Eugene Braunwald-5th ed. W.B. Saunders Company, Philadelphia, Vol. 1, Chap. 5 p. 153-176.
- Chaitman, B.R.; Stone, P.H.; Knatterud, G.L.; Henry, B.L. et al. (1995): Asymptomatic Cardiac Ischemic Pilot (ACIP) study: Impact of anti-ischemic therapy on 12-week rest ECG and exercise test outcomes. J. Am. Coll. Cardiol. 26: 585.
- Challapalli, Ligamnei, R. and Ehlert, F. (1997): Gender difference in QT dispersion. J. Am. Coll. Cardiol., 29(2) Suppl. A. : 148.
- Chetan P. Shah, Ranjan, K. Thakur, Earl J. Reisdorff, Ed. lone (1998): QT dispersion may be a useful adjunct for detection of myocardial infarction in the chest pain center. Am. Heart J., 136: 496-8.

- Chung, E.K. (1983): Principles of cardiac arrhythmias, third edition. Baltimore, Williams & Wilkins.
- Ciaroni, S.; Delonca, J. and Righetti, A. (1993): Early exercise testing after acute myocardial infarction in the elderly: clinical evaluation and prognostic significance. *Am. Heart J.*, 126: 304.
- Cobb, L.A. and Weaver, D.W. (1986): Exercise a risk for sudden death in patients with coronary heart disease, *J. Am. Coll. Cardiol.* 7: 215-9.
- Cobb, L.A. and Weaver, D.W. (1986): Exercise a risk for sudden death in patient coronary heart disease. *J. Am. Coll. Cardiol.* 7: 215-19.
- Corr, B. and Gillis, R.A. (1978): Autonomic neural influences on the dysrhythmias resulting from myocardial infarction. *Circ. Res.*, 43: 1.
- Cowan, J.C.; Yousoff, K. and Moore, M. (1988): Importance of lead selection in QT interval measurement. *Am. J. Cardiol.* 61: 83-7.
- Crenshaw, B.S., Ward, S.R. and Stebbins, A.L. (1995): Risk factors and outcomes in patients with atrial fibrillation following acute myocardial infarction. *Circulation*, 92 (Suppl.): 1-777.
- Creon, P.A.; Fox, K.M. (1987): Exercise electrocardiography in coronary artery diseases.
- Crimm, A., Severance, H.W., and Caffey, K. (1984): Prognostic significance of isolated sinus tachycardia during the first three days of acute myocardial infarction. *American Journal of Medicine*, 76: 983.
- Davey, P.; Bateman, J. and Mulligan, I.P. (1994): QT interval dispersion in chronic heart failure and left ventricular hypertrophy:

relation to autonomic nervous system and Holter tape abnormalities. *Br. Heart J.*, 71: 268-273.

Day, C.P.; Mc Comb, J.M.; Matthews, J. (1991): Reduction in QT disperison by sotalol following myocardial infraction. *Eur. heart.* 12: 423-427.

Day, C.P.; McComb, j.M. and Compbell, R.W. (1992): QT dispersion in sinus beats and ventricular extra systoles in normal hearts. *Br. Heart.* 67: 139-41.

Day, D.P., McComb, J.M.; Campbell, R.W.F. (1990): QT dispersion: an indication of arrhythmia risk in patient with long QT intervals. *Br. Heart J.*, 63: 342-344.

De Luna, A.B. and Vinolas, X. (1996): QT dispersion and heart rate variability. *Eur. Heart J.*, 17: 165-166.

DeSanctis, R.W., Block, P. and Hutter, A.M. (1972): Tachyarrhythmias in myocardial infarction. *circulation*, 45: 681.

Detry, J.M.R. and Fox, K.M. (1996): Exercise testing. In Julian textbook Diseases of the heart. 2nd ed. W.B. Saunders Company Ltd., London, vol. 1, Chap. 19, p. 372-385.

Dexter, L. and Whittenberger, J.L. (1951): Effect of exercise on circulatory dynamic of normal individual, *J. Appl. Physiol.*, 3: 439-453.

Doycep, M.C. Comb, J.M.; Campbell, R.W. (1990): QT dispersion on indication of arrhythmia risk in patients with long QT intervals. *Br. Heart*, 63: 342-4.

Eduard Homs, Vicens Marti, Ph.D. (1997): Automatic measurement of corrected QT interval in holter recording: comparison of its

- dynamic behavior in patients after myocardial infarction with and without life-threatening arrhythmias. *Am. Heart J.* 134: 181-7.
- Elder, M., Sievner, Z. and Goldbourt, U. (1992): Primary ventricular tachycardia in acute myocardial infarction: clinical characteristics and mortality: the sprint study group. *Ann. Intern. Med.*, 117: 30.
- Ellestad, M.H. (1986): *Stress testing: Principles and Practice*. 3rd ed. F.A. Davis Co. Philadelphia.
- El-Sherif, N., Myerburg, R.J. and Scherlag, B.J. (1976): Electrocardiographic antecedents of primary ventricular fibrillation: value of the R- on T phenomenon in myocardial infarction. *British Heart Journal*, 88: 415.
- Emanuela H. Locati, Wojciech Zareba, Arthur J. Moss, Peter J. Schwartz (1998): Age and sex related differences in clinical manifestation in patient with congenital long-QT syndrome. *Circulation* 97: 2237-2244.
- Ezz El-Din, H. and Khalil, M.T. (1978): The effect of B-blockers on exercise tolerance of patients with angina pectoris. *The Bulletin of Egyptian Society of Cardiology*, 17: 19.
- Farrell, T.G., Paul, V. and Gripps, T.R. (1991): Baroreflex sensitivity and electrophysiological correlates in patients after myocardial infarction. *Circulation*, 83: 945-952.
- Feil, H. and Siegel, M.L. (1928): Electrocardiographic changes during attacks of angina pectoris, *Am. J. Med. Sci.*, 157: 255.

- Fidela, L.I.; Moreno, Materesa, V.; BSMT., Labrose, A.; Koragounis and Jeffiey L. Anderson (1994): Anderson: Reduction QT interval dispersion by successful thrombolytic therapy in AMI. *Circulation* 90: 1-94.
- Finshenfeld, J., Desser, K.B. and Benchimol, A. (1973): Non-paroxysmal A-V junctional tachycardia associated with acute myocardial infarction. *American Heart Journal*, 86: 754.
- Fisch, C.D.; Sanctis, R.W.; Dodge, H.T.; Reeves, T.J. et al. (1986): Guidelines for exercise testing. A report of the American college of cardiology / American Heart Association: Task force on assessment of cardiovascular procedures. *J. Am. Coll. Cardiol.*, 8: 725-738.
- Fisher, M.I. and Nutter, D.O. (1973): Hemodynamic responses to isometric exercise (hand grip) in patients with heart disease, *Br. Heart J.*, 35: 422-432.
- Fleg, J.L., Schulman, S.O., Connor, F., Cerve, V. et al. (1994): Effects of acute B-adrenergic receptor blockade on age-associated change in cardiovascular performance during dynamic exercise *Circ.*, 90: 2333.
- Fletcher, G.F. and Schlant, R.C. (1994): The exercise test in: Harst's textbook of the heart, arteries and veins. 8th ed. by McGraw-Hill, Inc. p. 423-440.
- Fletcher, G.F.; Balady, G.; Froelicher, V.F.; Hartly, L.H. et al. (1995): Exercise standards: a statement for healthcare professionals from the American Heart Association Writing Group, Special Report, *Circ.* 91(2): 580-615.

- Frank Bogun, K. Kwok Chon, Mark Harvey, Bajiva Goyal, Mark Niebauer and Fred Morady (1996): QT dispersion in non sustained ventricular tachycardia and coronary artery diseases. *Am. J. Cardiol.* 77: 256.
- Froelicher, V.F. (1978): Exercise and the heart, clinical concepts, 2nd ed. Chicago, Year Book Medical Publishers, Inc.
- Froelicher, V.F., Wolthuis, R.A., Fisher, J. and Triebwasser, J.H. (1977): The response of healthy men to treadmill exercise. *Circ.*, 25: 153-160.
- Froelicher, V.F.; Myers, J., Follansbee, W.P. and Labovitz, A.J. (1993): Exercise and the heart. 3rd ed. St. Louis, Mosby-Year Book.
- Fronz, M.R.; Swerdlow, C.D.; Lein, L.B. and Schoefer, J. (1988): Cycle length dependence of human action potential duration in vivo. Effects of single extrastimuli, sudden sustained rate acceleration and deceleration and different steady-state frequencies. *J. Clin. Invest.* 82: 972-9.
- Ganz, L.I. and Friedman, P.L. (1995): Supraventricular tachycardia. *National English Journal of Medicine*, 332: 162.
- George, J.; Taylor, Richard, S.; Crampton, Robert S. Gibson, Pamela T. Stebbins, R.N., Maria T.G. Waldman and George A. Beller (1981): Prolonged QT interval at onset of AMI in predicting early phase ventricular tachycardia. *Am. Heart J.* 102: 1-16.
- Gerson, M.C.; Phillips, J.E.; Morris, S.N. and Mc Henry, P.L. (1979): Exercise induced U-wave inversion as a marker of stenosis of left anterior descending coronary artery, *Circ.*, 60: 1014-1020.

- Gilman, J.K., Jalal, S. and Naccarelli, G.V. (1994): Predicting and preventing sudden death from cardiac causes. *Circulation*, 90: 1082-1092.
- Glancy, J.M. Weston, P.J. and Bhullar, H.K. (1996): Reproducibility and automatic measurement of QT dispersion. *Eur. Heart J.*, 17: 1035-1039.
- Goldschlager, N. Cohn, K. and Goldschlager, A. (1979): Exercise related ventricular arrhythmias. *Mod. Concepts cardiovascular disease, Am. Int. Med.*, 85: 277.
- Goldschlager, N. and Sox, H.C. (1988): The diagnostic value of the treadmill exercise test in evaluation of chest pain, in patients with recent myocardial infarction and in asymptomatic individuals. *Am. Heart J.*, 116: 523-533.
- Gooch, A.S. (1972): Exercise testing for detecting changes in cardiac rhythm and conduction. *Am. J. Cardiol.*, 30: 741-747.
- Greeberg, P.S.; Friscia, D.A.; Ellsiead, M.H. (1979): Predictive accuracy of QT/QT ratio, QT_C interval, ST depression and R wave amplitude during stress testing. *Am. J. Cardiol.* 44: 18-23.
- Gressin, V., Gorgels, A.P. and Louvard, Y. (1993): Is arrhythmogenicity related to the speed of reperfusion during thrombolysis for myocardial infarction *European Heart Journal*, 14: 516.
- Hailer, B.; Van Leevwen, P.; Sallner, D. (2000): Changes of QT dispersion in patients with coronary artery disease dependent on different methods of stress induction. *Clin. Cardiol. Nor*; 23(3): 181-6.

- Hassanein, M.T. (1982): Study of systolic time intervals in some cardiac diseases at rest and after physical and pharmacological stress. Thesis (MD) Cardiology, Zagazig University, p. 43.
- Higham, P.D.; Hilton, C.J.; Aitcheson, D.A.; Furniss, S.S.; Bovrke, J.P.; Campbell, R.W.F. (1992): QT dispersion does reflect regional variation in ventricular recovery, *Circulation*, 86 (Suppl.) 392.
- Highom, P.D.; Campbell, R.W. (1994): QT dispersion. *Br. Heart J.*, 71: 508-10.
- Highom, p.D.; Furniss, S.S. and Campbell, R.W. (1995): QT dispersion and components of the QT interval in ischemia and infarction. *Br. Heart J.* 73: 32-36.
- Hii, J.T.Y.; Wyse, D.G.; Gillis, A.M. (1992): QT interval dispersion as a marker of Torsade de pointers. antiarrhythmic drugs and amiodaron. *Circulation*. 86: 1376-1382.
- Hii, J.T.Y.; Wyse, D.G.; Gallis, A.M.; Duff, H.J.; Solylo, M.A.; Mitchell, L.B. (1992): Precordial QT interval dispersion as a marker of torsades de pointes: dispete TS of class Ia ontiarsahythmic drug, and amiodorone. *Circualtion*, 86: 1376-1382.
- Irving, J.B., Bruce, R.A., Derouen, T.A. (1977): Variations in significance of systolic pressure during maximal exercise testing. Relation to severity of coronary artery disease and cardiac mortality. *Am. J. Cardiol.*, 39: 841.
- James M. Galancy, Clifford J. Garrat, Kent, L.; Woods and David, p. DeBono (1995): Use of lead adjustment formulas for QT dispersion after MI. *Br. Heart J.* 74: 676-679.

- Janse, M.J. and Opthof, T. (1995): Mechanisms of ischaemia induced arrhythmias. In Zipes, D.P. and Jalife, J. (eds.) Cardiac electrophysiology from cell to bedside . 2nd ed. Philadelphia, W.B. saunders Company, pp. 489-496.
- Kahan, K.A. and Simonson, E. (1957): Changes of man spatial QRS and vectors and of conventional electrocardiographic items in hom anaerobic work, *Circ. Res.*, 9: 629.
- Kambara, H. and Philips, J. (1976): Long term evaluation of ear repolarization syndrome, *Am. J. Cardiol.*, 58: 157-163.
- Kannel, W.B. (1982): Exercise and sudden death (editorial). *JAMA* 248: 3143-4.
- Kannel, W.B., Sprliep and Gordon, T. (1980): Labile hypertension faculty concept. *Circ.*, 61: 1183-1192.
- Karjolaenen, J.; Viitasolo, M.; Manttani M., et al. (1994): Relation between QT intervals and heart rates from 10 to 120 feets min in rest electrocardiograms of men and a simple method to adjust QT interval values *J. Am. Coll. Cardiol.*, 23: 1741-53.
- Katerina Hantkova, Mack Malika, Josef Kautzner, YiGang and A John Camm (1994): Adjustment of QT dispersion assessed from 12 lead electrocardiograms for different numbers of analyzed electrocardiographic leads: Comparison of stability of different methods. *Br. Heart J.* 72: 390-396.
- Kennedy, H.L., Whitlock, J.A. and Sprague, M.K. (1985): Long-term follow-up of asymptomatic healthy subjects with frequent and compelx ventricular ectopy. *National English. Journal of Medicine*, 312: 193-197.

- Khaching, Kchkhan, Janos, Monnar and John, J. (1997): Relation of left ventricular mass and QT dispersion in patients with systemic hypertension. *Excerpto, Medica-September*, vol. 23: 508-11.
- Kidwell, G.A. and Chung, U.K. (1996): Ischaemic ventricular arrhythmias. In: Fuster, V., Ross, R. and Topol, E.J. (eds.): *Atherosclerosis and coronary artery disease*. Philadelphia, Lippincott-Raven, 995-1012.
- Kink, J.W. and Ryan, T.J. (1994): Right ventricular infarction. *New England Journal of Medicine*, 330: 1211.
- Kirshenbaum, J.M., Kloner, R.F., McGowan, N. (1988): Use of an ultra-short-acting beta receptor blocker (Esmolol) in patients with acute myocardial ischemia and relative contraindications to beta-blockade therapy. *Journal of American College of Cardiology*, 12: 773.
- Kleiger, R.E., Miller, J.P. and Bigger, J.T. (1987): Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *American Journal of Cardiology*, 59: 256-262.
- Koide, Y.; Yotsukura, M.; Tajino, K. (1999): Use of QT dispersion measured on treadmill exercise electrocardiograms for detecting restenosis after percutaneous transluminal coronary angioplasty. *Clin. Cardiol.* 22(10): 369-48.
- Konnel, W.B. (1982): Exercise and sudden death (editorial). *JAMA* 248: 3143-4.
- Koppes, G.M., Kruyer, W., Beckmann, C.H., Hones, F.G. (1980): Response to exercise early after uncomplicated myocardial

- infarction in patients receiving no medication: Long-term follow up. *Am. J. Cardiol.* 46: 764.
- Kuchar, D.L., Thorburn, C.W. and Sammel, N.L. (1987): Prediction of serious arrhythmic events after myocardial infarction: signal-averaged electrocardiogram, Holter monitoring and radionuclide ventriculography. *Journal of American College of Cardiology*, 9: 531-538.
- Lary, D. and Goldschlager, N. (1974): Electrocardiographic changes during hyperventilation resembling myocardial ischemia in patients with normal coronary arteriograms. *Am. Heart J.*, 87: 383-390.
- Lee, H.S.; Cross, S.J.; Rawles, J. (1993): QT_c dispersion in patients with coronary artery disease. effect of exercise, dobutamine and dipyridamole myocardial stress 14: 210.
- Lee, J.H., Crump, R., and Ellestad, M.H. (1995): Significance of precordial T-wave increase during treadmill stress testing, *Am. J. Cardiol.*, 67: 1297-1299.
- Lehmann, M.H.; Timothy, K.; Frankovich, D.; Fromm, B. (1997): Age-sex influence on rate corrected QT interval and QT-heart rate relationship in families with genotypically characterized long QT syndrome. *J. Am. Coll. Cardiol.* 29: 93-99.
- Linhart, J.W. and Turnhoff, H.B. (1974): Maximal treadmill exercise test in patients with abnormal control ECG. *Circ.*, 50: 1179-1189.
- Lown, B., Vassan, X. and Hood, W.B. (1969): Unresolved problems in coronary care. *American Journal of Cardiology*, 20: 494-508.

- Macfarlane, P.W.; Veitch, Lawrie, T.D. eds (1998): Physiological aspects of cardiac arrhythmias. *Am. Heart, J.* 93: 645.
- Makoto Suzuki, Mitsuhiro Niishizaki, Masataka Arita, Takashi Ashikosae (1998): Increased QT dispersion in patient with vasospastic onsgina. *Circulation* , 98: 435-440.
- Marco, B., Robert, D., and Johann, B. (1992): Exercise induced ST depression and ST/HR index to predict triple or fifth main coronary disease, *J. am. Coll. Cardiol.*, 19(1): 11-18.
- Master, A.M. and Oppenheimer, E.T. (1929): A simple exercise tolerance test for circulatory efficiency with standard tables for normal individuals. *Am. J. Med. Sci.*, 177: 223.
- McHenry, P.L. (1977): The actual prevalence of false positive ST-segment responses to exercise in clinically normal subjects remains undefined. *Circulation*, 55: 683.
- Mclaoughlin, N.B., Campbell, R.W.L., Mvrrroy, A (1996): Comparison of automatic QT measurement techniques in the normal 12 lead electrocardiogram. *Br. Heart J.* 7H: 8H-9.
- Medias, J.E., Patel, D.C. and Singh, D. (1996): Atrial fibrillation in acute myocardial infarction. A prospective study based on data from a consecutive series of patients admitted to the coronary care unit. *Clin. Cardiol.*, 19: 180.
- Melandri, G.; Semprini, F.; Cervi, V.; Dube, F. et al. (1993): Benefit of adding low molecular weight heparin to the conventional treatment of stable angina pectoris: A double blind, randomized, placebocontrolle trial. *Circ.* 88: 2517.

- Meltzer, L.E. and Cohen, H.E. (1972): The incidence of arrhythmias associated with acute myocardial infarction. Textbook of coronary care unit. Philadelphia, Charles Press.
- Mirvis, D.M. (1985): Spatial variation of QT interval in normal persons and in patients with acute myocardial infarction, *J. Am. Coll. Cardiol.*, 3: 625-631.
- Mohamed, R.; Forsey, P.R. and Davis, M.K. (1996): Effect of liver transplantation on QT interval prolongation and autonomic dysfunction in end stage. Liver disease *Hepatology*, 23(5): 1128-1134.
- Molfatto G. Beria, G.; Sala, S.; Bonazzio, Schwartz, P.J. (1997): Quantitative analysis of T wave abnormalities and their prognostic implications in the idiopathic long QT syndrome. *J. Am. Coll. Cardiol.*, 23: 296-301.
- Moss, A.J. (1993): Measurement of interval and the risk associated with QT interval prolongation, a review, *Am. J. Card.* 72: 23-25.
- Musha, H.; So, T.; Hashimoto, N. (1999): Dynamic changes of QT dispersion as a predictor of myocardial ischemia on exercise testing in patients with angina pectoris. *Jpn Heart. J. Maur.*; 40(2): 119-26.
- Myerburg, R.J. and Castellanos, A. (1993): Cardiac arrest and sudden cardiac death. Braunwald: Heart Disease: A Textbook of Cardiovascular Medicine, Fifth ed., W.B. Saunders Company, Philadelphia, p. 746.

- Neil, B., McLaughlin, Ronald, W.F. Campbell and Alan Murray, (1995): Comparison of automatic QT measurement techniques in the normal 12-lead electrocardiogram. *Br. Heart J.* 74: 84-89.
- Nicolas, J.; Paolo, Colonna, Christopher, A. (1992): Assessment of QT dispersion syndrome in patients with congenital prolonged QT syndrome. *Am. J. Cardiol.* 69: 634-38.
- Niwa, H.; Meeda, T. and Shimoyama, N. (1993): Serial changes in QT interval during acute myocardial infarction: Comparison of QT interval changes of reperfused cases and non-reperfused cases. *J. Cardiol.*, 23(1): 14-50 (Abstr.).
- Osswald, S.; Trovton, T.G.; O'Nunain, S. et al. (1994): Patients with cardiac arrest and documented ventricular fibrillation. in: Akhtar M., Myerburg, R.J.; Ryskin, N.J., eds. sudden cardiac death. Prevalence, mechanisms, and approaches to diagnosis and management. Moleven: Williams and Wilkins, 465-85.
- Pantridge, J.F. and Adgey, A.A.J. (1969): Pre-hospital coronary care: the mobile coronary care unit. *American Journal of Cardiology*, 24: 666.
- Perkiomaki, J.; Koistinen, Linnal Uoto, M. (1996): Dispersion of QT interval at rest and during exercise in healthy subjects and patients with coronary artery disease. *Eur Heart J.* 14: 254.
- Perkiomaki, J.S., Koishinen, M.J., Yli Moyry S. and Huikuri, H.V. (1995): Dispersion of QT interval in patient with and without susceptibility to ventricular Tachyarrhythmia after previous myocardial infarction, *J. Am. Coll. Cardiol.* 26: 174-9.

- Pipberger, H.V.; Ishikawa, K. and Berson, A.S. (1972): QRS amplitude changes during heart failure and digitalisation. *Am. Heart J.*, 83: 202-297.
- Priori, S.G., Napolitano, C. and Diehl, L. (1994): Dispersion of the QT interval: A marker of therapeutic efficacy in the idiopathic long QT syndrome. *Circ.*, 89(4): 1681-1689.
- Pye, M., Quinn, A.C. and Cobbe, S.M. (1994): QT interval dispersion a non-invasive marker of susceptibility to arrhythmia in patients with sustained ventricular arrhythmia, *Br. Heart J.* 71: 511-4.
- Pye, M.; Quinn, A.C. and Cobbe, S.M. (1994): QT interval dispersion: a non-invasive marker of susceptibility to arrhythmia in patients with sustained ventricular arrhythmias *Br. Heart J.*, 71(6): 511-14.
- Raeder, E.A., Rosenbaum, D.S. and Bhasin, R. (1992): Alternating morphology of the QRST complex preceding sudden death. *national english Journal of Medicine*, 326: 271-272.
- Rochmis, P. and Blackburn, H. (1971): Exercise test. *J. Am. Med. Assoc.*, 216: 1061.
- Roden, D.M.; Lazzara, R.; Rosen, M.; Schwartz, P.J.; Towbin, J.; Vincent M. (1996): Multiple mechanism in the long QT syndrome: current knowledge, gaps and future directions. *Circulation*, 97: 2012.
- Romon, L.; Bellet, S. (1965): Significance of the QX/QT ratio and the QT ratio in the exercise electrocardiogram. *Circulation*, 32: 435-7.

- Rooke, G.A. and Feigl, E.O.: Work as a correlated of canine left ventricular oxygen consumption and the problem of catecholamine oxygen wasting, *Circ. Res.*, 50: 273-286, 1982.
- Rosenbaum, D.S., Jackson, L.E. and Smith, J.M. (1994): Electrical alternans and vulnerability to ventricular arrhythmias. *National English Journal of Medicine*, 330: 23-241.
- Rosenthal, H.A., Liu, H. and Komen, S. (1980): Exercise related R wave amplitude change in patients with chest pain syndromes. *Am. J. Cardiol.*, 46: 312-317.
- Rowell, L.B. (1986): *Human Circulation. Regulation During physical stress*, New York, Oxford University Press.
- Ruberman, W., Weinblatt, E. and Goldberg, J.D. (1977): Ventricular premature beats and mortality after myocardial infarction. *National English Journal of Medicine*, 297: 750-755.
- Salzman, S.H. (1969): Quantitative effects of physical conditioning on the exercise electrocardiograms of middle-aged subjects with arterosclerotic heart disease. In Blackburn. (4th ed.) *Measurements in exercise electrocardiography*. Charles C. Thomas, Spring Field, 111.
- Sannenblick, E.H., Ross, J. and Braunwald, E. (1968): Oxygen consumption of the heart. Newer concepts of its multifactorial determination. *Am. J. Cardiol.*, 22: 328-336.
- Sanz, G., Castaner, A. and Betriu, A. (1982): Determinants of prognosis in survivors of myocardial infarction: a prospective clinical angiographic study. *National english Journal of Medicine*, 306: 1065-1070.

- Schwartz, p.J., LaRovere, M.T. and Vanoli, E. (1992): Autonomic nervous system and sudden cardiac death: Experimental basis and clinical observation for post-myocardial infarction risk stratification. *Circulation*, 85: 177-191.
- Schwartz, P.J.; Locati, E.H.; Napolitano, C.; Priori, S.G. (1955): The long QT syndrome in: Zipes, D.P.; Jalife, J.; editors. *Cardioelectrophysiology: from cell to bedside*. 2nd ed. Philadelphia: W.B. Saunders; p. 788-811.
- Scant, R.C.; Blomqvist, G. and BrandEuburg, R.O. (1980): Guidelines for exercise testing. *Circ.*, 74: 653A - 667A.
- Scott, J.C. (1967): Physical activity and the coronary circulation. *Can. Med. Assoc. J.*, 96: 853.
- Seshadri Balaji, Yung R Lav, Paul C Gillette (1997): Effect of heart rate on QT interval in children and adolescents. 77: 128-129.
- Shaw, L.T., Younis, L.T., Stock, k.S., Wilken, K. et al. (1990): Effects of posture on metabolic and hemodynamic predischarge exercise response after the acute myocardial infarction. *Am. J. Cardiol.*, 66: 134.
- Sheffield, L.T., Holl, J.H. and Reeves, T.J. (1965): Exercise grade4d by heart rate in Electrocardiographic testing for angina pectoris, *Circ.*, 32: 622.
- Sheps, D.S.; Ernst, J.C. and Briese, W.G. (1977): Decreased frequency of exercise induced ventricular ectopic activity in the second of two consecutive treadmill tests. *Circ.*, 55: 892-898.

- Simon, S.S. Lo, Christopher J. Mathias and Martin St John Sutton (1996): QT interval and dispersion in primary autonomic failure. *Heart* 75: 498-501.
- Six, A.J., Louwerenburg, J.H. and Kingma, J.H. (1991): Predictive value of ventricular arrhythmias for patency of the infarct-related coronary artery after thrombolytic therapy. *British Heart Journal*, 66: 143.
- Skovick, D.S.; Weiss, N.S.; Fletcher, R.H.; Losky, T. (1986): The incidence of primary cardiac arrest during vigorous exercise. *N. Engl. J. Med.* 311: 874-7.
- Sporton, S.C., Taggart, p., Sutton, P.M.; Wolker, J.M. (1997): Acute ischemia a dynamic influence on QT dispersion, 349: 306-309.
- Sporton, S.C.; Taggart, p. Sutton, P.M. (1997): Acute ischemia: a dynamic influence on QT dispersion, 349: 306-309.
- Sullivan, M.; Atwood, J.E.; Myers, J.; Feuer, J. et al. (1989): Increased exercise capacity after digoxin administration in patients with heart failure. *J. Am. Coll. Cardiol.*, 13: 1138-1143.
- Tsug Ming Lee, Sheng-Fang Su, Ph.D.; Tzung-Dov wang (1998): Increased ventricular repolarization inhomogeneity during postural changes in patients with syndrome X. *Am. J. cardiol.* 82: 615-620.
- Tsung – Ming Lee, Sheng – Fong SV, Tzung Dav Wang, Wen-lin Wang (1998): Increased ventricular repolarization inhomogeneity during postural changes in patient with syndrome, *Am. J. Cardiol.*, 82: 615-620.

- Ulrich Stierle, Evangelos Giannitsis, Abdol hamid Sheikh zadeh, Dietmorkrugor (1998): Relation between QT dispersion and the extent of myocardial ischemia in patient with three vessel coronary artery disease. *Am. J. Cardiol.*, 81: 564-568.
- Volpi, A., Cavalli, A. and Franzosi, M.G. (1989): One-year pronosis of primary ventricular fibrillation complicating acute myocardial infarction. *American Journal of Cardiology*, 63: 1174.
- Wasserman, k., Hansen, J.E., Sue, D.Y. et al.: *Principles of Exercise Testing and Interpretation*, 2nd ed. Philadelphia, Lea and Febiger.
- Watanabe, Y.; Nishimuro, m. (1984): Clinical electrocardiographic studies of bifid T waves. *Br. Heart J.*, 52: 201-14.
- Wolfe, C.L., Nibley, C. and Bhandari, A. (1991): Polymorphous ventricular tachycardia associated with acute myocardial infarction. *Circulation*, 84: 1543.
- Wyns, W.; Melin, J.A., Van Butsele, R.J. et al. (1982): Assessment of right and left ventricular volumes during upright exercise in normal men. *Eur. H.J.*, 3: 329-536.
- Yoshimura, M.; Matsumoto, K.; Watanabe, M. (1999): Significance of exercise QT dispersion in patients with coronary artery disease who do not have exercise-induced ischemic ST-segment changes. *Jpn Circ. J.* 63(7): 517-21.
- Younis, L.T.; Melin, J.A.; Beckers, C. and Detry, J.M.R. (1990): Influence of age on the radionuclide left ventricular volumes and ejection fraction during maximal upright exercise in normal subjects. *Eur. H.J.*, 11: 916-924.

Yu, P.N.G. and Soffer, A. (1925): Studies of electrocardiographic changes during exercise (Modified double two-step test), *Circulation* 6: 183.

Zobel, M., Portony, S. and Franz, M.R. (1995): Electrocardiography indexes of dispersion of ventricular repolarization: an isoalted heart validation study, *J. Am. Coll. Cardiol.*, 25: 746-52.