
Abbott RD, Donahue RP, Kannel WB, Wilson PWF.
The impact of diabetes on survival following myocardial infarction in men vs women: the Framingham Study.
JAMA. 1988;260:3456.

Adams KF, Sueta CA, Gheorghiade M et al.
Gender differences in survival in advanced heart failure . Insights from the FIRST study.
Circulation 1999;99:1816.

Andrew F, Gray M, You-hui L, and John S.
Gender and age differences found in clotting mechanism of sudden cardiac death.
Circulation. 1998; 98:4891.

Antony I, Aptecar E, Lerebours G, et al.
Coronary artery constriction caused by the cold pressor test in human hypertension.
Hypertension 1994;24:212.

Assey ME.

The puzzle of normal coronary arteries in the patient with chest pain:
What to do?

Clin Cardiol 1993;16:170.

Austin MA.
Plasma triglycerides and coronary heart disease.
Arterioscler Thromb. 1991; 11:2.

Ayanian JZ, Epstein AM.

Differences in the use of procedures between women and men hospitalized for coronary heart disease.

N Engl J Med 1991;325:221.

Bairey Merz CN, Kelsey SF, Pepine CJ, et al.

The Women's Ischemia Syndrome Evaluation (WISE) study: protocol design, methodology, and feasibility report.

J Am Coll Cardiol. 1999;33:1453-1461.

Bateman TM, O'Keefe JH, Dorg VM, Bamhart C.

Prognostic significance of nonischemic SPECT stress myocardial perfusion scintigraphy in women.

J Nucl Med 1994;35:60P.

Barrett-Connor E, Cohn BA, Wingard DL, Edelstein SL.

Why is diabetes mellitus a stronger risk factor for fatal ischemic heart disease in women than in men ?

JAMA. 1991;265:627.

Barrett-Connor E, Wingard DL.

Sex differential in ischemic heart disease mortality in diabetics : a prospective population-based study.

Am J Epidemiol. 1983;118:489.

Beard C, Kottke T, Annegers J, Ballard D.

The Rochester Coronary Heart Disease Project: effect of cigarette smoking, hypertension, diabetes and steroidal estrogen use on coronary heart disease among 40-to-59-year-old women, 1960 through 1982.

Mayo Clin Proc. 1989;64:1471.

Bell MR, Grill DE, Garratt KN, et al.

Long-term Outcome of Women Compared With Men After Successful
Coronary Angioplasty
Circulation. 1995;91:2876.

Birdsall MA.

Association Between Polycystic Ovaries and Extent of Coronary
Artery Disease in Women Having Cardiac Catheterization.
Ann Intern Med. 1997;126:32.

Birdwell BG, Herbers JE, Kroenke K.

Evaluating chest pain.
Arch Intern Med. 1993;153:1991.

Bonow RO.

Cardiovascular Disease in Women: Viva la difference ? XXIst
Congress of the European society of Cardiology 1999.

Bugiardini R, Pozzati A, Ottani F, et al:

Vasotonic angina: A spectrum of ischemic syndromes involving
functional abnormalities of the epicardial and microvascular coronary
circulation.

J Am Coll Cardiol 1993;22:417.

Bullemer F, Graham KJ, Pankow J, et al.
Gender-related differences in risk factors of young patients with
symptomatic coronary artery disease (abstract).
J Am Coll Cardiol. 1995; February Special Issue : 267A.

Burt VL, Whelton P, Roccella EJ, et al.
Prevalence of hypertension in the US adult population. Results from
the Third National Health and Nutrition Examination Survey, 1988-
1991. *Hypertension* 1995;25:305.

Cannon RO , Camici PG, Epstein SE:
The pathophysiological dilemma of syndrome X.
Circulation 1992; 85:883.

Capron L, Jarret J, Kusandjian S, Honsait E.
Growth promoting effects of diabetes and insulin on arteries.
Diabetes. 1986;35:973.

Carmassi F, Morale M, Puccetti R, et al.
Coagulation and fibrinolytic system impairment in insulin dependent
diabetes mellitus.
Thromb Rev. 1992;67:643.

Carmen W, Barrett-Connor E, Sowers M, Khau K.
Higher risk of cardiovascular mortality among lean hypertensive
individuals in Tecumseh, Michigan.
Circulation. 1994;89:703.

Castelli WP. Epidemiology of triglycerides: A view from Framingham . *Am J Cardiol* 1992;70:3H-9H.

Chaitman BR, Bourassa MG, Davis K.
Angiographic prevalence of high-risk coronary artery disease in patient subsets (CASS).
Circulation 1981;269:2392.

Chauhan A.

Syndrome X : angina and normal coronary angiography.
Post Med J. 1995;71:341.

Chua TP, Saia F, Bhardwaj V, et al.
Are there gender differences in patients presenting with unstable angina ?
Intern J Cardiol. 2000;72:281.

Colditz GA, Willett WC, Stampfer MJ, et al .
Menopause and the risk of coronary heart disease in women.
N Eng J Med . 1987;316:1105.

Croft P, Hannaford PC.
Risk factors for acute myocardial infarction in women: evidence from the Royal College of General Practitioners' oral contraception study.
BMJ. 1989;298:165.

Dave TH, Wasir HS, Prabhakaran D, Dev V, Das G, Rajani M, Venugopal P, Tandon R.

Profile of coronary artery disease in Indian women: correlation of clinical, non invasive and coronary angiographic findings.
Indian Heart J 1991 ;43:25.

De Bacquer D, De Backer G, Kornitzer M, et al.
Prognostic value of ischaemic electrocardiographic findings for cardiovascular mortality in men and women.
J Am Coll Cardiol. 1998;32:680.

DeSanctis RW.
Clinical manifestations of coronary artery disease: chest pain in women.
In : *Cardiovascular Health and Disease In Women*, Wenger NK, Speroff L, Packard B.(Eds), Le Jacq Communications, Connecticut, 1993,p.67.

Dhawan J, Bennett DH, Morris J

Paucity of risk factors in women with angina and normal coronary angiograms. .

Int J Cardiol 1991 ; 31 : 181 .

Douglas PS and Ginsburg GS.
The evaluation of chest pain in women .
N Eng J Med. 1996;334:1311.

Donahue RP, Barrett-Connor E, Orchard TJ, Gutai JP.
Endogenous insulin and sex hormones in atherosclerosis and coronary heart disease.
Atherosclerosis. 1988;8:544.

Elhendy A, van Domburg RT, Bax JJ, Roelandt JR.
Gender differences in the relation between ST-T-wave abnormalities at
baseline electrocardiogram and stress myocardial perfusion
abnormalities in patients with suspected coronary artery disease.
Am J Cardiol. 1999;84:865.

Fleischmann KE, Hunink MG, Kuntz KM, Douglas PS.
Exercise echocardiography or exercise SPECT imaging? A meta-
analysis of diagnostic test performance.
J Am Med Assoc 1998;280:913.

Florence AT, Jean-Michel P, Collette C, and Claude R.
Coronary Heart Disease Risk Factors and Menopause :
a study in 1684 French women.
Atherosclerosis 1999;142:415.

✓ Florence HS, Braunwald E, Paul Canner, H, Dodge T, and
Joel G.
Effect of intravenous thrombolytic therapy on left ventricular function :
a report on tissue type plasminogen activator and streptokinase from
the thrombolysis in myocardial infarction (TIMI phase I) Trial.
Circulation 1987, 75; 4 : 817.

Fodor J, Frohlich J, Genest J, McPherson P.
Recommendations for the management and treatment of dyslipidemia
: report of the Working Group on Hypercholesterolemia and Other
Dyslipidemias.
Can Med Assoc J. 2000; 162: 1441.

-
- Frasure-Smith N, Lesperance F.**
Psychosocial risks and cardiovascular diseases.
Can J Cardiol. 1999;15[suppl G] : 93G-97G.
- Garcia MJ, McNamara PM, Gotdon T, Kannel WB.**
Morbidity and mortality in diabetics in the Framingham population:
sixteen-year follow-up study.
Diabetes Care. 1990;265:627.
- Gensini, G.G.:**
A more meaningful scoring system for determining the severity of
coronary artery disease.
Am. J. Cardiol. 1983; 51 : 606.
- Gillum RF.**
Sudden coronary death in the United States 1980-1985.
Circulation 1989;79:756.
- Ginsburg GS, Douglas PS , and Poppas A.**
Summary of the second report of the National Cholesterol Education
Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment
of High Blood Cholesterol in Adults (Adult Treatment Panel II).
JAMA 1993;269:3015.
- Glanz SA, Parmley WW.**
Passive smoking and heart disease: mechanisms and risk.
JAMA 1995;273:1047.
-

Grady D, Rubin SM, Petitti DB et al.

Hormone therapy to prevent disease and prolong life in postmenopausal women.

Ann Intern Med 1992;117:1016.

Grover-McKay M.

Gender-related imaging issues in assessment of coronary artery disease by nuclear techniques.

Am J Card Imaging 1996;10:54.

Grundy SM.

Second Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II).

Circulation 1994;89:1329.

Hachamovitch R, Berman DS, Kiat H, et al.

Effective risk stratification using exercise myocardial perfusion SPECT in women : gender related differences in prognostic nuclear testing.

J Am Coll Cardiol 1996;28:34.

Hanes DS, Weir MR, Sowers JR.

Gender considerations in hypertension pathophysiology and treatment.

Am J Med 1996;101 [suppl 3A] :10 S.

Hansen CL, Crabbe D, Rubin S.

Lower diagnostic accuracy of thallium-201 SPECT myocardial perfusion imaging in women: An effect of smaller chamber size.

J Am Coll Cardiol 1996;28:1214.

Hayashi T, Yamada K, Esaki T, et al.
 Estrogen increases endothelial nitric oxide by a receptor mediated
 system. *Biochem Biophys Res Commun*. 1995;214:847.

Haynes S, Feinleib M.
 Women, work and coronary heart disease: prospective findings from
 the Framingham Heart Study.
Am J Public Health. 1980;70:133.

Hitoshi S, Hajime W, Kiyotaka K, et al.
 Does Passive Smoking Impair Endothelium-dependent coronary Artery
 Dilation in Women?
J Am Coll Cardiol. 1998;31:811.

Hochman JS .
 Sex, Clinical Presentation, and Outcome in Patients with Acute
 Coronary Syndromes.
N Engl J Med. 1999;341:226.

Hochman JS, McCabe CH, Stone PH, Becker RC, Cannon
 CP, DeFeo-Fraulini T, Thompson B, , Steingart R,
 Knatterud G and Braunwald E. Outcome and Profile of Women
 and Men Presenting With Acute Coronary Syndromes: A Report From
 TIMI IIIB
J Am Coll Cardiol 1997;30:141.

Hodis HN, Mack WJ, Krauss RM, Alaupovic P.
Pathophysiology of triglyceride-Rich Lipoproteins in
Atherothrombosis : Clinical Aspects.
Atherosclerosis 1998;125:91.

Holdright DR.
Risk factors for cardiovascular disease in women .
J Hum Hypertens 1998;12:667.

Horwitz KB, Horwitz LD.
Canine vascular tissues are targets for androgens, estrogens,
progestins, and glucocorticoids.
J Clin Invest. 1982; 69: 750.

Hsueh WA, Anderson PW.
Hypertension, the endothelial cell, and the vascular complications of
diabetes mellitus.
Hypertension. 1992;20:253.

Hu FB, Willett WC, Manson JE, et al.
Snoring and risk of cardiovascular disease in women.
J Amer Coll Cardiol. 2000;35:308.

Isles CG, Hole DJ, Hawthorne VM, Lever AF.
Relation between coronary risk and coronary mortality in women of
the Renfrew and Paisley survey: comparison with men
Lancet 1992;339:702.

Jong P, Mohammed S, Sternberg L.

Sex differences in the features of coronary artery disease of patients undergoing coronary angiography.

Can J Cardiol 1996 ;12:671.

Jousilahti P, Tuomilehto J, Vartiainen E, et al.

Body weight, cardiovascular risk factors, and coronary mortality. 15 year follow-up of middle-aged men and women in eastern Finland.

Circulation.1996;93:1372.

Kannel WB, Wilson WF, D'Agostino RB, Janet Cobb.

Sudden Coronary Death in Women

Am Heart J 1998 ;136:205.

**Kennedy J.W., Killip T., Fisher L.D., Alderman E.L.,
Gillespie M.J. and Mock M.B.**

The clinical spectrum of coronary artery disease and its surgical and medical management, 1974–1979: The Coronary Artery Surgery Study.

Circulation 1982, 66:suppl.III:III-16.

King AC.

Physical Inactivity as a Contributor to Heart Disease in Women., The First International Conference on Women, Heart Disease and Stroke: Science and Policy in Action.

Conference summary 2000.

Kim C, Kwok YS, Saha S, Redberg RF.

Diagnosis of suspected coronary artery disease in women : a cost-effectiveness analysis.

Am Heart J . 1999;137:1019.

Kirchgatterer A, Weber T, Auer J, Wimmer L, Mayr H, Maurer E, Eber B.

Analysis of referral diagnoses of patients with normal coronary angiogram.

Wien Klin Wochenschr 1999 ;111:434.

Klein L.W., Agarwal J.B., Herlich M.B., Leavy T.M. and Helfant R.H.

Prognosis of symptomatic coronary artery disease in young adults aged 40 years or less.

Am J Cardiol 1987, 60:1269.

Knopp RH.

Estrogen, female gender, and heart disease. In: Topol EJ, ed. *Textbook of Cardiovascular Disease*. Philadelphia, Pa: Lippincott-Raven Publishers; 1998:195-218.

Kober L, Torp-Pedersen C, Ottesen M, et al.

On behalf of the TRACE study group Influence of gender on short- and long- term mortality after acute myocardial infarction.

Am J Cardiol 1996;77:1052.

Kokkinos PF, Holland JC, Pittaras AE, Narayan P, Dotson CO, Papademetriou V.

Cardiorespiratory fitness and coronary heart disease risk factor association in women.

Cardiovasc Rev Rep. 1995;11:49.

Kornowski R, Lansky AJ, Mintz GS, et al.

Comparison of men versus women in cross-sectional area, luminal narrowing, quantity of plaque, presence of calcium in plaque, and lumen location in coronary arteries by intravascular ultrasound in patients with stable angina pectoris.

Am J Cardiol. 1997;79:1601.

Kritz H, Schmidt P, Sinzinger H.

Passive smoking and cardiovascular risk.

Arch Intern Med. 1995;155:1942.

Krumholz HM, Douglas PS, Lauer MS, et al:

Selection of patients for coronary angiography and coronary revascularization early after myocardial infarction: Is there evidence for a gender bias?

Ann Intern Med 1992;116:785.

Kwok Y, Kim C, Grady D, et al.

Meta-analysis of exercise testing to detect coronary artery disease in women.

Am J Cardiol. 1999;83:660.

Kyriakidis J, Petropoulakis P, Androulakis A, et al.

Sex differences in anatomy of coronary artery disease .

J Clin Epidemiol 1995 ;48: 723.

Leaf DA, Sanmarco ME, Bahl RA.

Gender differences in coronary angiographic findings from 1972 through 1981 in Los Angeles, California.

Angiology 1990 ;41:609.

Lemaitre RN, Heckbert SR, Psaty BM, Siscovick DS.
Leisure-time physical activity and the risk of nonfatal myocardial
infarction in postmenopausal women.
Arch Intern Med. 1995;155:2302.

Linden W, Stossel C, Mourice J.
Psychosocial interventions for patients with coronary artery disease.
A meta-analysis.
Arch Intern Med. 1996;156:745.

MacDonald S, Joffres M, Stachenko S.
Multiple cardiovascular risk factors in Canadian adults.
Can Med Assoc J. 1992;146: 2021.

Malacrida R, Genoni M, Maggioni AP, et al.
A comparison of the early outcome of acute myocardial infarction in
women and men.
N Engl J Med. 1998;338:8.

Manson JE, Colditz GA, Stampfer MJ, et al.
A prospective of obesity and risk of coronary heart disease in women.
N Engl J Med. 1990; 322:882.

Manson JE, Stampfer MJ, Willet WC, et al.
Physical activity and incidence of coronary heart disease and stroke
in women.
Circulation. 1995;91;927.

Manson JE, Willet WC, Stampfer MJ, et al.
Body weight and mortality among women.
N Engl J Med. 1995;333:676.

Mark DB, Shaw LK, DeLong ER, et al:
Absence of sex bias in the referral of patients for cardiac
catheterization.
N Engl J Med 1994;330:1101.

Maynard C, Litwin PE, Martin JS, Weaver WD.
Gender differences in the treatment and outcome of acute myocardial
infarction: results from the Myocardial Infarction Triage and
Intervention Registry.
Arch Intern Med. 1992;152:972.

Maynard C, Weaver WD.
Treatment of women with acute MI: New findings from the MITI
registry.
J Myocardial Ischemia. 1992;4:27.

Mehta JL, Chen LY, Kone BC, Mehta P, Turner P.
Identification of constitutive and inducible forms of nitric oxide
synthase in human platelets.
J Lab Clin Med. 1995;25:370.

Meilahn EN, Becker RC, Corrao JM.
Primary prevention of coronary heart disease in women.
Cardiology. 1995;86:286.

Merkatz R, Temple R, Sobel S, Feiden K, Kessler D.
Women in clinical trials of new drugs.
N Engl J Med. 1993 ; 329: 292.

Miller VT.
Lipids, lipoproteins, women and cardiovascular disease.
Atherosclerosis. 1994;108:S73.

Milner KA, Funk M, Richards S, et al.
Gender differences in symptom presentation associated with coronary heart disease.
Am J Cardiol. 1999;341:217.

Moller J, Hallqvist J, Diderichsen F, Theorell T, Reuterwall C, Ahlbom A.
Do episodes of anger trigger myocardial infarction ? A case-cross over analysis in the Stockholm Heart Epidemiology program (SHEEP).
Psychosom Med. 1999;61:842.

Morise AP, Dalal JN, Duval RD.
Value of a simple measure of estrogen status for improving the diagnosis of coronary artery disease in women.
Am J Med. 1993; 94:491.

Morise AP, Dalal JN, Duval RD.
Value of a simple measure of estrogen status for improving the diagnosis of coronary artery disease in women .
N Engl J Med 1995;308:883.

Morise AP, Diamond GA, Detrano R, Bobbio M.
Incremental value of exercise electrocardiography and thallium- 201 testing in men and women for the presence and extent of coronary artery disease.
Am Heart J 1995;130:267.

Mosca L, Grundy S, Judelson D, et al.
Guide to preventive cardiology in women.
J Am Coll Cardiol. 1999;33:1751-1755.

Murabito JM, Evans JC, Larson MG, Levy D.
An investigation of differences in outcome between the sexes according to initial coronary disease presentation.
Circulation 1993;88:2548.

Nolan RP.
The threat of stress to the heart health of Canadians: results of the Foundation Health Survey.
Can J Cardiol. 2000;16:137.

Nolan RP.
Women ,Stress, and Cardiovascular Disease., The First International Conference on Women, Heart Disease and Stroke: Science and Policy in Action. Conference summary 2000.

Okin PM, Kligfield P.
Gender-specific criteria and performance of the exercise electrocardiogram.
Circulation 1995;92:1209.

Okin PM, Roman MJ, Devereux RB, Kligfield P.
 Gender differences and the electrocardiogram in left ventricular
 hypertrophy.
Hypertension. 1995;25:242.

Onaka H, Hirota Y, Shimada S, Kita Y, Sakai Y,
Kawakami Y, et al.

Clinical observation of spontaneous anginal attacks and multivessel
 spasm in variant angina pectoris with normal coronaries : Evaluation
 by 24-hour 12 lead electrocardiography with computer analysis.

J Am Coll Cardiol 1996;27:38.

Orth-Gomer K, Mittleman MA, Schenck-Gustafsson K,
et al.
 Lipoprotein-a as a determinant of coronary heart disease in young
 women. *Circulation*. 1997;95:329.

Ozaki Y, Keane D, Serruys P.

Fluctuation of spastic location in patients with vasospastic angina : A
 quantitative angiographic study.

J Am Coll Cardiol 1995;26:1606.

Palca J.
 Women left out at NIH.
Science. 1990 ; 248: 1601

Pepine CJ, El-Tamimi H, Lambert CR:

Prinzmetal's angina (variant angina).

Heart Dis Stroke 1992; 1:281.

Peto R, Lopez AD, Boreham J, Thun M, Heath C Jr, Doll R.
Mortality from smoking worldwide.

Br Med Bull. 1996;52:12-21.

Pinto RJ, Bhagwat AR, Loya YS, Sharma S.

Coronary artery disease in premenopausal Indian women: risk factors and angiographic profile.

Indian Heart J 1992 ;44:99.

Pramparo P.

Diagnosis, Treatment, and Access to Care., The First International Conference on Women, Heart Disease and Stroke: Science and Policy in Action.

Conference summary 2000.

Rabinowitz B.

Coronary heart disease in women.

Cardiol Rev 1997, 5:171.

Rexrode KM, Carey VJ, Hennekens CH, et al.

Abdominal Adiposity and Coronary Heart Disease in Women.

JAMA. 1998;280:1843.

Rich-Edwards JW, Manson JE, Hennekens CH, Buring JE.
The primary prevention of coronary heart disease in women.

N Engl J Med 1995;332:1758.

Robinson S.

Coronary heart disease in women.

Urban Cardiol. 1995; 17:23.

Roeters van Lennep JE, Zwinderman AH, Roeters van Lennep HW, Westerveld HE, Plokker HW, Voors AA, Bruschke AV, van der Wall EE.

Gender differences in diagnosis and treatment of coronary artery disease from 1981 to 1997. No evidence for the Yentl syndrome.

Eur Heart J 2000 ;21:911.

Roger VL, Jacobsen SJ, Pellikka PA, et al.

Gender differences in use of stress testing and coronary heart disease mortality: a population –based study in Olmsted County, Minnesota.

J Am Coll Cardiol. 1998;32:345.

Roger VL, Jacobsen SJ, Pellikka PA, et al.

Prognostic value of treadmill exercise testing: a population –based study in Olmsted County, Minnesota.

J Am Coll Cardiol. 1998;98:2836.

Romm PA, Green CE, Reagan K, Rackley CE.

Relation of serum lipoprotein cholesterol levels to presence and severity of angiographic coronary artery disease.

Am J Cardiol 1991 ;67:479.

Rosenberg L, Palmer JR, Shapiro S.

Decline in the risk of myocardial infarction among women who stop smoking.

N Engl J Med. 1990;322:213.

Ross J Jr, Brandenburg RO, Dinsmore RE, et al:
Guidelines for coronary angiography: A report of the American
College of Cardiology/American Heart Association Task Force on
assessment of diagnostic and therapeutic cardiovascular procedures:
Subcommittee on coronary angiography.

J Am Coll Cardiol 1987;10:935.

**Saito, T.; Date, H.; Taniguchi, I.; Nakamura, S.; Oka, H.;
Mizuno, Y.; Noda, K.; Yamashita, S.; Oshima, S. and
Yasue, H.**

Angiographic evaluation of culprit lesions in acute coronary syndrome
: relation to the original site on previous coronary angiography
(Abstract).

Jpn. Circ. J. 1998; 62: 359.

Saltzberg S, Stroh JA, Frishman WH.

Isolated systolic hypertension in the elderly: pathophysiology and
treatment.

Med Clin North Am. 1988;72:523.

Schrott H, Bittner B, Vittinghoff E, Herriington D, Hulley S,
For the HERS Research Group. Adherence to National Cholesterol
Education Program treatment goals in postmenopausal women with
heart disease.

JAMA. 1997;227:1281.

Secknus MA, Niedermaier On, Lauer MS, Marwick TH.
Diagnostic and prognostic implications of left ventricular cavity
obliteration response to dobutamine echocardiography.

Am J Cardiol 1998;81:1318.

Secknus MA, Marwick TH.

Influence of gender on physiologic response and accuracy of
dobutamine echocardiography.

Am J Cardiol 1997;80:721.

Sempos C, Cleeman J, Carroll M, et al.

Prevalence of high blood cholesterol among US adults: an update
based on guidelines from the Second Report of the National
Cholesterol Education Program Adult Treatment Panel.

JAMA. 1993; 269:3009.

Sheldon WC:

Indications for coronary arteriography.

Heart Dis Stroke 1993; 2:192.

Sowers JR.

Diabetes Mellitus and Cardiovascular Disease in Women.

Arch Intren Med. 1998;158:617.

Sowers JR.

Insulin and insulin-like growth factor in normal and pathological
cardiovascular physiology.

Hypertension. 1997;29:691.

Stone NJ and Kushner R

Effects of Dietary Modification and Treatment of Obesity :
Emphasis on Improving Vascular Outcomes.

Medical Clinics of North America 2000;84:111.

Sullivan AK, Holdright DR, Wright CA, et al.
Chest pain in women: Clinical, investigative, and prognostic features.
BMJ. 1994;308:883.

Taillefer R, DePuey EG, Udelson JE, et al.
Comparative diagnostic accuracy of Tl-201 and Tc-99m sestamibi
SPECT imaging [perfusion and ECG gated SPECT] in detecting
coronary artery disease in women.
J Am Coll Cardiol 1997;29:69.

Thomas JL; Braus PA.
Coronary artery disease in women, a historical perspective.
Arch Intern Med. 1998; 158: 333.

**Tunstall-Pedoe H, Kuulasmaa K, Amouyel P, Arveiler D,
Rajakangas A.M and Pajak A.**
Myocardial infarction and coronary deaths In the World Health
Organization MONICA Project.
Circulation 1994, 90:583.

Tunstall-Pedoe H, Kuulasmaa K, Mahonen M et al.
For the WHO MONICA [Monitoring Trends and Determinants in
Cardiovascular Disease] Project. Contributions of trends in survival
and coronary-event rates to changes in coronary heart disease
mortality: 10-year results from 37 WHO MONICA project populations.
Lancet 1999;353:1547.

Vaccarino V., Horwitz RI, Meehan TP, et al.
Sex Differences In Mortality After Myocardial Infarction : Evidence for
a Sex-Age Interaction.
Arch Intern Med. 1998;158:2054.

Vaccarino V, Krumholz Hm, Berkman LF, Horwitz RI.
Sex differences in mortality after myocardial infarction.
Circulation 1995;91:1861.

Vaccarino V, Parsons L, Every NR, et al.
Sex-based differences in early mortality after myocardial infarction.
N Engl J Med. 1999;341:217.

Vacek JL, Rosamond TL, Kramer PH, Crouse LJ, Porter CB, Robuck OW, White JL, Beauchamp GD.
Sex-related differences in patients undergoing direct angioplasty for acute myocardial infarction.
Am Heart J. 1994;126:521.

Valsania P.; Valsania P.; Zorio H.S.; Kowalschuk G.J. et al.
Severity of coronary artery disease in young patient with insulin dependent diabetes mellitus.
Am. H. J. 1991; 122 : 695.

Vigorito C, Russo P, Poto S, Genovese A, De Caprio L, Trimarco B, Condorelli M.
Coronary artery disease (CAD) in females: Coronary arteriographic findings in 34 women and comparison with 184 males with CAD.
G Ital Cardiol 1981;11:1228.

Walsh BW, Schiff I, Rosner B, Grienberg L, Ravinkar V, Sacks F.
Effects of postmenopausal estrogen replacement on the concentrations and metabolism of plasma lipoproteins.
N Engl J Med. 1991;325:1196.

Walsh JME, Grady D.

The treatment of hyperlipidemia in women.

JAMA. 1995; 274:1152.

Wang XL, Tam C, McCredie RM, Wilcken DEL.

Determinants of severity of coronary artery disease in Australian men and women.

Circulation 1994;89:1974.

Waters DD, Halphen C, Theroux P, David PR, Mizgala HF.

Coronary artery disease in young women: clinical and angiographic features and correlation with risk factors.

Am J Cardiol 1978 Jul;42:41.

Weber T, Kirchgatterer A, Auer J, Wimmer L, Lang G, Mayr H, Maurer E, Punzengruber C, Eber B.

Clinical presentation and coronary angiographic results in unstable angina pectoris

Acta Med Austriaca 1999;26:12.

Welch CC, Proudfit WL and Sheldon WC.

Coronary angiographic findings in 1000 women under age 50.

Am J Cardiol 1975;35:211.

Wenger NK.

Coronary heart disease in older women: prevention, diagnosis, management, and prognosis.

Clin Geriatr. 1996;4:53.

Wenger NK.

Coronary heart disease in women: gender differences in diagnostic evaluation.

JAMA. 1994;49:181.

Wenger NK, Speroff L, Parkard B.

Cardiovascular health and disease in women.

New Engl J Med. 1993;329:247.

Wenger NK, Speroff L, Packard B, eds.

Proceedings of NHLBI Conference: Cardiovascular Health and Disease in Women. Greenwich, Conn:

LeJacq Communications Inc; 1993.

Weiss K, Katzenschlager R, Egert H, Cajike P, Sinzinger H. Risk factors of atherosclerosis: results of screening in Southern Austria. *Wien Klin Wochenschr.* 1991;103:566.

Wexler LF.

Studies of acute coronary syndromes in women, lessons for everyone.

N Engl J Med 1999;341:275.

Whelton PK, He J, Appel LJ.

Treatment and prevention of hypertension. In: Manson JE, Ridker PM, Gaziano JM, Hennekens CH, eds. Prevention of Myocardial Infarction.

New York, NY: Oxford University Press, 1996:154-171.

Willet W, Green A, Stampfer M, et al.

Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes.

N Engl J Med. 1987;317:1303.

Williams Jk, Adams MR, Klopfenstein HS.

Estrogen modulates responses of atherosclerotic coronary arteries.

Circulation. 1990;81:1680.

Wilson E. ,

Young Women and Tobacco: Hope for Tomorrow. The First International Conference on Women, Heart Disease and Stroke: Science and Policy in Action.

Conference summary, 2000.

Wong CC; Froelicher ES; Bacchetti P; Barron HV; Gee L; Selby JV; Lundstrom R; Swain B; Truman A

Influence of gender on cardiovascular mortality in acute myocardial infarction patients with high indication for coronary angiography.

Circulation 1997 ;96(9 Suppl):II-51.

Yarzebski J, Col N, Pagley P, Savangeau J, Gore J, Goldberg R.

Gender differences and factors associated with the receipt of thrombolytic therapy in patients with acute myocardial infarction: a community-wide perspective.

Am Heart J. 1996;131:43.

Zhang F, Ram JL, Standley PR, Sowers JR.
17B estradiol attenuates voltage-dependent Ca^{2+} currents in vascular smooth muscle.

Am J Physiol. 1994; 193: C975.

Zuanetti G, Latini R, Maggioni AP, et al.

Influence of diabetes on mortality in acute myocardial infarction : Data from the GISSI-2 study.

J Am Coll Cardiol 1993;22:1788.