

References

- **Alderman E.; and Stadius M. (1992):**
The angiographic defintion of the bypass angioplacty revascularization investigation.
Coron Artery Dis. 3: 1189-1207.

- **Antonio s.; susanna s.; and Mauro s. (1997) :**
Detection of left ventricular systolic and diastolic abnormalities in patients with coronary artery disease by color kinesis.
Clin. Cardiol. 20, 927 – 933 .

- **Antman E.M.; Giugliano R.P.; and Gibson C.M. (1999):**
Abciximab facilitates the rate and extent of thrombolysis results of the thrombolysis in myocardial infarction (TIMI) 14.
Circulation; 99: 2720-2732.

- **Ambrose J.A .; winters S.L .; and stern A . (1985) :**
Angiographic morphology and the pathogenesis of unstable angina pectoris.
J. Am. coll cardiol;5:609-616.

- **Assmann P.; slager C.; and Borden S. (1990) :**
Quantitative echocardiographic analysis of global and regional left ventricular function: A problem revisited.
J. Am. Soc. Echocardiogr;3: 478 – 487.

- **Aygen M.; and popp R.L . (1987) :**
Influence of the orientation of myocardial fibers on echocardiographic images. Am.
J. cardiol; 60 : 147 – 152.

- **Becher H.;Tiemann K.; and Schlosser,T. (1998):**
Improvemeut in endocardial border delineation using tissue harmonic imaging.
Echocardiography ; 15:511-517.

References

- **Bednanz J. ; Vignon Ph.; Mor-avi V.; Weinert L.; Kock R.; et al. (1998):**
Color kinesis: principles of operation and technical guidelines. Echocardiography.
Am. jrnal. of CV ultrasound and Allied tech. 15 : 1. 21 - 23.

- **Borow K. M.; Green L. H.; Grossman W.; and Braunwald E. (1982) :**
Left ventricular end - systolic stress - shortening and stress length relations in humans : Normal values and sensitivity to inotropic state.
Am. J. Cardiol; 50 : 1301 : 1308 .

- **Bourdillon P.V.D.; Brooderick T.M.; Sawada S.G.; Armstrong W.F.; Dillon J.C.; et al (1989) :**
Regional wall motion index for infarct and non infarct regions after reperfusion in acute myocardial infarction : Comparison with global wall motion index.
J. Am.Soc. Echocardiogr.;2 : 398.

- **Chappuis F .; widmann t .; and Guth , B . (1988) :**
Quantitative assessment of regional left ventricular ejection fraction by densitometric analysis of intravenous digital subtraction ventriculograms: correlation with myocardial systolic shortening in dogs.
Circulation; 77:457-477.

- Chen C.H.; Nevo E.; Fetics B.; et al., (1997):**
Comparison of continuous left ventricular volumes by transthoracic 2-D digital echo quantification with simultaneous conductance catheter measurements in patients with cardiac disease.
Am. J. Cardiol; 80: 756-761.

- **Chesebro J.H.; Knatterud G.; and Roberts R.(1987) :**
Thrombolysis In Myocardial Infarction (TIMI) Trial, phase I: A comparison between Intravenous tissue plasminogen activator and intravenous streptokinase.
Circulation 76:142-154.

References

- **Clarkson P.B.M.; Wheeldon N.M.; and Iim P.O. (1995):**
left atrial size and function : Assessment using echo cardiographic – automated boundary detection. Br. Heart J ; 74:664-670.

- **Cowley M.J .; Disciascio G .; and Rehr R.B. (1989) :**
Angiographic observations and clinical relevance of coronary thrombus in unstable angina pectoris.
Am. J. Cardiol;63:E-108-E113.

- **Dangas G.; Mehran R .; and wallensteins (1997) :**
Correlation of angiographic morphology and clinical presentation in unstable angina.
J. Am. coll. cardiol;29:519 .

- **Davies R.F.; Goldberg A.D.; and Forman S. (1997) :**
A symptomatic cardiac Ischemia Pilot (ACIP) study two – year follow – up : outcomes of patients randomized to initial strategies of medical therapy versus revascularization.
Circulation; 95;2037.

- **Davies S.W .; Marchant B .; and Lyons J.P. (1991) :**
Irregular coronary lesion morphology after thrombolysis predicts early clinical instability .
J. Am. coll. cardiol;18:669-674.

- **Detrano R .; MacIntyre W.J.; and salcedo E.E .(1985):**
Videodensitometric ejection fraction from intravenous digital subtraction left ventriculograms: correlation with conventional direct contrast and radionuclide ventriculography.
Radiology.; 155:19-23.

References

- Dodge J.T.; Brown B.G.; and Bolson E.L . (1992) :

Lumen diameter of normal human coronary arteries: Influence of age, sex, anatomic variation, and left ventricular hypertrophy or dilation.
Circulation ; 86:232.

- Donald s ; Bain ; and William Grossmann . (2000) :

Coronary angiography. In Grossman cardiac catheterization, angiography and intervention, edited by D.S Baim and W. Grossmann. Sixth edition. Lippincott williams & Wilkins. Philadelphia . Chap 11. pp 211 - 256.

- D'cruz I.A.; Ialimalani G.G.; Sambasivan V.; Cohen H.C.; and Glick G.(1979) :

The superiority of mitral E point ventricular septum separation to other echo cardiographic indicators of left ventricular performance .
Clin. Cardiol.; 2:140.

- Ellis S.; Alderman E.I .; and Gaink. (1989) :

Morphology of left anterior descending coronary territory lesions as a predictor of anterior myocardial infarction. A CASS registry study.
J. Am. coll. cardiol ;13:1481-1491.

- Erbel R.; Schweizer P.; Meyer J.; Grenner H.; Krebs W.; et al. (1980) :

Left ventricular volume and ejection fraction determination by cross. Sectional echo cardiography in patients with coronary artery disease: A prospective study.
Clin . cardiol .; 3:377.

- Feigenbaum H. (1994) :

Coronary artery disease : In Echocardiography, edited by Fei-genbaum (ed). 5th edetion. A waverly company. Philadelphia - chap 8. pp 450 - 452.

References

- Felner J.M. (1990) :

Echocardiography . in the : Heart, Hurst, edited by J.W.; 7th edition , Mc Graw – Hill book company, New York , chap . 116. pp 325 - 334.

- Fentzke R.C.; Korcarz C.; and Mor-Avi V. (1997):

Non invasive assessment of left ventricular function in mice with genetically induced dilated cardiomyopathy using high frame rate two- dimensional contrast echocardiography. Abstract. Circulation; 96 (suppl 1): I-26.

- Fujii j.; sawada H.; Aizawa T.; Kato K.; and Onoe M. (1984)

Computer analysis of cross-sectional echocardiogram for quantitative evaluation of Left ventricular asnergy in myocardial infarction.

Br. Heart. J., 51:139.

- Gibson C.M . (1999) :

Primary angioplasty, rescue angioplasty and new devices. In Hennekens CH (ed): clinical Trial in cardiovascular disease: A companion to Braunwald's Heart disease Philadelphia, WB Saunders, , pp 185 – 197.

- Gibson C.M .; and Safian R.D . (1992) :

Limitations of cineangiography; impact of new technologies for image processing and quantitation.

Trends cardio vasc med ;2:156.

- Gibson C.M .; Murphy S.A.; and Men own I.B.(1999) :

Determinants of coronary blood flow after thrombolytic administration. TIMI study Group Thrombolysis In Myocardial Infarction.

J. Am. of coll. cardiol 34 : 1404 – 1413,

References

- **Gibson C.M .; cannon C.P. ; and Murphy S.A. (2000) :**
Relation ship of TIMI myocardial perfusion grade to mortality after administration of thrombolytic drugs.
circulation 101 : 124- 130,.

- **Gibson C.M .; Murphy S.A .; and Rizzo J.M .(1999) :**
Relation ship between TIMI frame count and clinical outcomes after thrombolytic administration. Thrombolysis in myocardial Infarction (TIMI) study Group.
Circulation 94:1945 – 1950.

- **Haft J.I .; and AL-Zarka A.M. (1993) :**
Comparison of the natural history of irregular and smooth coronary lesions: Insights into the pathogenesis, progression, and prognosis of coronary atherosclerosis.
Am. Heart. J. 126: 551-561.

- **Hartmann T.; kolev N.; and Blaicher A. (1997):**
Validity of acoustic quantification colour kinesis for detection of left ventricular regional wall motion abnormalities: A transesophageal echocardiographic study.
British. Journal of Anaesthesia; 79: 482 - 487.

- **Heger J.J.; Weyman A.E.; and wann L.s. (1980):**
Cross-sectional echocardiographic analysis of the extent of left ventricular asynergy in acute myocardial infarction.
Circulation; 61:1113 - 1118 .

- **Homans D.C.; Asinger R.; and Elsperger J.(1985) :**
Regional function and perfusion at the lateral border of ischemic myocardium .
circulation 71 : 1038- 1047.

References

- **Isaaz K.; Bruntz J.F.; and Ethevenot G. (1994) :**
Abnormal coronary flow velocity pattern in patients with left ventricular hypertrophy, angina pectoris, and normal coronary arteries: A transesophageal Doppler echo cardiographic study.
Am. Heart. J. 128 : 500-510 .

- **Jiang L.;Morrisey R.; and HandSchumacher M.D. (1996) :**
Quantitative three – dimensional reconstruction of left ventricular volume with complete borders detected by acoustic quantification underestimates volume.
Am. Heart. J. 131: 553 – 559.

- **Kimball T. R.; witt S. A.; and Daniels S. R. (1996) :**
The role of automatic endocardial edge detection in the evaluation of left ventricular diastolic function in children.
J. Am. Soc. Echocardiogr ; 9 : 18 - 26.

- **Lang R.; Vignon P.; Weinert L.; and Bednarz J. (1996) :**
Echocardiographic quantification of regional Left ventricular wall motion with color kinesis.
Circulation.; May 15-93 (10): 1877 – 1885.

- **Lau Y.S.; puryear J.V.; and Gan S.C. (1997) :**
Assessment of left ventricular wall motion abnormalities with the use of color kinesis, a valuable visual and training aid.
J. Am. Soc. Echocardiogr., 6 (10) : 665-672.

- **Lenderink T.; Simoons M.L .; and VAN ESG A.(1995) :**
Benefit of thrombolytic therapy is sustained throughout five years and is related to TIMI perfusion grade 3 but not grade 2 flow at discharge.
Circulation 92: 1110-1116.

References

- **Mintz G.S .; popma J.J.; and pichard A.D .(1996) :**
Limitations of angiography in the assessment of plaque distribution in coronary artery disease a systematic study of target lesion eccentricity in 1446 lesion.
Circulation;39; 924

- **Mor-Avi V.; Gillesberg I.E.; and korcarz C.(1995):**
Improved quantification effect of signal averaging.
J. Am. Soc. Echocardiogr .;11:792-802.

- **Mor – Avi V.; and Godoy I.. (1999) :**
Color kinesis: new technique or just another display of acoustic quantification? Vol. 16 , No. 1. Echocardioghy.
Am. Jrnl. of CV ultrasound & Allied tech. 16 : 95 - 103.

- **Mor- Avi V.; kirk T.; and Spencer M.D. (1999):**
Acoustic quantification today and its future horizons. Vol 16, no. 1,. Echocardiography.
Am. Jrnl. of CV ultrasound & Allied tech. 16 : 85 - 94.

- **Mor – Avi V.; lin R.M .; and Bednarz J. (1998) :**
Quantitative three-dimensional echocardiographic evaluation of regional systolic and diastolic LV function. Abstract.
J. Am. coll. cardiol; 31:297 A.

- **Mor-Avi V.; Vignon P.; Roch R.; Weinert L.; Maria J.; et al. (1997) :**
Segmental analysis of color kinesis images. New method for quantitative assessment of left ventricular contraction and relaxation.
Circulation : 95 : 2082 - 2097.

- **Mor- Avi V.; Vignon P.; and Bales A.S. (1998) :**
Acoustic quantification indexes of left ventricular size and function : Effects of signal averaging.
J. Am. Soc. Echocardiogr .; 11: 792 – 802.

Rferences

- Mottley J. G.; and Miller J.G. (1988):

Anisotropy of the ultrasonic backscatter of myocardial tissues. 1 : theory and measurements in vitro.

J. Acoust. Soc. Am; 766 – 771.

- Mulvagh S.I.; Foley D.A.; and Aeschbacher B.C. (1996):

Second harmonic imaging of an intravenously administered echo cardiographic contrast agent.

J.Am. coll. cardiol ;27: 1519-1525.

- Nidorf S.M.; Siu S.C.; and Galambos G. (1993) :

Benefit of late coronary reperfusion on ventricular morphology and function after myocardial infarction.

J. Am. coll. cardiol 21 : 683- 691 .

- Pandian N.G.; Kieso R.A.; and Kerber R.E. (1982) :

Two dimensional echo cardiography in experimental coronary stenosis. Relationship between systolic wall thinning and regional myocardial perfusion in sever coronary stenosis.

Circulation . 66:603.

- Parisi A.F.; Moynihan P.F.; and Folland E.D. (1981):

Quantitative detection of regional left ventricular contraction abnormalities by two dimensional echocardiography. II. Accuracy in coronary artery disease.

Circulation; 63 : 761 – 767.

- Perez J.E .; Woggoner A.D.; and Barzilai B. (1992):

On line assessment of ventricular function by automatic boundary detection and ultrasonic backscatter imaging.

J. Am. coll. cardiol,19: 313- 320.

Rferences

- **Popp R.L.; Agatston A.; and Armstrong W.F. (1998):**
Recommendations for training in performance and interpretation of stress echocardiography.
J. Am. Soc. Echocardiogr.; 11: 95 – 96.

- **Popp R.L. (1995) :**
"High time" for noninvasive assessment of regional ventricular diastolic ischemic dysfunction.
Circulation; 91 : 552 – 553.

- **Randall P.A . (1978) :**
Mach bands in cinecoronary arteriograph.
Radiology; 129:65.

- **Ryan T.J .; Anderson J.L .; and Antman E.M . (2000):**
ACC/AHA guidelines for the management of patients with acute myocardial infarction: Executive summary.
Circulation; 94:2341.

- **Ryan T.; Petrovic O.; Armstrong W.F.; and Feigenbaum H. (1986):**
Quantitative two-dimensional echocardiographic assessment of patients undergoing left ventricular aneurysmectomy .
Am. Heart. J., 111: 714.

- **Scanlon P.J.; Faxon D.P.; and Audet A. (1999) :**
AHA/ACC guidelines on coronary angiography; a report of the ACC/AHA Task force on practice guidelines.
J. Am. coll. cardiol :33: 1756.

- **Schwartz S.L.; Gao Q.L.; and Vannan M.A. (1996):**
Automatic backscatter analysis of regional left ventricular systolic function using color kinesis.
Am. J. cardiol; 77: 1345 -1350 .

References

- **Selien M.A.; Mc Williams E.T.; and Palileo M. (1996):**
Beat - to beat variability of left ventricular indexes measured by acoustic quantification : Influence of heart rate and respiration. Correlation with M-mode echocardiography.
J. Am. SOc. Echocardiogr; 9 : 221 - 230.

- **Sklenar. J.; Jayaweera A.R.; and kaul s. (1992) :**
A computer - aided approach for the quantitation of regional left ventricular function using two-dimensional echocardiography.
J. Am. Soc. Echcardiogr.; 5 : 33 - 40.

- **Spencer K.T.; Bednarz J.; and Rafter P.G . (1998) :**
Use of harmonic imaging without echocardiographic contrast to improve two-dimensional image quality.
Am. J. cardiol. 82 : 794 - 799.

- **Stadius M.L .; and Alderman E.L . (1990) :**
Coronary artery revascularization: critical need for and consequences of objective angiographic assessment of lesion severity.
Circulation;82;2231.

- **Starling M. R.; Walash R. A.; and Dell Italia L. J. (1987) :**
The relations ship of various measures of end - systole to left ventricular maximum time - varying elastance in man.
Circulation ; 76 : 32 - 43

- **Stefan G.; and bing R.J. (1972) :**
Echo cardiographic findings in experimental myocardial infarction of the posterior left ventricular wall.
Am. J. cardiol.; 30:629.

- **Stewart W.J.; Aurigemma G.P .; and Bierman F.Z . (1995):**
Task , force 4: Training in echo cardiology.
J. Am. coll. cardiol 25 : 16 – 19

References

- **Stewart W.; Rodkey S.M.; and Gunawarden A.S. (1993):**
left ventricular volume calculation with integrated backscatter from Echoardiography .
J. Am. Soc. Echocardiogr, 6: 553 - 563.
- **Stoddard M.F.; Pearson A.C .; and Kern M.J. (1989):**
Left ventricular diastolic function : Comparison of pulsed Doppler echo cardio graphic and hemo dynamic indexes in subjects with and without coronary artery disease .
J. Am. coll. cardial 13:327 – 336.
- **Swensen J.D.; Harkin C.; and Pace N.L. (1996):**
Transesophageal echo cardiography : An objective tool in defining maximum ventricular response to intravenous fluid therapy .
Anesth. Analg ; 38:1149-1153.
- **The TIMI III B Investigators . (1994) :**
Effects of tissue plasminogen activator and a comparison of early invasive and conservative strategies in unstable angina and non Q wave myocardial infarction: Results of the TIMI III B trial (Thrombolysis in myocardial Ischemia).
Circulation, 89: 1545.
- **TIMI study Group . (1985) :**
The thrombolysis In Myocardial Infarction (TIMI) Trial : phase I finding .
N. Eng. J. Med; 312:932–936.
- **Torry R.J.; Myers J.H.; Adler A.L.; liut C.L.; and Gallagher K.P. (1991) :**
Effects of nontransmural ischemia on inner and outer wall thickening in the canine left ventricle.
Am. Heart. J., 122 : 1292.

References

- Uren N.G . (1994) :

Relation between myocardial blood flow and the severity of coronary artery stenosis.
N. Engl. J. Med; 330:1782.

- Vermes E.; Guyon P.H.; Weingrod M.; et al., (2000):

Assessment of left ventricular regional wall motion by color kinesis technique.
Echocardiography: A Jrnl. of CV ultrasound & allied tech; 17.6: 521-527.

- Vignon P. ; mor-Avi V.; weinert L.; and Koch R. (1997) :

Quantitative evaluation of global and regional left ventricular diastolic function with color kinesis.
Circulation ; 11 : 1053 - 1061 .

- Vitarelli A.; Penco M.; and Ferro-luzzi M. (1996):

Assessment of diastolic left ventricular filling by echocardiographic automated border detection and comparison with radionuclide ventriculography.
J. Am. Soc. Echocardiography : 9:35 –146 .

- Vitarelli A.; Sciomer S.; and Schina M. (1997) :

Detection of left ventricular systolic and diastolic abnormalities in patients with coronary artery disease by color kinesis.
Clin. Cardiol.; 20 (11) : 927 – 933.

- Waggoner A.D.; Miller J.G.; and Perez J.E. (1994):

Two-dimensional echocardiographic automatic boundary detection for evaluation of left ventricular function in unselected adult patients.
J.Am. Soc. Echocardiogr; 7 : 459 – 464.