

REFERENCES

- Abou - Mourad , N.N ; Farah , F.S and Steel , D. (1979) :
Dermato pathic changes in hypozincemia . Arch -
Dermatol ; 115 : 956 .
- Ackerman , A.B. and Kligman , A.M. (1969) : Some
observations on dandruff . J Soc Cosmetic Chemists
; 20 : 81 .
- Agache, P. ; Blanc , D. Barrant , C. and Laurent , R . (1980)
:Sebum levels during the first year of life Br . J
Dermatol; 103 : 643 .
- AL - Doory . Y . (1980) : Text book of " Laboratory Medical
Mycology " . Philadelphia, Lea and Febiger .
- Aleen , H.B and Rippon , J.W (1985) : Seborrhoeic dermatitis
in the Textbook of Dermatology by Moschella S.L
and Hurely . H.J vol 2 2nd ed Saunders
Co.Philadelphia . London - Toronto p : 739 .
- Alexander ,S. (1967) . Loss of hair and dandruff . Br . J Derm-
tol, 79 : 549.

- Anthony, N. Domonkos, M.D,FACP(1971):
Seborrhoeic dermatitis and psoriasis . In : Textbook
of Clinical Dermatology . vol II 2 nd ed . Saunder's
Co . Philadelphia - London - Toronto p:204 .

- Arnold,H.L, Odom, R.B and James .W.D (Editors)(1990) :
Andrew's disease of the skin 8th ed. Saunder's Co.
Philadelphia - London - Toronto pp: 194 and 347 .

- Back O, Faergemann J and Hornqrist R(1985) :
Pityrosporum folliculitis : A common disease of the
young and middle - aged . JAm . Acad.Dermatol ;12
: 56 .

- Baker , H. (1972) : Drug reaction, in Rook , A. Wilkinson ,
D.S and Ebling , F.J.G(ed.) Text book of
Dermatology vol . I 2 nd ed . Oxford , England .
Blackwell scientific publications p . 1025 .

- Barba , A. Piubellow , Vantinl I . et al ; (1982) : Skin lesions
in chronic alcholic pancreatitis . Dermatologica 164 :
322 .

- Barnes , W .G, Sauer , G. C and Arnold . J . D (1973):
Scanning electron microscopy of *Tinea versicolor* organisms . Arch . Derm , 107 : 392 .
- Benham , R.W (1939) : The cultural characteristics of *pityrosporum ovale* , lipophilic fungus . Invest Dermatol - 2 : 187 .
- Bergbrant ,I.M , and Faergemann ,J.(1988) : Variations of *pityrosporum orbiculare* in middle - aged and elderly individuals . Acta Derm . Venereol (Stockh) 68 : 537 .
- Bergbrant I.M and Faergemann J (1989) :
Seborrhoeic dermatitis and *pityrosporum ovale* .
cultural and Immunological study . Acta . Derm .
Venereol . (Stockh) 69 : 332 .
- Bergbrant ,I.M. and Faergemann,J. (1990) : The role of *pityrosporum ovale* in seborrhoeic dermatitis . Semin - Dermatol . Dec ; 9 (4) ; P 262 .
- Bergbrant, I.M and Faergemann, J . (1991) ; Seborrhoeic dermatitis and *pityrosporum ovale* . A cultural and Immunological study . Acta Derm Venerol (Stockh) .

Berger R.S ,Stoner M.F , Hobbs E.R ; et al . (1988) : Cutaneous manifestations of early human immuno defeciency virus exposure .J Am . Acad . Dermatol , 19 : 298 .

Binder, R.L. and Jonelis, F.J (1983) : Seborrhoeic dermatitis in neuroleptic induced parkinsonism , Arch Dermatol 119 : 473 .

Boyum (1968) : Ficoll / Hypaque method for separation of mononuclear cells and granulocytes from human blood - Scand J ; Clin . Lab . Invest (suppl) 121 : 77 .

Brauer, E.W (1969) : Antiseborrhoeic qualities of zinc pyrithione in a cream vehicle . III clinical experience in pityriasis capitis (Dandruff , seborrhoeic dermatitis of scalp) 5 : 723 .

Brauner , G.J (1979) : Seborrhoeic dermatitis in Fitzpatrick , T.B ; Eisen , A.Z ; Wolf K ; Freedberg I.M ; Frank Austen K. (eds) Dermatology in general medicine , Textbook and Atlas 2 nd ed M.C Graw - Hill Book company New York , st . Louis p . 803 .

Broberg, A. and Faergemann, J. (1989) : Infantile seborrhoeic dermatitis and pityrosporum ovale. Br . J Dermatol 120 : 359 .

Bruneau, S.M. and Guinet, R.M.F (1983) : Quantitative immunoelectrophoretic study of genus pityrosporum. Sabouraud , Mykosen ; 27:123 .

Burton , J.L , Cartlidge, M. and Shuster, S. (1973) : Effect of L - dopa on the seborrhoea of parkinsonism. Br . J Dermatol 88 : 475 .

Burton , J.L . and Pye , R. J. (1983) : Seborrhoea is not a feature of seborrhoeic dermatitis . Br . Med . J ; 286 : 1169 .

Butterfield , W . Roberts, M . M and Dave, V . K (1987) : Sensitivities of pityrosporum species to selected commercial shampoos . Br J Dermatol ; 116 : 233 .

Byung, H . Park (1981) : Deficiencies in cell mediated immunity . In Text Book of Principles of Immunological diagnosis in medicine. chapter 51 p . 303 .

Caprilli, F . Mercantini, R . Nazarro-porro M , et al . (1973) :
Studies of the genus pityrosporum in submerged
culture . Mycopathol Mycol Appl ; 51 : 171 .

Carr,M. M . Pryee, D .M and Ive, F. A (1987) : Treatement of
seborrhoeic dermatitis with ketoconazole .vol I .
Responce of seborrhoeic dermatitis of the scalp to
topical ketoconazol . Br J Dermatol , 116 : 213 .

Castellani ,A. and Chalmers ,A . J (1913) : Manual of Tropical
Medicine 2nd ed , 1747 pp . Wm Wood and Co , New
York (cited by Gordon , 1951) .

Castellani, A. (1925) : Notes on three new yeast - like organisms
and a new bacillus . Jour . Trop Med ; 28 : 217 . (
cited by Gordon 1951) .

Clift ,D.C, Dodd, H . J , Kirby ,J.D.T , et al ; (1988) :
Seborrhoeic dermatitis and malignancy . Acta Derm
Venereol (Stockh) ; 68 : 48 .

Conant, N.F (1971) : Manual of Clinical Mycology ,W.B
Saunders . Philadelphia , p 595 .

Cowley, N. C , Farr, P. M and Shuster ,S.et al (1990) : The permissive effect of sebum in seborrhoeic dermatitis : an explanation of the rash in neurological disorders . Br J Dermatol ; 122 : 71 .

Cunningham, A.C , Leeming, J . P , Ingham et al ; (1990) : Differentiation of three serovars of *Malassezia furfur*. J Appl Bact ; 68 : 439 .

Domonkos,A.N. (1971) : Andrew's disease of the skin . Clinical Dermatology ed b . Philadelphia ,W.B Saunders , Co ; p 204 .

Eisenstat, B.A , Wormser,G.P (1984) : Seborrhoeic dermatitis and butterfly rash in AIDS , New Engl J Med 311 : 189 .

Emmons, C.W. (1940) : The isolation and pathogenicity of *pityrosporum ovale* . Publ . Health Rep . 55 ; 1306 .

Faergemann, J. and Bernander, S. (1979) : Tinea versicolor and *pityrosporum orbiculare*:A mycological investigation Sabouraudia ; 17:171 .

Faergemann ,J .and Fredriksson ,T.(1979) : Tinea versicolor with regard to seborrhoeic dermatitis . Arch Dermatol ; 115:966 .

Faergemann, J. and Fredriksson ,T.(1980) : Age incidence of pityrosporum orbiculare on human skin . Acta . Derm Venereol (Stockh) 60:531 .

Faergemann ,J.and Fredriksson,T.(1981) : Experimental infections in rabbits and humans with pityrosporum orbiculare and pityrosporum ovale . J Invest Dermatol ; 77:314 .

Faergemann,J. Tejernlund,U. Scheynius, A.et al; (1982) : Antigenic similarities and differences in genus pityrosporum J invest Dermatol ; 78:28 .

Faergemann ,J. (1983) : Antibodies to pityrosporum orbiculare in patients with Tinea versicolor and controls of various ages , J Invest Dermatol ; 80:133 .

Faergemann, J .(1984) : Quantitative culture of pityrosporum orbiculare . Int . J Dermatol ; 23:323 .

Faergemann ,J .(1986) : Seborrhoeic dermatitis and pityrosporum orbiculare : Treatement of seborrhoeic dermatitis of the scalp with miconazol-Hydrocortisone (Daktacort) , miconazol and Hydro - cortisone . Br J Dermatol ; 114 - 695 .

Faergemann,J . Johansson ,S. Bak, O. et al ; (1986) : An immunologic and cultural study of pityrosporum folliculitis . J Am Acad Dermatol ; 14:429 .

Faergemann, J.and Larko,O.(1987):The effect of U.V light on human skin microorganisms.Acta Derm Venereol (Stockh); 67:69.

Faergemann ,J. (1989) : A new method for growth and filament production of pityrosporum ovale (orbiculare) on human stratum corneum in vitro . J Invest . Dermatol ; 92:117 .

Faergemann ,J . (1990) : Treatement of seborrhoeic dermatitis of the scalp with ketoconazol shampoo . Acta Derm Venereol (Stockh); 70:171 .

Fishman ,H.C (1977) : Seborrhoeic dermatitis practical therapy .
Cutis . vol 20,724 .

Ford , G.P . Farr,P.M . Ive,F.A and Shuster,S.(1984) : The
response of seborrhoeic dermatitis to ketoconazol .
Br.J . Derm ; 111:603 .

Gloor, M. Wiegand,I. Friedrich ,H.C . (1972) : Uebermenge and
Jusammenst zung der Hautoberf la chenlipide beim
sogenannten seborrhois chen ckzem - Der Mschr 158:
759.

Goodman, D.S.Teplitz ,E.D . Wishner ,A. et al (1987) :
Prevelance of cutaneous disease in patients with
acquired immuno - deficiency syndrom (AIDS) or
AIDS releated complex . J Am Acad Dermatol 17:210 .

Gordon ,M.A(1951) : Lipophilic yeast like organisms associated
with Tinea versicolor. J. Invest . Derm , 17:267.

Gordon, M.A (1951) : The lipophilic mycoflora of the skin .
Mycologica ; 43:524 .

Goslen , J.B. and Kobayashi,G.S (1987) : Mycological infections
In : Dermatology in general medicine , by Fitzpatrick
T.B.Eisen.A.Z, Wolf . K. Freedberg .I.M and Austen
, KF. 3 rd ed Mc . Grow - Hill book company . New
York . st . Louis . Toronto . vol 3 ch . 179,p:2139 .

Gosse,R.M. Vanderwyk,R.W.(1969) : The relationship of a
nystatin resistant strain of pityrosporum oval to
dandruff . J Soc Cosmet ; 20 : 603 .

Green ,C.A , Far, P. M and Shuster,S.(1987) : Treatement of
seborrhoeic dermatitis with ketoconazol . II Response
of seborrhoeic dermatitis of the face , scalp and trunk
to topical ketoconazol . Br J Dermatol ; 116 : 217 .

Groisser,D. Bottone, E.J .Lebwohl,M.et al (1989) : Association of
pityrosporum orbiculare (Malassezia furfur) with
seborrhoeic dermatitis in patients with acquired
immuno - deficiency disease (AIDS) . J Am Acad .
Dermatal 20 , 770 .

Gueho ,E .Sinmons, R.B .Pruitt ,W.R , et al (1987) : Association
of Malassezia pachydermatis with systemic infections
of humans . J Clin Microbiol , 25 , 1789 .

Gueho, E. Meyer, S. (1989) : A reevaluation of the genus *Malassezia* by means of genome comparison . *Antonie Van Leeuwenhoek* ; 55:245.

Heng, M. C. Henderson, C. L. Barker, D. C. , et al (1990) : Correlation of *Pityrosporum ovale* density with the clinical severity of seborrhoeic dermatitis as assessed by a simplified technique . *J Am Acad. Dermatol* ; 23:82 .

Hodgson - Jones, I. S. Mackenna, R. M. B and Wheatley, V. R. (1953) : The surface skin fat in seborrhoeic dermatitis *Br J Dermatol*; 65:246 .

Honing, P. J and Smith , L. R (1979) : *Tinea capitis* masquerading as atopic or seborrhoeic dermatitis . *J. Pediatr* ; April . p 604.

Hudson, L. Frank , C. and Hay (1989) : In *Textbook of practical immunology* chapter 3 p:92 . Blackwell scientific publications - Oxford - London - Edinburgh - Boston - Melbourne .

Jacinto - Jamora ,S .Tanesis,J. and Katigbak,M.L , (1991) :
Pityrosporum folliculitis in the philppines : Diagnosis
, prevelance and management . J Am Acad Dermatol ;
24 : 693 .

Jawetz , E , Melnick , J . L . and Adelberg , E .A , et al (1989) :
Medical mycology In Text book of Medical
Microbiology 18th ed . Appleton and Lange California
p : 294 .

Johnson, M.L.T and Roberts,J.(1977): Prevelance of
dermatological diseases among persons 1 - 74 years of
age , United States publication No (P H S) 79 - 1660 ,
Washington D.C;U. S Departement of Health and
human services .

Jones, H .E ; Reinhart , J .H and Rinaldi , M .G (1974) :
Immunologic susceptibility to chronic
dermatophytosis . Arch Dermatol 110 : 213 .

Kanwar , A .J . Aboul Magid , Gary , M. P and Singh , G .(1981)
: Seborrhoeic dermatitis like eruption caused by
Cimetidine . Arch Dermatol , 117 : 65 .

- Katsambas, A .Antoniou, C.H , Frangouli ,E . et al (1989) : A double blind trial of treatment of seborrhoeic dermatitis with 2% ketoconazol cream compared with 1% Hydrocortison cream . Br . J Dermatol 121 : 353 .
- Keddie , F .M . and Barajas,I.(1972) : Quantitative ultra structural variations between pityrosporum ovale and pityrosporum orbiculare based on serial section electron microscopy . Int . J Dermatol ; 11 : 40 .
- Kieffer,M. Bergbrant,I. Faergemann,J.et al (1990) : Immune reactions to pityrosporum ovale in adult patients with atopic and seborrhoeic dermatitis . J Am . Acad Dermatol ; 22 : 739 .
- Kligman ,A.M .(1979) : Dandruff , its causes and treatment in Ortanos C , ed Haar and Haar kraukheiten . Stuttgart Fischer : 663 .
- Kligman,A.M. Leyden ,J.J (1983) : Seborrhoeic dermatitis . seminars in Dermatology , 2 : 57 .

Krestin, D. (1927) : The seborrhoeic facies as a manifestation of post encephalitic parkinsonism and allied disorders . Quarterly J Med ; 21 : 177 .

Kwon-Chung,K.J and Bennette ,J.E. (1992) : Infections caused by *Malassezia* species , In Text book of Medical Mycology . Lea and Febiger Philadelphia . London . chapter 8 p . 176 .

Langeron , M. (1987) : The family tree of fungi ; Quoted from the Textbook of mycology by Janssen research council . Belgium p . 30 .

Leeming ,J.P . Notman , F.H (1987) : Improved methods for isolation and enumeration of *Malassezia furfur* from human skin . J Clin Microbiol ; 25 : 2 017 .

Leeming, J.P. Notman, F. H and Holland, K .T (1989) : The distribution and etiology of *Malassezia furfur* and cutaneous bacteria on human skin .J Appl Bact .67 : 47 .

Lefkovits,I.and Pernis,B, (1979):In the Textbook of Immunological Methods . chapter 15 , 5 th ed

Academic Press New York- San Francisco- London .

Lever ,W.F and Schaumburg . Lever, G .(1990) : "
Histopathology of the skin " 7th ed Lippincot . J .B
company . Philadelphia London -
Toronto-Pennsylvania vol I . p . 110 , 218 and 364 .

Leyden ,J .J . Mc Ginley, K. J and Kligman ,A. M (1976): The
role of microorganisms in dandruff . Arch Dermatol ;
12 : 333 .

Mac Leod ,J.M.I.I and Dowling ,G .B (1928) : An experimental
study of Malassez: Its morphology , cultivation and
pathogenicity . Br J of Dermatol and syphilis XI . 139 .

Maietta ,G. Fornaro, P. Rongioletti ,F.et al (1990) : Patients with
mood depression have high prevelance of seborrhoeic
dermatitis . Acta Derm Venereol (Stockh) ; 70:432 .

Malassez, L .(1874) : Note sur le champignon du pityriasis
simplex Arch physiol . Norm . Pathol (Series 2) ; 1 :
451 (Cited by Gordon).

Marks,R.Pearse ,A.D . Walker, A.P (1985) : The effects of a shampoo containing zinc pyrithione on the control of dandruff . Br . J Dermatol ; 112 : 415 .

Mathes, B.M and Douglass ,M.C (1985) : Seborrhoeic dermatitis in patients with acquired immuno deficiency syndrom J Am Acad Dermatol 13 : 947 .

MC Ginley ,K.J . Lantis , L.R and Marples R .R (1970) : Microbiology of pityriasis versicolor .Arch Dermatol ; 102 : 168 .

MC Ginely ,K.J.Leyden J.J, Marples R .R, and Kligmann,A.M (1975) : Quantitative microbiology of the scalp in non dandruff , dandruff , and seborrhoeic dermatitis . J Invest Dermatol , 64 ;401 .

Midgley, G and Hay, R.J (1988) : Serological responses to pityrosporum (Malassezia) in seborrhoeic dermatitis demonstrated by ELISA and western blotting Bull. Soc Mycol . Med ; 17 : 267 .

- Midgley, G. (1989) : The diversity of pityrosporum (Malassezia) yeasts in vivo and in vitro Mycopathologica ; 106 : 143 .
- Monjardino, J .P . and Mac Gillivray ,A .j (1970) :RNA and histone metabolism in small lymphocytes stimulated by phytohaemagglutinin . Exp . Cell Res . 60 , 1 .
- Moore, M.Kile, R.L , Engman,M.et al (1926) : Pityrosporum ovale (bottle bacillus of Unna , spore of Malassez) . cultivation and possible role in seborrhoeic dermatitis . Arch Dermatol & Syph ; 33 : 457 .
- Moore,M.(1935) : Cultivation and study of pityrosporum ovale , the so called bottle bacillum of Unna . Arch Dermatol and Syph . 31 ; 661 .
- Moore ,G.S. and Jaciow , D.M. (Editors) (1979) : Mycology for the clinical laboratory . Reston Publishing Co . Virginia p:117 .
- Nazzaro - Porro, M . Passi ,S . Caprilli, F. et al (1976) : Growth requirements and lipid metabolism of pityrosporum ovale . J Invest . Dermatol ; 66 : 178 .

- Nazzaro - Porro ,M .Passi, S. Caprilli,F. et al (1977) : Induction of hyphae in cultures of pityrosporum by cholesterol and cholesterol esters . J Invest Dermatol ; 69 :531 .
- Noble,W.C and Midgley, G. (1978) : Scalp carriage of pityrosporum species : The effect of physiological maturity , sex and race .Sabouraudia ; 16 : 229 .
- Oberle ,A.D . Fowler, M .Grafton ,W.D (1981) : Isolate from the upper respiratory tract . Am Soc. Clin. Pathol ; 76 : 112 .
- Ota, M and Huang, P.T (1933) : Sur les champignons du genra pityrosporum sabauraud . Ann de parasitol ; II : 49 .
- Paker,H. and Wilkinson D .S (1979) : Psoriasis . In Rook A ; Wilkinson, D.S ; Ebling . F .J .G (eds) : Textbook of Dermatology vol II . 3rd ed Oxford England . Blackwell scientific publications p . 1326 .
- Panja ,G. (1927) : The Malassezia of the skin : cultivation , morphology and species . Trans Far Eastern Assoc . Trop . Med 1.422 .

- Parish , L.C . Witkowski , J .A.et al .(1974): Pityriasis, seborrhoeic dermatitis and dandruff : Evaluation of tar shampo cutis , 14:257 .
- Pierard ,J .and Dockx,P. (1972) : The ultra structure of *Tinea versicolor* and *Malassezia furfur* . Int J . Dermatol 11-116 Quoted from MC Daniel D.H and Welton W.A (1984) .
- Pierard - Franchimont C. Pierard ,G.E. Kligmam ,A. (1990) : Seasonal modulation on sebum excretion . Dermatologica ; 181:21 .
- Plewing,G.(1987) : Seborrhoeic dermatitis In the Text book of dermatology in general medicine Edited by Fitzpatrick T.B et al , 3 rd ed . M.C-Grow-Hill book company New york p978 .
- Provest,E.(1983) : The rise and fall of fluorescent *Tinea capitis* . Pediatr Dermatol 1,2:127.
- Pye ,R.J . Meyrich, G. Burton, G.L (1977) : Skin surface Lipids in seborrhoeic dermatitis .Br J Dermatol; 97(supple) 12 .

Ratzer, M.(1969) : The incidence of skin diseases in the west of Scotland . Br. J .Dermatol; 81:456 .

Richard,A.F. Clark and Terrence,T.and Hopkins(1992) : " The other eczemas " In the Text Book of Moschella, S.L .and Hurley ,H.J chapter 21 vol II 2nd ed Saunders Co Philadelphia London Toronto p:465 .

Rippon,J.H. (1982):The pathogenic fungi and the pathogenic Actinomycetes. In the Text book of Medical Mycology.2nd ed Philadelphia.W.B Saunders Company chapter 20p:59 .

Rivolta,S.(1873) : Parassiti vegetali 1st ed Torino Digiulo . Speirani F.Figglo; pp469(cited by Ive) .

Roberts, S.O.B(1969 a): Pityrosporum orbiculare incidence and distribution on clinically normal skin.Br.J. Derm : 81:264 .

Roberts, S.O.B and Lachapella,J.M (1969) : Confluent and reticulate papillomatosis (Gougerot - Carteaud) and pityrosporum orbiculare Br . J . Dermatol; 18:841 .

Roberts , S.O.B and Mackenzie, D.W.R (1979) ; Mycology in
Rook ,A .Wilkinson, D.S; Ebling , F.J.G (eds) : Text
book of Dermatology vol I 3rd ed .

Rocklin,R.A. (1981) : Invitro tests for cell-mediated immunity In
the Text book "Principles of Immunological diagnosis
in medicine " chapter 4 p:28 .

Roia ,F.C.Vander, W.Y.and Beal ,J.A(1963) : The human scalp
as a habitate for yeasts . J. Soc. Cosmetics Chemists ;
14:81.

Rook, A. and Wilkinson ,D.S (1979) : Seborrhoeic dermatitis in
Text book of Dermatology.Rook, A. and Wilkinson
,D.S Ebling ,F.J.G (eds):vol I 3rd ed Oxford England
Blackwell scientific publications p:308 .

Rook, A . Wilkinson, D.S.Ebling, F.J.G et al (1986) :Text Book
of Dermatology (ed4) Blackwell scientific publica-
tions Oxford , p:267 .

Rook,A. Wilkinson., D.S et al (1993): Text book of Dermatology
volII 5th ed Blackwell scientific publications. Oxford
.p445 .

- Rosenberg ,F.W(1980) : Effect of topical applications of heavy suspensions of killed *Malassezia ovalis* on rabbit skin *Mycopathologia*, 72:147.
- Sabouraud, R.(1904): *Maladies, du cuir chevelu II Les maladies desquamatives* Masson et cie,paris,1904 ;(ed1), p295 (cited by Ive) .
- Salkin,I.F .Gordon,M.A (1977) : Polymorphism of *Malassezia furfur*. *Can.J.Microbiol* ; 23:471 .
- Shuster ,S(1984) : The etiology of dandruff and the mode of action of therapeutic agents *Br.J. Dermatol* 111:23.
- Skinner,R.B.Noah,N.W. Taylor,R.M.M. et al (1985) : Double blind treatement of seborrhoeic dermatitis with 2% Ketoconazol cream *J.Am. Acad . Dermatol* 12:852 .
- Sohnle,P.G.and Lech,C.C (1978) : Cell - mediated immunity to *pityrosporum orbiculare* in tinea versicolor.*J Clin Invest* , 62:45.

pityrosporiasis in guinea pigs .J .Am .Acad. Dermatol
; 22:993.

Vander Wyk,R.W.and Hechemy ,K.E .(1967) : A comparison of
the bacterial and yeast flora of the human scalp and
their effect upon dandruff production . J Soc
Cosmetic Chemists ; 8 : 629 .

Vasarinsh,P. (1982) : Clinical dermatology - diagnosis and
therapy. p . 177 .

Wallace,M . Bangnali,H . Glen,D. et al ; (1983) : Isolation of
lipophilic yeast in " sterile " peritonitis - Lancet ; 2 :
262 .

Weary,P.E. (1968) : Pityrosporum ovale . Arch. Derm .98 :408.

Webster,G . (1991) : Seborrhoeic dermatitis . Int . J . Dermatol ,
Dec , 30 (12) ; p 843 .

Weidman,F.B .(1930) : Seborrhoeic dermatitis in a series of cas-
es of post epidemic encephalitis. A.M.A Arch Derm ;
21 : 690 .